

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 2,3-Diphenyl-1,3-thiazolidin-4-one

Hemant P. Yennawar,<sup>a</sup> John Tierney<sup>b</sup> and Lee J. Silverberg<sup>c\*</sup>

<sup>a</sup>Department of Chemistry, Pennsylvania State University, University Park, PA 16802, USA, <sup>b</sup>Pennsylvania State University, Brandywine Campus, 312 M Main Building, 25 Yearsley Mill Rd, Media, PA 19063, USA, and <sup>c</sup>Pennsylvania State University, Schuylkill Campus, 200 University Drive, Schuylkill Haven, PA 17972, USA  
Correspondence e-mail: ljs43@psu.edu


Received 5 March 2014; accepted 26 June 2014

Key indicators: single-crystal X-ray study;  $T = 298$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.049;  $wR$  factor = 0.147; data-to-parameter ratio = 19.5.

The title compound,  $\text{C}_{15}\text{H}_{13}\text{NOS}$ , is a chiral molecule crystallized as a racemate, with two molecules in the asymmetric unit. In each of the molecules, the five-membered thiazine ring has an envelope conformation, with the S atom forming the flap. In one molecule, the angle between the two phenyl-ring planes is  $82.77(7)^\circ$ , while in the other it is  $89.12(6)^\circ$ . In the crystal, molecules are linked into chains along the  $b$ -axis direction by  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonds.

## Related literature

For the preparation of the title compound, see: Tierney (1989). For the crystal structure of a tin complex of the title compound, see: Smith *et al.* (1995). For the synthesis and crystal structures of related compounds, see: Yennawar & Silverberg (2013, 2014); Fun *et al.* (2011). For reviews on 1,3-thiazolidin-4-ones, see: Brown (1961); Singh *et al.* (1981); Metally *et al.* (2006); Abhishek *et al.* (2012).



## Experimental

## Crystal data

$\text{C}_{15}\text{H}_{13}\text{NOS}$   
 $M_r = 255.32$   
Monoclinic,  $C2/c$

$a = 32.413(13)$  Å  
 $b = 6.196(3)$  Å  
 $c = 25.964(11)$  Å

$\beta = 100.258(7)^\circ$   
 $V = 5131(4)$  Å<sup>3</sup>  
 $Z = 16$   
Mo  $K\alpha$  radiation

$\mu = 0.24$  mm<sup>-1</sup>  
 $T = 298$  K  
 $0.14 \times 0.12 \times 0.08$  mm

## Data collection

Bruker SMART APEX CCD diffractometer  
Absorption correction: multi-scan (SADABS; Bruker, 2001)  
 $T_{\min} = 0.807$ ,  $T_{\max} = 0.981$

23146 measured reflections  
6334 independent reflections  
5015 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.028$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.049$   
 $wR(F^2) = 0.147$   
 $S = 1.01$   
6334 reflections

325 parameters  
H-atom parameters not refined  
 $\Delta\rho_{\text{max}} = 0.33$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.22$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C15}-\text{H15}\cdots\text{O1}^{\text{i}}$	0.93	2.58	3.470 (2)	160
$\text{C1}-\text{H1}\cdots\text{O1}^{\text{i}}$	0.98	2.49	3.466 (2)	172
$\text{C16}-\text{H16}\cdots\text{O2}^{\text{ii}}$	0.98	2.34	3.301 (3)	168
$\text{C17}-\text{H17B}\cdots\text{O2}^{\text{iii}}$	0.97	2.41	3.313 (3)	155

Symmetry codes: (i)  $x, y + 1, z$ ; (ii)  $x, y - 1, z$ ; (iii)  $-x, -y + 1, -z + 1$ .

Data collection: SMART (Bruker, 2001); cell refinement: SAINT (Bruker, 2001); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: XSELL (Bruker, 2001) and ORTEP-3 for Windows (Farrugia, 2012); software used to prepare material for publication: SHELXL97.

We acknowledge NSF funding (CHEM-0131112) for the X-ray diffractometer.

Supporting information for this paper is available from the IUCr electronic archives (Reference: FY2114).

## References

- Abhishek, K. J., Vaidya, A., Ravichandran, V., Kashaw, S. K. & Agrawal, R. K. (2012). *Bioorg. Med. Chem.* **20**, 3378–3395.  
Brown, F. C. (1961). *Chem. Rev.* **61**, 463–521.  
Bruker (2001). SMART, SAINT, XSELL and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.  
Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.  
Fun, H.-K., Hemamalini, M., Shanmugavelan, P., Ponnuswamy, A. & Jagatheesan, R. (2011). *Acta Cryst.* **E67**, o2706.  
Metally, M. A., Etman, H. A., Keshk, E. M. & Fekry, A. (2006). *Phosphorus, Sulfur Silicon Relat. Elem.* **181**, 1039–1058.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Singh, S. P., Parmar, S. S., Raman, K. & Stenberg, V. I. (1981). *Chem. Rev.* **81**, 175–203.  
Smith, F. E., Hynes, R. C., Tierney, J., Zhang, Y. Z. & Eng, G. (1995). *Can. J. Chem.* **73**, 95–99.  
Tierney, J. (1989). *J. Heterocycl. Chem.* **26**, 997–1001.  
Yennawar, H. P. & Silverberg, L. J. (2013). *Acta Cryst.* **E69**, o1659.  
Yennawar, H. P. & Silverberg, L. J. (2014). *Acta Cryst.* **E70**, o133.

## supporting information

*Acta Cryst.* (2014). E70, o847 [doi:10.1107/S1600536814015128]

**2,3-Diphenyl-1,3-thiazolidin-4-one**

**Hemant P. Yennawar, John Tierney and Lee J. Silverberg**

**1. Comment**


We have recently reported the syntheses and crystal structures of 6,7-diphenyl-5-thia-7-azaspiro[2.6]nonan-8-one, a seven-membered heterocycle (Yennawar and Silverberg, 2013), and 2,3-diphenyl-2,3,5,6-tetrahydro-4*H*-1,3-thiazin-4-one, a similar six-membered heterocycle (Yennawar and Silverberg, 2014). We report here the crystal structure of 2,3-diphenyl-1,3-thiazolidin-4-one (Tierney, 1989), the analogous five-membered heterocycle. The crystal structure of a tin complex of the title compound has been previously reported (Smith *et al.*, 1995), but the structure of the title compound has not. The crystal structure of similar compound 3-benzyl-2-phenyl-1,3-thiazolidin-4-one has been reported (Fun *et al.*, 2011). The 1,3-thiazolidin-4-ones are an important class of compounds with a wide range of biological activity (Brown, 1961; Singh, *et al.*, 1981; Metally *et al.*, 2006; Abhishek *et al.*, 2012).

**2. Experimental**


A sample of the title compound, prepared according to Tierney (1989), was recrystallized from ethanol.  $R_f = 0.54$  (50% EtOAc/hexanes). m.p.: 131–133°C (lit. 131–132°C). Crystals for X-ray crystallography were grown by slow evaporation from toluene.

**3. Refinement**


The C-bound H atoms were geometrically placed with C—H = 0.93–0.97 Å, and refined as riding with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ .

**Figure 1**

ORTEP view of the title compound. Thermal ellipsoids are drawn at 50% probability.

**Figure 2**

View of *b-c* plane with C—H...O interactions shown as dashed lines.



**Figure 3**  
Unit-cell contents.

### 2,3-Diphenyl-1,3-thiazolidin-4-one

#### Crystal data

$C_{15}H_{13}NO$

$M_r = 255.32$

Monoclinic,  $C2/c$

Hall symbol:  $-C\ 2yc$

$a = 32.413\ (13)\ \text{\AA}$

$b = 6.196\ (3)\ \text{\AA}$

$c = 25.964\ (11)\ \text{\AA}$

$\beta = 100.258\ (7)^\circ$

$V = 5131\ (4)\ \text{\AA}^3$

$Z = 16$

$F(000) = 2144$

$D_x = 1.322\ \text{Mg m}^{-3}$

Melting point:  $405(1)\ \text{K}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 7831 reflections

$\theta = 2.2\text{--}28.3^\circ$

$\mu = 0.24\ \text{mm}^{-1}$

$T = 298\ \text{K}$

Block, colorless

$0.14 \times 0.12 \times 0.08\ \text{mm}$

#### Data collection

Bruker SMART APEX CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

Detector resolution:  $8.34\ \text{pixels mm}^{-1}$

$\phi$  and  $\omega$  scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2001)

$T_{\min} = 0.807$ ,  $T_{\max} = 0.981$

23146 measured reflections

6334 independent reflections

5015 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.028$

$\theta_{\max} = 28.3^\circ$ ,  $\theta_{\min} = 1.6^\circ$

$h = -42 \rightarrow 42$

$k = -8 \rightarrow 8$

$l = -34 \rightarrow 30$

*Refinement*

Refinement on  $F^2$   
 Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.049$   
 $wR(F^2) = 0.147$   
 $S = 1.01$   
 6334 reflections  
 325 parameters  
 0 restraints  
 Primary atom site location: structure-invariant  
 direct methods

Secondary atom site location: difference Fourier  
 map  
 Hydrogen site location: inferred from  
 neighbouring sites  
 H-atom parameters not refined  
 $w = 1/[\sigma^2(F_o^2) + (0.0897P)^2 + 1.5154P]$   
 where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.33 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.22 \text{ e } \text{\AA}^{-3}$

*Special details*

**Experimental.** The data collection nominally covered a full sphere of reciprocal space by a combination of 4 sets of  $\omega$  scans each set at different  $\varphi$  and/or  $2\theta$  angles and each scan (30 s exposure) covering  $-0.300^\circ$  degrees in  $\omega$ . The crystal to detector distance was 5.82 cm.

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.16111 (4)	0.4561 (2)	0.38426 (6)	0.0375 (3)
H1	0.1730	0.5960	0.3771	0.045*
C2	0.18453 (7)	0.1206 (3)	0.44221 (7)	0.0596 (5)
H2A	0.2083	0.0676	0.4671	0.072*
H2B	0.1601	0.0348	0.4455	0.072*
C3	0.19346 (5)	0.1053 (3)	0.38717 (6)	0.0426 (3)
C4	0.11402 (4)	0.4644 (2)	0.36653 (6)	0.0384 (3)
C5	0.09284 (6)	0.6505 (3)	0.37624 (7)	0.0524 (4)
H5	0.1078	0.7693	0.3913	0.063*
C6	0.04944 (6)	0.6605 (4)	0.36362 (8)	0.0656 (5)
H6	0.0355	0.7856	0.3706	0.079*
C7	0.02702 (6)	0.4875 (4)	0.34099 (8)	0.0682 (6)
H7	-0.0021	0.4951	0.3324	0.082*
C8	0.04765 (6)	0.3011 (4)	0.33086 (8)	0.0651 (5)
H8	0.0325	0.1834	0.3154	0.078*
C9	0.09108 (5)	0.2897 (3)	0.34371 (7)	0.0514 (4)
H9	0.1048	0.1639	0.3370	0.062*
C10	0.18622 (4)	0.3123 (3)	0.30503 (5)	0.0373 (3)
C11	0.17313 (5)	0.1509 (3)	0.26884 (7)	0.0511 (4)
H11	0.1613	0.0246	0.2790	0.061*
C12	0.17791 (7)	0.1806 (4)	0.21694 (8)	0.0693 (6)
H12	0.1690	0.0736	0.1923	0.083*

C13	0.19558 (7)	0.3659 (4)	0.20187 (8)	0.0704 (6)
H13	0.1989	0.3836	0.1673	0.084*
C14	0.20842 (6)	0.5253 (4)	0.23787 (7)	0.0629 (5)
H14	0.2203	0.6510	0.2275	0.075*
C15	0.20371 (5)	0.4996 (3)	0.28963 (6)	0.0471 (4)
H15	0.2123	0.6083	0.3139	0.057*
C16	0.08432 (5)	0.0210 (3)	0.54421 (6)	0.0402 (3)
H16	0.0716	-0.1127	0.5541	0.048*
C17	0.05773 (6)	0.3511 (3)	0.48483 (7)	0.0577 (5)
H17A	0.0809	0.4348	0.4764	0.069*
H17B	0.0321	0.3998	0.4628	0.069*
C18	0.05479 (5)	0.3792 (3)	0.54178 (6)	0.0426 (3)
C19	0.13159 (4)	-0.0004 (2)	0.55810 (5)	0.0371 (3)
C20	0.14969 (5)	-0.1974 (3)	0.55010 (6)	0.0455 (4)
H20	0.1327	-0.3128	0.5370	0.055*
C21	0.19270 (5)	-0.2246 (3)	0.56134 (7)	0.0513 (4)
H21	0.2044	-0.3573	0.5554	0.062*
C22	0.21819 (5)	-0.0559 (3)	0.58124 (7)	0.0521 (4)
H22	0.2471	-0.0746	0.5890	0.063*
C23	0.20071 (5)	0.1409 (3)	0.58958 (7)	0.0536 (4)
H23	0.2179	0.2552	0.6031	0.064*
C24	0.15749 (5)	0.1688 (3)	0.57782 (7)	0.0469 (4)
H24	0.1459	0.3024	0.5833	0.056*
C25	0.07029 (4)	0.1912 (3)	0.62627 (6)	0.0406 (3)
C26	0.08743 (5)	0.3596 (3)	0.65831 (7)	0.0531 (4)
H26	0.0975	0.4818	0.6439	0.064*
C27	0.08940 (6)	0.3447 (4)	0.71192 (8)	0.0694 (6)
H27	0.1007	0.4575	0.7336	0.083*
C28	0.07459 (7)	0.1627 (5)	0.73317 (8)	0.0745 (6)
H28	0.0758	0.1535	0.7692	0.089*
C29	0.05807 (7)	-0.0052 (4)	0.70153 (8)	0.0704 (6)
H29	0.0485	-0.1283	0.7162	0.084*
C30	0.05565 (5)	0.0088 (3)	0.64786 (7)	0.0537 (4)
H30	0.0442	-0.1042	0.6264	0.064*
N1	0.18173 (3)	0.28663 (19)	0.35879 (4)	0.0351 (3)
N2	0.06799 (4)	0.2018 (2)	0.57083 (5)	0.0397 (3)
O1	0.20931 (4)	-0.0527 (2)	0.37092 (5)	0.0597 (3)
O2	0.04255 (4)	0.54461 (19)	0.55954 (5)	0.0579 (3)
S1	0.175220 (14)	0.39920 (8)	0.454463 (16)	0.05330 (15)
S2	0.065806 (16)	0.06983 (9)	0.474346 (18)	0.06317 (17)

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0426 (7)	0.0352 (7)	0.0370 (7)	-0.0009 (6)	0.0132 (6)	-0.0069 (6)
C2	0.0781 (12)	0.0579 (11)	0.0449 (9)	0.0001 (9)	0.0168 (9)	0.0069 (8)
C3	0.0441 (8)	0.0392 (8)	0.0443 (8)	-0.0014 (6)	0.0074 (6)	0.0008 (6)
C4	0.0418 (7)	0.0411 (8)	0.0348 (7)	0.0036 (6)	0.0132 (6)	0.0014 (6)

C5	0.0591 (10)	0.0454 (9)	0.0545 (10)	0.0119 (8)	0.0146 (8)	-0.0003 (8)
C6	0.0604 (11)	0.0722 (13)	0.0668 (12)	0.0306 (10)	0.0180 (9)	0.0060 (10)
C7	0.0427 (9)	0.1013 (17)	0.0607 (11)	0.0169 (10)	0.0097 (8)	0.0031 (11)
C8	0.0455 (9)	0.0795 (14)	0.0704 (13)	-0.0051 (9)	0.0104 (8)	-0.0143 (11)
C9	0.0426 (8)	0.0533 (10)	0.0592 (10)	0.0017 (7)	0.0114 (7)	-0.0104 (8)
C10	0.0342 (6)	0.0433 (8)	0.0352 (7)	0.0057 (6)	0.0079 (5)	-0.0031 (6)
C11	0.0538 (9)	0.0532 (10)	0.0463 (9)	-0.0014 (7)	0.0085 (7)	-0.0137 (8)
C12	0.0784 (13)	0.0829 (15)	0.0450 (10)	0.0099 (11)	0.0065 (9)	-0.0251 (10)
C13	0.0863 (14)	0.0896 (16)	0.0386 (9)	0.0114 (12)	0.0202 (9)	0.0010 (10)
C14	0.0737 (12)	0.0718 (13)	0.0473 (10)	-0.0001 (10)	0.0222 (9)	0.0105 (9)
C15	0.0515 (9)	0.0504 (9)	0.0408 (8)	-0.0016 (7)	0.0122 (7)	-0.0012 (7)
C16	0.0413 (7)	0.0379 (8)	0.0423 (8)	-0.0002 (6)	0.0096 (6)	-0.0062 (6)
C17	0.0537 (9)	0.0693 (12)	0.0489 (10)	0.0055 (9)	0.0055 (8)	0.0070 (9)
C18	0.0365 (7)	0.0410 (8)	0.0500 (9)	-0.0020 (6)	0.0067 (6)	-0.0005 (7)
C19	0.0411 (7)	0.0376 (8)	0.0336 (7)	0.0021 (6)	0.0093 (5)	0.0006 (6)
C20	0.0508 (8)	0.0392 (8)	0.0456 (8)	0.0029 (6)	0.0065 (7)	-0.0051 (7)
C21	0.0535 (9)	0.0496 (10)	0.0504 (9)	0.0174 (7)	0.0079 (7)	0.0010 (8)
C22	0.0414 (8)	0.0668 (11)	0.0474 (9)	0.0074 (8)	0.0057 (7)	0.0061 (8)
C23	0.0443 (8)	0.0546 (10)	0.0611 (11)	-0.0075 (7)	0.0073 (7)	-0.0021 (8)
C24	0.0471 (8)	0.0391 (8)	0.0553 (10)	0.0004 (6)	0.0116 (7)	-0.0029 (7)
C25	0.0351 (6)	0.0466 (9)	0.0405 (8)	0.0064 (6)	0.0079 (6)	-0.0038 (6)
C26	0.0509 (9)	0.0543 (10)	0.0531 (10)	0.0032 (7)	0.0069 (7)	-0.0110 (8)
C27	0.0635 (11)	0.0880 (15)	0.0518 (11)	0.0155 (11)	-0.0033 (9)	-0.0251 (11)
C28	0.0700 (12)	0.1132 (19)	0.0399 (10)	0.0204 (13)	0.0089 (9)	0.0033 (11)
C29	0.0716 (13)	0.0898 (16)	0.0521 (11)	0.0036 (11)	0.0171 (9)	0.0159 (11)
C30	0.0528 (9)	0.0592 (11)	0.0500 (10)	-0.0004 (8)	0.0114 (7)	0.0029 (8)
N1	0.0375 (6)	0.0343 (6)	0.0348 (6)	0.0018 (5)	0.0099 (5)	-0.0036 (5)
N2	0.0405 (6)	0.0379 (7)	0.0413 (7)	0.0023 (5)	0.0094 (5)	-0.0027 (5)
O1	0.0756 (8)	0.0412 (7)	0.0634 (8)	0.0151 (6)	0.0151 (6)	0.0017 (6)
O2	0.0582 (7)	0.0416 (7)	0.0725 (9)	0.0086 (5)	0.0076 (6)	0.0006 (6)
S1	0.0549 (2)	0.0697 (3)	0.0357 (2)	0.0024 (2)	0.00903 (17)	-0.01146 (19)
S2	0.0619 (3)	0.0795 (4)	0.0439 (3)	0.0126 (2)	-0.0019 (2)	-0.0167 (2)

*Geometric parameters (Å, °)*

C1—N1	1.4641 (18)	C16—N2	1.4638 (19)
C1—C4	1.515 (2)	C16—C19	1.516 (2)
C1—S1	1.8328 (17)	C16—S2	1.8315 (17)
C1—H1	0.9800	C16—H16	0.9800
C2—C3	1.511 (2)	C17—C18	1.509 (2)
C2—S1	1.791 (2)	C17—S2	1.790 (2)
C2—H2A	0.9700	C17—H17A	0.9700
C2—H2B	0.9700	C17—H17B	0.9700
C3—O1	1.216 (2)	C18—O2	1.219 (2)
C3—N1	1.360 (2)	C18—N2	1.359 (2)
C4—C5	1.387 (2)	C19—C24	1.383 (2)
C4—C9	1.386 (2)	C19—C20	1.386 (2)
C5—C6	1.388 (3)	C20—C21	1.383 (2)

C5—H5	0.9300	C20—H20	0.9300
C6—C7	1.368 (3)	C21—C22	1.375 (3)
C6—H6	0.9300	C21—H21	0.9300
C7—C8	1.383 (3)	C22—C23	1.378 (3)
C7—H7	0.9300	C22—H22	0.9300
C8—C9	1.389 (2)	C23—C24	1.391 (2)
C8—H8	0.9300	C23—H23	0.9300
C9—H9	0.9300	C24—H24	0.9300
C10—C11	1.386 (2)	C25—C30	1.382 (2)
C10—C15	1.382 (2)	C25—C26	1.387 (2)
C10—N1	1.4378 (18)	C25—N2	1.430 (2)
C11—C12	1.396 (3)	C26—C27	1.385 (3)
C11—H11	0.9300	C26—H26	0.9300
C12—C13	1.371 (3)	C27—C28	1.379 (4)
C12—H12	0.9300	C27—H27	0.9300
C13—C14	1.373 (3)	C28—C29	1.374 (4)
C13—H13	0.9300	C28—H28	0.9300
C14—C15	1.389 (2)	C29—C30	1.384 (3)
C14—H14	0.9300	C29—H29	0.9300
C15—H15	0.9300	C30—H30	0.9300
N1—C1—C4	113.81 (12)	C19—C16—H16	108.6
N1—C1—S1	104.96 (10)	S2—C16—H16	108.6
C4—C1—S1	111.61 (10)	C18—C17—S2	107.33 (13)
N1—C1—H1	108.8	C18—C17—H17A	110.2
C4—C1—H1	108.8	S2—C17—H17A	110.2
S1—C1—H1	108.8	C18—C17—H17B	110.2
C3—C2—S1	107.19 (12)	S2—C17—H17B	110.2
C3—C2—H2A	110.3	H17A—C17—H17B	108.5
S1—C2—H2A	110.3	O2—C18—N2	124.19 (16)
C3—C2—H2B	110.3	O2—C18—C17	123.34 (16)
S1—C2—H2B	110.3	N2—C18—C17	112.46 (14)
H2A—C2—H2B	108.5	C24—C19—C20	118.54 (14)
O1—C3—N1	124.89 (15)	C24—C19—C16	122.89 (14)
O1—C3—C2	122.89 (15)	C20—C19—C16	118.57 (14)
N1—C3—C2	112.22 (14)	C21—C20—C19	120.90 (15)
C5—C4—C9	118.82 (15)	C21—C20—H20	119.5
C5—C4—C1	118.53 (14)	C19—C20—H20	119.5
C9—C4—C1	122.56 (13)	C20—C21—C22	120.17 (15)
C4—C5—C6	120.52 (18)	C20—C21—H21	119.9
C4—C5—H5	119.7	C22—C21—H21	119.9
C6—C5—H5	119.7	C23—C22—C21	119.70 (15)
C7—C6—C5	120.33 (18)	C23—C22—H22	120.2
C7—C6—H6	119.8	C21—C22—H22	120.2
C5—C6—H6	119.8	C22—C23—C24	120.14 (16)
C6—C7—C8	119.88 (17)	C22—C23—H23	119.9
C6—C7—H7	120.1	C24—C23—H23	119.9
C8—C7—H7	120.1	C19—C24—C23	120.55 (15)



C7—C8—C9	120.04 (19)	C19—C24—H24	119.7
C7—C8—H8	120.0	C23—C24—H24	119.7
C9—C8—H8	120.0	C30—C25—C26	120.09 (16)
C4—C9—C8	120.41 (17)	C30—C25—N2	119.07 (14)
C4—C9—H9	119.8	C26—C25—N2	120.84 (15)
C8—C9—H9	119.8	C27—C26—C25	119.63 (19)
C11—C10—C15	120.12 (15)	C27—C26—H26	120.2
C11—C10—N1	120.44 (15)	C25—C26—H26	120.2
C15—C10—N1	119.43 (13)	C28—C27—C26	120.0 (2)
C10—C11—C12	119.10 (18)	C28—C27—H27	120.0
C10—C11—H11	120.5	C26—C27—H27	120.0
C12—C11—H11	120.5	C29—C28—C27	120.43 (19)
C13—C12—C11	120.64 (18)	C29—C28—H28	119.8
C13—C12—H12	119.7	C27—C28—H28	119.8
C11—C12—H12	119.7	C28—C29—C30	120.0 (2)
C12—C13—C14	119.99 (18)	C28—C29—H29	120.0
C12—C13—H13	120.0	C30—C29—H29	120.0
C14—C13—H13	120.0	C25—C30—C29	119.84 (19)
C13—C14—C15	120.30 (19)	C25—C30—H30	120.1
C13—C14—H14	119.9	C29—C30—H30	120.1
C15—C14—H14	119.9	C3—N1—C10	123.39 (12)
C10—C15—C14	119.85 (16)	C3—N1—C1	116.99 (12)
C10—C15—H15	120.1	C10—N1—C1	119.50 (12)
C14—C15—H15	120.1	C18—N2—C25	123.43 (13)
N2—C16—C19	112.93 (12)	C18—N2—C16	117.51 (13)
N2—C16—S2	105.00 (10)	C25—N2—C16	118.80 (12)
C19—C16—S2	112.89 (10)	C2—S1—C1	91.65 (8)
N2—C16—H16	108.6	C17—S2—C16	92.37 (8)
S1—C2—C3—O1	164.39 (14)	N2—C25—C26—C27	-179.78 (15)
S1—C2—C3—N1	-15.65 (18)	C25—C26—C27—C28	0.4 (3)
N1—C1—C4—C5	-162.23 (14)	C26—C27—C28—C29	0.4 (3)
S1—C1—C4—C5	79.20 (16)	C27—C28—C29—C30	-0.9 (3)
N1—C1—C4—C9	21.5 (2)	C26—C25—C30—C29	0.1 (3)
S1—C1—C4—C9	-97.08 (15)	N2—C25—C30—C29	179.28 (16)
C9—C4—C5—C6	0.4 (3)	C28—C29—C30—C25	0.7 (3)
C1—C4—C5—C6	-176.01 (16)	O1—C3—N1—C10	0.9 (2)
C4—C5—C6—C7	-0.6 (3)	C2—C3—N1—C10	-179.02 (14)
C5—C6—C7—C8	0.3 (3)	O1—C3—N1—C1	176.89 (15)
C6—C7—C8—C9	0.1 (3)	C2—C3—N1—C1	-3.07 (19)
C5—C4—C9—C8	0.0 (3)	C11—C10—N1—C3	47.6 (2)
C1—C4—C9—C8	176.27 (17)	C15—C10—N1—C3	-132.45 (15)
C7—C8—C9—C4	-0.3 (3)	C11—C10—N1—C1	-128.26 (15)
C15—C10—C11—C12	0.1 (2)	C15—C10—N1—C1	51.69 (18)
N1—C10—C11—C12	-179.99 (15)	C4—C1—N1—C3	-102.68 (15)
C10—C11—C12—C13	0.5 (3)	S1—C1—N1—C3	19.64 (15)
C11—C12—C13—C14	-0.6 (3)	C4—C1—N1—C10	73.44 (16)
C12—C13—C14—C15	0.2 (3)	S1—C1—N1—C10	-164.24 (10)

C11—C10—C15—C14	-0.5 (2)	O2—C18—N2—C25	-2.1 (2)
N1—C10—C15—C14	179.58 (15)	C17—C18—N2—C25	176.81 (13)
C13—C14—C15—C10	0.4 (3)	O2—C18—N2—C16	-176.11 (14)
S2—C17—C18—O2	-167.49 (13)	C17—C18—N2—C16	2.75 (19)
S2—C17—C18—N2	13.64 (17)	C30—C25—N2—C18	135.69 (16)
N2—C16—C19—C24	-21.5 (2)	C26—C25—N2—C18	-45.1 (2)
S2—C16—C19—C24	97.45 (16)	C30—C25—N2—C16	-50.32 (19)
N2—C16—C19—C20	159.33 (14)	C26—C25—N2—C16	128.84 (15)
S2—C16—C19—C20	-81.74 (16)	C19—C16—N2—C18	106.20 (15)
C24—C19—C20—C21	-0.2 (2)	S2—C16—N2—C18	-17.21 (15)
C16—C19—C20—C21	178.98 (15)	C19—C16—N2—C25	-68.14 (17)
C19—C20—C21—C22	0.7 (3)	S2—C16—N2—C25	168.44 (10)
C20—C21—C22—C23	-0.5 (3)	C3—C2—S1—C1	22.65 (14)
C21—C22—C23—C24	-0.1 (3)	N1—C1—S1—C2	-23.76 (11)
C20—C19—C24—C23	-0.4 (2)	C4—C1—S1—C2	99.98 (12)
C16—C19—C24—C23	-179.55 (15)	C18—C17—S2—C16	-19.83 (12)
C22—C23—C24—C19	0.5 (3)	N2—C16—S2—C17	20.77 (11)
C30—C25—C26—C27	-0.6 (2)	C19—C16—S2—C17	-102.66 (12)

Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ )

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
C15—H15 $\cdots$ O1 <sup>i</sup>	0.93	2.58	3.470 (2)	160
C1—H1 $\cdots$ O1 <sup>i</sup>	0.98	2.49	3.466 (2)	172
C16—H16 $\cdots$ O2 <sup>ii</sup>	0.98	2.34	3.301 (3)	168
C17—H17B $\cdots$ O2 <sup>iii</sup>	0.97	2.41	3.313 (3)	155

Symmetry codes: (i)  $x, y+1, z$ ; (ii)  $x, y-1, z$ ; (iii)  $-x, -y+1, -z+1$ .