

Review Article

Chondrosarcoma and Peroxisome Proliferator-Activated Receptor

K. Nishida,^{1,2} T. Kunisada,² Z. N. Shen,² Y. Kadota,² K. Hashizume,² and T. Ozaki²

¹ Department of Human Morphology, Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama University, 2-5-1 Shikata-cho, Okayama 700-8558, Japan

² Department of Orthopaedic Surgery, Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama University, 2-5-1 Shikata-cho, Okayama 700-8558, Japan

Correspondence should be addressed to K. Nishida, knishida@md.okayama-u.ac.jp

Received 1 March 2008; Accepted 17 July 2008

Recommended by Dipak Panigrahy

Induction of differentiation and apoptosis in cancer cells by ligands of PPAR γ is a novel therapeutic approach to malignant tumors. Chondrosarcoma (malignant cartilage tumor) and OUMS-27 cells (cell line established from grade III human chondrosarcoma) express PPAR γ . PPAR γ ligands inhibited cell proliferation in a dose-dependent manner, and induced apoptosis of OUMS-27. The higher-grade chondrosarcoma expressed a higher amount of antiapoptotic Bcl-xL in vivo. The treatment of OUMS-27 by 15d-PGJ₂, the most potent endogenous ligand for PPAR γ , downregulated expression of Bcl-xL and induced transient upregulation of proapoptotic Bax, which could accelerate cytochrome c release from mitochondria to the cytosol, followed by induction of caspase-dependent apoptosis. 15d-PGJ₂ induced the expression of CDK inhibitor p21 protein in human chondrosarcoma cells, which appears to be involved in the mechanism of inhibition of cell proliferation. These findings suggest that targeted therapy with PPAR γ ligands could be a novel strategy against chondrosarcoma.

Copyright © 2008 K. Nishida et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. INTRODUCTION

Cancers are associated with dysregulation of differentiation and apoptotic cell death. Recent investigations have demonstrated that induction of these cellular events by targeted therapy with ligands of nuclear hormone receptors could be a novel strategy against cancers [1]. Peroxisome proliferator-activated receptor (PPAR) γ , a member of the nuclear receptor superfamily, acts as a ligand-activated transcription factor, and is involved in many processes important for homeostasis of cells and tissues, including metabolism, immune and inflammatory controls, cell proliferation and apoptotic cell death [2–6]. Because PPAR γ is expressed by many malignant tumors, activation of PPAR γ by 15-deoxy- $\Delta^{12,14}$ -prostaglandin J₂ (15d-PGJ₂), the most potent endogenous ligand for PPAR γ [7], and the synthetic PPAR γ ligands (e.g., rosiglitazone, pioglitazone, troglitazone, and indomethacin) have been regarded as a novel therapeutic approach for certain human malignancies through growth inhibition, induction of apoptosis and terminal differentiation, and inhibition of angiogenesis [8]. This review will

outline the inhibitory effects of synthetic and endogenous PPAR γ ligands and discuss their potential therapeutic effects on chondrosarcoma.

2. CLINICAL FEATURES OF CHONDROSARCOMA

Chondrosarcoma is a malignant tumor of cartilage; the matrix formed by tumor cells is uniformly and entirely chondroid in nature [9]. Human chondrosarcoma is a rare bone tumor, accounting for <10% of primary malignant bone tumors. Chondrosarcoma also arises in pre-existing benign lesions (e.g., osteochondromatosis, enchondromatosis) and is termed “secondary chondrosarcoma.”

Primary (conventional) chondrosarcoma arises centrally in a previously normal bone, and mostly grows slowly through the diaphyseal cortex. Most patients are aged >50 years. Conventional chondrosarcoma is more common in men. The commonest sites are the bones of the pelvis, followed by the femur and the humerus. Recognizable histologic variants are clear cell, mesenchymal, and dedifferentiated chondrosarcomas. On the basis of histologic

features (nuclear atypia and cellularity), conventional chondrosarcoma is further subdivided into three grades: I, II, and III [10]. The histologic grade of chondrosarcoma indicates the differentiation status of tumor cells, and is one of the most important factors for prognosis [11]. Progression of a locally aggressive low-grade chondrosarcoma to a metastasizing high-grade chondrosarcoma is associated with loss of cartilaginous phenotype, genomic instability, and aneuploidy [12]. The grading of chondrosarcoma correlates well with clinical behavior, although chondrosarcoma is one of the most difficult malignant tumors of bone to diagnose [13].

Most conventional chondrosarcomas are grade I or II. For low-grade chondrosarcoma, surgical treatment with adequate marginal resection is reported to be associated with better clinical outcomes [14]. Only 5–10% of conventional chondrosarcomas are grade-III lesions, which have definite metastatic potential. The prognosis for high-grade chondrosarcoma is poor, despite adequate surgery, because they are highly resistant to conventional chemotherapy and radiotherapy [15]. These facts, that the differentiation status of chondrosarcoma is predictive of clinical outcomes, suggest the favorable effects of the modification of the differentiation status on clinical behavior. Recent advances in understanding the progression or development of chondrosarcoma have suggested several molecular targets for future development of new adjuvant therapy [16], such as chondrocyte differentiation factors (PTHrP, CTGF) [17, 18], antiapoptotic gene (*Bcl-2*) [19, 20], tumor suppressor gene (*p16*, *p53*) [21, 22], and others (PDGF- α , VEGF, *MDR-1*) [23–25].

3. OUMS-27, A CHONDROSARCOMA CELL LINE

A cell line derived from chondrosarcoma, particularly from high-grade chondrosarcoma, can provide a useful model for the investigation of cell development and treatment of chondrosarcoma [26–30]. The OUMS-27 cell line has been established from grade III human chondrosarcoma [31]. The cells do not show contact inhibition after reaching confluence, grow rapidly in multiple layers, and express proteoglycan, as well as collagen type I, II, III, IX, and XI after 120 passages, showing stable maintenance of the differentiated chondrocytic properties. The transplantation of OUMS-27 cells into athymic mice resulted in formation of grade II chondrosarcoma at the injection site. There have been many studies on the etiology and treatment of chondrosarcoma using this cell line [32–34].

4. EXPRESSION OF PPAR γ IN HUMAN CHONDROSARCOMA AND OUMS-27

Subramanian et al. [35] investigated gene expression profiling of ten extraskeletal myxoid chondrosarcomas (EMCs) using 42000 spot cDNA microarrays. Eighty-six genes that distinguished EMC from the other sarcomas were identified by significance analysis of microarrays with 0.25% likelihood of false significance. Of these, PPAR γ and PARGC1A, an interacting protein with PPAR γ and also a coactivator, were highly expressed in EMCs.

TABLE 1: Summary of immunohistochemical study for PPAR γ in human chondrosarcoma tissues.

Positive cell ratios (%)	Pathological grade of chondrosarcoma (%)		
	I (<i>n</i> = 20)	II (<i>n</i> = 6)	III (<i>n</i> = 2)
<10	35	17	50
10–40	40	33	50
>40	25	50	0

In vivo PPAR γ protein content was examined in conventional chondrosarcoma specimens from 28 patients undergoing surgery [36]. Immunohistochemical study revealed that human chondrosarcoma cells frequently express PPAR γ protein. The positivity (cutoff positivity of 10%) of chondrosarcoma cells were 65.0% in grade I, 83.3% in grade II, and 50.0% in grade III; overall positivity was 67.9% (see Table 1). Expression of PPAR γ in OUMS-27 cells at protein and mRNA levels was confirmed by immunocytochemistry and reverse transcriptase-polymerase chain reaction (RT-PCR) analysis, respectively [36]. These data indicated that PPAR γ is frequently expressed in primary chondrosarcomas and chondrosarcoma cell line OUMS-27, and led the authors to test the effect of PPAR γ activators on cell proliferation and survival of OUMS-27.

5. EVIDENCE OF APOPTOTIC CELL DEATH OF OUMS-27 CELLS AFTER TREATMENT BY PPAR γ LIGANDS

In our previous report [36], OUMS-27 cells were treated with increasing concentrations of pioglitazone (synthetic PPAR γ ligand) and 15d-PGJ₂ for up to 48 hours. The results of immunostain for Ki-67 (cell proliferation marker) and colorimetric MTT assay showed that treatment with both pioglitazone and 15d-PGJ₂ for 24 hours inhibited cell growth and reduced cell viability in a dose-dependent manner, respectively. 15d-PGJ₂ had more noticeable effects on OUMS-27 cell growth than pioglitazone. It was unclear whether the effects of ligands on OUMS-27 cells were strictly due to PPAR γ activation. When cells were treated with 15d-PGJ₂ doses of $\geq 5 \mu\text{g/mL}$, they showed relatively round shapes and some cells no longer adhered to the dish.

Semithin sectioned, LR White-embedded cells stained by toluidine blue revealed that many OUMS-27 cells treated with 15d-PGJ₂ show apoptotic appearances with cell shrinkage and nuclear condensation (see Figure 1). DNA fragmentations in OUMS-27 cells treated by 15d-PGJ₂ (10 $\mu\text{g/mL}$) for 24 hours were confirmed by DNA ladder formation and TUNEL staining. Transmission electron microscopic study revealed sections of OUMS-27 cells treated with 15d-PGJ₂ contained many cells consistent with morphological apoptosis with condensed chromatin, many vacuoles in cytoplasm, and membrane budding. Early apoptotic change and the translocation of phosphatidylserine (PS) on the outer leaflet of the cell membrane were demonstrated by FACS analysis. The population of apoptotic cells with PS at the outer membrane of the cells (annexin-V-positive,




FIGURE 1: Cell morphology of chondrosarcoma cell line OUMS-27 after incubation with (b) or without (a) 15d-PGJ₂. Cells were treated with 10 μg/mL of 15d-PGJ₂ for 8 hours, and the cell pellet embedded in hydrophilic resin. Semithin sections stained by toluidine blue show more apoptotic cells with cell shrinkage and nuclear condensation (arrows) after treatment with 15d-PGJ₂.

PI-negative) was ~53.9% and 67.6% at 4 hours and 24 hours after coincubation with 15d-PGJ₂, respectively.

6. MECHANISM OF APOPTOTIC CELL DEATH OF OUMS-27 CELLS BY PPAR_γ LIGANDS

cDNA microarray analysis was carried out to comprehensively explore the changes in gene expression pattern during OUMS-27 cell growth inhibition and possible cell cycle arrest caused by treatment with 15d-PGJ₂ [37]. Among the 1081 genes analyzed, 52 genes were upregulated and 81 genes were downregulated significantly in OUMS-27 cells after 8-hour treatment with 15d-PGJ₂ (10 μg/mL). Microarray analysis is shown in Table 2. Interestingly, the proapoptotic gene Bax was upregulated, and the antiapoptotic gene Bcl-xL was downregulated. The other Bcl-2 members were unchanged. These results were further confirmed by RT-PCR and real-time PCR analysis.

Upregulation of Bax, concurrent with the downregulation of Bcl-xL, can destabilize mitochondria, leading to the release of several mitochondrial intermembrane space proteins such as cytochrome c, AIF, Smac/DIABLO, Endo G, and Omi/HtrA2 into the cytosol, where they are actively involved in apoptotic cell death [38]. This hypothesis is supported by our observations of the release of cytochrome c from mitochondria into the cytosol, and the activation of caspase-3 in 15d-PGJ₂-treated OUMS-27 cells. Coincubation

TABLE 2: Changes in apoptosis-related gene expression in OUMS-27 cells after treatment with 15d-PGJ₂.

Gene name	Fold
Clusterin	+3.7
Defender against cell death 1	+3.8
Tumor necrosis factor receptor 1	+3.2
State-induced state inhibitor 2	+2.5
Heat shock protein 60	+2.4
V-akt murine thymoma viral oncogene homolog 1	+2.3
Apoptosis regulator bax	+2.3
Interferon-induced RNA-dependent protein kinase	+2.1
Apoptosis regulator bcl-xl	-2.2
Calpain, small subunit 1	-2.3
Heat shock 70 KD protein 1	-2.4
Endothelin 2	-2.4
Insulin-like growth factor 1 receptor	-2.5
Death-associated protein 6	-2.8

of cells with the broad-spectrum caspase inhibitor Z-VAD-FMK completely inhibited caspase activity, and prevented the cell death induced by 15d-PGJ₂. These results indicate that 15d-PGJ₂ induced apoptosis in OUMS-27 cells through a caspase-dependent signal transduction pathway which, at least in part, was triggered by cytosolic release of cytochrome c [37].

The decreased expression of antiapoptotic Bcl-xL in OUMS-27 treated by 15d-PGJ₂ led us to examine the expression in the tissue of human chondrosarcoma samples to study the clinical application of differentiation therapy by PPAR_γ activation. The result of immunohistochemical study demonstrated that Bcl-xL was expressed in all three grades of chondrosarcoma; the expression was strongest in grade III. These results indicated that higher-grade chondrosarcoma cells may be resistant to apoptosis by overexpression of Bcl-xL, and 15d-PGJ₂ might induce apoptotic cell death by downregulation of Bcl-xL and transient upregulation of Bax [37]. Similar results were reported in renal cell carcinoma cells (786-O and A498 cells) showing the thiazolidinedione (TZD) induction of apoptosis with increased Bax expression and decreased Bcl-2 expression [39].

6.1. Genetic and epigenetic alterations in chondrosarcoma

Little is known about the role of genetic or epigenetic alterations in tumor progression from low-malignant chondroblastic to highly malignant anaplastic chondrosarcoma. The appearance of de novo aberrant DNA methylation is the commonest molecular change in the cancer cell, which inactivates many cellular pathways [40]. The most studied change of DNA methylation in neoplasms is the silencing of tumor suppressor genes by deoxy-cytidylatephosphate-deoxy-guanylate (CpG) island promoter hypermethylation, which targets genes and molecules associated in cell differentiation, such as p16(INK4a), BRCA1, and hMLH1

[41–43]. Röpke et al. reported the p16 and E-cadherin promoter methylation in low-grade chondroid compartment of dedifferentiated chondrosarcoma. Van Beerendonk et al. found p16 promoter methylation by methylation-specific PCR in 5 of 30 tumors, but this did not correlate with protein expression, or with loss of heterozygosity (LOH) at 9p21 region, one of the few consistent genetic aberrations found in conventional chondrosarcoma [44]. In OUMS-27, methylation was not detectable in the promoter of p16 gene (unpublished data).

Some reports suggested that p53 mutation and p53 loss of heterozygosity are involved [43, 45]. In OUMS-27, we have previously shown that the p53 gene is mutated [31]. Asp et al. analyzed p16 and p53 in cartilaginous tumor tissues and showed that the p16 gene was found to be partly methylated in 5 high-grade chondrosarcomas and homozygously deleted in 1 chondrosarcoma, whereas the p53 gene revealed an unchanged structure in all 22 chondrosarcoma samples [46].

7. INDUCTION OF CELL CYCLE ARREST BY 15d-PGJ₂ IN OUMS-27

Ligands for PPAR γ reportedly induce cell cycle arrest in various cancer cells [39, 47–54]. 15d-PGJ₂ induces G₁ arrest and inhibits cell growth of human anaplastic thyroid carcinoma through a p53-independent, but p21- and p27-dependent, manner [55]. Activation of PPAR γ by troglitazone inhibited cell growth and induced G₁ arrest through the increase of cyclin-dependent kinase (CDK) inhibitor p27 in several cell lines, including human pancreatic carcinoma cells, gastric cancer cells, and hepatocellular carcinoma cells [56–58]. The effect of troglitazone on the proliferation of cancer cells was inhibited by antisense for p27. Yang et al. also showed TZD decreased the protein levels of proliferating cell nuclear antigen, pRb, cyclin D, and Cdk4, but increased the levels of p21 and p27, in RCC cells [39].

In OUMS-27, 15d-PGJ₂ induced the expression of the CDK inhibitor p21 protein, and it was increased within 24 hours. Expression of the other CDK inhibitors, p16 and p27 proteins, were detected at time zero, and were not significantly influenced by 15d-PGJ₂ treatment [37]. 15d-PGJ₂-induced p21 may exert cell cycle arrest in a p53-independent manner.

Whether 15d-PGJ₂ induces p21 expression in OUMS-27 cells through a PPAR γ -dependent or -independent pathway is unclear. It is possible that p21 expression is directly regulated by PPAR γ activation because p21 gene contains a potentially conserved consensus PPAR γ response element in the promoter region [59]. Copland et al. reported [60] that RS5444, a novel high-affinity PPAR γ agonist, inhibits anaplastic thyroid carcinoma (ATC) tumor growth and angiogenesis in mice. In DRO cells derived from ATC tumor, they demonstrated that upregulation of p21 by RS5444 is PPAR γ dependent, and might be the major mechanism by which RS5444 inhibits DRO cell proliferation. Han et al. demonstrated the link of PPAR γ activation and p21 signaling to cell growth inhibition in human lung cell carcinoma cells using p21 antisense oligonucleotides [61]. They also indicated the induction of p21 expression by

PPAR γ ligands might be mediated through increased Sp-1 and NF-interleukin 6 (IL6) CAAT/enhancer binding protein (C/EBP)-dependent transcriptional activation.

8. CLINICAL APPLICATION OF PPAR γ AGONIST FOR CHONDROSARCOMA SUPPRESSION

Accumulating evidence suggests that PPAR γ activators might have clinical therapeutic benefit in the treatment of cancers. Although initial clinical trials with troglitazone reported promising results in liposarcomas [62] and prostate cancers [63], recent studies failed to show the expected therapeutic values of rosiglitazone in liposarcomas [64] and early-stage breast cancers [65], and troglitazone in chemotherapy-resistant metastatic colorectal cancers [66]. However, a single study of a phase-I clinical trial of LY293111 in patients with advanced solid tumors reported a potential efficacy of PPAR γ agonist for chondrosarcoma [67]. LY293111 is an orally stable leukotriene B₄ (LTB₄) receptor antagonist, as well as a PPAR γ agonist, as demonstrated by activity in the rat ZDF diabetes model, and the induction of adipocyte differentiation. One patient with progressive chondrosarcoma had stable disease lasting ~336 days of LY293111 administration at the dose of 200 mg bd.

9. FUTURE DIRECTION

In chondrosarcoma, whether the cell death and growth inhibitory effects induced by 15d-PGJ₂ are PPAR γ -dependent or -independent is unknown. As 15d-PGJ₂ at high doses is toxic for most of cell types independent of PPAR γ activation, we examined the effects of the caspase inhibitor Z-VAD-FMK, and the PPAR γ antagonist GW9662, on caspase-3 activation and cell viability of OUMS-27 cells treated by 15d-PGJ₂ [37]. 15d-PGJ₂ alone clearly increased cell death; the addition of GW9662 partially inhibited cell death. Cell death was inhibited almost to control level when Z-VAD-FMK was added to 15d-PGJ₂. The activity of caspase-3 was attenuated, though not completely, by stimulation of 15d-PGJ₂ together with GW9662. These data indicate that the greater proapoptotic effects of 15d-PGJ₂ on chondrosarcoma cells may result from the cumulative effects of PPAR γ -dependent and -independent pathways. Detailed analysis of the effects of ligands on cells transfected with PPAR γ siRNA should provide important clues to understanding this phenomenon. Whether endogenous or synthetic PPAR γ ligands can also induce tumor cell death in an experimentally transplanted chondrosarcoma model remains to be examined before human trial.

ACKNOWLEDGMENT

The authors wish to thank Mrs. Aki Yoshida (Department of Orthopedic Surgery, Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama University) for technical assistance.

REFERENCES

- [1] B. H. Park, B. Breyer, and T.-C. He, "Peroxisome proliferator-activated receptors: roles in tumorigenesis and chemoprevention in human cancer," *Current Opinion in Oncology*, vol. 13, no. 1, pp. 78–83, 2001.
- [2] B. Desvergne and W. Wahli, "Peroxisome proliferator-activated receptors: nuclear control of metabolism," *Endocrine Reviews*, vol. 20, no. 5, pp. 649–688, 1999.
- [3] G. Chinetti, J.-C. Fruchart, and B. Staels, "Peroxisome proliferator-activated receptors (PPARs): nuclear receptors at the crossroads between lipid metabolism and inflammation," *Inflammation Research*, vol. 49, no. 10, pp. 497–505, 2000.
- [4] E.-Z. Amri, F. Bonino, G. Ailhaud, N. A. Abumrad, and P. A. Grimaldi, "Cloning of a protein that mediates transcriptional effects of fatty acids in preadipocytes. Homology to peroxisome proliferator-activated receptors," *The Journal of Biological Chemistry*, vol. 270, no. 5, pp. 2367–2371, 1995.
- [5] T. Lemberger, B. Desvergne, and W. Wahli, "Peroxisome proliferator-activated receptors: a nuclear receptor signaling pathway in lipid physiology," *Annual Review of Cell and Developmental Biology*, vol. 12, pp. 335–363, 1996.
- [6] S. Kersten, B. Desvergne, and W. Wahli, "Roles of PPARs in health and disease," *Nature*, vol. 405, no. 6785, pp. 421–424, 2000.
- [7] B. M. Forman, P. Tontonoz, J. Chen, R. P. Brun, B. M. Spiegelman, and R. M. Evans, "15-deoxy- $\Delta^{12,14}$, 14-prostaglandin J_2 is a ligand for the adipocyte determination factor PPAR γ ," *Cell*, vol. 83, no. 5, pp. 803–812, 1995.
- [8] T. Wang, J. Xu, X. Yu, R. Yang, and Z. C. Han, "Peroxisome proliferator-activated receptor γ in malignant diseases," *Critical Reviews in Oncology/Hematology*, vol. 58, no. 1, pp. 1–14, 2006.
- [9] H. D. Dorfman and B. Czerniak, "Malignant cartilage tumors," in *Bone Tumors*, pp. 353–440, Mosby, Saint Louis, Mo, USA, 1998.
- [10] F. Bertoni, P. Bacchini, and P. C. Hogendoorn, "Chondrosarcoma," in *World Health Organization Classification of Tumours: Pathology and Genetics, Tumours of Soft Tissue and Bone*, C. D. M. Fletcher, K. K. Unni, and F. Mertens, Eds., IARC Press, Lyon, France, 2002.
- [11] H. L. Evans, A. G. Ayala, and M. M. Romsdahl, "Prognostic factors in chondrosarcoma of bone. A clinicopathologic analysis with emphasis on histologic grading," *Cancer*, vol. 40, no. 2, pp. 818–831, 1977.
- [12] J. V. M. G. Bovée, A.-M. Cleton-Jansen, A. H. M. Taminiau, and P. C. W. Hogendoorn, "Emerging pathways in the development of chondrosarcoma of bone and implications for targeted treatment," *The Lancet Oncology*, vol. 6, no. 8, pp. 599–607, 2005.
- [13] K. K. Unni, "Chondrosarcoma (primary, secondary, dedifferentiated, and clear cell)," in *Dahlin's Bone Tumors: General Aspects and Data on 11,087 Cases*, pp. 71–108, Lippincott-Raven, Philadelphia, Pa, USA, 1996.
- [14] T. Ozaki, N. Lindner, A. Hillmann, R. Rödl, S. Blasius, and W. Winkelmann, "Influence of intralesional surgery on treatment outcome of chondrosarcoma," *Cancer*, vol. 77, no. 7, pp. 1292–1297, 1996.
- [15] V. O. Lewis, "What's new in musculoskeletal oncology," *The Journal of Bone & Joint Surgery*, vol. 89, no. 6, pp. 1399–1407, 2007.
- [16] R. M. Terek, "Recent advances in the basic science of chondrosarcoma," *Orthopedic Clinics of North America*, vol. 37, no. 1, pp. 9–14, 2006.
- [17] T. Kunisada, J. M. Moseley, J. L. Slavin, T. J. Martin, and P. F. M. Choong, "Co-expression of parathyroid hormone-related protein (PTHrP) and PTH/PTHrP receptor in cartilaginous tumours: a marker for malignancy?" *Pathology*, vol. 34, no. 2, pp. 133–137, 2002.
- [18] T. Shakunaga, T. Ozaki, N. Ohara, et al., "Expression of connective tissue growth factor in cartilaginous tumors," *Cancer*, vol. 89, no. 7, pp. 1466–1473, 2000.
- [19] J. V. M. G. Bovée, L. J. C. M. van den Broek, A.-M. Cleton-Jansen, and P. C. W. Hogendoorn, "Up-regulation of PTHrP and Bcl-2 expression characterizes the progression of osteochondroma towards peripheral chondrosarcoma and is a late event in central chondrosarcoma," *Laboratory Investigation*, vol. 80, no. 12, pp. 1925–1933, 2000.
- [20] M. Amling, M. Pösl, M. W. Hentz, M. Priemel, and G. Delling, "PTHrP and Bcl-2: essential regulatory molecules in chondrocyte differentiation and chondrogenic tumors," *Verhandlungen der Deutschen Gesellschaft für Pathologie*, vol. 82, pp. 160–169, 1998.
- [21] R. M. Terek, J. H. Healey, P. Garin-Chesa, S. Mak, A. Huvois, and A. P. Albino, "p53 mutations in chondrosarcoma," *Diagnostic Molecular Pathology*, vol. 7, no. 1, pp. 51–56, 1998.
- [22] B. Coughlan, A. Feliz, T. Ishida, B. Czerniak, and H. D. Dorfman, "p53 expression and DNA ploidy of cartilage lesions," *Human Pathology*, vol. 26, no. 6, pp. 620–624, 1995.
- [23] I. Sulzbacher, P. Birner, K. Trieb, M. Mühlbauer, S. Lang, and A. Chott, "Platelet-derived growth factor- α receptor expression supports the growth of conventional chondrosarcoma and is associated with adverse outcome," *The American Journal of Surgical Pathology*, vol. 25, no. 12, pp. 1520–1527, 2001.
- [24] T. Furumatsu, K. Nishida, A. Kawai, M. Namba, H. Inoue, and Y. Ninomiya, "Human chondrosarcoma secretes vascular endothelial growth factor to induce tumor angiogenesis and stores basic fibroblast growth factor for regulation of its own growth," *International Journal of Cancer*, vol. 97, no. 3, pp. 313–322, 2002.
- [25] R. N. Rosier, R. J. O'Keefe, L. A. Teot, et al., "P-glycoprotein expression in cartilaginous tumors," *Journal of Surgical Oncology*, vol. 65, no. 2, pp. 95–105, 1997.
- [26] M. Takigawa, H.-O. Pan, M. Enomoto, et al., "A clonal human chondrosarcoma cell line produces an anti-angiogenic antitumor factor," *Anticancer Research*, vol. 10, no. 2A, pp. 311–315, 1990.
- [27] T. Chano, H. Okabe, Y. Saeki, M. Ishizawa, K. Matsumoto, and S. Hukuda, "Characterization of a newly established human chondrosarcoma cell line, CS-OKB," *Virchows Archiv*, vol. 432, no. 6, pp. 529–534, 1998.
- [28] H. Chansky, J. R. Robbins, S. Cha, W. H. Raskind, E. U. Conrad, and L. J. Sandell, "Expression of cartilage extracellular matrix and potential regulatory genes in a new human chondrosarcoma cell line," *Journal of Orthopaedic Research*, vol. 16, no. 5, pp. 521–530, 1998.
- [29] R. Gil-Benso, C. Lopez-Gines, J. A. López-Guerrero, et al., "Establishment and characterization of a continuous human chondrosarcoma cell line, ch-2879: comparative histologic and genetic studies with its tumor of origin," *Laboratory Investigation*, vol. 83, no. 6, pp. 877–887, 2003.
- [30] I. Kudawara, N. Araki, A. Myoui, Y. Kato, A. Uchida, and H. Yoshikawa, "New cell lines with chondrocytic phenotypes from human chondrosarcoma," *Virchows Archiv*, vol. 444, no. 6, pp. 577–586, 2004.

- [31] T. Kunisada, M. Miyazaki, K. Mihara, et al., "A new human chondrosarcoma cell line (OUMS-27) that maintains chondrocytic differentiation," *International Journal of Cancer*, vol. 77, no. 6, pp. 854–859, 1998.
- [32] K. Demircan, S. Hirohata, K. Nishida, et al., "ADAMTS-9 is synergistically induced by interleukin-1 β and tumor necrosis factor α in OUMS-27 chondrosarcoma cells and in human chondrocytes," *Arthritis & Rheumatism*, vol. 52, no. 5, pp. 1451–1460, 2005.
- [33] T. Furumatsu, N. Yamaguchi, K. Nishida, et al., "Endostatin inhibits adhesion of endothelial cells to collagen I via $\alpha_2\beta_1$ integrin, a possible cause of prevention of chondrosarcoma growth," *The Journal of Biochemistry*, vol. 131, no. 4, pp. 619–626, 2002.
- [34] T. Hayami, C. Shukunami, K. Mitsui, et al., "Specific loss of chondromodulin-I gene expression in chondrosarcoma and the suppression of tumor angiogenesis and growth by its recombinant protein in vivo," *FEBS Letters*, vol. 458, no. 3, pp. 436–440, 1999.
- [35] S. Subramanian, R. B. West, R. J. Marinelli, et al., "The gene expression profile of extraskelatal myxoid chondrosarcoma," *The Journal of Pathology*, vol. 206, no. 4, pp. 433–444, 2005.
- [36] K. Nishida, T. Furumatsu, I. Takada, et al., "Inhibition of human chondrosarcoma cell growth via apoptosis by peroxisome proliferator-activated receptor- γ ," *British Journal of Cancer*, vol. 86, no. 8, pp. 1303–1309, 2002.
- [37] Z.-N. Shen, K. Nishida, H. Doi, et al., "Suppression of chondrosarcoma cells by 15-deoxy- $\Delta^{12,14}$ -prostaglandin J₂ is associated with altered expression of Bax/Bcl-xL and p21," *Biochemical and Biophysical Research Communications*, vol. 328, no. 2, pp. 375–382, 2005.
- [38] Y. Tsujimoto, "Cell death regulation by the Bcl-2 protein family in the mitochondria," *Journal of Cellular Physiology*, vol. 195, no. 2, pp. 158–167, 2003.
- [39] F. Yang, Z. Zhang, D. Xin, et al., "Peroxisome proliferator-activated receptor γ ligands induce cell cycle arrest and apoptosis in human renal carcinoma cell lines," *Acta Pharmacologica Sinica*, vol. 26, no. 6, pp. 753–761, 2005.
- [40] M. Röpke, C. Boltze, H. W. Neumann, A. Roessner, and R. Schneider-Stock, "Genetic and epigenetic alterations in tumor progression in a dedifferentiated chondrosarcoma," *Pathology, Research and Practice*, vol. 199, no. 6, pp. 437–444, 2003.
- [41] M. Esteller, "Aberrant DNA methylation as a cancer-inducing mechanism," *Annual Review of Pharmacology and Toxicology*, vol. 45, pp. 629–656, 2005.
- [42] M. Esteller, "CpG island hypermethylation and tumor suppressor genes: a booming present, a brighter future," *Oncogene*, vol. 21, no. 35, pp. 5427–5440, 2002.
- [43] M. Esteller, M. F. Fraga, M. Guo, et al., "DNA methylation patterns in hereditary human cancers mimic sporadic tumorigenesis," *Human Molecular Genetics*, vol. 10, no. 26, pp. 3001–3007, 2001.
- [44] H. M. van Beerendonk, L. B. Rozeman, A. H. M. Taminiau, et al., "Molecular analysis of the INK4A/INK4A-ARF gene locus in conventional (central) chondrosarcomas and enchondromas: indication of an important gene for tumour progression," *The Journal of Pathology*, vol. 202, no. 3, pp. 359–366, 2004.
- [45] H. J. Grote, R. Schneider-Stock, W. Neumann, and A. Roessner, "Mutation of p53 with loss of heterozygosity in the osteosarcomatous component of a dedifferentiated chondrosarcoma," *Virchows Archiv*, vol. 436, no. 5, pp. 494–497, 2000.
- [46] J. Asp, L. Sangiorgi, S. E. Inerot, et al., "Changes of the p16 gene but not the p53 gene in human chondrosarcoma tissues," *International Journal of Cancer*, vol. 85, no. 6, pp. 782–786, 2000.
- [47] K. Hashimoto, R. T. Ethridge, and B. M. Evers, "Peroxisome proliferator-activated receptor γ ligand inhibits cell growth and invasion of human pancreatic cancer cells," *International Journal of Gastrointestinal Cancer*, vol. 32, no. 1, pp. 7–22, 2002.
- [48] N. G. Nikitakis, H. Siavash, C. Hebert, M. A. Reynolds, A. W. Hamburger, and J. J. Sauk, "15-PGJ₂, but not thiazolidinediones, inhibits cell growth, induces apoptosis, and causes downregulation of Stat3 in human oral SCCa cells," *British Journal of Cancer*, vol. 87, no. 12, pp. 1396–1403, 2002.
- [49] C. Ward, I. Dransfield, J. Murray, S. N. Farrow, C. Haslett, and A. G. Rossi, "Prostaglandin D₂ and its metabolites induce caspase-dependent granulocyte apoptosis that is mediated via inhibition of κ B α degradation using a peroxisome proliferator-activated receptor- γ -independent mechanism," *The Journal of Immunology*, vol. 168, no. 12, pp. 6232–6243, 2002.
- [50] R. Butler, S. H. Mitchell, D. J. Tindall, and C. Y. F. Young, "Nonapoptotic cell death associated with S-phase arrest of prostate cancer cells via the peroxisome proliferator-activated receptor γ ligand, 15-deoxy- $\Delta^{12,14}$ -prostaglandin J₂," *Cell Growth & Differentiation*, vol. 11, no. 1, pp. 49–61, 2000.
- [51] M. A. K. Rumi, H. Sato, S. Ishihara, et al., "Peroxisome proliferator-activated receptor γ ligand-induced growth inhibition of human hepatocellular carcinoma," *British Journal of Cancer*, vol. 84, no. 12, pp. 1640–1647, 2001.
- [52] C. Wang, M. Fu, M. D'Amico, et al., "Inhibition of cellular proliferation through κ B kinase-independent and peroxisome proliferator-activated receptor γ -dependent repression of cyclin D1," *Molecular and Cellular Biology*, vol. 21, no. 9, pp. 3057–3070, 2001.
- [53] C. E. Clay, G. Atsumi, K. P. High, and F. H. Chilton, "Early *de Novo* gene expression is required for 15-deoxy- $\Delta^{12,14}$ -prostaglandin J₂-induced apoptosis in breast cancer cells," *The Journal of Biological Chemistry*, vol. 276, no. 50, pp. 47131–47135, 2001.
- [54] S. Laurora, S. Pizzimenti, F. Briatore, et al., "Peroxisome proliferator-activated receptor ligands affect growth-related gene expression in human leukemic cells," *Journal of Pharmacology and Experimental Therapeutics*, vol. 305, no. 3, pp. 932–942, 2003.
- [55] S. H. Chung, N. Onoda, T. Ishikawa, et al., "Peroxisome proliferator-activated receptor gamma activation induces cell cycle arrest via the p53-independent pathway in human anaplastic thyroid cancer cells," *Japanese Journal of Cancer Research*, vol. 93, no. 12, pp. 1358–1365, 2002.
- [56] W. Motomura, N. Takahashi, M. Nagamine, et al., "Growth arrest by troglitazone is mediated by p27^{Kip1} accumulation, which results from dual inhibition of proteasome activity and Skp2 expression in human hepatocellular carcinoma cells," *International Journal of Cancer*, vol. 108, no. 1, pp. 41–46, 2004.
- [57] W. Motomura, T. Okumura, N. Takahashi, T. Obara, and Y. Kohgo, "Activation of peroxisome proliferator-activated receptor γ by troglitazone inhibits cell growth through the increase of p27^{Kip1} in human pancreatic carcinoma cells," *Cancer Research*, vol. 60, no. 19, pp. 5558–5564, 2000.
- [58] S. Takeuchi, T. Okumura, W. Motomura, M. Nagamine, N. Takahashi, and Y. Kohgo, "Troglitazone induces G1 arrest by p27^{Kip1} induction that is mediated by inhibition of proteasome

- in human gastric cancer cells," *Japanese Journal of Cancer Research*, vol. 93, no. 7, pp. 774–782, 2002.
- [59] R.F. Morrison and S.R. Farmer, "Role of PPAR in regulating a cascade expression of cyclin-dependent kinase inhibitors, p18(INK4c) and p21(Waf1/Cip1), during adipogenesis," *The Journal of Biological Chemistry*, vol. 274, no. 24, pp. 17088–17097, 1999.
- [60] J. A. Copland, L. A. Marlow, S. Kurakata, et al., "Novel high-affinity PPAR γ agonist alone and in combination with paclitaxel inhibits human anaplastic thyroid carcinoma tumor growth via p21^{WAF1/CIP1}," *Oncogene*, vol. 25, no. 16, pp. 2304–2317, 2006.
- [61] S. Han, N. Sidell, P. B. Fisher, and J. Roman, "Up-regulation of p21 gene expression by peroxisome proliferator-activated receptor γ in human lung carcinoma cells," *Clinical Cancer Research*, vol. 10, no. 6, pp. 1911–1919, 2004.
- [62] G. D. Demetri, C. D. M. Fletcher, E. Mueller, et al., "Induction of solid tumor differentiation by the peroxisome proliferator-activated receptor- γ ligand troglitazone in patients with liposarcoma," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 96, no. 7, pp. 3951–3956, 1999.
- [63] E. Mueller, M. Smith, P. Sarraf, et al., "Effects of ligand activation of peroxisome proliferator-activated receptor γ in human prostate cancer," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 97, no. 20, pp. 10990–10995, 2000.
- [64] G. Debrock, V. Vanhentenrijk, R. Sciot, M. Debiec-Rychter, R. Oyen, and A. Van Oosterom, "A phase II trial with rosiglitazone in liposarcoma patients," *British Journal of Cancer*, vol. 89, no. 8, pp. 1409–1412, 2003.
- [65] L. D. Yee, N. Williams, P. Wen, et al., "Pilot study of rosiglitazone therapy in women with breast cancer: effects of short-term therapy on tumor tissue and serum markers," *Clinical Cancer Research*, vol. 13, no. 1, pp. 246–252, 2007.
- [66] M. H. Kulke, G. D. Demetri, N. E. Sharpless, et al., "A phase II study of troglitazone, an activator of the PPAR γ receptor, in patients with chemotherapy-resistant metastatic colorectal cancer," *Cancer Journal*, vol. 8, no. 5, pp. 395–399, 2002.
- [67] G. K. Schwartz, A. Weitzman, E. O'Reilly, et al., "Phase I and pharmacokinetic study of LY293111, an orally bioavailable LTB $_4$ receptor antagonist, in patients with advanced solid tumors," *Journal of Clinical Oncology*, vol. 23, no. 23, pp. 5365–5373, 2005.