

Title: The neurotrophic factor artemin and its receptor GFR α 3 mediate migraine-like pain via the ion channel TRPM8.

Chenyu Yang^{1,2}, Chao Wei^{1,3}, Sanaa Alam¹, Xunyang Chen¹, David D. McKemy^{1,2}

¹Neurobiology Section, Department of Biological Sciences, University of Southern California, Los Angeles, CA 90089; ²Molecular and Computational Biology Graduate Program, University of Southern California, Los Angeles, CA 90089; ³Department of Anesthesia and Critical Care, Beth Israel Deaconess Medical Center and Harvard Medical School, Boston, MA 02215.

Corresponding author:

David D. McKemy, Ph.D.

3641 Watt Way / HNB 201

University of Southern California

Los Angeles, CA 90089

(213) 821-5724

mckemy@usc.edu

ABSTRACT

Background

Migraine has a strong genetic foundation, including both monogenic and polygenic types. The former are rare, with most migraine considered polygenic, supported by genome-wide association studies (GWAS) identifying numerous genetic variants associated with migraine risk. Surprisingly, some of the most common mutations are associated with TRPM8, a non-selective cation channel that is the primary sensor of cold temperatures in primary afferent neurons of the somatosensory system. However, it is unlikely that the temperature sensitivity of TRPM8 underlies its role in migraine pathogenesis. To define the basis of the channel's involvement, we reasoned that cellular processes that increase cold sensitivity in the skin, such as the neurotrophic factor artemin, via its receptor GFR α 3, also mediate TRPM8-associated migraine-like pain in the meninges.

Methods

To investigate the role of artemin and GFR α 3 in preclinical rodent migraine models, we infused nitroglycerin acutely and chronically, and measured changes in periorbital and hind paw mechanical sensitivity in male and female mice lacking GFR α 3, after neutralization of free artemin with specific monoclonal antibodies, or by systemic treatment with a TRPM8-specific antagonist. Further, in wildtypes and mice lacking either GFR α 3 or TRPM8, we tested the effects of supradural infusions of a mix of inflammatory mediators, artemin, and a TRPM8-specific agonist on migraine-related pain in mice.

Results

We find that mechanical allodynia induced by systemic nitroglycerin, or supradural infusion of inflammatory mediators, involves GFR α 3. In addition, neutralization of circulating artemin reduces the nitroglycerin phenotype, demonstrating the importance of this neurotrophic pathway. Further, we show TRPM8 expression in the meninges and that direct supradural infusion of either a TRPM8-specific agonist or artemin itself produces mechanical allodynia, the latter dependent on TRPM8 and ameliorated by concurrent treatment with sumatriptan.

Conclusions

These results indicate that neuroinflammatory events in the meninges can produce migraine-like pain in mice via artemin and GFR α 3, likely acting upstream of TRPM8, providing a novel pathway that may contribute to migraine pathogenesis.

Keywords

TRPM8, Artemin, GFR α 3, trigeminal

INTRODUCTION

Migraine is a strongly heritable neurovascular disorder, yet little is known of the underlying genetic basis for common migraine^{1, 2}. To this end, multiple genome-wide association studies (GWAS) have searched for single-nucleotide polymorphisms (SNPs) linked to either protection from migraine or associated with increased migraine susceptibility³⁻⁷. Surprisingly, some of the most common mutations reside in or near the genetic locus of the menthol receptor, *transient receptor potential melastatin 8 (trpm8)*³⁻⁸, a cold-gated ion channel highly expressed in trigeminal ganglia (TG)^{9, 10}. While its role in cold somatosensation and nociception in the skin is well-studied¹¹, how TRPM8 channels might contribute to migraine pathogenesis is unclear¹².

The SNPs identified in *trpm8* reside in noncoding regions, suggesting they do not alter function but may influence channel expression^{5, 7, 13-20}. For example, the reduced migraine risk allele rs10166942[C] is located 5' of the start codon^{3, 5} and homozygous individuals express less TRPM8 and are less sensitive to cold than those with a T at this position²¹. Intriguingly, this T allele, which is prevalent in individuals from higher latitudes and proposed to arise from positive selection for a thermoregulatory advantage in colder climates²², is associated with increased migraine predisposition^{23, 24}. Thus, increased TRPM8 expression may correlate with increased migraine risk. What is also intriguing is that cold triggers headaches in humans (headache-induced by cold stimuli (HICS)), suggesting a link between headaches and TRPM8 activity²⁵.

Given its relative novelty as a migraine susceptibility gene, only a few preclinical studies have examined TRPM8 function related to migraine²⁶⁻³⁰. Previously, using both mouse genetics and TRPM8-specific pharmacology, we showed that TRPM8 is required

for both chronic and acute migraine-like phenotypes induced by systemic application of either nitroglycerin (NTG) or calcitonin gene-related peptide (CGRP)³¹. However, the molecular mechanisms whereby TRPM8 channels and cells mediate migraine-like pain are not fully understood. To investigate this, we reasoned that processes underlying TRPM8-mediated cold allodynia and hyperalgesia in the skin could also facilitate the channel's sensitization in migraine-relevant tissues. Specifically, the glial cell-line derived neurotrophic factor (GDNF) artemin and its receptor GFR α 3 are required for TRPM8-dependent cold allodynia and hyperalgesia in multiple pain models³²⁻³⁶. Further, artemin expression is present in meningeal vascular smooth muscle and GFR α 3 expressing nerve fibers are found throughout the dura mater^{37, 38}. Expression of both meningeal GFR α 3 and vascular artemin in the rat increases after nitroglycerin (NTG) treatment³⁸, and artemin is proposed to increase expression of the inducible form of nitric oxide synthase (iNOS) in trigeminal neurons³⁹. Thus, there is circumstantial evidence for a role of artemin/GFR α 3 signaling in the meninges that may correlate with migraine pathogenesis.

Here, we report that, like TRPM8, GFR α 3 is required for NTG-induced acute and chronic tactile allodynia, a phenotype that is also attenuated by neutralization of artemin. Further, GFR α 3 is involved in allodynia produced by a supradural infusion of a mixture of inflammatory mediators. Lastly, application of either a TRPM8 agonist or artemin directly onto the mouse dura matter leads to similar phenotypes that are TRPM8-dependent, as well as sensitive to the abortive migraine agent sumatriptan. Our findings suggest a conserved mechanism underlies both TRPM8-dependent cold allodynia and migraine-like pain, highlighting a novel molecular pathway that may, in part, contribute to migraine headaches.

MATERIALS & METHODS

Animals

All experiments were approved by University of Southern California Institutional Animal Care and Use Committee and in compliance with ARRIVE guidelines for Animal Research: Reporting of In Vivo Experiments ⁴⁰. Mice aged from 8 weeks to 16 weeks were used in all experiments and unless otherwise specified, equal numbers of male and female mice were used. All mice were housed in a temperature-controlled environment on a 12-hour light and dark cycle with ad libitum access to food and water. All experiments were performed during the light cycle from 8 AM to 5 PM. Trpm8-null mice (Trpm8^{-/-}, Strain # 008198) ⁴¹ were purchased from Jackson Laboratories and were in the C57Bl/6 background. Wildtype C57Bl/6 and Trpm8^{GFP} ¹⁰ mice were maintained in-house, as were Gfrα3-null (Gfrα3^{-/-}) mice ⁴² that were bred from Gfrα3^{+/-} animals to obtain Gfrα3-nulls and wildtype littermate controls, as described ³⁴.

Supradural Injections

Supradural injections were performed as previously described ⁴³. Briefly, mice were anesthetized with 3%-4% isoflurane and then maintained by 1.5%-2% isoflurane through a nose cone. The various agents used were injected in a volume of 5 µl using an internal cannula (Invivo1, part #C313IS-5/SPC, Internal Canula, 28 gauge) connected to plastic tubing (Perkin Elmer, Cat#N0695476, 2-stop, I.D. 0.19 mm) and a 10 µl glass syringe (Hamilton, Cat# 80300). The inner projection of the cannula was adjusted to be 0.6 mm to inject through the soft tissue at the intersection of lambdoidal and sagittal sutures but

not puncture the dura. Post-mortem analysis was done to confirm that the dura remained intact after each injection.

Retrograde Labeling of Trigeminal Neurons Innervating Mouse Dura

Trpm8^{GFP} mice were anesthetized and 5 μ l supradural injections of the retrograde tracer wheat germ agglutinin (1 mg/ml, WGA-Alexa-350; Cat#W11263, ThermoFisher) were performed as described ⁴³. After recovery from anesthesia, mice were housed in their cages for 14 days to allow for the transport of the tracers to the dural afferent somata in the trigeminal ganglia (TG).

Tissue Preparation and Immunohistochemistry

TG were dissected from Trpm8^{GFP} mice injected with WGA tracer as previously described ¹⁰. Briefly, the mice were euthanized and perfused transcardially with Phosphate Buffer Solution (PBS) followed by fixative (cold 4% paraformaldehyde solution, PFA). Tissues were carefully removed and post-fixed in the same fixative for another 2 hours at 4°C. After cryoprotection in 30% sucrose solution for at least 24 hours, the tissues were fast-frozen in OCT and sectioned with a cryostat at 16 μ m. The tissue sections were mounted on Superfrost Plus slides and kept at -80°C. On the day of immunostaining, frozen slides were dried at 25°C for 30 mins, washed with deionized water and PBS for three times, and permeabilized in 0.1% Triton + PBS (PBST) for 30 mins. The mounted sections were blocked in 0.1% PBST solution with 20% Normal Goat Serum (NGS) for 1 hour. The slides were incubated at 4°C overnight with primary antibodies in 0.1% PBST solution with 0.5% BSA + 0.5% NGS (Chicken anti-GFP antibody, 1:500, Cat# GFP-1020, Aves Lab; Rabbit

anti-CGRP antibody, 1:1000, Cat# C8198, ThermoFisher). The slides were washed in incubation buffer 6x5 mins and incubated in secondary antibodies for another 2 hours at room temperature (Goat anti-Chicken-IgY antibody, Alexa-488, 1:2000, Cat# A11039, ThermoFisher; Goat anti-Rabbit-IgG antibody, CF594, 1:2000, Cat# SAB4600234, Sigma Aldrich). The slides were washed in incubation buffer 6x5 mins followed by 2x5 mins PBS and covered by media and coverslips. The slides were dried and kept at 4°C overnight before imaging.

The meninges (including dura mater) were dissected from Trpm8^{GFP} mice using a protocol adapted from previous reports ⁴⁴. Briefly, mice were euthanized and perfused transcardially with ice-cold PBS. Skulls with meninges attached were removed and fixed in 2% PFA solution at 4°C overnight. The meninges were carefully detached from skulls in PBS solution the next day and transferred to a 24-well plate. The tissues were blocked for 1 hour in 0.1% PBST solution with 2% NGS + 1% BSA and then incubated overnight at 4°C in 0.1% PBST solution with 1% BSA and primary antibodies. After 3x5 mins wash with PBS, the tissues were incubated in 0.1% PBST solution with 1% BSA and secondary antibodies at room temperature for 2 hours. The tissues, after 3x5 mins wash with PBS, were flattened and dried in the dark on Superfrost Slides for 10-15 mins, covered with appropriate media and coverslips and imaged immediately.

Image Acquisition and Analysis

Digital images were acquired using a Zeiss Axio Imager.M2 (Oberkochen, Germany) with Apotome attachment and Zeiss Zen Blue software. Quantification of overlap between expression of neuronal markers, retrograde tracers, and GFP was obtained per field and

expressed as percentage overlap with the mean \pm standard error of the mean between fields.

Pharmacological agents and migraine-like pain models

Nitroglycerin (NTG) was purchased from VWR (Cat# 101095-632) and supplied as 1% solution in Propylene Glycol (PG) and further diluted to the desired working concentration (1 mg/kg or 10 mg/kg) using a 0.9% saline solution. NTG or vehicle was administered intraperitoneally (i.p.) every other day for 9 days, as described^{31, 45}, with mechanical sensitivity measured before and 2 hours after each injection. Icilin was purchased from Tocris Bioscience (Cat# 1531), dissolved, and diluted to 1 nmol / 5 μ l in PG and injected supradurally as described above. PBMC was purchased from Focus Biomolecules (Cat# 10-1413), dissolved, and diluted to working concentration (4 mg/kg) using 10% DMSO in a 0.9% saline solution and injected i.p. as described^{32, 46}. Artemin was purchased from R&D Systems (Cat# 1085-AR), dissolved, and diluted to 1 pg / 5 μ l using a Hank's Balances Salt Solution (HBSS: 140 mM NaCl, 5 mM KCl, 1 mM CaCl₂, 0.4 mM MgSO₄, 0.5 mM MgCl₂, 0.3 mM Na₂HPO₄, 0.4 mM KH₂PO₄, 6 mM D-glucose, 4 mM NaHCO₃) and injected supradurally. The inflammatory soup (IS) components were purchased from Sigma Aldrich, consisting of 1 mM Bradykinin (Cat# 05-23-0500), 1 mM 5-HT (Cat# H9523), 1 mM Histamine (Cat# 59964), 200 μ M PGE₂ (Cat# P5640) and at pH 5.0 in Synthetic Interstitial Fluid (SIF: 10 mM HEPES, 5 mM KCl, 135 mM NaCl, 1 mM MgCl₂, 2 mM CaCl₂, 10 mM glucose). Vehicle solutions were used as control injections in experiments accordingly. The IS or vehicle was injected once supradurally and mechanical sensitivity was measured from one to six hours post-injection, then again

twenty-four hours later. The anti-artemin monoclonal and control IgG Isotype monoclonal antibodies were purchased from R&D Systems (Artemin: MAB1085, Control: MAB002) and reconstituted in sterile PBS solution. The antibodies were injected intraperitoneally at a concentration of 10 mg/kg. Sumatriptan was purchased from Sigma Aldrich (Cat# S1198), diluted using 0.9% Saline solution, and injected subcutaneously (s.c.) at a concentration of 0.6 mg/kg.

von Frey Assay

The von Frey assay was used to assess the sensitivity to mechanical stimulation as previously described^{31, 43}. Mechanical sensitivity (50% withdrawal threshold) was determined at the timepoints indicated with calibrated von Frey monofilaments (Semmes-Weinstein) using the up-and-down method. Briefly, the mice tested were conditioned by handling for 5 minutes and habituated to plexiglass chambers on metal mesh (hindpaw) or horizontal paper cups (periorbital assay) for 3 consecutive days before experiments. The filaments (for hind paw assay: starting from 0.4 g filament; flexible force range of 0.008-2 g; for periorbital assay: starting from 0.07 g filament, flexible force range of 0.008g-0.6 g) were applied to the plantar surface of the hind paw or the center of the face between eyes for up to 5-6 seconds until the filament bent or positive response was recorded. The next heaviest (if no response) or lightest (if a response) filament was used in the next measurement and 4 additional measurements were performed. The 50% withdrawal threshold was calculated using the SUDO method previously described⁴⁷. The mice were returned to their home cage after measurements and retested on the next treatment/test day.

Statistical Analysis

The experimenters were blinded to either genotype of the mice receiving treatments or substances injected. Unless otherwise specified, eight to ten mice were tested in each group and equally balanced between male and female mice. No sex differences were observed in the data. All data are presented as mean \pm standard error of the mean (SEM) with the statistical assay used described in the figure legend or associated text, along with sample numbers. Time course measurements and comparisons between different injection conditions such as treatment or genotype were made with were determined via a two-way ANOVA with either Bonferroni's, Dunnett's, or Tukey's multiple comparison tests. Two-tailed paired t-tests were used to compare pre- versus post-injection conditions with the same injection conditions or genotype.

RESULTS

TRPM8 afferents are present in the mouse dura.

To further investigate the role of TRPM8 in migraine-like pain, we confirmed that TRPM8-positive afferents are present in the mouse meninges, first using retrograde tracer back-labeling⁴³. Naïve adult *Trpm8*^{GFP} mice¹⁰ received a supradural injection of the retrograde tracer wheat germ agglutinin (1%; WGA-Alexa-dye) and TG were processed for immunolabeling 14 days post-injection as described^{10, 34, 35, 48}. We found that $15.7 \pm 2.7\%$ of WGA-labeled cells were GFP positive (Fig. 1A, n=4 mice; 805 WGA-positive cells), consistent with the proportion of TG neurons per ganglia known to express TRPM8^{9, 10, 34, 49, 50}. Next, we performed immunolabeling of dura dissected from adult *Trpm8*^{GFP} mice as described^{10, 34, 51}, observing GFP expression in *Trpm8*^{GFP} animals (Fig. 1B), labeling that was absent in wildtype animals (Fig. 1C). These results are consistent with prior studies showing TRPM8 expressed in this tissue⁵². Of note, a proportion of GFP-positive axons also showed immunoreactivity for CGRP, consistent with soma co-expression of TRPM8 and CGRP in the trigeminal ganglia^{9, 10, 34, 49, 50}.

GFR α 3 is required for nitroglycerin-induced tactile allodynia.

Next, we asked how activation of TRPM8 channels in the meninges, which are likely not responding to temperature in this tissue, might lead to migraine-like pain. We reasoned that mechanisms known to induce TRPM8-dependent cold allodynia in the skin might also be relevant in the dura matter. The neurotrophic factor artemin, by binding to its cognate receptor GDNF Family Receptor α 3 (GFR α 3), can induce TRPM8-dependent cold hypersensitivity, and GFR α 3 is also required for or involved in inflammatory and

neuropathic cold pain of various etiologies, including migraine³²⁻³⁵. To test this premise, we measured acute periorbital mechanical allodynia in wildtype and $GFR\alpha3$ -null mice ($Gfr\alpha3^{-/-}$)^{34, 42} 2 hours after they were given an intraperitoneal injection (i.p.) of 1 mg/kg NTG or vehicle, repeated every 2 days^{31, 45} (Fig. 2A). As expected, wildtype animals exhibited mechanical allodynia after each NTG injection that was significantly different than baseline (BL) at all days tested, as well as compared to vehicle injected mice by the 2nd injection on Day 3 ($p < 0.01$, 2-way ANOVA with Tukey's multiple comparisons test). In contrast, $Gfr\alpha3^{-/-}$ mice were insensitive to NTG treatment as their mechanical sensitivity was not different than baseline or compared to vehicle injected animals ($p > 0.05$, 2-way ANOVA with Tukey's multiple comparisons test) at all days tested. To ensure that the lack of a phenotype in $Gfr\alpha3^{-/-}$ mice wasn't due to an NTG dosage effect, we also tested their acute response to 10 mg/kg NTG, again finding that the robust increase in mechanical allodynia in wildtype mice 2 hours after a single injection was absent in $Gfr\alpha3^{-/-}$ animals (Fig. 2B). Lastly, to determine if $GFR\alpha3$ was also required for extracephalic allodynia induced by NTG, we measured hind paw mechanical responses in wildtype and $Gfr\alpha3^{-/-}$ mice before and 2 hrs after an i.p. injection of 10 mg/kg NTG, observing no differences in $Gfr\alpha3^{-/-}$ mice given NTG (Fig. 2C).

Repeated NTG dosing leads to an increase in basal mechanical sensitivity that is considered a potential model of chronic migraine^{31, 45}. To test the role of $GFR\alpha3$ in chronic migraine, we compared baseline behaviors at each injection day prior to treatment with 1 mg/kg NTG or vehicle, observing the expected increase in basal allodynia in wildtypes, an effect that was absent in $Gfr\alpha3^{-/-}$ animals (Fig. 3A). We again assessed 10 mg/kg NTG, also observing that basal mechanical allodynia was absent in $Gfr\alpha3^{-/-}$ mice 2 days after a

single NTG injection (Day 3, Fig. 3B). Similarly, hindpaw mechanical allodynia did not develop in $Gfr\alpha3^{-/-}$ mice given 10 mg/kg NTG (Fig. 3C). Thus, $GFR\alpha3$ is required for both cephalic and extracephalic tactile allodynia in both acute and chronic models of rodent migraine-like pain, a phenotype like that observed in TRPM8-null mice ³¹.

Neutralizing artemin reduces nitroglycerin-induced tactile allodynia.

Our results show that $GFR\alpha3$ is required for NTG-induced tactile allodynia, suggesting that its specific ligand artemin might also play a role in this phenotype. To test this, we employed monoclonal antibodies (mAb) raised against artemin that we have shown can specifically ameliorate pathological cold allodynia. ^{32-34, 53}. Wildtypes were tested for basal cephalic mechanical sensitivity then injected with either an anti-artemin mAb or an isotype control. The mice were re-tested at 1 hour post-injection, demonstrating that mAb treatment did not alter basal mechanosensation (Fig. 4A). Immediately after testing, mice treated with the two mAbs were injected with 1 mg/kg NTG and re-tested 2 hours post-injection. Mice given the isotype control showed robust mechanical allodynia, as expected, whereas mice injected with the anti-artemin mAb exhibited normal sensitivity (Fig. 4A).

Next, we asked if chronic allodynia induced by NTG was sensitive to artemin neutralization. Animals were tested for basal mechanical sensitivity (Day 1) and given a single injection of 1 mg/kg NTG. Their basal sensitivity was remeasured 2 days later (Day 3), with all animals exhibiting mechanical allodynia (Fig 4B). These mice were then given either an injection of the isotype control or anti-artemin mAbs and tested 1 hour later. As with acute sensitization, artemin neutralization decreased mechanical allodynia, bringing

the 50% withdrawal threshold back to basal levels. These data indicate that both acute and the initial phase of chronic mechanical allodynia induced by NTG are dependent on free, circulating artemin, as well as suggest that established allodynia can be alleviated by artemin neutralization.

Dura artemin treatment induces TRPM8-dependent tactile allodynia.

Our results suggest a key role for artemin and GFR α 3 signaling in the NTG model of migraine-like pain but doesn't directly determine if GFR α 3 is acting at sites involved in migraine, such as the meninges. To address this, we first asked if periorbital mechanical allodynia induced by stimulation of the meninges via direct supradural injection of a cocktail of inflammatory mediators required GFR α 3⁴³. Wildtype and Gfr α 3^{-/-} mice were tested for basal (BL) mechanical sensitivity as above, then given a 5 μ l injection of either vehicle (SIF; synthetic interstitial fluid) or an inflammatory soup (IS) mixture (in SIF at pH 5.0). Both wildtype and Gfr α 3^{-/-} mice given IS showed robust mechanical allodynia, compared to vehicle injected mice (wildtypes: p<0.05 at 1 hour, p<0.0001 at 2-4 hours post-injection; Gfr α 3^{-/-}: p<0.01 at 2 hours, p<0.05 at 3hrs, two-way ANOVA with Tukey's multiple comparison tests) and basal sensitivity (wildtypes: p<0.01 at 1 hour, p<0.0001 at 2 to 4 hours post-injection; Gfr α 3^{-/-}: p<0.01 at 2 hours, p<0.001 at 3 hours, p<0.05 at 4 hours post-injection; two ANOVA with Dunnett's multiple comparison test), for several hours post-injection. However, the level of sensitization after IS injection was significantly lower in Gfr α 3^{-/-} mice compared to wildtypes at 2-4hrs post-injection (Fig. 5A), indicating that GFR α 3 mediates, in part, migraine-like pain produced by meningeal stimulation with inflammatory mediators.

Next, we asked if direct application of artemin to the meninges in wildtype mice could induce a similar migraine-like phenotype. Naïve wildtype mice were tested for baseline periorbital mechanical sensitivity, given a supradural injection of either vehicle (HBSS) or 1 pg (5 μ l) artemin, then retested for mechanical allodynia over the next 96 hours. Consistent with our $Gfr\alpha3^{-/-}$ and artemin neutralization results, we observed robust allodynia in mice receiving artemin that lasted over 24 hours post-injection (Fig. 5B). Thus, like other inflammatory agents, artemin directly induces tactile allodynia when applied to the dura. Next, as artemin induces TRPM8-dependent cold hypersensitivity in the skin^{32, 34, 35}, we reasoned that artemin and $GFR\alpha3$ signaling underlies the TRPM8-dependent migraine-like phenotypes we observed³¹. To test this hypothesis, we repeated the dura artemin injections in TRPM8-null mice ($Trpm8^{-/-}$)⁴¹, finding that this treatment did not induce mechanical allodynia in the absence of TRPM8 (Fig. 5B). Thus, as with cold allodynia in the skin, artemin-induced sensitization in the meninges is TRPM8 dependent. Lastly, to further define this artemin phenotype, we concurrently treated wildtype mice with supradural artemin and 0.6 mg/kg (subcutaneously) sumatriptan, the latter a traditional migraine treatment in humans and an agent used in animal studies to determine if a phenotype is migraine-like pain⁵⁴⁻⁵⁶. Mice given artemin and the vehicle for sumatriptan (0.9% Saline) again showed robust mechanical allodynia starting at 2 hours post-injection ($p < 0.5$ at 2 and 6 hours, $p < 0.0001$ at 3 and 4 hours, $p < 0.01$ at 5 hours post-injection, $n = 8$, two-way ANOVA with Tukey's multiple comparisons test). In contrast, mice given artemin and sumatriptan exhibited reduced mechanical allodynia from 2 to 4 hours post-injection, compared to artemin/vehicle treated mice (Fig. 5C), results suggesting that the artemin phenotype is consistent with migraine-like pain.

Activation of meningeal TRPM8 induces cephalic tactile allodynia independent of GFR α 3.

As we find that artemin neutralization alleviates both acute and chronic tactile allodynia induced by NTG, and artemin-induced mechanical allodynia is TRPM8-dependent, we posit that this effect of artemin is via TRPM8 sensitization. Previously, we showed that acute extracephalic allodynia induced by 10 mg/kg NTG could be prevented by pre-treatment with the TRPM8 antagonist PBMC, but that established chronic NTG measured after 4 NTG treatments given every two days was insensitive to TRPM8 inhibition³¹. Although inconsistent with our results in Fig. 4B, these prior results were with a substantially higher dose of NTG (10mg/kg versus 1mg/kg) given repeatedly. Thus, to determine if TRPM8 underlies the basal sensitization shown in Fig. 4B, wildtype mice were given a single 1 mg/kg NTG injection, then tested for basal allodynia 2 days later (BL-D3) with all animals exhibiting sensitization (n=13, paired two-tailed t-test, $p < 0.0001$). These mice were randomly separated into two groups with one receiving an injection of 4 mg/kg PBMC and the other vehicle, then tested for mechanical sensitivity every hour for 5 hours. Mice given vehicle remained allodynic over this time period as there were no significant differences in mechanical sensitivity compared to the Day 3 baseline (D3-BL; $p > 0.05$, two-way ANOVA with Dunnett's multiple comparison test) and remained significantly different than the responses measured pretreatment (BL) at all time points tested ($p < 0.01$ two-way ANOVA with Dunnett's multiple comparison test). Conversely, mice given PBMC showed transient relief from this allodynia by 2 hours post-PBMC compared to the Day 3 baseline and to vehicle injected animals (Fig. 6A). Thus, inhibition of TRPM8 systemically relieved the initial NTG induced basal mechanical allodynia in this

chronic migraine model. Further, to contrast with our prior results, mice were similarly given 4 injections of 1 mg/kg NTG at 2-day intervals, then their baseline (BL-D9) was measured 2 days after the 4th injection (Day 9), with all animals exhibiting robust allodynia (Fig. 6A, paired two-tailed t-test, $p < 0.0001$). In contrast to its effects on Day 3, mice given 4 mg/kg PBMC on Day 9 remained allodynic with periorbital mechanical allodynia that was like vehicle treated animals (Fig. 6A). These results are consistent with previous findings and demonstrate that, while the initial basal allodynia induced with 1 dose of NTG is TRPM8 and artemin dependent, the robust phenotype developed by repeated NTG treatments is independent of TRPM8.

These data, and evidence that mouse meningeal afferents express TRPM8 (Fig. 1), suggest that sensitization of TRPM8 channels by artemin and GFR α 3 signaling in this tissue leads to migraine-like phenotypes in the mouse, but it is unclear if these are functional channels. Thus, we next asked if direct application of the potent and selective TRPM8 agonist icilin^{9,57} to the mouse dura could induce behaviors consistent with mouse migraine-like phenotypes. Previously, Burgos-Vega et al. (2016) demonstrated that supradural application of icilin in the rat induced both cephalic and extracephalic mechanical allodynia that was inhibited by co-application of sumatriptan and L-NAME²⁶. While this effect was sensitive to TRPM8 antagonism, the genetic dependence of this icilin phenotype in mice has not been established. Thus, we directly applied icilin (1 nmol in 5 μ l) to the mouse dura and measured changes in periorbital mechanical sensitivity pre- and post-injection, observing robust allodynia that was significantly different than vehicle injected mice by 2 hours, lasting up to 5 hours before returning to baseline (BL) by 6 hours post-injection (Fig. 6B). This time course is consistent with previous findings

in the rat ²⁶. We also tested extracephalic sensitivity in the mouse hindpaw, observing mechanical allodynia at 3 hours post-injection, sensitization that was absent in *Trpm8*^{-/-} mice (Fig. 6C). Lastly, in a final test of this hypothetical pathway, we predict that artemin and GFR α 3 act upstream of TRPM8 and thus the dura icilin induced allodynia should occur independent of GFR α 3, a hypothesis supported by clear periorbital tactile allodynia in *Gfra3*^{-/-} mice treated with supradural icilin (Fig. 6D). Thus, TRPM8 expressed in the mouse meninges is functional and activation of these channels leads to both cephalic and extracephalic mechanical allodynia that is consistent with preclinical rodent migraine models.

DISCUSSION

TRPM8 is the principal detector of environmental cold in the skin but, like the heat-sensitive channel TRPV1, is also present in afferents innervating tissues not involved in thermosensation^{58, 59}. For example, TRPM8-positive afferents extensively innervate the rodent bladder and contain nociceptor markers such as CGRP at a higher percentage than the overall TRPM8-positive somas per ganglion, consistent with the channel's role in bladder dysfunction⁶⁰⁻⁶². Similarly, TRPV1 is expressed in visceral and muscle afferents in larger proportions compared to those innervating the skin⁶³. Thus, while present and functional, the thermosensitivity of these receptors is likely not relevant in these tissues, indicating that endogenous agonists or modulators sensitize or activate these channels to assert their physiological roles in regions other than the skin^{64, 65}. For example, agonists like menthol and capsaicin sensitize the temperature sensitivity of thermoTRPs like TRPM8⁹ and TRPV1⁶⁶, respectively, such that the channels gate at physiological temperatures⁶⁷. Similarly, several signal transduction cascades induced by inflammation sensitize TRPV1 such that the channel is active at skin temperatures^{68, 69}. Thus, their sensitivity to endogenous and exogenous stimuli likely underlies their potential contribution to migraines⁶⁴. For TRPM8, artemin is the only known endogenous agent that leads directly to TRPM8-dependent hypersensitivity, a pathway that has only been studied in relation to cutaneous cold sensation¹¹. Here, we provide compelling evidence that this pathway also underlies a mechanism whereby TRPM8 sensitization may lead to migraine-like pain.

Artemin is a ligand of the glial cell line-derived neurotrophic factor (GDNF) family and plays crucial roles in nervous system development, nociception, and the

pathogenesis of neurological disorders ⁷⁰. At the transcriptional level, artemin mRNA is expressed in developing nerve roots, Schwann cells, vascular and visceral smooth muscle cells, endothelial cells, skin fibroblasts, and immune cells, indicating its potential release from these sites ^{71, 72}. The GDNF Family Receptor Alpha 3 (GFR α 3) is a high affinity receptor for artemin and shows a similar elevated expression pattern in sensory ganglia, smooth muscle cells, skin fibroblasts, and endothelial cells ⁷². Notably, both artemin and GFR α 3 are expressed in dura afferents, with elevated levels in rodent models of migraine, suggesting a significant role in migraine pathogenesis ^{37, 38}. Intriguingly, our previous work has demonstrated that artemin, through GFR α 3, induces nociception, including cold hypersensitivity, in vivo ³²⁻³⁵. This effect is TRPM8-dependent, with GFR α 3 expressed in approximately half of TRPM8 neurons in the trigeminal ganglion (TG), and both inflammatory and neuropathic cold pain are absent in GFR α 3-null mice ^{32, 34}. Peripheral ARTN neutralization alleviates established inflammatory and neuropathic cold pain, highlighting a potential therapeutic intervention ^{32, 34, 53}. These findings suggest similarities in ARTN/GFR α 3 signaling in both cold and trigeminovascular pain.

Nitroglycerin (NTG), a nitric oxide (NO) donor, can induce headaches in both migraineurs and non-migraineurs by sensitizing the trigeminovascular system, causing vasodilation of meningeal blood vessels, and promoting the release of proinflammatory factors ⁷³⁻⁷⁵. In this study, we administered a moderate dose of NTG (1 mg/kg, one-tenth of the commonly used dosage ⁴⁵) to induce a significant tactile allodynic phenotype in the cephalic region of mice, minimizing the risk of overwhelming sensitization. We found that the absence of GFR α 3 and the neutralization of artemin via a monoclonal antibody prevented acute migraine-like pain triggered by NTG, suggesting that systemic NTG

administration could induce artemin release and interaction with GFR α 3 at the meninges. Additionally, a chronic migraine-like phenotype established by repeated NTG treatment was also attenuated by artemin neutralization or the absence of GFR α 3, indicating sustained increased artemin levels facilitated baseline hypersensitivity in the rodent model of chronic migraine⁴⁵. These data compellingly demonstrate that the NTG-induced migraine-like phenotype is highly dependent on artemin release and signaling through GFR α 3. This significant correlation is underscored by the observed elevation of artemin and GFR α 3 levels in the dura mater of NTG-induced rodent migraine models and trigeminal ganglion cultures³⁹. Artemin, which is abundantly expressed in vascular smooth muscle cells, promotes vasodilation by relaxing these cells, and its level positively correlates with inducible nitric oxide synthase (iNOS) levels³⁹. Consequently, neutralizing artemin could mitigate NTG-induced vasodilation, a primary contributor to nociception in this model. Furthermore, the excessive production of nitric oxide in the NTG model can trigger an inflammatory response and subsequent neurogenic inflammation⁷⁶. The precise molecular mechanisms linking NTG effects and artemin release remain to be elucidated, and uncovering these relationships is crucial for developing future treatments.

A limitation of the NTG model is the systemic nature of its effects, making it difficult to determine if the phenotype is primarily produced in the meninges. However, we found that direct application of artemin to the mouse dura mater triggers cephalic tactile allodynia, a TRPM8-dependent phenotype. Artemin induced allodynia was also inhibited by concurrent treatment with sumatriptan, further supporting this phenotype to be migraine-like. As shown here and in other reports, TRPM8-positive afferents are functionally expressed in mouse meninges, and activation of this channel in this tissue

leads to both cephalic and extracephalic tactile allodynia^{26, 27, 52, 77}. This phenotype is reminiscent of our prior work showing that peripheral nociceptor activation by endogenous proalgesics leads to artemin-dependent cold allodynia³² and that subcutaneous artemin application leads to TRPM8-dependent cold pain³⁵. It is also essential to recognize that artemin might not account for all migraine episodes. For instance, in the inflammatory soup migraine model, deleting the GFR α 3 receptor only partially alleviates cephalic tactile allodynia induced by inflammatory factors. Hence, the development of migraine-like pain likely involves multiple molecular pathways, necessitating a therapeutic approach that targets multiple pathogenic pathways for effective treatment.

Our results suggest that activating TRPM8 leads to migraine-like pain, but cooling is also reported to have a short-term effect alleviating headaches, and menthol modestly relieves migraines when applied either intranasally or to the forehead^{28-30, 78-80}. Further, one report suggests TRPM8 has a protective effect and contributes to a faster recovery in the chronic migraine model in males but not females⁸¹. TRPM8 is the predominant cold sensor in mammals, mediating innocuous cool and noxious cold sensation both acutely and in pathological settings^{9, 33, 41, 46, 48, 82-84}. Conversely, modest thermal or chemical cooling of the skin inhibits both inflammatory and neuropathic pain^{48, 85-87}. These same stimuli also alleviate acute and chronic itch, airway irritation, and aversion to oral nicotine⁸⁸⁻⁹¹. These processes are TRPM8-dependent and likely occur via TRPM8 afferent activation of spinal cord dorsal horn inhibitory interneurons, suggesting a similar mechanism in the trigeminocervical complex (TCC) may provide headache relief^{89, 92}. As cephalic and extracephalic cutaneous allodynia with migraines is proposed to result from spinal and supraspinal central sensitization, respectively⁹³⁻⁹⁶, stimulation of cephalic

cutaneous TRPM8 afferents, which converge with meningeal afferents in the TCC, may provide an inhibitory input that reduces sensitization produced in the meninges. Therefore, in the context of migraine-like pain, there may be a distinction between the activity of TRPM8-expressing afferents in the facial skin and those in the meninges, an intriguing posit that requires further investigation.

How can TRPM8 activation lead to migraine-like pain when its only known role in nociception is related to cold sensation? The prevailing model for the sequence of events leading to migraine suggests that a variety of factors alter cortical excitability that eventually leads to activation of meningeal trigeminal nociceptors which induce sensitization of 2nd order neurons in the brainstem ^{97, 98}. Our results suggest that artemin/GFR α 3 sensitization of TRPM8 in the meninges is one such pathway that can result in migraine and headaches. Further, when considered in the context of the TRPM8 variants identified in GWAS on migraine ^{21, 23}, individuals with elevated TRPM8 expression would be more susceptible to an increase in artemin or other endogenous agents that sensitize TRPM8 channels, perhaps underlying their predisposition to migraines ²¹. This hypothesis requires further study, but our results now implicate artemin/GFR α 3/TRPM8 as novel players in the molecular mechanisms of migraine.

Acknowledgements

The authors wish to thank current and former members of the McKemy Laboratory for their assistance in the completion of this work.

Declaration of conflicting interests

The authors declare no financial or conflicts of interest.

Funding

The present study was funded by the National Institutes of Health - National Institute of Neurological Disorders and Stroke Awards #R01NS106888 and #R21NS118852.

References

1. Anttila V, Wessman M, Kallela M, et al. Genetics of migraine. *Handb Clin Neurol* 2018; 148: 493-503. 2018/02/27. DOI: 10.1016/B978-0-444-64076-5.00031-4.
2. Peck KR, Johnson YL and Smitherman TA. Migraine. *Handb Clin Neurol* 2016; 138: 283-293. 2016/09/18. DOI: 10.1016/B978-0-12-802973-2.00016-1.
3. Chasman DI, Schurks M, Anttila V, et al. Genome-wide association study reveals three susceptibility loci for common migraine in the general population. *Nat Genet* 2011; 43: 695-698. 2011/06/15. DOI: 10.1038/ng.856.
4. Gormley P, Anttila V, Winsvold BS, et al. Meta-analysis of 375,000 individuals identifies 38 susceptibility loci for migraine. *Nat Genet* 2016; 48: 856-866. 2016/06/21. DOI: 10.1038/ng.3598.
5. Freilinger T, Anttila V, de Vries B, et al. Genome-wide association analysis identifies susceptibility loci for migraine without aura. *Nat Genet* 2012; 44: 777-782. 2012/06/12. DOI: 10.1038/ng.2307.
6. Chasman DI, Anttila V, Buring JE, et al. Selectivity in genetic association with sub-classified migraine in women. *PLoS Genet* 2014; 10: e1004366. 2014/05/24. DOI: 10.1371/journal.pgen.1004366.
7. Esserlind AL, Christensen AF, Le H, et al. Replication and meta-analysis of common variants identifies a genome-wide significant locus in migraine. *Eur J Neurol* 2013; 20: 765-772. 2013/01/09. DOI: 10.1111/ene.12055.
8. Key FM, Abdul-Aziz MA, Mundry R, et al. Human local adaptation of the TRPM8 cold receptor along a latitudinal cline. *PLoS Genet* 2018; 14: e1007298. 2018/05/04. DOI: 10.1371/journal.pgen.1007298.

9. McKemy DD, Neuhausser WM and Julius D. Identification of a cold receptor reveals a general role for TRP channels in thermosensation. *Nature* 2002; 416: 52-58. 2002/03/08. DOI: 10.1038/nature719.
10. Takashima Y, Daniels RL, Knowlton W, et al. Diversity in the neural circuitry of cold sensing revealed by genetic axonal labeling of transient receptor potential melastatin 8 neurons. *J Neurosci* 2007; 27: 14147-14157. 2007/12/21. DOI: 10.1523/JNEUROSCI.4578-07.2007.
11. McKemy DD. Molecular basis of peripheral innocuous cold sensitivity. *Handb Clin Neurol* 2018; 156: 57-67. 2018/11/21. DOI: 10.1016/B978-0-444-63912-7.00003-5.
12. Dussor G and Cao YQ. TRPM8 and Migraine. *Headache* 2016; 56: 1406-1417. 2016/09/17. DOI: 10.1111/head.12948.
13. Kaur S, Ali A, Ahmad U, et al. rs2651899 variant is associated with risk for migraine without aura from North Indian population. *Mol Biol Rep* 2019; 46: 1247-1255. 2019/01/13. DOI: 10.1007/s11033-019-04593-1.
14. Fu X, Yang J, Wu X, et al. Association between PRDM16, MEF2D, TRPM8, LRP1 gene polymorphisms and migraine susceptibility in the She ethnic population in China. *Clin Invest Med* 2019; 42: E21-E30. 2019/03/25. DOI: 10.25011/cim.v42i1.32389.
15. Chen SP, Fuh JL, Chung MY, et al. Genome-wide association study identifies novel susceptibility loci for migraine in Han Chinese resided in Taiwan. *Cephalalgia* 2018; 38: 466-475. 2017/09/28. DOI: 10.1177/0333102417695105.
16. Sintas C, Fernandez-Morales J, Vila-Pueyo M, et al. Replication study of previous migraine genome-wide association study findings in a Spanish sample of migraine with aura. *Cephalalgia* 2015; 35: 776-782. 2014/11/13. DOI: 10.1177/0333102414557841.

17. Fan X, Wang J, Fan W, et al. Replication of migraine GWAS susceptibility loci in Chinese Han population. *Headache* 2014; 54: 709-715. 2014/03/29. DOI: 10.1111/head.12329.
18. Ghosh J, Pradhan S and Mittal B. Genome-wide-associated variants in migraine susceptibility: a replication study from North India. *Headache* 2013; 53: 1583-1594. 2013/11/26. DOI: 10.1111/head.12240.
19. An XK, Ma QL, Lin Q, et al. PRDM16 rs2651899 variant is a risk factor for Chinese common migraine patients. *Headache* 2013; 53: 1595-1601. 2013/09/12. DOI: 10.1111/head.12212.
20. Schurks M. Genetics of migraine in the age of genome-wide association studies. *J Headache Pain* 2012; 13: 1-9. 2011/11/11. DOI: 10.1007/s10194-011-0399-0.
21. Gava NR, Sandrock R, Arnold GE, et al. Reduced TRPM8 expression underpins reduced migraine risk and attenuated cold pain sensation in humans. *Sci Rep* 2019; 9: 19655. 2019/12/25. DOI: 10.1038/s41598-019-56295-0.
22. Edvinsson L. Role of CGRP in Migraine. *Handb Exp Pharmacol* 2019; 255: 121-130. 2019/02/07. DOI: 10.1007/164_2018_201.
23. Ling YH, Chen SP, Fann CS, et al. TRPM8 genetic variant is associated with chronic migraine and allodynia. *J Headache Pain* 2019; 20: 115. 2019/12/16. DOI: 10.1186/s10194-019-1064-2.
24. Vigano A, Manica A, Di Piero V, et al. Did Going North Give Us Migraine? An Evolutionary Approach on Understanding Latitudinal Differences in Migraine Epidemiology. *Headache* 2019; 59: 632-634. 2019/04/09. DOI: 10.1111/head.13520.

25. Chebini A and Dilli E. Cold Stimulus Headache. *Curr Neurol Neurosci Rep* 2019; 19: 46. 20190606. DOI: 10.1007/s11910-019-0956-5.
26. Burgos-Vega CC, Ahn DD, Bischoff C, et al. Meningeal transient receptor potential channel M8 activation causes cutaneous facial and hindpaw allodynia in a preclinical rodent model of headache. *Cephalalgia* 2016; 36: 185-193. 20150505. DOI: 10.1177/0333102415584313.
27. Ren L, Dhaka A and Cao YQ. Function and postnatal changes of dural afferent fibers expressing TRPM8 channels. *Mol Pain* 2015; 11: 37. 2015/06/27. DOI: 10.1186/s12990-015-0043-0.
28. Lopresti AL, Smith SJ and Drummond PD. Herbal treatments for migraine: A systematic review of randomised-controlled studies. *Phytother Res* 2020; 34: 2493-2517. 20200420. DOI: 10.1002/ptr.6701.
29. Borhani Haghighi A, Motazedian S, Rezaii R, et al. Cutaneous application of menthol 10% solution as an abortive treatment of migraine without aura: a randomised, double-blind, placebo-controlled, crossed-over study. *Int J Clin Pract* 2010; 64: 451-456. DOI: 10.1111/j.1742-1241.2009.02215.x.
30. Rafieian-Kopaei M, Hasanpour-Dehkordi A, Lorigooini Z, et al. Comparing the Effect of Intranasal Lidocaine 4% with Peppermint Essential Oil Drop 1.5% on Migraine Attacks: A Double-Blind Clinical Trial. *Int J Prev Med* 2019; 10: 121. 20190705. DOI: 10.4103/ijpvm.IJPVM_530_17.
31. Wei C, Kim B and McKemy DD. Transient receptor potential melastatin 8 is required for nitroglycerin- and calcitonin gene-related peptide-induced migraine-like pain

behaviors in mice. *Pain* 2022; 163: 2380-2389. 20220329. DOI: 10.1097/j.pain.0000000000002635.

32. Yang C, Yamaki S, Jung T, et al. Endogenous Inflammatory Mediators Produced by Injury Activate TRPV1 and TRPA1 Nociceptors to Induce Sexually Dimorphic Cold Pain That Is Dependent on TRPM8 and GFRalpha3. *J Neurosci* 2023; 43: 2803-2814. 20230310. DOI: 10.1523/JNEUROSCI.2303-22.2023.

33. Yamaki S, Chau A, Gonzales L, et al. Nociceptive afferent phenotyping reveals that transient receptor potential ankyrin 1 promotes cold pain through neurogenic inflammation upstream of the neurotrophic factor receptor GFRalpha3 and the menthol receptor transient receptor potential melastatin 8. *Pain* 2021; 162: 609-618. 2020/08/23. DOI: 10.1097/j.pain.0000000000002043.

34. Lippoldt EK, Ongun S, Kusaka GK, et al. Inflammatory and neuropathic cold allodynia are selectively mediated by the neurotrophic factor receptor GFRalpha3. *Proc Natl Acad Sci U S A* 2016; 113: 4506-4511. 2016/04/07. DOI: 10.1073/pnas.1603294113.

35. Lippoldt EK, Elmes RR, McCoy DD, et al. Artemin, a glial cell line-derived neurotrophic factor family member, induces TRPM8-dependent cold pain. *J Neurosci* 2013; 33: 12543-12552. 2013/07/26. DOI: 10.1523/JNEUROSCI.5765-12.2013.

36. Su L, Shu R, Song C, et al. Downregulations of TRPM8 expression and membrane trafficking in dorsal root ganglion mediate the attenuation of cold hyperalgesia in CCI rats induced by GFRalpha3 knockdown. *Brain Res Bull* 2017; 135: 8-24. 20170901. DOI: 10.1016/j.brainresbull.2017.08.002.

37. McIlvried LA, Albers K and Gold MS. Distribution of artemin and GFRalpha3 labeled nerve fibers in the dura mater of rat: artemin and GFRalpha3 in the dura. *Headache* 2010; 50: 442-450. 2009/10/23. DOI: 10.1111/j.1526-4610.2009.01548.x.
38. Shang HQ, Wang Y, Mao YY, et al. Expression of artemin and GFRalpha3 in an animal model of migraine: possible role in the pathogenesis of this disorder. *J Headache Pain* 2016; 17: 81. 2016/09/08. DOI: 10.1186/s10194-016-0673-2.
39. Shang H, Wang Y, Chao X, et al. Artemin transiently increases iNOS expression in primary cultured trigeminal ganglion neurons. *Neurosci Lett* 2017; 660: 34-38. 2017/09/14. DOI: 10.1016/j.neulet.2017.09.016.
40. Percie du Sert N, Ahluwalia A, Alam S, et al. Reporting animal research: Explanation and elaboration for the ARRIVE guidelines 2.0. *PLoS Biol* 2020; 18: e3000411. 2020/07/14. DOI: 10.1371/journal.pbio.3000411.
41. Bautista DM, Siemens J, Glazer JM, et al. The menthol receptor TRPM8 is the principal detector of environmental cold. *Nature* 2007; 448: 204-208. 2007/06/01. DOI: 10.1038/nature05910.
42. Honma Y, Araki T, Gianino S, et al. Artemin is a vascular-derived neurotropic factor for developing sympathetic neurons. *Neuron* 2002; 35: 267-282. 2002/08/06. DOI: 10.1016/s0896-6273(02)00774-2.
43. Burgos-Vega CC, Quigley LD, Trevisan Dos Santos G, et al. Non-invasive dural stimulation in mice: A novel preclinical model of migraine. *Cephalalgia* 2019; 39: 123-134. 2018/06/01. DOI: 10.1177/0333102418779557.

44. Louveau A, Filiano AJ and Kipnis J. Meningeal whole mount preparation and characterization of neural cells by flow cytometry. *Curr Protoc Immunol* 2018; 121: e50. 20180416. DOI: 10.1002/cpim.50.
45. Pradhan AA, Smith ML, McGuire B, et al. Characterization of a novel model of chronic migraine. *Pain* 2014; 155: 269-274. 2013/10/15. DOI: 10.1016/j.pain.2013.10.004.
46. Knowlton WM, Daniels RL, Palkar R, et al. Pharmacological blockade of TRPM8 ion channels alters cold and cold pain responses in mice. *PLoS One* 2011; 6: e25894. 2011/10/11. DOI: 10.1371/journal.pone.0025894.
47. Bonin RP, Bories C and De Koninck Y. A simplified up-down method (SUDO) for measuring mechanical nociception in rodents using von Frey filaments. *Mol Pain* 2014; 10: 26. 20140416. DOI: 10.1186/1744-8069-10-26.
48. Knowlton WM, Palkar R, Lippoldt EK, et al. A sensory-labeled line for cold: TRPM8-expressing sensory neurons define the cellular basis for cold, cold pain, and cooling-mediated analgesia. *J Neurosci* 2013; 33: 2837-2848. 2013/02/15. DOI: 10.1523/JNEUROSCI.1943-12.2013.
49. Dhaka A, Earley TJ, Watson J, et al. Visualizing cold spots: TRPM8-expressing sensory neurons and their projections. *J Neurosci* 2008; 28: 566-575. DOI: 10.1523/JNEUROSCI.3976-07.2008.
50. Peier AM, Moqrich A, Hergarden AC, et al. A TRP channel that senses cold stimuli and menthol. *Cell* 2002; 108: 705-715. DOI: 10.1016/s0092-8674(02)00652-9.

51. Avona A, Mason BN, Burgos-Vega C, et al. Meningeal CGRP-Prolactin Interaction Evokes Female-Specific Migraine Behavior. *Ann Neurol* 2021; 89: 1129-1144. 20210422. DOI: 10.1002/ana.26070.
52. Ren L, Chang MJ, Zhang Z, et al. Quantitative Analysis of Mouse Dural Afferent Neurons Expressing TRPM8, VGLUT3, and NF200. *Headache* 2018; 58: 88-101. 2017/09/20. DOI: 10.1111/head.13188.
53. Jeong SL, Zhang H, Yamaki S, et al. Immunoglobulin somatic hypermutation in a defined biochemical system recapitulates affinity maturation and permits antibody optimization. *Nucleic Acids Res* 2022; 50: 11738-11754. DOI: 10.1093/nar/gkac995.
54. Edelmayer RM, Le LN, Yan J, et al. Activation of TRPA1 on dural afferents: a potential mechanism of headache pain. *Pain* 2012; 153: 1949-1958. 20120717. DOI: 10.1016/j.pain.2012.06.012.
55. Edelmayer RM, Vanderah TW, Majuta L, et al. Medullary pain facilitating neurons mediate allodynia in headache-related pain. *Ann Neurol* 2009; 65: 184-193. DOI: 10.1002/ana.21537.
56. De Felice M, Ossipov MH, Wang R, et al. Triptan-induced latent sensitization: a possible basis for medication overuse headache. *Ann Neurol* 2010; 67: 325-337. DOI: 10.1002/ana.21897.
57. Wei ET and Seid DA. AG-3-5: a chemical producing sensations of cold. *J Pharm Pharmacol* 1983; 35: 110-112. DOI: 10.1111/j.2042-7158.1983.tb04279.x.
58. Stein RJ, Santos S, Nagatomi J, et al. Cool (TRPM8) and hot (TRPV1) receptors in the bladder and male genital tract. *J Urol* 2004; 172: 1175-1178. DOI: 10.1097/01.ju.0000134880.55119.cf.

59. Harrington AM, Hughes PA, Martin CM, et al. A novel role for TRPM8 in visceral afferent function. *Pain* 2011; 152: 1459-1468. 20110413. DOI: 10.1016/j.pain.2011.01.027.
60. Hayashi T, Kondo T, Ishimatsu M, et al. Expression of the TRPM8-immunoreactivity in dorsal root ganglion neurons innervating the rat urinary bladder. *Neurosci Res* 2009; 65: 245-251. 20090719. DOI: 10.1016/j.neures.2009.07.005.
61. Lashinger ES, Steinginga MS, Hieble JP, et al. AMTB, a TRPM8 channel blocker: evidence in rats for activity in overactive bladder and painful bladder syndrome. *Am J Physiol Renal Physiol* 2008; 295: F803-810. 20080618. DOI: 10.1152/ajprenal.90269.2008.
62. Tsukimi Y, Mizuyachi K, Yamasaki T, et al. Cold response of the bladder in guinea pig: involvement of transient receptor potential channel, TRPM8. *Urology* 2005; 65: 406-410. DOI: 10.1016/j.urology.2004.10.006.
63. Christianson JA, McIlwrath SL, Koerber HR, et al. Transient receptor potential vanilloid 1-immunopositive neurons in the mouse are more prevalent within colon afferents compared to skin and muscle afferents. *Neuroscience* 2006; 140: 247-257. 20060329. DOI: 10.1016/j.neuroscience.2006.02.015.
64. Benemei S and Dussor G. TRP Channels and Migraine: Recent Developments and New Therapeutic Opportunities. *Pharmaceuticals (Basel)* 2019; 12 2019/04/12. DOI: 10.3390/ph12020054.
65. Spekker E, Kortesi T and Vecsei L. TRP Channels: Recent Development in Translational Research and Potential Therapeutic Targets in Migraine. *Int J Mol Sci* 2022; 24 20221231. DOI: 10.3390/ijms24010700.

66. Caterina MJ, Schumacher MA, Tominaga M, et al. The capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature* 1997; 389: 816-824. 1997/12/31 23:16. DOI: 10.1038/39807.
67. Julius D. TRP channels and pain. *Annu Rev Cell Dev Biol* 2013; 29: 355-384. DOI: 10.1146/annurev-cellbio-101011-155833.
68. Chuang HH, Prescott ED, Kong H, et al. Bradykinin and nerve growth factor release the capsaicin receptor from PtdIns(4,5)P₂-mediated inhibition. *Nature* 2001; 411: 957-962. DOI: 10.1038/35082088.
69. Premkumar LS and Ahern GP. Induction of vanilloid receptor channel activity by protein kinase C. *Nature* 2000; 408: 985-990. DOI: 10.1038/35050121.
70. Forrest SL, Osborne PB and Keast JR. Characterization of axons expressing the artemin receptor in the female rat urinary bladder: a comparison with other major neuronal populations. *J Comp Neurol* 2014; 522: 3900-3927. 2014/07/22. DOI: 10.1002/cne.23648.
71. Zhu S, Li Y, Bennett S, et al. The role of glial cell line-derived neurotrophic factor family member artemin in neurological disorders and cancers. *Cell Prolif* 2020; 53: e12860. 2020/06/24. DOI: 10.1111/cpr.12860.
72. Uhlen M, Fagerberg L, Hallstrom BM, et al. Proteomics. Tissue-based map of the human proteome. *Science* 2015; 347: 1260419. 2015/01/24. DOI: 10.1126/science.1260419.
73. Akerman S, Romero-Reyes M, Karsan N, et al. Therapeutic targeting of nitroglycerin-mediated trigeminovascular neuronal hypersensitivity predicts clinical

outcomes of migraine abortives. *Pain* 2021; 162: 1567-1577. DOI: 10.1097/j.pain.0000000000002142.

74. Christiansen I, Thomsen LL, Daugaard D, et al. Glyceryl trinitrate induces attacks of migraine without aura in sufferers of migraine with aura. *Cephalalgia* 1999; 19: 660-667; discussion 626. 1999/10/19. DOI: 10.1046/j.1468-2982.1999.019007660.x.

75. Demartini C, Greco R, Zanaboni AM, et al. Nitroglycerin as a comparative experimental model of migraine pain: From animal to human and back. *Prog Neurobiol* 2019; 177: 15-32. 20190213. DOI: 10.1016/j.pneurobio.2019.02.002.

76. Chen SP, Hsu CL, Wang YF, et al. Genome-wide analyses identify novel risk loci for cluster headache in Han Chinese residing in Taiwan. *J Headache Pain* 2022; 23: 147. 2022/11/21. DOI: 10.1186/s10194-022-01517-6.

77. Huang D, Li S, Dhaka A, et al. Expression of the transient receptor potential channels TRPV1, TRPA1 and TRPM8 in mouse trigeminal primary afferent neurons innervating the dura. *Mol Pain* 2012; 8: 66. 2012/09/14. DOI: 10.1186/1744-8069-8-66.

78. Hsu YY, Chen CJ, Wu SH, et al. Cold intervention for relieving migraine symptoms: A systematic review and meta-analysis. *J Clin Nurs* 2022 20220520. DOI: 10.1111/jocn.16368.

79. Ugurlu YK and Enc N. The effect of local cold compresses for nitroglycerin-induced headache: An observational pretest-posttest study. *Nurs Crit Care* 2022 20220715. DOI: 10.1111/nicc.12823.

80. St Cyr A, Chen A, Bradley KC, et al. Efficacy and Tolerability of STOPAIN for a Migraine Attack. *Front Neurol* 2015; 6: 11. 2015/02/24. DOI: 10.3389/fneur.2015.00011.

81. Alarcon-Alarcon D, Cabanero D, de Andres-Lopez J, et al. TRPM8 contributes to sex dimorphism by promoting recovery of normal sensitivity in a mouse model of chronic migraine. *Nat Commun* 2022; 13: 6304. 20221022. DOI: 10.1038/s41467-022-33835-3.
82. Knowlton WM, Bifolck-Fisher A, Bautista DM, et al. TRPM8, but not TRPA1, is required for neural and behavioral responses to acute noxious cold temperatures and cold-mimetics in vivo. *Pain* 2010; 150: 340-350. 2010/06/15. DOI: 10.1016/j.pain.2010.05.021.
83. Pogorzala LA, Mishra SK and Hoon MA. The cellular code for mammalian thermosensation. *J Neurosci* 2013; 33: 5533-5541. DOI: 10.1523/JNEUROSCI.5788-12.2013.
84. Ongun S, Sarkisian A and McKemy DD. Selective cold pain inhibition by targeted block of TRPM8-expressing neurons with quaternary lidocaine derivative QX-314. *Commun Biol* 2018; 1: 53. 2018/10/03. DOI: 10.1038/s42003-018-0062-2.
85. Dhaka A, Murray AN, Mathur J, et al. TRPM8 is required for cold sensation in mice. *Neuron* 2007; 54: 371-378. 2007/05/08. DOI: 10.1016/j.neuron.2007.02.024.
86. Proudfoot CJ, Garry EM, Cottrell DF, et al. Analgesia mediated by the TRPM8 cold receptor in chronic neuropathic pain. *Curr Biol* 2006; 16: 1591-1605. 2006/08/22. DOI: 10.1016/j.cub.2006.07.061.
87. Liu B, Fan L, Balakrishna S, et al. TRPM8 is the principal mediator of menthol-induced analgesia of acute and inflammatory pain. *Pain* 2013; 154: 2169-2177. 20130629. DOI: 10.1016/j.pain.2013.06.043.

88. Palkar R, Ongun S, Catich E, et al. Cooling Relief of Acute and Chronic Itch Requires TRPM8 Channels and Neurons. *J Invest Dermatol* 2018; 138: 1391-1399. 2017/12/31. DOI: 10.1016/j.jid.2017.12.025.
89. Kardon AP, Polgar E, Hachisuka J, et al. Dynorphin acts as a neuromodulator to inhibit itch in the dorsal horn of the spinal cord. *Neuron* 2014; 82: 573-586. 20140410. DOI: 10.1016/j.neuron.2014.02.046.
90. Fan L, Balakrishna S, Jabba SV, et al. Menthol decreases oral nicotine aversion in C57BL/6 mice through a TRPM8-dependent mechanism. *Tob Control* 2016; 25: ii50-ii54. 20161003. DOI: 10.1136/tobaccocontrol-2016-053209.
91. Ha MA, Smith GJ, Cichocki JA, et al. Menthol attenuates respiratory irritation and elevates blood cotinine in cigarette smoke exposed mice. *PLoS One* 2015; 10: e0117128. 20150213. DOI: 10.1371/journal.pone.0117128.
92. Pergolizzi JV, Jr., Taylor R, Jr., LeQuang JA, et al. The role and mechanism of action of menthol in topical analgesic products. *J Clin Pharm Ther* 2018; 43: 313-319. 20180310. DOI: 10.1111/jcpt.12679.
93. Nosedá R and Burstein R. Migraine pathophysiology: anatomy of the trigeminovascular pathway and associated neurological symptoms, cortical spreading depression, sensitization, and modulation of pain. *Pain* 2013; 154 Suppl 1: S44-53. 20130725. DOI: 10.1016/j.pain.2013.07.021.
94. Burstein R, Yamamura H, Malick A, et al. Chemical stimulation of the intracranial dura induces enhanced responses to facial stimulation in brain stem trigeminal neurons. *J Neurophysiol* 1998; 79: 964-982. DOI: 10.1152/jn.1998.79.2.964.

95. Burstein R, Yarnitsky D, Goor-Aryeh I, et al. An association between migraine and cutaneous allodynia. *Ann Neurol* 2000; 47: 614-624. 2000/05/11.
96. Burstein R, Cutrer MF and Yarnitsky D. The development of cutaneous allodynia during a migraine attack clinical evidence for the sequential recruitment of spinal and supraspinal nociceptive neurons in migraine. *Brain* 2000; 123 (Pt 8): 1703-1709. 2000/07/25. DOI: 10.1093/brain/123.8.1703.
97. Ashina M, Hansen JM, Do TP, et al. Migraine and the trigeminovascular system-40 years and counting. *Lancet Neurol* 2019; 18: 795-804. 20190531. DOI: 10.1016/S1474-4422(19)30185-1.
98. Charles A and Brennan KC. The neurobiology of migraine. *Handb Clin Neurol* 2010; 97: 99-108. DOI: 10.1016/S0072-9752(10)97007-3.

FIGURE LEGENDS

Figure 1: TRPM8-positive afferents innervate mouse meninges.

(A) Representative images of GFP (TRPM8) and WGA (Alex-350) co-labeling in trigeminal ganglia in adult *Trpm8^{GFP}* mice that received an intradural injection of WGA. Arrows highlight co-labeled cells. Immunoreactivity for GFP (TRPM8) and calcitonin gene-related peptide (CGRP) in the meninges of *Trpm8^{GFP}* mice (B) versus that in wildtype mice (C). Arrowheads denote axons that co-label for GFP and CGRP.

Figure 2: GFR α 3 is required for nitroglycerin-induced acute cephalic and extracephalic tactile allodynia.

(A) Left: Schematic of acute injections and mechanical testing timeline. Right: Wildtype mice exhibit periorbital mechanical allodynia measured 2 hrs after each systemic NTG injection (1 mg/kg) compared to baseline (BL) sensitivity (n=8 for each condition, two-way ANOVA with Dunnett's multiple comparison test, *p<0.05, ***p<0.001, ****p<0.0001). In contrast, GFR α 3-null mice (*Gfra3^{-/-}*) are insensitive to 1 mg/kg NTG injections on each day tested (n=6 for each condition, two-way ANOVA with Dunnett's multiple comparison test compared to BL, p>0.05). Wildtypes receiving a 10 mg/kg dose of NTG show robust periorbital (B) and hindpaw (C) mechanical allodynia 2 hr post-injection compared to baseline (BL), a phenotype absent in *Gfra3^{-/-}* mice (n=6 and 8 for periorbital and hindpaw, respectively, paired two-tailed t-test, ^{ns}p>0.05, **p<0.01, ***p<0.001).

Figure 3: GFR α 3 is required for chronic allodynia induced by repeated nitroglycerin treatments.

(A) Left: Schematic of chronic injections and mechanical testing timeline. Right: Wildtype and Gfr α 3^{-/-} mice received 1 mg/kg NTG injections every two days, with only wildtypes displaying enhanced basal periorbital mechanical allodynia, compared to baseline at Day 1 (n=6-8 for each condition, two-way ANOVA with Dunnett's multiple comparison test, *p<0.05, **p<0.01, ***p<0.001). Wildtypes receiving a 10 mg/kg dose of NTG show robust basal periorbital (B) and hindpaw (C) mechanical allodynia 2 days post-injection compared to baseline (BL), a phenotype is absent in Gfr α 3^{-/-} mice (n=6 each condition for periorbital and hindpaw, respectively, paired two-tailed t-test, ^{ns}p>0.05, **p<0.01, ****p<0.0001).

Figure 4: Neutralizing artemin ameliorates nitroglycerin-induced tactile allodynia.

(A) Top: Schematic of the experimental approach. Wildtype mice treated with either an anti-artemin or isotype control mAb exhibit normal periorbital mechanical sensitivity 1hr post-injection compared to baseline (BL). A subsequent injection of 1 mg/kg NTG induced allodynia in mice given the isotype control, but not the anti-artemin mAb (n=6 for each condition, one-way ANOVA with Tukey's multiple comparisons tests for each time point and condition, ^{ns}p>0.05, **p<0.01). (B) Top: Schematic of the experimental approach. Wildtype mice treated with 1 mg/kg NTG to induce basal periorbital mechanical allodynia 2 days post-injection compared to BL, a phenotype that remained in mice subsequently treated with the isotype control mAb. However, allodynia was reduced in animals treated

with the anti-artemin mAb (n=10 for each condition, one-way ANOVA with Tukey's multiple comparisons test for different time point and condition, **p<0.01, ^{ns}p>0.05).

Figure 5: Necessity of GFR α 3 and artemin in tactile allodynia induced by direct stimulation of the meninges.

(A) Direct supradural infusion of an inflammatory soup (IS) mixture induces periorbital tactile allodynia that is partially dependent on GFR α 3. Gfr α 3^{-/-} mice given IS exhibited a reduced level of mechanical allodynia compared to wildtypes at 2-4hrs post-injection (n=10 each genotype, two-way ANOVA with Tukey's multiple comparisons test, *p<0.05, **p<0.01). (B) Supradural artemin induces robust periorbital tactile allodynia in wildtype but not Trpm8^{-/-} mice (n=6 for each condition and genotype, two-way ANOVA with a Tukey's multiple comparisons test for vehicle vs. artemin at each time point; *p<0.05, ***p<0.001). (C) Wildtype mice treated with artemin and sumatriptan exhibit a partial reduction in periorbital mechanical allodynia compared to those given artemin and vehicle (n=8 for each condition, *p<0.05, **p<0.01, two-way ANOVA with a Tukey's multiple comparison test).

Figure 6: Activation of TRPM8 afferents in the mouse dura induces tactile allodynia.

(A) Wildtype mice treated with 1 mg/kg NTG to induce basal periorbital mechanical allodynia at 2 days post-injection (Day 3), as well as after 4 injections repeated every 2 days, measured 9 days after the initial injections, compared to BL, a phenotype that remained in mice subsequently treated with vehicle (n=8, two-way ANOVA with Dunnett's multiple comparisons test, compared to baseline, p<0.01 on Day 3, p<0.001 on Day 9).

However, basal allodynia on Day 3, but not Day 9, was reduced in animals treated with the PBMC at 2 hrs post-injection compared to vehicle treated mice (n=8, two-way ANOVA with Dunnett's multiple comparisons test, ***p<0.001). (B) Periorbital mechanical sensitivity measured in wildtype mice that received an intradural injection of either icilin or vehicle. Mechanical allodynia was evident by 2 hrs post-injection and lasted for up to 5 hrs (n=6 for each condition, two-way ANOVA with a Bonferroni's multiple comparisons test for vehicle vs. icilin at each time point; *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001). (C) TRPM8-null (*Trpm8*^{-/-}) mice displayed normal mechanical sensitivity in their hindpaws 3 hrs after an intradural injection of icilin, in contrast to wildtype animals (n=10 for each condition, one-way ANOVA with a Tukey's multiple comparisons test for vehicle vs. icilin in each condition; ^{ns}p>0.05, **p<0.01, ****p<0.0001). (D) Intradural icilin induces periorbital tactile allodynia in *Gfra3*^{-/-} mice (3 hrs post-injection, n=8, paired two-tailed t-test, **p<0.01).

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6

