

Smoking and COVID-19: A Scoping Review

Chadia Haddad^{1,2*} , Sandrella Bou Malhab^{3*}, Hala Sacre⁴ 
and Pascale Salameh^{4,5,6}

¹Research Department, Psychiatric Hospital of the Cross, Jal Eddib, Lebanon. ²INSERM, Univ. Limoges, CH Esquirol, IRD, U1094 Tropical Neuroepidemiology, Institute of Epidemiology and Tropical Neurology, GEIST, Limoges, France. ³Ecole Doctorale des Sciences et Technologie, Lebanese University, Hadath, Lebanon. ⁴INSPECT-LB: Institut National de Santé Publique, Épidémiologie Clinique et Toxicologie-Liban, Beirut, Lebanon. ⁵Faculty of Pharmacy, Lebanese University, Hadat, Lebanon. ⁶School of Medicine, University of Nicosia, Cyprus.

Tobacco Use Insights
Volume 14: 1–9
© The Author(s) 2021
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/1179173X21994612



ABSTRACT: Smoking affects a person's overall health and damages nearly every organ of the body. Since smoking tobacco affects and damages the lungs, it increases the risk of respiratory infections and makes it easier for the coronavirus disease (COVID-19) to invade the lung tissue, causing more severe symptoms and increasing the risk of death. However, debates are still ongoing as to the effect of cigarette smoking on vulnerability to COVID-19. Some studies, where active smokers were underrepresented among patients with COVID-19, claimed that a “smoker's paradox” may exist in COVID-19 and that smokers are protected from infection and severe complications of COVID-19. However, other studies reported the opposite trend. The objective of this study is to review the findings of epidemiological and in vitro studies about the association between smoking and the risk of contracting COVID-19, taking into account disease severity. Several epidemiological studies have found a higher smoking prevalence among COVID-19 infected patients. Also, studies had shown that people with respiratory diseases caused by tobacco use are at higher risk of developing severe COVID-19 symptoms. Studies have shown that in vitro, the acute exposure allows for more severe proximal airway epithelial disease from SARS-CoV-2 by reducing the mucosal innate immune response and the proliferation of airway basal stem cells and has implications for disease spread and severity in people exposed to cigarette smoke, with a more severe viral infection and cell death. Smoker patients with different comorbidities are at higher risk of contracting the COVID-19 virus and have a worse prognosis for the virus as well as for their comorbidities. Further investigations of the interaction between smoking and COVID-19 are warranted to accurately assess the risk of contracting COVID-19 among smokers, and the progression to mechanical ventilation or death in patients who suffer from it.

KEYWORDS: Smoking, tobacco waterpipe, coronavirus disease, epidemiologic studies, cigarette smoking

RECEIVED: September 9, 2020. **ACCEPTED:** January 25, 2021.

TYPE: Review

FUNDING: The author(s) received no financial support for the research, authorship, and/or publication of this article.

DECLARATION OF CONFLICTING INTERESTS: The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

CORRESPONDING AUTHORS: Chadia Haddad, Research Department, Psychiatric Hospital of the Cross, P.O. Box 60096, Jall-Eddib, Lebanon. Email: Chadia_9@hotmail.com

Pascale Salameh, Faculty of Pharmacy, Lebanese University, Hadat, Lebanon. Email: pascalesalameh1@hotmail.com

Introduction: Smoking in the Context of COVID-19 Pandemic

The COVID-19 pandemic has imposed on humans a new way of living: restricted movements, confinement in closed spaces, fear of being infected, and strict hygiene measures for endless periods. Psychologically, this era is considered as a “transition” phase and seems to require a search for a new balance. While boredom and confinement might have stimulated smoking among some, the threat of contracting COVID-19 and becoming severely ill might have motivated others to improve their health by quitting smoking; this is particularly true for individuals with moderate to severe stress.¹

Studies have reported mixed findings regarding changes in tobacco use during the pandemic. Indeed, in the United States, far more participants reported increasing their tobacco use since COVID-19 started (40.9%) versus decreasing their tobacco use (17.8%).² In Italy, most exclusive cigarette smokers have considered quitting, but most exclusive e-cigarette users

have not considered stopping the use of e-cigarettes. In the group of former smokers, about one-third of participants reported having thoughts about smoking again, and in the group of never smokers, few participants declared the intention to start smoking.³ In China, relapses from smoking abstinence were relatively common during the pandemic (25%). Similarly, 20% of regular smokers increased their usage amount.⁴ In England, a study have found that increased smoking among smokers and relapsed smoking among former smokers were a result of social isolation and potential stress.⁵ A study done by Chertok⁶ among 810 adults from Ohio have found a change in smoking behavior among current smokers since COVID-19 were 36.7% having attempted to quit since the outbreak.

Debates are still ongoing as to the effect of cigarette smoking on vulnerability to COVID-19. Although it is well established that smoking is associated with morbidity and mortality in several respiratory infections, some studies, where active smokers were underrepresented among patients with COVID-19, claimed that a “smoker's paradox” may exist in COVID-19 and that smokers are protected from infection and severe

*First Co-authors



complications of COVID-19. However, other studies reported the opposite trend. Therefore, the need of a scoping review to provide an overview of the available research evidence. The objective of this study is to review the findings of epidemiological and in vitro studies about the association between smoking and the risk of contracting COVID-19, taking into account disease severity.

Method and research design

A comprehensive literature review was conducted using the MEDLINE/PubMed database and references cited in relevant articles were also searched. Included studies were recently published about the effect of COVID-19 on smoking from January 1st, up till the end of September, 2020. Also, studies about the mechanisms by which smoking affects the body were included from December 2001 up till May 2018. No restrictions were placed on language of publication or study origin. Limitations were placed on the type of publication when using the PubMed database and included all of the following: case reports, clinical studies, clinical trials, comparative studies, controlled clinical trials, journal articles, meta-analyses, observational studies, systematic reviews, and books and documents. Whenever possible, peer-reviewed randomized clinical trials (RCTs) and placebo controlled trials were preferred over case reports, case studies and commentaries, while review articles and meta-analyses were also included whenever relevant. Similarly, books and book chapters on the topic of smoking were also reviewed for eligibility.

To ensure retrieval of the largest number of articles covering the topic of smoking, the following key search terms were used: “smoking,” “tobacco,” “nicotine,” “covid-19,” “coronavirus disease,” “hookah” and “waterpipe,” “cigarette,” “smoker,” “epidemic,” “SARS-CoV-2,” “virus,” and “respiratory function”. The search term “smoking” was included in the initial run of the online databases but resulted in over 315 133 articles, the majority of which were irrelevant. Consequently “smoking” as a search term was eliminated from the primary searches conducted.

Then we used the term “Smoking and covid-19” and “effect of smoking on the body” and we obtained a total of 20188 articles using the MEDLINE/PubMED database. Of the 20188 article, 557 articles were about ‘smoking and COVID-19’ and 19631 were about “harmful effect of smoking.” Title and abstract screening resulted in the elimination of articles that were not relevant or did not cover the topic of smoking; did not meet the initial inclusion criteria related to article type; focused primarily on other disease and disorders other than smoking; or primarily focused on psycho-social aspects and risk factors related to smoking. This resulted in 200 articles that required in-depth analysis, of which 50 were eliminated due to irrelevance, and 48 were not found to be focused on smoking and COVID-19. This resulted in 107 articles which were included in the final version of the article.

Modes of COVID-19 Transmission Using a Tobacco Smoking

Viral transmission through tobacco smoking might first occur through contact since smoking involves regular contact between saliva, hands, and devices.⁷ Moreover, smoking tobacco (cigarettes, e-cigarettes, or waterpipe) produces exhaled smoke and aerosols, and causes coughing or sneezing, thus contaminating surroundings and surfaces with SARS-CoV-2 (severe acute respiratory syndrome coronavirus 2).

Additionally, SARS-CoV-2 survives in aerosols and surfaces (plastic, paper, and steel) for several hours to days⁸ after oral/nasal release (coughing, sneezing, smoking).^{9,10} Also, the virus is actively found for 4 hours up to 2 to 3 days on shared surfaces of smoking devices.⁷ Thus, touching contaminated surfaces of smoking devices can actively harm smokers and non-smokers, even when people practice social distancing.⁷

The act of cigarette smoking and virus transmission

The COVID-19 is transmitted from hand to mouth as smoking cigarettes implies that fingers and contaminated cigarettes are in contact with the lips or comes close to the mouth, which increases the risk of transmission.^{11,12} The latter also depends on whether hands are contaminated and how the packet is prepared and handled.¹² Indeed, infected hands can contaminate a clean packet and the cigarettes it contains, thus it is no longer clean during the remaining time of its use.¹²

The act of waterpipe smoking and virus transmission

Waterpipe smoking is the process of inhaling tobacco smoke after it has passed through a chilled water chamber, with a hose capped with a plastic mouthpiece.¹³ Virus transmission through the waterpipe is high since smokers might cough and contaminate the inner side of the hose. The moisture in tobacco smoke and the use of cold water in the water chamber promote the survival of viruses and bacteria.^{13,14} Also, uncontrolled manual preparation of waterpipe could spread infectious diseases.¹⁴ Another factor of contamination is that most cafés prefer not to clean the waterpipe upon each smoking session, since it is labor-intensive and time-consuming to wash and clean pieces, including the cold water jar, ideal for virus transmission.^{13,15} Moreover, the waterpipe tobacco smoke contains many dangerous chemicals that damage the respiratory tract and predispose the smoker to viral infection.¹⁵

On the other hand, smoking waterpipe is an activity typically performed within groups in public settings.¹⁶ Social gatherings encourage the transmission of the COVID -19 virus.¹⁷ The risk may be higher when waterpipe is used indoors.¹⁸ Participants usually pass the hose from person to person and may use the same mouthpiece¹⁶ during smoking sessions that last from 45 to 60 minutes but can also be up to several hours.¹⁴ Even if only 1 customer uses the pipe and mouthpiece at a

time, other customers might reuse uncleaned waterpipes chambers and hoses on the same day.¹⁴

It is noteworthy to mention that until now no studies have directly examine the mode of transmission of COVID-19 through cigarette or waterpipe use. The abovementioned studies are most often prospective envisioning or predicting the COVID-19 transmission based on facts about waterpipe and cigarette use. Further studies and evidence are needed in order to understand the transmission way of COVID-19 through smoking and measures should be implemented within the community to limit its direct and indirect effects.⁷

Mechanisms Through Which Smoking may Increase the Risk of Infection with COVID-19

Mechanistic studies indicate that the enhanced susceptibility to infection could be attributed to angiotensin converting enzyme 2 (ACE2) receptor upregulation, the key receptor used to gain entrance into the host mucosa and induce active infection by the acute respiratory syndrome coronavirus 2 (SARS-CoV-2), an apparently unique mechanism for this virus. Smoking contributes to a worse disease state and increased vulnerability to SARS-CoV-2 infection via the activation of peripheral nicotinic acetylcholine receptors (nAChRs) expressed in many organ systems. Nicotine affects homeostasis of the renin angiotensin system (RAS)¹⁹ and contributes to the up-regulation of the angiotensin-converting enzyme (ACE)/angiotensin (ANG)-II/ANG II type 1 receptor axis, leading to the development of cardiovascular and pulmonary diseases.¹⁹

In parallel, coronaviruses have distinctive crown-shaped features due to the presence of large type 1 transmembrane spike (S1 and S2) glycoproteins mediating entry of the virus into the host cell. The S1 domain is responsible for first stage host cell entry and contains the angiotensin-converting enzyme-2 (ACE2) receptor. Recent evidence suggests that the SARS-CoV-2 modified S protein exhibited a significantly higher affinity for ACE2 and is 10- to 20-fold more likely to bind to ACE2 in human cells, enabling an easy person-to-person spread of the virus.²⁰ So the action of nicotine on nAChRs facilitates the SARS-CoV-2 host entry.²¹

Smoking and the respiratory function

The lung RAS is involved in the pathogenesis of different lung diseases other than blood pressure and fluid balance, like pulmonary arterial hypertension (PAH)²² lung infection/inflammation,²³ acute respiratory distress syndrome,²⁴ Chronic Obstructive Pulmonary Disease (COPD),²⁵ and pulmonary fibrosis.²⁶

Bronchial epithelial cells absorb inhaled nicotine and smoke components. The gas mixture generated by smoking contains hundreds of components including carbon monoxide, nicotine, oxidants, fine particulate matters (PM), and aldehydes associated with the onset of COPD.²⁷ Inhaled smoke is responsible for the destruction of the matrix, shortage of blood supply, and

death of epithelial cells. It leads to the infiltration of inflammatory cells to mucosa and glandular tissue.²⁷

Furthermore, acrolein, a smoke component, impairs the alveolar-capillary barrier function, resulting in lung inflammation and increased susceptibility to acute respiratory distress disorder (ARDS).²⁸

Chronic smoking significantly decreases pulmonary function, that is, Forced Expiratory Volume (FEV1), Forced Vital Capacity (FVC), Forced Expiratory Flow (FEF), and Fractional Exhaled Nitric Oxide (FeNO).²⁹ The latter is associated with decreased NO and its conversion to peroxynitrite (ONOO) due to oxidative stress.³⁰

Smoking and the immune system

Smoking is a cause of the release of many inflammatory mediators (pro-inflammatory and anti-inflammatory cytokines).³¹ Many studies have demonstrated the association of smoking with chronic systemic inflammation and autoimmune diseases (rheumatoid arthritis, psoriasis, COPD, and systemic lupus erythematosus).³²

Nicotine interacts with nAChRs $\alpha 7$ on the surface of immune cells like macrophages and lymphocytes T and B,³³ and induces an immunosuppressive effect, modulating innate and adaptive immune responses.^{34,35} Therefore, smoking alters the development and function of innate (dendritic cells, macrophages, and natural killer cells) and adaptive immune cells (cytotoxic CD8+ T cells, CD4+ T helper cells (Th), regulatory T cells, and B cells), leading to pro-inflammatory responses, thus worsening smoking-associated COPD.³²

Smoking and oxidative stress

Oxidative stress is the result of an imbalance between the production of reactive oxygen species (ROS) and the antioxidant capacity of the organism. ROS are essential for some signaling pathways, but their excess leads to proteins, lipids, and DNA oxidation, which is involved in some diseases.³⁶ Tobacco smoke constituents induce oxidative stress via different mechanisms, including the direct effects of nicotine (neutralized by N-acetylcysteine) and the abundance of PM.³⁷ Oxidative stress and inflammation lead to the increased production of superoxide anion ($O_2^{\cdot -}$) via the NADPH oxidase and nitric oxide synthase (iNOS) as principal sources of ROS.³⁸ Furthermore, carbon monoxide released by smoking is directly absorbed in the lungs. It forms a tight but reversible bond with hemoglobin and leads to tissue hypoxia. But once eliminated, oxygen reperfusion occurs, thus increasing ROS.³⁹

Waterpipe is also a source of heavy metals such as Pb, Cr and arsenic that lead to free radicals formation.⁴⁰ Acrolein, a toxic unsaturated aldehyde produced by cigarette and waterpipe, binds to proteins and DNA, and induce oxidative stress and inflammation. Acrolein decreases antioxidant enzyme promoting ROS production, oxidizes lipids and produces more

acrolein in a vicious circle.⁴¹ Acute and high dose exposures to acrolein suppresses immune response while in the case for repetitive waterpipe smoking, (chronic exposure with low doses) enhances inflammatory response.⁴²

Smoking and the cardiovascular function

Chronic exposure to nicotine (through the activation of nAChRs, widely expressed in the heart)⁴³ exacerbates pathological changes in the heart. Cigarette smoke accelerates the progression of cardiac hypertrophy and heart failure.⁴⁴ Previous findings demonstrated a possible cardiotoxic interaction between nicotine and the RAS.⁴⁵ Nicotine was shown to exacerbate cardiovascular remodeling induced by ANG II, which included increased heart rate, and increased thickening of the aortic wall.⁴⁵ Both nicotine and RAS are involved in the pathogenesis of vascular calcification and abdominal aortic aneurysms (AAA).⁴¹ Thus, smoking is a considerable risk factor associated with the development, expansion, and rupture of AAA.⁴⁶

Smoking and the endocrine system

The activation of nAChRs affects many organ systems, including the endocrine system. Prolactin levels are directly related to dopamine secretion, while the hypothalamic-pituitary-adrenal (HPA) axis stimulation is associated with the stress level controlled by nicotine administration.⁴⁷ Furthermore, smoking worsens insulin resistance and diabetes mellitus (DM) via many proposed mechanisms, including but not limited to the activation by the nicotine of a macrophage $\alpha 7$ -nAChR associated anti-inflammatory pathway and the exacerbation of nicotine-associated inflammation and insulin resistance.⁴⁸ Another mechanism may be the increased ROS due to the increased expression of tumor necrosis factor- α (TNF- α),⁴⁹ in addition to maternal smoking during pregnancy and breastfeeding, causing insulin resistance later in childhood.⁵⁰

A Known Fact: Harmful Effects of Smoking

In addition to the increased risk of virus transmission, smoking affects a person's overall health and damages nearly every organ of the body.^{51,52} When tobacco toxins enter the body through the mouth and nose, they cause tissue and cells damage up to the lungs,⁵³ including upper lung airways and alveoli, resulting in lung inflammation and reduced lung and immune function.⁵³ Consequently, cigarette smoking causes lung diseases such as bronchiolitis, chronic bronchitis, pulmonary emphysema, tuberculosis, and lung cancers, in addition to an increased risk of respiratory infections.⁵³ While the lung has initial ways to protect itself from damage caused by inhalation agents, these defenses are gradually overwhelmed as tobacco smoke is repeatedly inhaled over time.⁵³ After years of smoking, the lung tissue becomes severely damaged, loses its elasticity, and can no longer easily exchange air.⁵³ Similar to cigarettes, waterpipe

smoke harms pulmonary cells and damages alveoli, causing tissue inflammation and undermining the capacity of the lungs to fight off infection.⁵⁴ Indeed, after a single session, cell damage and lung inflammation have been detected among waterpipe users.⁵⁴ Thus, smoking reduces the ability of the lung to absorb oxygen and release carbon dioxide, storing up mucus, which leads to excessive coughing and difficulties in breathing.⁵³

Since smoking tobacco affects and damages the lungs, it makes it easier for the SARS-CoV-2 to invade the lung tissue, causing more severe symptoms and increasing the risk of death.¹⁸ Smoking also requires high respiratory volumes, deep inhalation, progressive cooling and drying of the mucous respiratory tract, and changes in nose-to-mouth breathing. It reduces the mobility of ciliated cells and raises the viscosity of the mucous membrane, which impairs the filtration of microorganisms from the upper respiratory tract system.⁵³

Smoking and COVID-19: Current Clinical Findings

Based on these mechanisms, smoking is hypothesized to increase risks associated with COVID-19. Several epidemiological and clinical studies conducted to confirm this hypothesis found conflicting results.

Lower prevalence of smoking in COVID-19 infected patients

A spatial analysis involving 175 countries showed that the percentage of the total smoking population was inversely associated with COVID-19 at the global level.⁵⁵ In Mexico, among 89 756 laboratory-confirmed positive cases (retrospective case series used a publicly available nation-level dataset released on May 31, 2020, by the Mexican Ministry of Health), current smokers were 23% less likely to be diagnosed with COVID-19 compared to non-smokers. However, having COPD (among other comorbidities) was associated with hospitalization and adverse outcomes. Current smoking was not linked to unfavorable outcomes.⁵⁶ Similarly, an ecological study found a statistically significant negative association between smoking prevalence and the prevalence of COVID-19 across the 38 European nations, after controlling for confounding factors. However, it could not demonstrate a direct association between smoking prevalence and COVID-19 mortality ($P = .626$).⁵⁷ A preliminary study that analyzed the data available in published articles showed a lower prevalence of smoking in COVID-19 patients compared to the regional average.⁵⁸

It is noteworthy that statistical associations do not imply causation: they might be explained by several facts, such as an interfering dose-effect relationship (whenever the number of packs, the duration, and the timing of smoking, are not measured), reverse causation (some people had stopped smoking during the pandemic), or a residual confounding (additional confounding factors not taken into account). Likely, the low prevalence of smokers among COVID-19 hospitalized patients is partly due to the fact that many smokers were misclassified

as nonsmokers.⁵⁹ Researchers have also identified information and selection biases, in addition to knowledge gaps, which may give the false impression that smoking is protective against COVID-19.⁶⁰

Plausible biologic mechanisms by which smoking might be protective against COVID-19 include an anti-inflammatory effect of nicotine, a blunted immune response among smokers (reducing the risk of a cytokine storm in COVID-19), and increased nitric oxide in the respiratory tract (which may inhibit replication of SARS-CoV-2 and its entry into cells),⁶⁰ in addition to the possible role of squamous cell metaplasia, which is commonly associated with smoking.⁶¹ As of now, however, the data supporting the smoker's paradox claims are limited and questionable. The claims of a protective effect must be viewed with extreme caution by both the general population and clinicians.⁶⁰

Higher prevalence of smoking in COVID-19 infected patients

Several epidemiological studies have found a higher smoking prevalence among COVID-19 infected patients. In a systematic review including thirteen studies the prevalence of current smoking among 5960 hospitalized COVID-19 patients in China ranged from 1.4% to 12.6%.⁶¹ The random effects pooled prevalence of current smoking was 6.5% (95% confidence interval [CI]: 4.9%–8.2%).⁶¹ Another, meta-analysis done by Emami et al.⁶² analyzed data for 2986 patients and found a pooled prevalence of smoking of 7.6% (CI: 3.8%–12.4%). In a cross sectional study done among 53 002 adults from United Kingdom found that the prevalence of confirmed COVID-19 was higher among current smokers (0.56% [CI: 0.41%–0.75%]) as compared with never smokers.⁶³ Data from a hospital in France also indicate that while prevalence of current smoking was lower in hospitalized COVID-19 patients than the general population, prevalence of former smoking was much higher.⁶⁴ In a cross-sectional study among adolescents and young adults, COVID-19 diagnosis was 5 times more likely among ever-users of e-cigarettes only (95% CI: 1.82–13.96), 7 times more likely among ever-dual-users (95% CI: 1.98–24.55), and 6.8 times more likely among past 30-day dual-users (95% CI: 2.40–19.55). Symptoms were also 4.7 times more likely among past 30-day dual-users (95% CI: 3.07–7.16).⁶⁵ Furthermore, a systematic review of reviews also showed a significant association between COVID-19 infection and current or ever smoking.⁶⁶

At this time, it is impossible to estimate the prevalence of smoking status among COVID-19 patients in a population-representative sample.⁶¹ The accuracy and consistency of the reported smoking status is also unknown. Given the emergency of the outbreak, the patient's smoking status could not have been properly reported or certain patients have been unable to register their smoking status.⁶¹ It is also possible that some patients may also have been in critical condition at the time

they were hospitalized, affecting their communication capacity and their capacity to disclose their smoking status.⁶¹

Smoking and COVID-19 progression, severity, and mortality

Numerous studies had shown that people with respiratory diseases caused by tobacco use are at higher risk of developing severe COVID-19 symptoms.^{67–69} A study among 1099 Chinese COVID-19 patients found that 32% of patients with a history of smoking (smokers and ex-smokers) had a severe form of COVID-19 pneumonia at the time of hospitalization, compared with 15% of patients who had never smoked.⁶⁷ Another study among 78 patients with COVID-19-induced pneumonia found that 27% of smokers worsened within 2 weeks of hospitalization, compared to 3% of non-smokers.⁶⁸ A meta-analysis based on 15 studies showed that 22% of current smokers and 46% of ex-smokers had severe complications of the COVID-19.⁶⁹

During a follow up of 28 days among 720 patients, 71.50% had recovered or were symptomatically stable, COPD was a risk factor of progression, while smoking was inversely related to it.⁷⁰ A meta-analysis found that smoking is a risk factor for COVID-19 progression, with smokers having higher odds of COVID-19 progression than never smokers (OR = 1.91).⁷¹ In Kuwait, among 1096 consecutive COVID-19 patients, the risk factors found to be significantly associated with admission to intensive care were age above 50 years old, a qSOFA score above 0, smoking, elevated CRP, and elevated procalcitonin levels. Smoking and elevated procalcitonin levels also correlated significantly with mortality.⁷² In a study to characterize a large cohort of patients with COVID-19 (n = 4536) and their outcomes, several factors were involved with increased hospitalization risk, including former smoking history, shortness of breath, and certain medications (NSAIDs, immunosuppressive treatment).⁷³ A Brazilian study on 10 713 COVID-19 patients found that both smoking and pulmonary disease increased the risk of hospitalization.⁷⁴ Other smaller studies showed similar results.⁷⁵ Among patients with lung cancer, determinants of COVID-19 severity got largely patient-specific features, including smoking status and COPD (odds ratio for severe COVID-19 = 2.9).⁷⁶ Moreover, there is a possible role of smoking and vaping on cerebrovascular and neurological dysfunction in COVID-19 patients, along with potential pathogenic mechanisms associated with it.⁷⁷

When assessing clinical characteristics and risk factors for mortality of patients with COVID-19 from Mexico, smoking was found to increase the risk of death. Less frequent comorbidities, such as COPD, chronic kidney disease, and immunosuppression, also showed a significant risk for death ($P < .0001$).⁷⁸ Common independent factors associated with an increased risk of death were older age, history of smoking status, number of comorbidities, more advanced performance status, and active cancer.⁷⁹ Moreover, a pooled meta-analysis

showed that the prevalence of respiratory diseases, specifically COPD (OR 4.21; 95% CI, 2.9-6.0), and smoking (current smoking OR 1.98; 95% CI, 1.16-3.39 and former smoking OR 3.46; 95% CI, 2.46-4.85) were significantly associated with severe COVID-19 outcomes.⁸⁰ Another meta-analysis also showed that current smokers were more likely to have an adverse outcome compared with non-current smokers (odds ratio [OR]: 1.53, 95%CI: 1.06-2.20, $P=.022$) but less likely compared with former smokers (OR: 0.42, 95% CI: 0.27-0.74, $P=.003$).⁸¹ An older meta-analysis also found similar results: current smokers were 1.45 times more likely (95% CI: 1.03-2.04) to have severe complications compared to former and never smokers. Current smokers also had a higher mortality rate of 38.5%.⁶⁹

COVID-19 disease state worsening by smoking is evidence based, despite some studies claiming a reduced SARS-CoV-2 infection risk in smoker population. In fact, a review assessing the relationship between smoking status and some COVID-19 disease outcomes (infection, hospitalization, disease severity, and mortality), showed no important association between being a smoker and hospitalization or mortality and that current smokers were at lower risk of getting infected with SARS-CoV-2 compared to never smokers. But the authors related these results to study specific factors that complicated the interpretation, including inverse causality.⁸² In all cases, it is impossible to prove causality through observational studies or to conduct interventional studies, except for smoking cessation trials in the general population and among patients after COVID.

Confirmation of Smoking as a Risk Factor for COVID-19 in vitro

Even if smoking may not necessarily increase one's risk for contracting COVID-19, the biological and inflammatory cascade that occurs upon SARS-CoV-2 infection may be particularly devastating for a smoker.⁸³

Smoking and demographics suggested mechanisms

Evidence has shown that smoker patients with different comorbidities (DM, cardiovascular diseases (CVD), COPD, cancer etc.) are at higher risk of contracting SARS-CoV-2. They also have a worse prognosis for COVID-19 as well as for their comorbidities, especially for higher age patients. Older men are in particular at higher risk of testing positive for COVID-19 because smoking rates are highest for men, and older male smokers will have smoked for most of their lives (causing maximum damage to their respiratory and cardiovascular systems). In fact the SARS-CoV-2 viral capsid binds to surface ACE2 and activates a cellular serine protease *TMPRSS2* (an androgen-responsive gene highly expressed in men⁸⁴) for protein priming.⁸⁵ Therefore, the higher expression of *TMPRSS2* in men could increase their vulnerability to COVID-19,⁸⁴ thus explaining the higher morbidity and mortality associated with older age and male gender.⁸⁶

Smoking and comorbidities suggested mechanisms

DM may aggravate the risk of severe COVID-19⁸⁷ and mortality because it increases the surface expression of ACE2 through hyperinsulinemia,⁸⁸ in addition to other synergistic mechanisms. DM is associated with immunodeficiency due to reduced TNF- α and impaired Treg cells, which facilitates infections.^{88,89}

CVD patients are also predisposed to an increased risk of more severe disease for many reasons. Hypertension is associated with a dysregulation in the immune system, manifested by elevated IL-17 levels and abnormal T-cell function.⁹⁰ Other factors include overactive sympathetic drive and elevations in angiotensin II, a pro-inflammatory peptide.⁹¹ ACE2 modulators (ACE1 inhibitors, angiotensin receptor blockers, and thiazolidinediones), frequently used to treat hypertension, upregulate ACE2 expression.²⁹

Furthermore, the expression of ACE-2 receptors is increased in COPD, leading to severe COVID-19 symptoms, including structural damage to lungs, lower immunity, and increased mucous production and airway blocking.⁹² Consequently, higher morbidity and mortality are associated with chronic lung diseases.⁸⁶

Patients suffering from any malignancy, exacerbated by smoking, are also at a higher risk of developing COVID-19 infection due to the weak immunity providing an efficient viral replication environment.⁹³

Additional smoking suggested mechanisms

The impact of the toxic constituents of tobacco products on endothelial function has been thoroughly discussed. Studies have shown that smoking (cigarettes, waterpipe, and e-cigarette) may have contributed to an increased burden of COVID-19 symptoms and severe health consequences.⁹⁴ In vitro, the acute exposure allows for more severe proximal airway epithelial disease from SARS-CoV-2 by reducing the mucosal innate immune response and the proliferation of airway basal stem cells and has implications for disease spread and severity in people exposed to cigarette smoke, with a more severe viral infection and cell death.⁹⁵ Furthermore, the viral entry has been shown to cause cytokine storm involving excessive production of pro-inflammatory cytokines/chemokines, including IL-6, TNF- α , IFN- γ , IL-2, IL-7, IP-10, MCP-3, or GM-CSF, which is increased by smoking.⁹⁶ Also, the expression levels of *TMPRSS4*, a gene coding for a protease that primes SARS-CoV-2 for cell entry similarly to *TMPRSS2*, were elevated in bronchial epithelial cells from current smokers compared with never smokers, suggesting that higher bronchial *TMPRSS4* levels in smokers might put them at higher risk for SARS-Cov-2 infection.⁹⁷

Chronic smoke exposure triggers the expansion of this cell population and a concomitant increase in ACE2 expression. In contrast, quitting smoking decreases the abundance of these

secretory cells and reduces ACE2 levels. Thus, ACE2 expression is responsive to inflammatory signaling and can be upregulated by viral infections or interferon treatment. Taken together, these results may partly explain why smokers are particularly susceptible to severe SARS-CoV-2 infections. Furthermore, ACE2 was identified as an interferon-stimulated gene in lung cells, suggesting that SARS-CoV-2 infections could create positive feedback loops that increase ACE2 levels and facilitate viral dissemination.⁹⁸ This biological phenomenon could also apply to new electronic smoking devices such as electronic cigarettes and “heat-not-burn” IQOS devices.²⁰ In the same line, smoking may upregulate ACE2 expression through the upregulation of the androgen pathway. These data provide a potential model for the increased susceptibility of smoking patients to COVID-19.⁹⁹

Finally, increased age, comorbidities, smoking status, and recent chemotherapy were associated with higher viral load. In turn, a high viral load (OR=6) was linked to an increased need for intubation (OR=3) and in-hospital mortality.¹⁰⁰

Future Perspectives

Further investigations of the interaction between smoking and COVID-19 are warranted to accurately assess the risk of contracting COVID-19 among smokers, and the progression to mechanical ventilation or death in patients who suffer from it. Also, further systematic review and meta-analysis studies are needed that could provide a critical analyses of the available data suggesting Covid19 susceptibilities amongst smokers. Risk factors for COVID-19, such as cardiovascular disease, chronic obstructive pulmonary disease, and diabetes, are all strongly associated with smoking habits.¹⁰¹ Thus, the probability of a beneficial effect of smoking on COVID-19 outcomes is manifestly unlikely. Future investigation should focus on homogenizing variables' definitions, leading to less heterogeneity and easier results interpretation.

Authors of potentially controversial articles should thus restrain from promoting their results before the peer-review process is complete,¹⁰² given all the limitations and biases that they could include, and the false message they could convey. This may be particularly exploited by tobacco industry to promote smoking as not only safe activity but as preventive agent from COVID-19 complications. Furthermore, given the likely benefits of smoking cessation in vulnerable populations, the current pandemic should provide an opportunity to enforce smoking cessation in health facilities (including mental health) and encourage nicotine replacement therapy alongside smoking cessation counseling, to increase the rate of quitting.¹⁰³⁻¹⁰⁵ For instance, studies aiming at assessing smokers' attitude toward quitting smoking during the pandemic, showed increased motivation to quit and decreased quantity of smoking.¹⁰⁶ This may offer for policy makers a strong justification for implementing strong antismoking campaigns while making quitting practices available and affordable for the general public. Moreover, nicotine, nicotinic agonists, or modulators of

nicotinic receptors may play a potential therapeutic role in COVID-19, owing to their varied effects, including mood regulation, anti-inflammatory, as well as purported interference with SARS-CoV-2 entry and/or replication.¹⁰⁷ The hypothesis that nicotine may have a protective effect against COVID-19, partially masked by smoking-related toxicity and abrupt cessation of nicotine consumption when smokers are hospitalized, need to be explored in laboratory studies using nicotine-based pharmaceutical products.⁸¹ Additionally, its role in the prevention and treatment of COVID-19 should be evaluated in appropriate placebo-controlled trials.¹⁰⁴

Limitations of epidemiological data

Smoking contributes to COVID-19 morbidity and mortality as explained by several biological mechanisms, while epidemiological data is not conclusive whether smoking is associated with increased odds of infection as well as worse disease state. The inconclusive results may be due to some limitations. First data was not collected directly from the participants (due to the applied social distancing policies). These were rather provided by data banks (like the European bank or the WHO data. . .), so questions about smoking habits may not have adequately reported. Also some hospitalized COVID-19 patients may also have been unable to communicate and disclose their smoking habits because of their critical condition. Some inflated results may be due to the fact that current smokers, negative for COVID-19, confusing their chronic smoking symptoms (cough, increased sputum, altered sense of smell or taste) with those of COVID-19 with those of COVID-19, and may present for testing more frequently than nonsmokers.⁸² Studies taking into consideration potential confounders, selection and information biases are recommended. Finally, tobacco industry may have exploited some investigations drawbacks to diffuse falsified studies about the benefits of smoking as a protective agent against COVID-19.

In conclusion, physicians and public health professionals should adequately collect detailed data from trusted sources on smoking as part of clinical management to draw useful conclusions for policy makers to adopt. This will promote smoking cessation as a lead practice, contributing to the mitigation of the COVID-19 pandemic.⁷¹ Further studies, including systematic reviews and meta-analyses, and studies with appropriate assessment of potential confounders and bias minimization, are necessary to conclude about the association of smoking with COVID-19 severity and fatality.

ORCID iDs

Chadia Haddad  <https://orcid.org/0000-0003-2413-2684>

Hala Sacre  <https://orcid.org/0000-0001-6547-6251>

REFERENCES

1. Bommele J, Hopman P, Walters BH, et al. The double-edged relationship between COVID-19 stress and smoking: implications for smoking cessation. *Tob Induc Dis.* 2020;18:63.

2. Kowitz SD, Cornacchione Ross J, Jarman KL, et al. Tobacco quit intentions and behaviors among cigar smokers in the United States in response to COVID-19. *Int J Environ Res Public Health*. 2020;17:5368.
3. Caponnetto P, Inguscio L, Saitta C, Maglia M, Benfatto F, Polosa R. Smoking behavior and psychological dynamics during COVID-19 social distancing and stay-at-home policies: a survey. *Health Psychol Res*. 2020;8:9124.
4. Sun Y, Li Y, Bao Y, et al. Brief report: increased addictive internet and substance use behavior during the COVID-19 pandemic in China. *Am J Addict*. Published online June 4, 2020. doi:10.1111/ajad.13066
5. Patwardhan P. COVID-19: risk of increase in smoking rates among England's 6 million smokers and relapse among England's 11 million ex-smokers. *BJGP open*. 2020;4:bjgpopen20X101067.
6. Chertok IRA. Perceived risk of infection and smoking behavior change during COVID-19 in Ohio. *Public Health Nurs*. Published online September 27, 2020. doi:10.1111/phn.12814
7. Ahmed N, Maqsood A, Abduljabbar T, Vohra F. Tobacco smoking a potential risk factor in transmission of COVID-19 infection. *Pak J Med Sci*. 2020;36:S104-S107.
8. Van Doremalen N, Bushmaker T, Morris DH, et al. Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1. *N Engl J Med*. 2020;382:1564-1567.
9. Zhu N, Zhang D, Wang W, et al. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med*. 2020;382:727-733.
10. World Health Organization. *Modes of Transmission of Virus Causing COVID-19: Implications for IPC Precaution Recommendations: Scientific Brief, 27 March 2020*. World Health Organization; 2020.
11. Sabino-Silva R, Jardim ACG, Siqueira WL. Coronavirus COVID-19 impacts to dentistry and potential salivary diagnosis. *Clin Oral Investig*. 2020;24:1619-1621.
12. Ranieri M. COVID-19 self-infection in cigarette smoking: description of the mechanism. Preprint. doi:10.13140/RG.2.2.15310.00321 2020
13. Alagaili AN, Briese T, Amor NM, Mohammed OB, Lipkin WI. Waterpipe smoking as a public health risk: potential risk for transmission of MERS-CoV. *Saudi J Biol Sci*. 2019;26:938-941.
14. World Health Organization. Knowledge hub for waterpipe tobacco smoking. Increased risk of COVID-19 infection amongst smokers and amongst waterpipe users. 2020. Accessed August 25, 2020. <https://untobaccocontrol.org/kh/waterpipes/covid-19/#:~:text=Increased%20risk%20of%20Coronavirus%20infection%20amongst%20smokers&text=Among%20Chinese%20patients%20diagnosed%20with,those%20who%20did%20not%20smoke>
15. Shekhar S, Hannah-Shmouni F. Hookah smoking and COVID-19: call for action. *CMAJ*. 2020;192:E462-E462.
16. Maziak W, Taleb ZB, Bahelah R, et al. The global epidemiology of waterpipe smoking. *Tob control*. 2015;24(suppl 1):i3-i12.
17. Centers for Disease Control and Prevention. Coronavirus disease 2019 (COVID-19). Community mitigation resources. 2020. Accessed August 25, 2020. <https://www.cdc.gov/coronavirus/2019-ncov/php/open-america/community-mitigation-concepts.html>
18. World Health Organization. Tobacco Free initiative. Tobacco and waterpipe use increases the risk of COVID-19. 2020. Accessed August 25, 2020. <http://www.emro.who.int/tfi/known-the-truth/tobacco-and-waterpipe-users-are-at-increased-risk-of-covid-19-infection.html>
19. Oakes JM, Fuchs RM, Gardner JD, Lazartigues E, Yue X. Nicotine and the renin-angiotensin system. *Am J Physiol Regul Integr Comp Physiol*. 2018;315:R895-R906.
20. Brake SJ, Barnsley K, Lu W, McAlinden KD, Eapen MS, Sohal SS. *Smoking Upregulates Angiotensin-Converting Enzyme-2 Receptor: A Potential Adhesion Site for Novel Coronavirus SARS-CoV-2 (Covid-19)*. Multidisciplinary Digital Publishing Institute; 2020.
21. Changeux J-P, Amoura Z, Rey FA, Miyara M. A nicotinic hypothesis for Covid-19 with preventive and therapeutic implications. *C R Biol*. 2020;343:33-39.
22. de Man FS, Tu L, Handoko ML, et al. Dysregulated renin-angiotensin-aldosterone system contributes to pulmonary arterial hypertension. *Am J Respir Crit Care Med*. 2012;186:780-789.
23. Jia H. Pulmonary angiotensin-converting enzyme 2 (ACE2) and inflammatory lung disease. *Shock*. 2016;46:239-248.
24. Imai Y, Kuba K, Penninger JM. The discovery of angiotensin-converting enzyme 2 and its role in acute lung injury in mice. *Exp Physiol*. 2008;93:543-548.
25. Han S-X, He G-M, Wang T, et al. Losartan attenuates chronic cigarette smoke exposure-induced pulmonary arterial hypertension in rats: possible involvement of angiotensin-converting enzyme-2. *Toxicol Appl Pharmacol*. 2010;245:100-107.
26. Bruce E, Shenoy V, Rathinasabapathy A, et al. Selective activation of angiotensin AT 2 receptors attenuates progression of pulmonary hypertension and inhibits cardiopulmonary fibrosis. *Br. J. Pharmacol*. 2015;172:2219-2231.
27. Alwis KU, deCastro BR, Morrow JC, Blount BC. Acrolein exposure in US tobacco smokers and non-tobacco users: NHANES 2005-2006. *Environ Health Perspect*. 2015;123:1302-1308.
28. Borgas D, Chambers E, Newton J, et al. Cigarette smoke disrupted lung endothelial barrier integrity and increased susceptibility to acute lung injury via histone deacetylase 6. *Am J Respir Cell Mol Biol*. 2016;54:683-696.
29. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*. 2020;395:1054-1062.
30. Barua RS, Ambrose JA, Srivastava S, DeVoe MC, Eales-Reynolds L-J. Reactive oxygen species are involved in smoking-induced dysfunction of nitric oxide biosynthesis and upregulation of endothelial nitric oxide synthase: an in vitro demonstration in human coronary artery endothelial cells. *Circulation*. 2003;107:2342-2347.
31. Gonçalves R, Coletta R, Silvério K, et al. Impact of smoking on inflammation: overview of molecular mechanisms. *Inflamm Res*. 2011;60:409-424.
32. Qiu F, Liang C-L, Liu H, et al. Impacts of cigarette smoking on immune responsiveness: up and down or upside down? *Oncotarget*. 2017;8:268.
33. Cloëz-Tayarani I, Changeux JP. Nicotine and serotonin in immune regulation and inflammatory processes: a perspective. *J Leukoc Biol*. 2007;81:599-606.
34. Sopori M. Effects of cigarette smoke on the immune system. *Nat Rev Immunol*. 2002;2:372-377.
35. Cui W-Y, Li MD. Nicotinic modulation of innate immune pathways via $\alpha 7$ nicotinic acetylcholine receptor. *J Neuroimmune Pharmacol*. 2010;5:479-488.
36. Steven S, Frenis K, Oelze M, et al. Vascular inflammation and oxidative stress: major triggers for cardiovascular disease. *Oxid Med Cell Longev*. 2019;2019:7092151.
37. Qasim H, Alarabi AB, Alzoubi KH, Karim ZA, Alshbool FZ, Khasawneh FT. The effects of hookah/waterpipe smoking on general health and the cardiovascular system. *Environ Health Prev Med*. 2019;24:1-17.
38. Sun L, Liu C, Xu X, et al. Ambient fine particulate matter and ozone exposures induce inflammation in epicardial and perirenal adipose tissues in rats fed a high fructose diet. *Part Fibre Toxicol*. 2013;10:1-20.
39. Piantadosi CA. Carbon monoxide, reactive oxygen signaling, and oxidative stress. *Free Radic Biol Med*. 2008;45:562-569.
40. Ercal N, Gurer-Orhan H, Aykin-Burns N. Toxic metals and oxidative stress part I: mechanisms involved in metal-induced oxidative damage. *Curr Top Med Chem*. 2001;1:529-539.
41. Sui Y-B, Chang J-R, Chen W-J, et al. Angiotensin-(1-7) inhibits vascular calcification in rats. *Peptides*. 2013;42:25-34.
42. Moghe A, Ghare S, Lamoreau B, et al. Molecular mechanisms of acrolein toxicity: relevance to human disease. *Toxicol Sci*. 2015;143:242-255.
43. Dvorakova M, Lips KS, Brüggmann D, Slavikova J, Kuncova J, Kummer W. Developmental changes in the expression of nicotinic acetylcholine receptor α -subunits in the rat heart. *Cell Tissue Res*. 2005;319:201-209.
44. Meurrens K, Ruf S, Ross G, Schlee R, von Holt K, Schlüter K-D. Smoking accelerates the progression of hypertension-induced myocardial hypertrophy to heart failure in spontaneously hypertensive rats. *Cardiovasc Res*. 2007;76:311-322.
45. Colombo ES, Davis J, Makvandi M, et al. Effects of nicotine on cardiovascular remodeling in a mouse model of systemic hypertension. *Cardiovasc Toxicol*. 2013;13:364-369.
46. Daugherty A, Cassis LA. Mouse models of abdominal aortic aneurysms. *Arterioscler Thromb Vasc Biol*. 2004;24:429-434.
47. Tweed JO, Hsia SH, Lutfy K, Friedman TC. The endocrine effects of nicotine and cigarette smoke. *Trends Endocrinol Metab*. 2012;23:334-342.
48. Wang X, Yang Z, Xue B, Shi H. Activation of the cholinergic anti-inflammatory pathway ameliorates obesity-induced inflammation and insulin resistance. *Endocrinology*. 2011;152:836-846.
49. Tatebe J, Morita T. Enhancement of TNF- α expression and inhibition of glucose uptake by nicotine in the presence of a free fatty acid in C2C12 skeletal myocytes. *Horm Metab Res*. 2011;43:11-16.
50. Bruin JE, Gerstein HC, Holloway AC. Long-term consequences of fetal and neonatal nicotine exposure: a critical review. *Toxicol Sci*. 2010;116:364-374.
51. US Department of Health and Human Services. *How Tobacco Smoke Causes Disease: What It Means To You*. US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2010.
52. US Department of Health and Human Services. *The Health Consequences of Smoking—50 Years of Progress: A Report of the Surgeon General*. US Department of Health and Human Services, Centers for Disease Control and Prevention; 2014.
53. Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health. *How Tobacco Smoke Causes Disease: The Biology and Behavioral Basis for Smoking-Attributable Disease: A Report of the Surgeon General*. Centers for Disease Control Prevention; 2010.

54. Nemmar A, Al-Salam S, Yuvaraju P, Beegam S, Yasin J, Ali BH. Chronic exposure to water-pipe smoke induces alveolar enlargement, DNA damage and impairment of lung function. *Cell Physiol Biochem*. 2016;38:982-992.
55. Iyanda AE, Adeleke R, Lu Y, et al. A retrospective cross-national examination of COVID-19 outbreak in 175 countries: a multiscale geographically weighted regression analysis (January 11-June 28, 2020). *J Infect Public Health*. 2020;13:1438-1445.
56. Yannouchos T, Sussman R, Mier JM, Poulas K, Farsalinos K. Characteristics and risk factors for COVID-19 diagnosis and adverse outcomes in Mexico: an analysis of 89,756 laboratory-confirmed COVID-19 cases. *Eur Respir J*. 2020;2002144.
57. Tsigaris P, Teixeira da Silva JA. Smoking prevalence and COVID-19 in Europe. *Nicotine Tob Res*. 2020;22:1646-1649.
58. Tajlil A, Ghaffari S, Pourafkari L, Mashayekhi S, Roshanravan N. Nicotine and smoking in the COVID-19 era. *J Cardiovasc Thorac Res*. 2020;12:136.
59. Cattaruzza M, Zagà V, Gallus S, D'Argenio P, Gorini G. Tobacco smoking and COVID-19 pandemic: old and new issues. A summary of the evidence from the scientific literature. *Acta Biomed*. 2020;91:106-112.
60. Usman MS, Siddiqi TJ, Khan MS, et al. Is there a smoker's paradox in COVID-19? *BMJ Evid Based Med*. 2020.
61. Farsalinos K, Barbouni A, Niaura R. Systematic review of the prevalence of current smoking among hospitalized COVID-19 patients in China: could nicotine be a therapeutic option? *Intern Emerg Med*. 2020;15:845-852.
62. Emami A, Javanmardi F, Pirbonyeh N, Akbari A. Prevalence of underlying diseases in hospitalized patients with COVID-19: a systematic review and meta-analysis. *Arch Acad Emerg Med*. 2020;8:e35.
63. Jackson SE, Brown J, Shahab L, Steptoe A, Fancourt D. COVID-19, smoking and inequalities: a study of 53 002 adults in the UK. *Tob Control*. 2020.
64. Miyara M, Tubach F, Pourcher V, Morelot-Panzini C, Pernet J, Haroche J. Low rate of daily active tobacco smoking in patients with symptomatic COVID-19. *Qeios*. Published online May 2020.
65. Gaiha SM, Cheng J, Halpern-Felsher B. Association between youth smoking, electronic cigarette use, and Coronavirus Disease 2019. *J Adolesc Health*. 2020;67:519-523.
66. Grundy EJ, Suddek T, Filippidis FT, Majeed A, Coronini-Cronberg S. Smoking, SARS-CoV-2 and COVID-19: a review of reviews considering implications for public health policy and practice. *Tob Induc Dis*. 2020;18:58.
67. Guan W-j, Ni Z-y, Hu Y, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med*. 2020;382:1708-1720.
68. Liu W, Tao Z-W, Wang L, et al. Analysis of factors associated with disease outcomes in hospitalized patients with 2019 novel coronavirus disease. *Chin Med J*. 2020;133:1032-1038.
69. Alqahtani JS, Oyelade T, Aldahir AM, et al. Prevalence, severity and mortality associated with COPD and smoking in patients with COVID-19: a rapid systematic review and meta-analysis. *PLoS One*. 2020;15:e0233147.
70. Cen Y, Chen X, Shen Y, et al. Risk factors for disease progression in mild to moderate COVID-19 patients—a multi-center observational study. *Clin Microbiol Infect*. 2020;26:1242-1247.
71. Patanavanich R, Glantz SA. Smoking is associated with COVID-19 progression: a meta-analysis. *Nicotine Tob Res*. 2020;22:1653-1656.
72. Almazeedi S, Al-Youha S, Jamal MH, et al. Characteristics, risk factors and outcomes among the first consecutive 1096 patients diagnosed with COVID-19 in Kuwait. *EclinicalMedicine*. 2020;24:100448.
73. Jehi L, Ji X, Milinovich A, et al. Development and validation of a model for individualized prediction of hospitalization risk in 4,536 patients with COVID-19. *PLoS One*. 2020;15:e0237419.
74. Soares RdCM, Mattos LR, Raposo LM. Risk factors for hospitalization and mortality due to COVID-19 in Espírito Santo State, Brazil. *Am J Trop Med Hyg*. 2020;103:1184-1190.
75. Zhao Z, Chen A, Hou W, et al. Prediction model and risk scores of ICU admission and mortality in COVID-19. *PLoS One*. 2020;15:e0236618.
76. Luo J, Rizvi H, Preeshagul IR, et al. COVID-19 in patients with lung cancer. *Ann Oncol*. 2020;31:1386-1396.
77. Archie SR, Cucullo L. Cerebrovascular and neurological dysfunction under the threat of COVID-19: is there a comorbid role for smoking and vaping? *Int J Mol Sci*. 2020;21:3916.
78. Parra-Bracamonte G, Lopez-Villalobos N, Parra-Bracamonte F. Clinical characteristics and risk factors for mortality of patients with COVID-19 in a large dataset from Mexico. *Ann Epidemiol*. 2020;52:93-98.e2.
79. Curigliano G. Cancer patients and risk of mortality for COVID-19. *Cancer Cell*. 2020;38:161-163.
80. Sanchez-Ramirez DC, Mackey D. Underlying respiratory diseases, specifically COPD, and smoking are associated with severe COVID-19 outcomes: a systematic review and meta-analysis. *Respir Med*. 2020;171:106096.
81. Farsalinos K, Barbouni A, Poulas K, Polosa R, Caponnetto P, Niaura R. Current smoking, former smoking, and adverse outcome among hospitalized COVID-19 patients: a systematic review and meta-analysis. *Ther Adv Chronic Dis*. 2020;11:2040622320935765.
82. Simons D, Shahab L, Brown J, Perski O. The association of smoking status with SARS-CoV-2 infection, hospitalisation and mortality from COVID-19: a living rapid evidence review. *Addiction*. 2020.
83. Leung JM, Sin DD. Smoking, ACE-2, and COVID-19: ongoing controversies. *Eur Respir J*. 2020;56:2001759.
84. Asselta R, Paraboschi EM, Mantovani A, Duga S. ACE2 and TMPRSS2 variants and expression as candidates to sex and country differences in COVID-19 severity in Italy. *Aging*. 2020;12:10087-10098.
85. Hoffmann M, Kleine-Weber H, Schroeder S, et al. SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell*. 2020;181:271-280.e8.
86. Porzionato A, Emmi A, Barbon S, et al. Sympathetic activation: a potential link between comorbidities and COVID-19. *FEBS J*. 2020;287:3681-3688.
87. Apicella M, Campopiano MC, Mantuano M, Mazoni L, Coppelli A, Del Prato S. COVID-19 in people with diabetes: understanding the reasons for worse outcomes. *Lancet Diabetes Endocrinol*. 2020;8:782-792.
88. Li B, Yang J, Zhao F, et al. Prevalence and impact of cardiovascular metabolic diseases on COVID-19 in China. *Clin Res Cardiol*. 2020;109:531-538.
89. Sardu C, D'Onofrio N, Balestrieri ML, et al. Outcomes in patients with hyperglycemia affected by Covid-19: can we do more on glycemic control? *Diabetes Care*. 2020;43:1408-1415.
90. Scheen AJ, Marre M, Thivolet C. Prognostic factors in patients with diabetes hospitalized for COVID-19: findings from the CORONADO study and other recent reports. *Diabetes Metab*. 2020;46:265-271.
91. Iacobellis G, Penaherrera CA, Bermudez LE, Mizrachi EB. Admission hyperglycemia and radiological findings of SARS-CoV2 in patients with and without diabetes. *Diabetes Res Clin Pract*. 2020;164:108185.
92. Wan Y, Shang J, Graham R, Baric RS, Li F. Receptor recognition by the novel coronavirus from Wuhan: an analysis based on decade-long structural studies of SARS coronavirus. *J Virol*. 2020;94:e00127-20.
93. Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*. 2020;395:507-513.
94. Münzel T, Hahad O, Kuntic M, Keaney JF, Deanfield JE, Daiber A. Effects of tobacco cigarettes, e-cigarettes, and waterpipe smoking on endothelial function and clinical outcomes. *Eur Heart J*. 2020;41:4057-4070.
95. Purkayastha A, Sen C, Garcia G, et al. Direct exposure to SARS-CoV-2 and cigarette smoke increases infection severity and alters the stem cell-derived airway repair response. *Cell Stem Cell*. 2020;27:869-875.e4.
96. Kaur G, Lungarella G, Rahman I. SARS-CoV-2 COVID-19 susceptibility and lung inflammatory storm by smoking and vaping. *J Inflamm*. 2020;17:21.
97. Voinsky I, Gurwitz D. Smoking and COVID-19: similar bronchial ACE2 and TMPRSS2 expression and higher TMPRSS4 expression in current versus never smokers. *Drug Dev Res*. Published online August 5, 2020.doi:10.1002/ddr.21729
98. Smith JC, Sausville EL, Girish V, et al. Cigarette smoke exposure and inflammatory signaling increase the expression of the SARS-CoV-2 receptor ACE2 in the respiratory tract. *Dev Cell*. 2020;53:514-529.e3.
99. Chakladar J, Shende N, Li WT, Rajasekaran M, Chang EY, Ongkeko WM. Smoking-mediated upregulation of the androgen pathway leads to increased SARS-CoV-2 susceptibility. *Int J Mol Sci*. 2020;21:3627.
100. Magleby R, Westblade LF, Trzebecki A, et al. Impact of SARS-CoV-2 viral load on risk of intubation and mortality among hospitalized patients with coronavirus disease 2019. *Clin Infect Dis*. Published online June 30, 2020. doi:10.1093/cid/ciaa851
101. Komiya M, Hasegawa K. Smoking cessation as a public health measure to limit the coronavirus disease 2019 pandemic. *Eur Cardiol*. 2020;15:e16.
102. Kamiński M, Muth A, Bogdański P. Smoking, vaping, and tobacco industry during COVID-19 pandemic: Twitter data analysis. *Cyberpsychol Behav Soc Netw*. 2020;23:811-817.
103. Seiler NK, Swamy R, Xiao J, Yun Y. Tobacco smoking cessation in mental health services during the COVID-19 pandemic. *J Addict Dis*. 2020;38:582-584.
104. Boutou AK, Pitsiou G, Kontakiotis T, Kioumis I. Nicotine treatment and smoking cessation in the era of COVID-19 pandemic: an interesting alliance. *ERJ Open Res*. 2020;6:00306-2020.
105. Tetik BK, Tekinemre IG, Taş S. The effect of the COVID-19 pandemic on smoking cessation success. *J Community Health*. Published online July 8, 2020. doi:10.1007/s10900-020-00880-2
106. Klemperer EM, West JC, Peasley-Miklus C, Villanti AC. Change in tobacco and electronic cigarette use and motivation to quit in response to COVID-19. *Nicotine Tob Res*. 2020;22:1662-1663.
107. Tizabi Y, Getachew B, Copeland RL, Aschner M. Nicotine and the cholinergic system in COVID-19. *FEBS J*. 2020;287:3656-3663.