

# The immunometabolite itaconate inhibits heme synthesis and remodels cellular metabolism in erythroid precursors

Jason R. Marcero,<sup>1</sup> James E. Cox,<sup>2,3</sup> Hector A. Bergonia,<sup>4</sup> Amy E. Medlock,<sup>1,5</sup> John D. Phillips,<sup>4</sup> and Harry A. Dailey Jr<sup>1,6</sup>

<sup>1</sup>Department of Biochemistry and Molecular Biology, University of Georgia, Athens, GA; <sup>2</sup>Department of Biochemistry; <sup>3</sup>Metabolomics, Proteomics and Mass Spectrometry Cores, University of Utah, Salt Lake City, UT; <sup>4</sup>Division of Hematology, Department of Medicine, University of Utah School of Medicine, Salt Lake City, UT; <sup>5</sup>Augusta University/University of Georgia Medical Partnership, Athens, GA; and <sup>6</sup>Department of Microbiology, University of Georgia, Athens, GA

## Key Points

- The immunometabolite itaconate is taken up by erythroid precursors and converted to itaconyl-CoA by the CoA transferase SUGCT.
- Itaconyl-CoA is a competitive inhibitor of ALAS2 and inhibits erythropoietic heme synthesis.

As part of the inflammatory response by macrophages, *Irg1* is induced, resulting in millimolar quantities of itaconate being produced. This immunometabolite remodels the macrophage metabolome and acts as an antimicrobial agent when excreted. Itaconate is not synthesized within the erythron but instead may be acquired from central macrophages within the erythroid island. Previously, we reported that itaconate inhibits hemoglobinization of developing erythroid cells. Herein we show that this action is accomplished by inhibition of tetrapyrrole synthesis. In differentiating erythroid precursors, cellular heme and protoporphyrin IX synthesis are reduced by itaconate at an early step in the pathway. In addition, itaconate causes global alterations in cellular metabolite pools, resulting in elevated levels of succinate, 2-hydroxyglutarate, pyruvate, glyoxylate, and intermediates of glycolytic shunts. Itaconate taken up by the developing erythron can be converted to itaconyl-coenzyme A (CoA) by the enzyme succinyl-CoA:glutarate-CoA transferase. Propionyl-CoA, propionyl-carnitine, methylmalonic acid, heptadecanoic acid, and nonanoic acid, as well as the aliphatic amino acids threonine, valine, methionine, and isoleucine, are increased, likely due to the impact of endogenous itaconyl-CoA synthesis. We further show that itaconyl-CoA is a competitive inhibitor of the erythroid-specific 5-aminolevulinic acid synthase (ALAS2), the first and rate-limiting step in heme synthesis. These findings strongly support our hypothesis that the inhibition of heme synthesis observed in chronic inflammation is mediated not only by iron limitation but also by limitation of tetrapyrrole synthesis at the point of ALAS2 catalysis by itaconate. Thus, we propose that macrophage-derived itaconate promotes anemia during an inflammatory response in the erythroid compartment.

## Introduction

Induction of heme biosynthesis is a critical process in differentiating erythroid cell precursors and is ultimately responsible for the production of >400 billion hemoglobin molecules per second in adult human bone marrow.<sup>1</sup> In metazoans, heme production in the erythroid compartment is a committed pathway that consists of 8 enzymatic steps. Erythroid heme synthesis begins with production of 5-aminolevulinic acid (ALA) via the rate-limiting enzyme ALA synthase 2 (ALAS2) and concludes with the insertion of ferrous iron into protoporphyrin IX (PPIX) by ferrochelatase (FECH)<sup>2,3</sup> (Figure 1). *ALAS2* is upregulated by

Submitted 15 March 2021; accepted 7 June 2021; prepublished online on *Blood Advances* First Edition 7 September 2021; final version published online 24 November 2021.  
DOI 10.1182/bloodadvances.2021004750.

The full-text version of this article contains a data supplement.


© 2021 by The American Society of Hematology. Licensed under Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0), permitting only noncommercial, nonderivative use with attribution. All other rights reserved.

erythroid-specific factors such as GATA binding factor 1 (GATA1), nuclear factor, erythroid 2 (NFE2), and erythroid Krüppel-like factor (EKLF)<sup>4,5</sup> in the proerythroblast and basophilic stages of development in mice and humans, respectively,<sup>6</sup> and it subsequently governs the remaining steps of erythroid differentiation.<sup>7-9</sup> Notably, *ALAS2* induction occurs after intimate contacts between a red cell progenitor and central macrophage (CM) are established in an erythroblastic island.<sup>10-12</sup> The CM assists in erythroblast development by inhibiting apoptosis,<sup>13</sup> facilitating iron delivery,<sup>14-16</sup> and enhancing enucleation by pyrenocytolysis.<sup>12,17</sup>

The inability to produce sufficient numbers of normal erythrocytes results in anemia. Although anemia is the most prevalent of all blood disorders, it is generally underappreciated and undertreated despite its impact on quality of life and the economy.<sup>18</sup> Iron-deficiency anemia (IDA), followed in frequency by anemia of inflammation (AI), are major contributors to the morbidity and mortality of chronic illnesses.<sup>19</sup> IDA is characterized as a hypochromic, microcytic anemia, whereas AI is usually a normochromic, normocytic anemia.<sup>20</sup> Microcytosis and loss of *ALAS2* activity have been observed in the bone marrow of patients with rheumatoid arthritis (RA).<sup>21</sup> It has been postulated that systemic hypoferrremia common to AI is causative. However, ferrokinetic studies in patients with RA do not indicate reduced iron supply to bone marrow,<sup>22</sup> and *FECH* activity in patients with RA is normal.<sup>21</sup> This finding contrasts with what is found in IDA,<sup>23</sup> in mouse models of IDA in which *IRP2* is knocked out with decreased transferrin receptor expression,<sup>24</sup> and in low cellular iron conditions in which *FECH* activity is proposed to be limited by decreased iron-sulfur cofactor

assembly.<sup>25</sup> In samples of whole bone marrow from a patient with RA, *ALA* synthesis is reduced to a far greater magnitude than in purified erythroblasts,<sup>21</sup> suggesting that resident immunoreactive leukocytes contribute to the loss of *ALAS2* activity and corresponding anemia. Together, these observations strongly suggest that factors other than the availability of iron for erythropoiesis must be involved to limit heme synthesis in AI.

We hypothesize that alterations in the erythropoietic program during inflammation are mediated by the CM in erythroblastic islands. Lipopolysaccharide (LPS)-mediated inflammation in mice results in anemia via inhibition of erythropoiesis in bone marrow at an early stage, with an observed depletion of proerythroblasts and basophilic erythroblasts,<sup>26</sup> the stages during which *ALAS2* is first induced.<sup>6</sup> A variety of factors are produced and excreted by macrophages as part of their inflammatory response. In the last decade, multiple studies have revealed that macrophages synthesize itaconate (supplemental Figure 1A), an  $\alpha,\beta$ -unsaturated carboxylic acid and close analogue of succinate (methylenesuccinic acid) (supplemental Figure 1B), at the outset of an inflammatory response.<sup>27-32</sup> Itaconate is generated in millimolar quantities from the TCA cycle intermediate *cis*-aconitate via immune-response gene 1 (*IRG1*) protein, which can be induced by proinflammatory factors such as LPS and interferon- $\gamma$  in bone marrow-derived macrophages (BMDMs).<sup>29,31</sup> Importantly, itaconate is secreted by macrophages,<sup>28,30,32</sup> including BMDMs, and has recently been identified in local tissues during sepsis<sup>30</sup> and in the plasma of patients with RA.<sup>33</sup> Thus, it is believed that itaconate has characteristics that affect metabolism via autocrine, paracrine, and endocrine processes.<sup>34</sup>



**Figure 1. Heme biosynthesis and the erythroblastic island.** The terminal steps of erythroid development occur while erythroblasts are closely associated with a CM via  $\alpha\beta$  integrins, VCAM1, erythroblast macrophage protein (EMP), and ICAM4. Heme synthesis in this context is fully activated by the basophilic erythroblast (BasoEB) stage of development. CD163 and CD169 are macrophage-specific cell surface markers. Heme synthesis enzymes are in bold green text. CFU-E, colony forming unit–erythroid; CPOX, coproporphyrinogen III oxidase; HMBS, hydroxymethylbilane synthase; HSC, hematopoietic stem cell; OrthoEB, orthochromatic erythroblast; PBGS, porphobilinogen synthase; PolyEB, polychromatic erythroblast; PPOX, protoporphyrinogen oxidase; ProEB, proerythroblasts; RBC, red blood cell; SCoA, succinyl-CoA; UROD, URO decarboxylase; UROS, uroporphyrinogen III synthase.

Itaconate inhibits succinate dehydrogenase (SDH)<sup>31,32,35</sup> and has been associated with succinate-driven stabilization of hypoxia-inducible factor-1 $\alpha$  and proinflammatory interleukin 1 $\beta$  (IL1 $\beta$ ) production.<sup>36</sup> It also attenuates liver phosphofructokinase 2 (PFK2) activity in glycolysis<sup>37</sup> and further disrupts the TCA cycle by blocking substrate-level phosphorylation.<sup>38</sup> Older publications proposed that itaconate is converted to itaconyl-CoA (supplemental Figure 1C) by reverse succinyl CoA synthetase (SCS) activity with adenosine triphosphate (ATP) or guanosine triphosphate (GTP) supplementation,<sup>39,40</sup> although direct evidence for this theory was lacking. Itaconate has been shown to have antimicrobial activity by inhibition of the glyoxylate shunt<sup>41-43</sup> and as itaconyl-CoA by inactivation of vitamin B<sub>12</sub>.<sup>44,45</sup> B<sub>12</sub> can be scavenged by microbes in their host organisms,<sup>46</sup> and the suicide inhibition of the B<sub>12</sub>-dependent methylmalonic acid mutase (MUT) by itaconyl-CoA leads to the accumulation of toxic propionyl-CoA and may curtail cholesterol-dependent growth of *Mycobacterium tuberculosis*.<sup>45</sup>

Herein we report that the immunometabolite itaconate taken up by the developing erythron is converted to itaconyl-CoA and controls porphyrin synthesis at the point of ALAS2 catalysis. We also show global metabolic changes that occur in developing erythrocytes upon itaconate treatment suggest alterations in carbon flux for heme synthesis. Moreover, we provide evidence that itaconyl-CoA is produced by a

CoA transferase and not SCS in red cell progenitors. We believe this mechanism as a whole is crucial during acute and/or chronic conditions of anemia when itaconate has been shown to accumulate.


## Methods

### Reagents

All reagents were obtained from Sigma unless otherwise noted. <sup>13</sup>C<sub>5</sub>-itaconate was synthesized by the Metabolite Standards Synthesis Core at SRI International, arranged through the National Institutes of Health Common Funds Metabolomics initiative.<sup>47</sup> Details regarding the preparation of itaconyl-CoA are provided in the supplemental Methods.

### Cell culture

DS19 murine erythroleukemia (MEL) cells<sup>48,49</sup> were used as erythroblast models and propagated in Dulbecco's modified Eagle medium/F12 media (Corning) containing 25 mM glucose and 1 mM sodium pyruvate supplemented with 2 mM glutamine, 10% fetal bovine serum, and 1% penicillin-streptomycin under a humidified 5% carbon dioxide atmosphere. Details of the culture methods are presented in the supplemental Methods.



**Figure 2. Porphyrin and heme analysis of MEL cells treated with itaconate.** Heme (A), PPIX (B), and total carboxyl porphyrins (C,D) were determined by high-performance liquid chromatography for wild-type MEL cells differentiated in dimethyl sulfoxide media for 72 hours. (E) Relative fluorescence of cyclic porphyrins in media from MEL cell cultures overexpressing *Homo sapiens* ALAS2 (hsALAS2) over 72 hours without dimethyl sulfoxide induction. Itaconate concentrations were 2.5 mM in all experiments shown here. Bars represent means  $\pm$  1 standard deviation ( $n = 3-4$  biological replicates as indicated red dots). Multiple Student  $t$  tests combined with Bonferroni analysis produced  $P$  values  $<.01$  for all phosphate-buffered saline (PBS) vs itaconate comparisons shown. RFU, relative fluorescence units.

## Heme and porphyrin analysis

MEL cells treated with or without 2.5 mM unlabeled itaconate and 1.5% dimethyl sulfoxide were analyzed in quadruplicate ( $n = 4$  biological replicates) for heme, PPIX, and pathway intermediates 8-, 7-, 6-, 5-, and 4-carboxylporphyrins (for isomers I and III) by using high-performance liquid chromatography, as reported previously.<sup>50</sup> Details are presented in the supplemental Methods.

## Expression and purification of recombinant enzymes

The design and preparation of expression plasmids, recombinant protein expression in *Escherichia coli*, and purification are detailed in the supplemental Methods.

## Spectrophotometric assays and kinetics

ALAS activity assays were conducted either in coupled assays or with derivatized ALA detection via Ehrlich's modified reagent, with micro-scale modifications as detailed in the supplemental Methods. For inhibition studies, inhibitory constant and mode of inhibition were determined by titrating 10 to 200  $\mu\text{M}$  succinyl-CoA in the presence of 50 mM glycine substrate and 0, 200, and 1000  $\mu\text{M}$  itaconyl-CoA. Michaelis-Menten nonlinear regression analysis was subsequently conducted on data for 3 separate protein preparations ( $n = 3$  biological replicates) with Prism software (GraphPad 8.4).

Pig heart SCS-G (Sigma) and recombinant SCS-A were assayed for reverse activity (GTP or ATP consumption as appropriate) with succinate and/or itaconate as substrate(s) using a direct spectrophotometric assay for CoA ester bond formation at 225 to 235  $\text{nm}$ <sup>51</sup> as detailed in the supplemental Methods.


## High-performance liquid chromatography assays of CoA transferase activity

SUGCT (succinyl-CoA:glutarate-CoA transferase) activity was determined as reported in the literature,<sup>52</sup> with minor changes as detailed in the supplemental Methods.

## Results

### Itaconate decreases heme and porphyrin synthesis in erythroid precursors

Previously, it was shown that when medium from RAW264.7 macrophages treated with LPS was added to differentiating MEL cells, the level of heme production decreased.<sup>53</sup> We have verified these experimental results knowing that these cells produce and excrete itaconate under these conditions (supplemental Figure 2). Previously, we established that the inflammatory metabolite itaconate attenuates hemoglobin production in MEL cells undergoing erythroid differentiation and that this block is bypassed by providing the cells with ALA.<sup>54</sup> Here



**Figure 3. Liquid chromatography-mass spectrometry analysis of differentiating MEL cells treated with labeled itaconate.** MEL cultures were induced with dimethyl sulfoxide and treated with phosphate-buffered saline (PBS) or 1 mM labeled itaconate for 72 hours. Select metabolites targeted by liquid chromatography-mass spectrometry analysis are given on the y-axes. Analyses were conducted as described in the Methods section. Bars represent means  $\pm 1$  standard deviation ( $n = 4$  biological replicates). Multiple Student  $t$  tests combined with Bonferroni analysis produced  $P$  values  $< .01$  for all PBS vs  $^{13}\text{C}_5$ -itaconate for the metabolites (A)  $^{13}\text{C}_5$ -itaconate, (B)  $^{13}\text{C}_5$ -itaconyl-CoA, (D) succinyl-CoA, (E) propionyl-CoA, (F) propionyl-carnitine, and (G) inositol, comparisons except (C) succinate (0.024) and (H) AMP (0.034).

we show that heme, PPIX, and all carboxyl porphyrin intermediates are significantly decreased during MEL cell differentiation in the presence of itaconate (Figure 2A-D). These data indicate that inhibition of heme synthesis occurs upstream of uroporphyrinogen decarboxylase (Figure 1). To better target the point of inhibition, we examined the effects of itaconate on transgenic MEL cells overexpressing hsALAS2. Normally, these cells overproduce porphyrins because pathway regulation is bypassed by the expression of excess ALAS2. Treatment of these cells with itaconate reduces porphyrin-induced fluorescence (Figure 2E), suggesting that blockage of the pathway occurs before tetrapyrrole formation.

In agreement with our MEL cell data,<sup>54</sup> hemoglobin production in EPO-induced human TF1 cells is diminished with itaconate and can be rescued by ALA (supplemental Figure 3A). However, we also noted a significant increase in cell proliferation with itaconate. Unlike MEL cells<sup>55</sup> or erythroid-differentiated CD34<sup>+</sup> cells,<sup>56</sup> TF1 cells express the succinate cognate G protein-coupled receptor 91 (GPR91, or SUCNR1).<sup>57</sup> Because SUCNR1 is sensitive to itaconate<sup>58</sup> and known to stimulate ERK signaling, we added combinations of itaconate and the SUCNR1 antagonist compound 4c (gift of Xin Li) or MEK inhibitor U0126 to differentiating TF1 cultures to assess the impact of the SUCNR1-MAPK/ERK pathway on differentiation. The inhibitors independently rescued hemoglobin levels and proliferation rates measured in control cells, suggesting that in TF-1 cells, the observed decrease in hemoglobinization is at least partially due to nonspecific itaconate binding to SUCNR1 and the associated increase in TF1 cell proliferation via the SUCNR1-ERK signaling axis<sup>57</sup> (supplemental Figure 3B-C).

### Itaconate is imported by erythroid precursors and converted to itaconyl-CoA


Although it is known that itaconate is actively secreted by macrophages and that itaconate added to cultures of MEL cells results in decreased hemoglobinization, it was necessary to show that itaconate can be imported by developing erythroid cells. Mass spectrometry-based metabolomic tracing of dimethyl sulfoxide-induced MEL cells cultured with uniformly labeled (<sup>13</sup>C<sub>5</sub>)-itaconate exhibited the

intracellular accumulation of the same species (Figure 3A). In support of this observation, mitochondrial itaconate transporters, including SLC25A10 (succinate) and SLC25A1 (citrate), have been identified *in vitro*,<sup>59</sup> making movement of exogenously supplied itaconate into the mitochondrial compartment feasible. These transporters are expressed at significant levels in erythroid-differentiating MEL<sup>55</sup> and CD34<sup>+</sup> cells.<sup>56</sup> With respect to plasmalemmal counterpart(s), a cell surface itaconate transporter has not yet been identified in any mammalian cell type, although our data clearly show that a mechanism for itaconate uptake by cells must exist.<sup>60</sup>

Further tracing analysis of the differentiating MEL cell samples indicates that one metabolic fate of internalized <sup>13</sup>C<sub>5</sub>-itaconate is <sup>13</sup>C<sub>5</sub>-itaconyl-CoA (Figure 3B). This result was unsurprising based on 50-year-old data suggesting that mammalian mitochondrial SCS may use itaconate as an alternative substrate for succinate in ATP- or GTP-driven synthesis of itaconyl-CoA.<sup>39,40</sup>

### Itaconate affects flux through the erythroblast TCA cycle and glycolytic shunts

Liquid chromatography-mass spectrometry analysis of differentiating MEL cells treated with 1 mM <sup>13</sup>C<sub>5</sub>-itaconate or 2.5 mM unlabeled itaconate showed significant differences in TCA cycle metabolite levels from cells differentiated in the absence of itaconate. Specifically, succinate (Figure 3C; supplemental Figures 4, 5A, and 6) and succinyl-CoA (Figure 3D) were elevated in itaconate-treated cultures, although neither was labeled by <sup>13</sup>C<sub>5</sub>-itaconate. The finding of elevated succinate levels was not surprising given the established direct inhibitory action of itaconate on SDH in other cell types.<sup>31,32</sup> Fumarate increased as well with itaconate (supplemental Figure 4A) despite putative SDH inhibition. The  $\alpha$ -ketoglutarate derivative 2-hydroxyglutarate (supplemental Figures 4A, 5A, and 6) also accumulated, and increased glutamic acid was found in the media (supplemental Figure 6B). Both these observations may be due to product inhibition of  $\alpha$ -ketoglutarate dehydrogenase caused by the elevated levels of succinyl-CoA.<sup>61</sup>



**Figure 4. Inhibition of the reverse SCS reaction by itaconate.** Succinate but not itaconate at 10 mM concentrations is metabolized by pig heart SCS-GTP (SCS-G) (A) and recombinant SCS-ATP (SCS-A) (B). (C) Itaconate inhibits SCS-A at the indicated ratios using 50 mM succinate in all reactions. Summary bar plots represent mean rates of acyl-CoA formation at 225 to 235 nm. The corresponding UV scans for panels A and B are provided in supplemental Figure 7. Bars represent means and error bars = 1 standard deviation (n = 2-3 biological replicates). Panel A: Student *t* test *P* value = .0001. Panel C: one-way analysis of variance *P* value = .0022 with Bonferroni post hoc test *P* values = .026, .0022, and .15 (not significant [ns]) for control vs 1:200, control vs 1:50, and 1:200 vs 1:50, respectively.



Our results also suggest that glycolysis is disrupted by itaconate during terminal erythropoiesis. Intermediates of glycolytic shunts such as the pentose phosphate and serine-glycine synthesis pathways increased compared with controls, including AMP (Figure 3E), adenosine and thymine (supplemental Figure 4A), serine (supplemental Figures 4, 5, and 6), and glycine (supplemental Figure 6A). Unexpectedly, levels of glutathione in the culture media were also elevated (supplemental Figures 4B and 6B), as itaconate has been reported to directly alkylate glutathione in LPS-activated macrophages.<sup>62</sup> Myoinositol (supplemental Figures 4A, 5A, and 6A) and inositol (Figure 3F), products of glucose-6-phosphate metabolism, increased with itaconate as well. Allosteric inhibition of the glycolysis-activating PFK2 by itaconate, a phosphoenolpyruvate analogue, has been previously described in the liver<sup>37</sup> and could explain these observations in our differentiating erythroid progenitors.

### Itaconate impairs metabolism of amino acids and fatty acid oxidation compounds


It is well established that aliphatic amino acids and odd-chain fatty acids are ultimately catabolized to propionyl-CoA and then methylmalonyl-CoA by propionyl-CoA carboxylase and B<sub>12</sub>-dependent methylmalonyl-CoA mutase (MUT), respectively. Further metabolomic data gathered here indicate that flux through these pathways is attenuated in differentiating erythroblast cultures treated with itaconate, likely via itaconyl-CoA and its ability to inactivate B<sub>12</sub> and MUT activity as previously discussed in adipocytes<sup>44</sup> and macrophages.<sup>45</sup> Namely, we found increased levels of propionyl-CoA (Figure 3G), propionyl-carnitine (Figure 3H), and methylmalonic acid (supplemental

Figures 4, 5, and 6) with itaconate treatment. Upstream of propionyl-CoA, we measured elevations in the odd-chain fatty acids heptadecanoic acid (supplemental Figures 4A and 6A) and nonanoic acid (supplemental Figure 6B), as well as the aliphatic amino acids threonine, valine, methionine, and isoleucine (supplemental Figure 6). Notably, itaconyl-CoA accumulations may also explain increased pyruvate and glyoxylate levels (supplemental Figures 4B, 5B, and 6B). Itaconyl-CoA is an alternative substrate of methylglutaconase and is converted by this enzyme to citramalyl-CoA.<sup>40,44</sup> It is therefore reasonable to believe that citramalyl-CoA lyase (CLYBL) catalyzes the cleavage of citramalyl-CoA to pyruvate and acetyl-CoA<sup>40</sup> in lieu of metabolizing glyoxylate.<sup>63</sup>

### Itaconyl-CoA is produced from itaconate via SUGCT but not SCS

Indirect evidence from approximately one-half century ago has been the basis for the proposal that itaconate is converted to itaconyl-CoA by SCS,<sup>39,40</sup> and in the absence of definitive data, it has been assumed by others to occur in this fashion.<sup>38,64</sup> However, we observed that neither ATP- nor GTP-specific SCS is able to use itaconate under the same conditions as succinate in vitro (Figure 4A-B; supplemental Figure 7A-B). Furthermore, we observed that SCS activity is inhibited by itaconate in a dose-dependent manner (Figure 4C).

In an effort to identify a mammalian enzyme for itaconyl-CoA production, we first turned to bacterial enzymes capable of itaconyl-CoA synthesis. It is known that *Pseudomonas aeruginosa* itaconyl-CoA transferase (Palct) is an enzyme capable of forming itaconyl-CoA




**Figure 5. SUGCT converts itaconate to itaconyl-CoA.** (A) Human SUGCT is homologous to the *P. aeruginosa* itaconyl-CoA transferase (Palct). Black and gray boxes highlight identical residues and conservative substitutions, respectively. (B) Discontinuous high-performance liquid chromatography (HPLC) assays showed that SUGCT generates itaconyl-CoA from succinyl-CoA and itaconate in vitro. (C) Immunoblotting of MEL mitochondria from cells treated with and without 2.5 mM itaconate and dimethyl sulfoxide (DMSO) for 72 hours. MEL mitochondrial lysates were probed with anti-ALAS2, anti-FECH, and anti-SUGCT primary antibodies. HPLC assays were performed on a C18 reverse-phase column with UV detection (mAU = milli-absorbance units at 254 nm). Optimal autoexposure within the linear dynamic range was performed for each immunoblot on the ChemiDoc imaging system (Bio-Rad). The HPLC chromatogram (B) and western blot (C) shown are representative of n = 2 biological replicates.

from succinyl-CoA and itaconate without added nucleotides.<sup>64</sup> Using a bioinformatics approach, we found that mammalian SUGCT (encoded by the *C7ORF10* gene) is a homologue of the *P aeruginosa* Palct (Figure 5A). Others have reported SUGCT activity with the canonical substrate glutaryl-CoA.<sup>52</sup> We cloned, expressed, and purified SUGCT and determined that it effectively catalyzes the production of itaconyl-CoA using succinyl-CoA (Figure 5B). In agreement with transcript data in MEL cells,<sup>55</sup> immunoblotting showed that SUGCT protein is present in mitochondria isolated from induced


and uninduced MEL cells (Figure 5C). Thus, SUGCT is likely responsible for itaconyl-CoA synthesis in developing erythrocytes.

### Itaconyl-CoA inhibits, but is not a substrate of, ALAS2

Itaconyl-CoA (supplemental Figure 1C) is a close analogue of succinyl-CoA (supplemental Figure 1D) and may affect ALAS2 enzyme activity. Although itaconate does not affect ALAS2 activity (supplemental Figure 8A), discontinuous kinetic assays revealed



**Figure 6. ALAS2 inhibition by itaconyl-CoA.** Kinetic progress curves (A) and Lineweaver-Burk plot (B) indicating competitive inhibition of ALAS2 with an inhibitory constant =  $100 \pm 20 \mu\text{M}$  (mean  $\pm 1$  standard deviation) for itaconyl-CoA. Michaelis constant =  $10 \pm 2 \mu\text{M}$  (mean  $\pm 1$  standard deviation) for succinyl-CoA. Discontinuous colorimetric assays were conducted for  $n = 3$  biological replicates (separate protein preparations).



**Figure 7. Model of the erythroblastic island during an inflammatory response.** Itaconate is produced via IRG1 in immunoactivated blood island macrophages and excreted to the surrounding milieu. Subsequent import and metabolism of itaconate to itaconyl-CoA by proximal red cell precursors result in inhibition of ALA synthesis and cellular hemoglobinization at the point of ALAS2 catalysis. Interleukin 6 (IL6), tumor necrosis factor- $\alpha$  (TNF $\alpha$ ), and interleukin 1 $\beta$  (IL1 $\beta$ ) are proinflammatory cytokines. GSH, glutathione; IFNGR, interferon- $\gamma$  (IFN $\gamma$ ) receptor; TLR4, Toll-like receptor 4.

itaconyl-CoA to be a competitive inhibitor of recombinant ALAS2 with an inhibitory constant of  $100 \pm 20 \mu\text{M}$  (Figure 6). We measured the Michaelis constant of succinyl-CoA to be  $10 \pm 2 \mu\text{M}$  for the uninhibited enzyme, which agrees with the literature for the mammalian enzyme.<sup>65-69</sup> Thus, itaconyl-CoA could be partially responsible for the observed increase in succinyl-CoA in induced MEL cells (Figure 3D). Continuous coupled ALAS assays were also performed because ALA derivatives synthesized from other acyl-CoA substrates have shown unpredictable stability in the  $\alpha$ -aminoketone formed via reaction with acetylacetone in the discontinuous assay protocol.<sup>66</sup> The results of the coupled assays verify that itaconyl-CoA is not an alternative ALAS2 substrate even at millimolar concentrations (supplemental Figure 8B). This finding explains the alterations in heme precursor levels (Figure 2) and the global alterations in metabolites.

## Discussion

Itaconate is an antimicrobial compound whose impact on mammalian liver metabolism was first explored in the 1940s,<sup>39,70,71</sup> and we provide evidence here that itaconate has a significant impact on the erythroid compartment (Figure 7; supplemental Figure 9). Immunoactivated macrophages have been identified as the source of itaconate via IRG1 catalysis.<sup>29-32</sup> Importantly, measurable quantities of itaconate have been found in bronchial lavages of patients with severe lung infections<sup>30</sup> and in blood plasma of patients with RA.<sup>33</sup> Experiments examining the impact of RA on the erythroid compartment have shown diminished ALAS2 activity in microcytic red cell precursors, most significantly in the context of whole bone marrow samples containing immunoactivated leukocytes such as BMDMs.<sup>21</sup> Based on the results of this study, we propose that itaconate negatively alters heme biosynthesis and erythroid differentiation at the point of ALAS2 catalysis in the context of an inflammatory erythroblastic island. Specifically, we have determined that heme, PPIX, and total carboxyl porphyrin levels are decreased in MEL cells differentiated in the presence of itaconate or media from immunoactivated, itaconate-generating macrophages. Addition of ALA to itaconate-treated cultures rescues hemoglobinization, as described previously.<sup>54</sup> <sup>13</sup>C tracing data show that itaconate is imported by differentiating MEL cells and is subsequently metabolized to itaconyl-CoA. Levels of succinyl-CoA are also increased in these cells with itaconate supplementation, suggesting that ALAS2 is directly affected. Indeed, our kinetic assays showed that itaconyl-CoA is a competitive inhibitor of ALAS2 without acting as an alternative substrate.

The widely accepted explanation for decreased ALAS2 activity in inflammatory states such as AI is that systemic iron is sequestered via hepcidin<sup>72</sup> and could therefore downregulate iron-dependent ALAS2 synthesis. Mature ALAS2 messenger RNA contains a 5' iron-responsive element that is lacking in the housekeeping ALAS1 enzyme,<sup>73</sup> and this element is bound by apo iron-regulatory proteins IRP1 and IPR2 when erythroid iron levels are low, effectively blocking ALAS2 translation.<sup>24,74</sup> The impact of IRP1 on ALAS2 protein levels in particular has been described in cases of sideroblastic anemia caused by loss of glutaredoxin 5<sup>75</sup> and mitoferrin-1-deficient porphyria.<sup>76</sup> However, others have reported that erythroblast iron delivery during terminal differentiation bypasses the cytosolic iron pool for IRP2, and IRP numbers in general appear to be overwhelmed late in erythroid differentiation when heme synthesis is massively upregulated.<sup>77</sup> A more recent study has verified low levels of translated *Irp1* and *Irp2* relative to *Alas2* at the basophilic erythroblast stage in mice and has

provided evidence that heme-regulated eukaryotic initiation factor 2  $\alpha$  (eIF2 $\alpha$ ) kinase is the predominant regulator of hemoglobin synthesis in iron-replete and iron-deficient conditions in these cells.<sup>78</sup> Perhaps most significantly, ferrokinetic data from RA patient samples do not indicate reduced iron supply to the marrow,<sup>22</sup> and iron-dependent FECH activity is unaffected.<sup>21</sup> Thus, we postulate that itaconate is a significant factor in AI by downregulating ALAS2 activity and porphyrin synthesis in the bone marrow during an inflammatory response.

Additional data also support many of the previously substantiated effects of itaconate on glycolysis,<sup>37</sup> B<sub>12</sub> metabolism,<sup>44,45</sup> and the TCA cycle<sup>29,31,32</sup> in other tissues (supplemental Figure 9). For example, we see increases in compounds such as inositol as well as products of the pentose phosphate and serine-glycine synthesis pathways, possibly due to allosteric inhibition of PFK2.<sup>37</sup> Elevated levels of aliphatic amino acids (eg, valine), odd-chain fatty acids (eg, heptadecanoic acid), and propionyl-CoA metabolites suggest that itaconyl-CoA compromises B<sub>12</sub> constitution in erythroblasts and irreversibly inhibits the MUT enzyme as seen in adipocytes and immunoactivated macrophages.<sup>44,45</sup> In addition to succinyl-CoA, elevations in succinate and fumarate were observed with itaconate treatment. It is tempting to postulate that succinate accumulation is caused by itaconate-driven inhibition of SDH and accounts for succinyl-CoA production via the reverse SCS reaction. However, this does not justify the increase in fumarate. We also showed in vitro that itaconate inhibits reverse SCS activity and is not converted to itaconyl-CoA by SCS as proposed by others.<sup>39,40</sup> Our results fit with those of a more recent study showing that itaconate impairs substrate-level phosphorylation in LPS-stimulated macrophages in respiratory distress and behaves similarly to a specific SCS inhibitor in these cells.<sup>38</sup>

Bioinformatics and immunoblotting data support the hypothesis that exogenous itaconate is derivatized to itaconyl-CoA by SUGCT in red cell progenitors. In addition, succinate contributes little to the supply of heme carbon during erythroid differentiation, and succinyl-CoA used for heme synthesis is supplied primarily through a separate pool of  $\alpha$ -ketoglutarate dehydrogenase acting outside of the TCA cycle.<sup>54</sup> ALAS2 inhibition thus provides a valid explanation for both increased succinyl-CoA and decreased porphyrin synthesis when erythroid precursors are exposed to itaconate. The global metabolic changes that occurred from treatment of itaconate to developing erythroid cells show a direct effect of itaconate on multiple metabolic processes, independent of additional cytokines and other immunoregulated molecules. Although some of these metabolites are clearly upstream and linked to heme synthesis in the developing erythron, others may be downstream of the process. Follow-up experiments to determine the metabolic connections and temporal response will be necessary to determine the impact of itaconate in the erythroblastic island.

## Acknowledgments

The authors thank D. Bishop (Mount Sinai), A. Cantor (Boston Children's Hospital), and X. Li (New York University) for providing some reagents used in this study.

This work was supported by National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases grants DK11653 and DK110858 via Pilot and Feasibility Grants from the University of Utah Center for Iron and Heme Disorders (A.E.M.) and U54DK110858 and U24DK126127 (J.D.P.). Mass



spectrometry equipment was obtained through 1S10OD016232-01 and 1S10OD021505-01 (J.E.C.).

## Authorship

Contribution: J.R.M. designed experiments, analyzed data, and wrote the paper; J.E.C. and H.A.B. performed research and analyzed data; J.E.C., A.E.M., and J.D.P. analyzed data and edited the paper; and H.A.D. designed experiments, analyzed data, and wrote the paper.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

ORCID profiles: J.E.C., 0000-0002-5977-2350; A.E.M., 0000-0001-9402-3593; J.D.P., 0000-0003-1567-1678.

Correspondence: Harry A. Dailey, Department of Microbiology, 230 Coverdell Center, University of Georgia, Athens, GA 30602; e-mail: hdailey@uga.edu.

## References

1. Harris JW. X-linked, pyridoxine-responsive sideroblastic anemia. *N Engl J Med.* 1994;330(10):709-711.
2. Dailey HA, Meissner PN. Erythroid heme biosynthesis and its disorders. *Cold Spring Harb Perspect Med.* 2013;3(4):a011676.
3. Phillips JD. Heme biosynthesis and the porphyrias. *Mol Genet Metab.* 2019;128(3):164-177.
4. Ferreira GC, Andrew TL, Karr SW, Dailey HA. Organization of the terminal two enzymes of the heme biosynthetic pathway. Orientation of protoporphyrinogen oxidase and evidence for a membrane complex. *J Biol Chem.* 1988;263(8):3835-3839.
5. Kramer MF, Gunaratne P, Ferreira GC. Transcriptional regulation of the murine erythroid-specific 5-aminolevulinate synthase gene. *Gene.* 2000;247(1-2):153-166.
6. An X, Schulz VP, Li J, et al. Global transcriptome analyses of human and murine terminal erythroid differentiation. *Blood.* 2014;123(22):3466-3477.
7. Lake-Bullock H, Dailey HA. Biphasic ordered induction of heme synthesis in differentiating murine erythroleukemia cells: role of erythroid 5-aminolevulinate synthase. *Mol Cell Biol.* 1993;13(11):7122-7132.
8. Nakajima O, Takahashi S, Harigae H, et al. Heme deficiency in erythroid lineage causes differentiation arrest and cytoplasmic iron overload. *EMBO J.* 1999;18(22):6282-6289.
9. Brownlie A, Donovan A, Pratt SJ, et al. Positional cloning of the zebrafish sauternes gene: a model for congenital sideroblastic anaemia. *Nat Genet.* 1998;20(3):244-250.
10. Chasis JA, Mohandas N. Erythroblastic islands: niches for erythropoiesis. *Blood.* 2008;112(3):470-478.
11. Astwood E, Vora A. Erythroblastic islands. *Blood.* 2011;117(1):10.
12. Yeo JH, Lam YW, Fraser ST. Cellular dynamics of mammalian red blood cell production in the erythroblastic island niche. *Biophys Rev.* 2019;11(6):873-894.
13. Rhodes MM, Kopsombut P, Bondurant MC, Price JO, Koury MJ. Adherence to macrophages in erythroblastic islands enhances erythroblast proliferation and increases erythrocyte production by a different mechanism than erythropoietin. *Blood.* 2008;111(3):1700-1708.
14. Iacopetta BJ, Morgan EH. The kinetics of transferrin endocytosis and iron uptake from transferrin in rabbit reticulocytes. *J Biol Chem.* 1983;258(15):9108-9115.
15. Sheftel AD, Zhang AS, Brown C, Shirihai OS, Ponka P. Direct interorganellar transfer of iron from endosome to mitochondrion. *Blood.* 2007;110(1):125-132.
16. Kafina MD, Paw BH. Intracellular iron and heme trafficking and metabolism in developing erythroblasts. *Metallomics.* 2017;9(9):1193-1203.
17. Yoshida H, Kawane K, Koike M, Mori Y, Uchiyama Y, Nagata S. Phosphatidylserine-dependent engulfment by macrophages of nuclei from erythroid precursor cells. *Nature.* 2005;437(7059):754-758.
18. Smith RE Jr. The clinical and economic burden of anemia. *Am J Manag Care.* 2010;16(Suppl Issues):S59-S66.
19. Madu AJ, Ughasoro MD. Anaemia of chronic disease: an in-depth review. *Med Princ Pract.* 2017;26(1):1-9.
20. Davis SL, Littlewood TJ. The investigation and treatment of secondary anaemia. *Blood Rev.* 2012;26(2):65-71.
21. Houston T, Moore M, Porter D, Sturrock R, Fitzsimons E. Abnormal haem biosynthesis in the chronic anaemia of rheumatoid arthritis. *Ann Rheum Dis.* 1994;53(3):167-170.
22. Cavill I, Bentley DP. Erythropoiesis in the anaemia of rheumatoid arthritis. *Br J Haematol.* 1982;50(4):583-590.
23. Sharma DC, Mathur R. Some biochemical changes in heme synthesis in iron deficiency. *Indian J Physiol Pharmacol.* 2000;44(4):491-494.
24. Cooperman SS, Meyron-Holtz EG, Olivier-Wilson H, Ghosh MC, McConnell JP, Rouault TA. Microcytic anemia, erythropoietic protoporphyria, and neurodegeneration in mice with targeted deletion of iron-regulatory protein 2. *Blood.* 2005;106(3):1084-1091.
25. Crooks DR, Ghosh MC, Haller RG, Tong WH, Rouault TA. Posttranslational stability of the heme biosynthetic enzyme ferrochelatase is dependent on iron availability and intact iron-sulfur cluster assembly machinery. *Blood.* 2010;115(4):860-869.
26. Brendt P, Rehfeld I, Kamphausen A, Kreissig C, Peters J. Lipopolysaccharide interference in erythropoiesis in mice. *Anaesthesia.* 2012;67(5):493-500.
27. Strelko CL, Lu W, Dufort FJ, et al. Itaconic acid is a mammalian metabolite induced during macrophage activation. *J Am Chem Soc.* 2011;133(41):16386-16389.
28. Sugimoto M, Sakagami H, Yokote Y, et al. Non-targeted metabolite profiling in activated macrophage secretion. *Metabolomics.* 2012;8(4):624-633.

29. Michelucci A, Cordes T, Ghelfi J, et al. Immune-responsive gene 1 protein links metabolism to immunity by catalyzing itaconic acid production. *Proc Natl Acad Sci USA*. 2013;110(19):7820-7825.
30. Meiser J, Kraemer L, Jaeger C, et al. Itaconic acid indicates cellular but not systemic immune system activation. *Oncotarget*. 2018;9(63):32098-32107.
31. Cordes T, Wallace M, Michelucci A, et al. Immunoresponsive gene 1 and itaconate inhibit succinate dehydrogenase to modulate intracellular succinate levels. *J Biol Chem*. 2016;291(27):14274-14284.
32. Lampropoulou V, Sergushichev A, Bambouskova M, et al. Itaconate links inhibition of succinate dehydrogenase with macrophage metabolic remodeling and regulation of inflammation. *Cell Metab*. 2016;24(1):158-166.
33. Daly R, Blackburn G, Best C, et al. Changes in plasma itaconate elevation in early rheumatoid arthritis patients elucidates disease activity associated macrophage activation. *Metabolites*. 2020;10(6):E241.
34. Zaslona Z, O'Neill LAJ. Cytokine-like roles for metabolites in immunity. *Mol Cell*. 2020;78(5):814-823.
35. Ackermann WW, Potter VR. Enzyme inhibition in relation to chemotherapy. *Proc Soc Exp Biol Med*. 1949;72(1):1-9.
36. Tannahill GM, Curtis AM, Adamik J, et al. Succinate is an inflammatory signal that induces IL-1 $\beta$  through HIF-1 $\alpha$ . *Nature*. 2013;496(7444):238-242.
37. Sakai A, Kusumoto A, Kiso Y, Furuya E. Itaconate reduces visceral fat by inhibiting fructose 2,6-bisphosphate synthesis in rat liver. *Nutrition*. 2004;20(11-12):997-1002.
38. Németh B, Doczi J, Csete D, et al. Abolition of mitochondrial substrate-level phosphorylation by itaconic acid produced by LPS-induced Irg1 expression in cells of murine macrophage lineage. *FASEB J*. 2016;30(1):286-300.
39. Adler J, Wang SF, Lardy HA. The metabolism of itaconic acid by liver mitochondria. *J Biol Chem*. 1957;229(2):865-879.
40. Wang SF, Adler J, Lardy HA. The pathway of itaconate metabolism by liver mitochondria. *J Biol Chem*. 1961;236(1):26-30.
41. McFadden BA, Purohit S. Itaconate, an isocitrate lyase-directed inhibitor in *Pseudomonas indigofera*. *J Bacteriol*. 1977;131(1):136-144.
42. Patel TR, McFadden BA. *Caenorhabditis elegans* and *Ascaris suum*: inhibition of isocitrate lyase by itaconate. *Exp Parasitol*. 1978;44(2):262-268.
43. Williams JO, Roche TE, McFadden BA. Mechanism of action of isocitrate lyase from *Pseudomonas indigofera*. *Biochemistry*. 1971;10(8):1384-1390.
44. Shen H, Campanello GC, Flicker D, et al. The human knockout gene CLYBL connects itaconate to vitamin B<sub>12</sub>. *Cell*. 2017;171(4):771-782.e11.
45. Ruetz M, Campanello GC, Purchal M, et al. Itaconyl-CoA forms a stable biradical in methylmalonyl-CoA mutase and derails its activity and repair. *Science*. 2019;366(6465):589-593.
46. Gopinath K, Venclovas C, Ioerger TR, et al. A vitamin B<sub>12</sub> transporter in *Mycobacterium tuberculosis*. *Open Biol*. 2013;3(2):120175.
47. Sud M, Fahy E, Cotter D, et al. Metabolomics Workbench: an international repository for metabolomics data and metadata, metabolite standards, protocols, tutorials and training, and analysis tools. *Nucleic Acids Res*. 2016;44(D1):D463-D470.
48. Ohta Y, Tanaka M, Terada M, et al. Erythroid cell differentiation: murine erythroleukemia cell variant with unique pattern of induction by polar compounds. *Proc Natl Acad Sci USA*. 1976;73(4):1232-1236.
49. Singer D, Cooper M, Maniatis GM, Marks PA, Rifkin RA. Erythropoietic differentiation in colonies of cells transformed by Friend virus. *Proc Natl Acad Sci USA*. 1974;71(7):2668-2670.
50. Yien YY, Robledo RF, Schultz IJ, et al. TMEM14C is required for erythroid mitochondrial heme metabolism. *J Clin Invest*. 2014;124(10):4294-4304.
51. Cha S, Parks RE Jr. Succinic thiokinase. I. Purification of the enzyme from pig heart. *J Biol Chem*. 1964;239(6):1961-1967.
52. Marlaire S, Van Schaftingen E, Veiga-da-Cunha M. C7orf10 encodes succinate-hydroxymethylglutarate CoA-transferase, the enzyme that converts glutarate to glutaryl-CoA. *J Inher Metab Dis*. 2014;37(1):13-19.
53. Sassa S, Wolpe S, Cerami A. Inhibition of erythroid differentiation of mouse erythroleukemia cells by a macrophage product(s). *Blood Cells*. 1987;13(1-2):161-169.
54. Burch JS, Marcero JR, Maschek JA, et al. Glutamine via  $\alpha$ -ketoglutarate dehydrogenase provides succinyl-CoA for heme synthesis during erythropoiesis. *Blood*. 2018;132(10):987-998.
55. Zhen R, Moo C, Zhao Z, et al. Wdr26 regulates nuclear condensation in developing erythroblasts. *Blood*. 2020;135(3):208-219.
56. Gautier EF, Ducamp S, Leduc M, et al. Comprehensive proteomic analysis of human erythropoiesis. *Cell Rep*. 2016;16(5):1470-1484.
57. Hakak Y, Lehmann-Bruinsma K, Phillips S, et al. The role of the GPR91 ligand succinate in hematopoiesis. *J Leukoc Biol*. 2009;85(5):837-843.
58. Trauelsen M, Rexen Ulven E, Hjorth SA, et al. Receptor structure-based discovery of non-metabolite agonists for the succinate receptor GPR91. *Mol Metab*. 2017;6(12):1585-1596.
59. Mills EL, Ryan DG, Prag HA, et al. Itaconate is an anti-inflammatory metabolite that activates Nrf2 via alkylation of KEAP1. *Nature*. 2018;556(7699):113-117.
60. O'Neill LAJ, Artyomov MN. Itaconate: the poster child of metabolic reprogramming in macrophage function. *Nat Rev Immunol*. 2019;19(5):273-281.
61. Smith CM, Bryla J, Williamson JR. Regulation of mitochondrial  $\alpha$ -ketoglutarate metabolism by product inhibition at  $\alpha$ -ketoglutarate dehydrogenase. *J Biol Chem*. 1974;249(5):1497-1505.
62. Bambouskova M, Gorvel L, Lampropoulou V, et al. Electrophilic properties of itaconate and derivatives regulate the I $\kappa$ B $\zeta$ -ATF3 inflammatory axis. *Nature*. 2018;556(7702):501-504.

63. Strittmatter L, Li Y, Nakatsuka NJ, Calvo SE, Grabarek Z, Mootha VK. CLYBL is a polymorphic human enzyme with malate synthase and  $\beta$ -methylmalate synthase activity. *Hum Mol Genet.* 2014;23(9):2313-2323.
64. Sasikaran J, Ziemski M, Zadora PK, Fleig A, Berg IA. Bacterial itaconate degradation promotes pathogenicity. *Nat Chem Biol.* 2014;10(5):371-377.
65. Hunter GA, Ferreira GC. A continuous spectrophotometric assay for 5-aminolevulinate synthase that utilizes substrate cycling. *Anal Biochem.* 1995;226(2):221-224.
66. Shoolingin-Jordan PM, LeLean JE, Lloyd AJ. Continuous coupled assay for 5-aminolevulinate synthase. *Methods Enzymol.* 1997;281:309-316.
67. Bishop DF, Tchaikovskii V, Nazarenko I, Desnick RJ. Molecular expression and characterization of erythroid-specific 5-aminolevulinate synthase gain-of-function mutations causing X-linked protoporphyria. *Mol Med.* 2013;19(1):18-25.
68. Bishop DF, Tchaikovskii V, Hoffbrand AV, Fraser ME, Margolis S. X-linked sideroblastic anemia due to carboxyl-terminal ALAS2 mutations that cause loss of binding to the  $\beta$ -subunit of succinyl-CoA synthetase (SUCLA2). *J Biol Chem.* 2012;287(34):28943-28955.
69. Bailey HJ, Bezerra GA, Marcero JR, et al. Human aminolevulinate synthase structure reveals a eukaryotic-specific autoinhibitory loop regulating substrate binding and product release. *Nat Commun.* 2020;11(1):2813.
70. Booth AN, Taylor J, Wilson RH, Deeds F. The inhibitory effects of itaconic acid in vitro and in vivo. *J Biol Chem.* 1952;195(2):697-702.
71. Finkelstein M, Gold H, Paterno CA. Pharmacology of itaconic acid and its sodium, magnesium, and calcium salts. *J Am Pharm Assoc Am Pharm Assoc.* 1947;36(6):173-179.
72. Ganz T, Nemeth E. Hepcidin and iron homeostasis. *Biochim Biophys Acta.* 2012;1823(9):1434-1443.
73. Cox TC, Bawden MJ, Martin A, May BK. Human erythroid 5-aminolevulinate synthase: promoter analysis and identification of an iron-responsive element in the mRNA. *EMBO J.* 1991;10(7):1891-1902.
74. Melefors O, Goossen B, Johansson HE, Stripecke R, Gray NK, Hentze MW. Translational control of 5-aminolevulinate synthase mRNA by iron-responsive elements in erythroid cells. *J Biol Chem.* 1993;268(8):5974-5978.
75. Ye H, Jeong SY, Ghosh MC, et al. Glutaredoxin 5 deficiency causes sideroblastic anemia by specifically impairing heme biosynthesis and depleting cytosolic iron in human erythroblasts. *J Clin Invest.* 2010;120(5):1749-1761.
76. Chung J, Anderson SA, Gwynn B, et al. Iron regulatory protein-1 protects against mitoferrin-1-deficient porphyria. *J Biol Chem.* 2014;289(11):7835-7843.
77. Schranzhofer M, Schiffrer M, Cabrera JA, et al. Remodeling the regulation of iron metabolism during erythroid differentiation to ensure efficient heme biosynthesis. *Blood.* 2006;107(10):4159-4167.
78. Zhang S, Macias-Garcia A, Ulirsch JC, et al. HRI coordinates translation necessary for protein homeostasis and mitochondrial function in erythropoiesis. *eLife.* 2019;8:8.