



Article

Anti-Apoptotic and Antioxidant Activities of the Mitochondrial Estrogen Receptor Beta in N2A Neuroblastoma Cells

Ioannis Tsialtas , Achilleas Georgantopoulos, Maria E. Karipidou , Foteini D. Kalousi, Aikaterini G. Karra, Demetrios D. Leonidas and Anna-Maria G. Psarra *

Department of Biochemistry and Biotechnology, University of Thessaly, 81500 Larissa, Greece; tsialtasj@gmail.com (I.T.); ageorgant@bio.uth.gr (A.G.); mkaripidou@uth.gr (M.E.K.); fokalous@bio.uth.gr (F.D.K.); aikaterini.g.karra@gmail.com (A.G.K.); ddleonidas@bio.uth.gr (D.D.L.)

* Correspondence: ampsarra@bio.uth.gr; Tel.: +30-2410-565221

Abstract: Estrogens are steroid hormones that play a crucial role in the regulation of the reproductive and non-reproductive system physiology. Among non-reproductive systems, the nervous system is mainly affected by estrogens due to their antioxidant, anti-apoptotic, and anti-inflammatory activities, which are mediated by membranous and nuclear estrogen receptors, and also by non-estrogen receptor-associated estrogen actions. Neuronal viability and functionality are also associated with the maintenance of mitochondrial functions. Recently, the localization of estrogen receptors, especially estrogen receptor beta, in the mitochondria of many types of neuronal cells is documented, indicating the direct involvement of the mitochondrial estrogen receptor beta (mtER β) in the maintenance of neuronal physiology. In this study, cell lines of N2A cells stably overexpressing a mitochondrial-targeted estrogen receptor beta were generated and further analyzed to study the direct involvement of mtER β in estrogen neuroprotective antioxidant and anti-apoptotic actions. Results from this study revealed that the presence of estrogen receptor beta in mitochondria render N2A cells more resistant to staurosporine- and H₂O₂-induced apoptotic stimuli, as indicated by the reduced activation of caspase-9 and -3, the increased cell viability, the increased ATP production, and the increased resistance to mitochondrial impairment in the presence or absence of 17- β estradiol (E2). Thus, the direct involvement of mtER β in antioxidant and anti-apoptotic activities is documented, rendering mtER β a promising therapeutic target for mitochondrial dysfunction-associated degenerative diseases.

Keywords: estrogen receptor beta; mitochondria; apoptosis; oxidative stress; neuroprotection



Citation: Tsialtas, I.; Georgantopoulos, A.; Karipidou, M.E.; Kalousi, F.D.; Karra, A.G.; Leonidas, D.D.; Psarra, A.-M.G. Anti-Apoptotic and Antioxidant Activities of the Mitochondrial Estrogen Receptor Beta in N2A Neuroblastoma Cells. *Int. J. Mol. Sci.* **2021**, *22*, 7620. <https://doi.org/10.3390/ijms22147620>

Academic Editor: Tuuli Käämbre

Received: 7 June 2021

Accepted: 13 July 2021

Published: 16 July 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Estrogens play a central role in the regulation of the biochemical processes related to female reproduction [1] and non-reproductive tissue physiology [2], but they also act on male reproductive tissue [3]. Thus, a wide variety of biological actions, including differentiation, cell proliferation, apoptosis, antioxidant defense, and inflammation, are regulated by the main estrogenic hormone 17- β estradiol (E2) [4]. The classical mechanism of estrogen actions is mainly mediated by the nuclear estrogen receptors (ERs), estrogen receptor alpha (ER α) and estrogen receptor beta (ER β), which function as ligand-dependent transcription factors, regulating the transcription of target genes containing the consensus estrogen response element (ERE) in their promoter regions. Estrogen binding to ERs triggers their nuclear translocation, homo- or heterodimerization, and binding to specific EREs in the nuclear DNA. ERs can also be found in the nucleus independently of the presence of the hormone [5,6]. Estrogen-activated nuclear ERs can also regulate non-ERE dependent transcription via interactions with other transcription factors, such as AP-1, affecting positively or negatively the expression of their target genes. In addition to the classical nuclear actions of estrogen, rapid membrane-initiated actions could also take place, which are mediated via plasma membrane-associated mER α , mER β and G-protein coupled ERs (GPER, GPER1) [7–9]. Estrogen actions that proceed independently of estrogen receptors

have also been reported [9]. Moreover, mitochondrial localization of ERs has been observed in a variety of cell types, including neuronal cells [10–12]. The biochemical mechanisms of the direct actions of estrogen in mitochondria, via their cognate receptors, is under investigation and many aspects of these actions remain to be verified and further explored. Mitochondrial estrogen receptors, especially estrogen receptor beta, is proposed to act as a direct target of estrogens, contributing to the estrogen-dependent orchestration and coordination of nuclear and mitochondrial functions, affecting mitochondrial transcription, reactive oxygen species (ROS) production, and apoptosis [6,13,14].

Protective actions of estrogens for human physiology are well documented [15–18]. Especially for the nervous system, a significant amount of evidence revealed estrogen's neuroprotective actions. Thus, estrogen is proposed to promote neuronal survival, to control neuronal excitability and synaptic plasticity, and to regulate neuronal differentiation, brain development, neuronal homeostasis, and astroglial and microglial inflammatory responses [19,20]. Both nuclear and membrane estrogen receptors participate in estradiol neuroprotective actions, among others, via the control of aromatase, b-cell lymphoma 2 (BCL-2) family member and cytokine synthesis, glycogen synthase kinase-3 β (GSK3 β) inhibition, glutamate uptake, and calcium and ROS level regulation [19–21]. ERs actions also involve interactions with alternative pathways orchestrated by other neuroprotective factors, namely insulin growth factor-1, brain derived neurotrophic growth factor, and regulatory factors of WNT (Wingless-Int) and Notch signaling [19]. Thus, estradiol is proposed as a neuroprotective factor in many neurodegenerative diseases such as Alzheimer's disease, also affecting amyloid beta synthesis and toxicity [22].

It is also widely accepted that neurons critically depend on mitochondrial function to ensure membrane excitability and to execute the complex processes of neurotransmission and plasticity [23]. Thus, mitochondrial dysfunction which is accompanied with increased ROS production and oxidative damage is associated with numerous neurodegenerative disorders [24].

To explore whether mitochondrial estrogen receptor beta is directly involved in neuroprotection against apoptotic and oxidative stress stimuli, neuroblastoma N2A cells stably overexpressing a mitochondrial targeted ER β fused with the green fluorescence protein were generated (N2AmtGFPER β). N2A cells stably overexpressing a mitochondrial-targeted GFP were also produced and used as control cells. Comparative studies to investigate the mtER β effect on staurosporine- and H₂O₂-induced apoptosis and oxidative stress, in the presence or absence of E₂, were performed, applying MTT assay, mitochondrial staining assessment via confocal microscopy, Western blot analysis of mitochondrial dependent apoptosis caspases, and ATP measurements. Our results showed that mtER β expression renders N2A neuroblastoma cells more resistant against apoptotic and oxidative stress stimuli. Thus, the direct antioxidant and anti-apoptotic actions of mtER β are documented, rendering mitochondrial ER β a promising therapeutic target for mitochondrial dysfunction-associated degenerative diseases.

2. Results

2.1. Generation of N2A Cells Stably Overexpressing a Mitochondrial-Targeted GFPER β Protein

Focusing on the elucidation of the role of the mitochondrial ER β in mitochondrial functions, and particularly in the regulation of the mitochondrial-dependent apoptosis and antioxidant stress, N2A cells stably overexpressing a mitochondrial-targeted human ER β fused with the green fluorescence protein (GFP) at its *n* terminus was generated (N2AmtGFPER β), as described in the experimental section. N2A cells stably overexpressing a mitochondrial-targeted GFP (N2AmtGFP) were also produced and used as a control. Characterization of single colonies of the N2AmtGFPER β and N2AmtGFP cells was performed as follows.

Confocal microscopy single section images of colonies of N2A cells stably overexpressing a mitochondrial targeted GFP (mtGFP) or a mitochondrial targeted GFPER β protein (mtGFPER β) are shown in Figure 1A,B, respectively. As is shown in Figure 1, staining of

GFP in N2AmtGFP and N2AmtGFPER β cells (Figure 1A,B) exhibited the same pattern and colocalized with the mitochondrial marker MitoTracker Red CMXRos (CMX), verifying the mitochondrial localization of the expressed mtGFPER β and mtGFP proteins in the N2AmtGFP and N2AmtGFPER β cells, respectively. Quantification of the relative expression levels of the fluorescent mtGFP and mtGFPER β proteins in the respective cell colonies was performed as described in the experimental section and the results are presented in Figure 1C,D. Quantitative colocalization analysis of the GFP fusion proteins and the CMX mitochondrial staining in N2AmtGFP and N2AmtGFPER β cells revealed a Pearson's correlation coefficient of 0.75 and 0.62, respectively, verifying the colocalization of the GFP and the mitochondrial CMX staining. Similarly, quantitative colocalization analysis of the GFP fusion proteins and the Hoechst 33342 nuclear staining in N2AmtGFP and N2AmtGFPER β cells revealed a Pearson's correlation coefficient of -0.25 and -0.23 , respectively, indicating no colocalization of the GFP and the nuclear Hoechst staining (see also Supplementary Figure S1). The pattern of GFPER β staining did not indicate plasma membrane localization of the receptor. Verification and assessment of the relative expression levels of the mtGFP and mtGFPER β proteins in colonies of the respective stable cell lines of N2A cells was also achieved by Western blot analysis, using specific anti-GFP antibodies (Figure 1E,F). Quantification of the results is presented in Figure 1G,H. Thus, as is shown in Figure 1D,H, colony 1 (Col. 1) of the N2AmtGFPER β cells exhibited the highest mtGFPER β expression levels and showed approximately two- to five-fold higher mtGFPER β expression compared to the other positive selected colonies (Figure 1D,H). As regards N2AmtGFP cells, colony 1 of the N2AmtGFP cells showed two- to eight-fold higher expression levels of mtGFP, compared to the respective positive selected colonies (Figure 1C,G).

To further confirm the expression and the mitochondrial localization of the GFPER β protein, immunofluorescence, and Western blot analysis using antibodies against ER β were performed. Thus, as is shown in Figure 2A, immunofluorescence analysis using antibodies against human ER β (hER β) verified the colocalization of the stably expressed hER β with the mitochondrial GFP staining. Quantitative colocalization analysis of the CMX and ER β staining revealed a Pearson's correlation coefficient of 0.84 (see also supplementary Figure S1). Moreover, via Western blot analysis using antibodies against human ER β , the detection of the GFPER β fusion protein in colonies of the N2AmtGFPER β cells, but not in N2A and N2AmtGFP cells, was verified (Figure 2C). Endogenous levels of ER β in N2A cells (Figure 2A,C) are limited and thus undetectable in accordance with previously reported observations [25–27]. The overexpression of mtER β in N2A mtGFPER β stable cell lines was also confirmed by assessment of human ER β mRNA levels. For that purpose, specific primers for the human ER β gene (ER β forward: CGGAAGCTGGCTCACTTGCT; ER β reverse: ATGCCTGACGTGGGACAGGA) (Figure 2B) and primers for β -actin (β -actin forward: TGTGACGTTGACATCCGTAA; β -actin reverse: GCTAGGAGCCAGAGCAGTAA) as a housekeeping gene were applied in real-time PCR analysis. An approximately two- to three-fold increase in mRNA levels of the human ER β levels in colony 1, compared to that of colony 3 and 2, respectively, was observed (Figure 2B). The differential expression of mRNA levels in colonies of N2AmtGFPER β cells is in accordance with the differential protein expression levels, presented in Figure 1D,H.




Figure 1. Characterization of the N2AmtGFP, N2AmtGFPERβ stable cell lines. Confocal microscopy analysis for the assessment of the GFP expression levels in colonies of the N2AmtGFP (A) and N2AmtGFPERβ, (B) cells and its colocalization

with the CMX mitochondrial dye. Hoechst staining was applied for nuclear staining. Bars indicate 10 μm . Quantification of the relative expression levels of the mitochondrial green fluorescence protein in colonies of the N2AmtGFP and N2AmtGFPER β cells is presented in (C,D), respectively. The lowest expression level was set as 1. Data are presented as the mean \pm SD; *** $p < 0.001$ ($n = 30\text{--}50$), compared to the respective colony 1 of each cell line. Representative images of Western blot analysis of GFP and β -actin (E,F), and quantification of the GFP expression levels in colonies of the N2AmtGFP and N2AmtGFPER β cells (G,H) are presented. Western blot analysis of β -actin was applied for the normalization of the results (G,H). The lowest GFP expression level, in the respective colony of each cell line was set as 1. Data are expressed as mean \pm S.D. ($n = 3$), * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, compared to the respective colony 1 of each cell line.




Figure 2. Verification of the mitochondrial ER β expression in N2AmtGFPER β cells. **(A)** Confocal colocalization studies in colony 1 of the N2AmtGFP and N2AmtGFPER β cells. Anti-ER β antibody and anti-mouse Alexa 568 secondary antibodies were applied for the ER β detection. Mitochondrial GFP fusion proteins are depicted in green. Nuclear Hoechst staining is depicted in blue. Bars indicate 10 μm . **(B)** Results from real-time PCR analysis for the assessment of the relative human mRNA ER β levels in colonies of N2AmtGFPER β cells. β -actin was used as reference gene. The highest mRNA expression level in the respective colony of the N2AmtGFPER β was set as 1. Data are presented as the mean \pm SD; *** $p < 0.001$ ($n = 4\text{--}5$), compared to colony 1. **(C)** Western blot analysis of mtGFPER β , in N2A and in colonies of N2AmtGFP and N2AmtGFPER β cells. Anti-ER β and anti- β -actin antibodies were used for ER β and β -actin detection.

The results from Figures 1 and 2 show that colonies of number 1 (colony 1) of the N2AmtGFPER β and N2AmtGFP cells showed the highest expression of the mtGFPER β and mtGFP proteins, exhibiting four- to eight-fold higher expression of the mtGFPER β and mtGFP proteins, respectively, compared to the respective ones in the others selected colonies. Thus, colonies of number 1 were selected for further analysis as described below.

2.2. Effect of mtER β on Staurosporine-Induced Apoptosis

To assess the possible anti-apoptotic activities of mtER β , comparative studies applying MTT cell viability assay, as well as Western blot analysis, of caspase-9 and -3 were performed in N2AmtGFPER β and N2AmtGFP cells, pre-cultured in a hormone-free medium and subsequently treated, or not, with various concentrations of staurosporine, in the presence or absence of 10^{-9} M E2. As is shown in Figure 3A, no statistically significant differences in cell viability were observed between N2AmtGFPER β and N2AmtGFP cells treated with staurosporine at a range concentration of 0.05 μ M to 1 μ M for 12 h. Incubation of the cells with 0.5–1 μ M staurosporine in the presence or absence of E2 caused a 30% to 60% reduction in the cell viability of both cell lines, compared to the respective vehicle-treated cells. No statistically significant differences were observed between E2-treated and E2-non treated cells. According to the literature, treatment of N2A cells with 1 μ M for 24 h triggers the induction of apoptosis [28]. Thus, further assessment of caspase activation to investigate the effect of the mitochondrial ER β in mitochondrial-dependent apoptosis was performed. Interestingly, a statistically significant increased reduction in procaspase-9 and procaspase-3 protein levels in N2AmtGFP cells, compared to the respective ones in N2AmtGFPER β cells, was observed upon treatment of the cells with 0.5 μ M or 2 μ M staurosporine (Figure 3B,C and Figure 3D,E, respectively) for 12 h. Accordingly, a statistically significant increased production of activated cleaved caspase-9 in 0.5 μ M or 2 μ M staurosporine-treated N2AmtGFP cells, compared to that in N2AmtGFPER β cells, was observed (Figure 3B,C and Figure 3D,E, respectively). Similarly, assessment of cleaved caspase-3 protein levels revealed an increased resistance in cleaved caspase-3 production in staurosporine-treated N2AmtGFPER β cells compared to the respective ones in N2AmtGFP cells. Thus, a statistically significant reduction in cleaved caspase-3 production was observed in N2AmtGFPER β , compared to that in N2AmtGFP cells, upon treatment with the relative low concentrations (0.5 μ M) of staurosporine (Figure 3B,C) and upon co-treatment with 0.5 or 2 μ M staurosporine/ 10^{-9} M E2 (Figure 3B,C and Figure 3D,E, respectively). As it was expected, the presence of E2 led to suppression of the staurosporine-induced apoptosis in both cell lines only when the cells were treated with relatively low concentrations of staurosporine. Thus, reduced cleaved caspase-9 protein levels in E2/0.5 μ M staurosporine treated cells compared to those in E2-non treated/0.5 μ M staurosporine-treated cells (Figure 3B,C) was observed.

2.3. Effect of mtER β on H₂O₂-Induced Apoptosis

In the same context, the role of mtER β in H₂O₂-induced apoptosis was assessed in N2AmtGFP and N2AmtGFPER β cells. Comparative studies on cell viability and caspase activation were performed for that purpose (Figure 4). MTT cell viability assay revealed a higher resistance (by approximately 20–30%) to the H₂O₂-induced reduction in cell viability in N2AmtGFPER β cells treated with 10 to 500 μ M H₂O₂ for 12 h, compared to the H₂O₂-treated N2AmtGFP cells (Figure 3A,B). No statistically significant differences between the N2AmtGFPER β and N2AmtGFP cells were observed upon treatment with H₂O₂ at concentrations higher than 0.5 mM. Treatment with 1 and 2 mM H₂O₂ for 12 h caused a severe reduction in cell viability that reached 90% compared to control treated cells (Figure 4B). The presence of E2 did not significantly alter H₂O₂ effects on cell viability (Figure 4A,B). Western blot analysis of caspase activation upon treatment with 2mM H₂O₂ for 2.5 h was also performed. Treatment conditions were chosen based on results from pilot experiments (Supplementary Figure S2). Thus, Western blot analysis of caspase-9 and -3 in 2 mM H₂O₂ treated cells for 2.5 h showed a statistically significant increase in cleaved

caspase-9 protein levels and a statistically significant increase in the reduction of the pro-caspase-9 and -3 protein levels in N2AmtGFP cells, compared to that in N2AmtGFPER β cells. These results further substantiate the mtER β -associated resistance of N2A cells to H₂O₂-induced apoptosis (Figure 4C,D). The protective actions of mtER β are strengthened in the presence of E₂, as indicated by the statistically significant reduced production of cleaved caspase-9 in the presence of E₂ in H₂O₂-treated N2AmtGFPER β cells compared to that in the N2AmtGFP cells (Figure 4C,D).




Figure 3. Protective effect of mtER β on staurosporine-induced apoptosis. **(A)** Assessment of the relative cell viability of N2AmtGFP and N2AmtGFPER β cells upon treatment with 0.05 to 1 μ M staurosporine for 12 h, in the presence or absence of 10⁻⁹M E₂, via MTT analysis. Data were analyzed by two-way ANOVA and expressed as mean \pm S.D. ($n = 4-6$), *** $p < 0.001$, compared to the respective vehicle-treated cells of each cell line. Western blot analysis of procaspase-9 and -3 and cleaved caspase-9 and -3, in N2AmtGFP and N2AmtGFPER β cells, treated with 0.5 μ M **(B)** and 2 μ M **(D)** staurosporine, in the presence or absence of 10⁻⁹M E₂, for 12 h. Western blot analysis of β -actin was applied for the normalization of the results. Quantification of the results is presented in **(C)** and **(E)** respectively. Data were analyzed by two-way ANOVA and expressed as mean \pm S.D. ($n = 3-4$), * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.




Figure 4. Protective effect of mtER β on H_2O_2 -induced apoptosis. Assessment of the relative cell viability of N2AmtGFP and N2AmtGFPER β cells upon treatment with 10 to 100 μM H_2O_2 (A) and 0.2 mM to 2 mM H_2O_2 (B) for 12 h, in the presence or absence of $10^{-9} M$ E_2 , via MTT analysis. Data were analyzed by two-way ANOVA and expressed as mean \pm S.D. ($n = 4-6$), ** $p < 0.01$, *** $p < 0.001$. Vertical asterisks represent statistically significant differences compared to the respective vehicle-treated cells of each cells line, while horizontal lines and the respective asterisks corresponds to statistically significant differences between the two types of cells subjected to the same treatment, as indicated. (C) Western blot analysis of procaspase-9 and -3 and cleaved caspase-9 in N2AmtGFP and N2AmtGFPER β cells, treated with 2 mM H_2O_2 , in the presence or absence of $10^{-9} M$ E_2 , for 2.5 h. Western blot analysis of β -actin was applied for the normalization of the results. Quantification of the results is presented in (D). Band density of the highest expression level of each analyzed molecule, in the respective set of experiment, was set at 1. Data were analyzed by two-way ANOVA and expressed as mean \pm S.D. ($n = 3-5$), * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

2.4. Effect of mtER β on H_2O_2 -Induced Mitochondrial Impairment in N2A Cells

The effect of mtER β on H_2O_2 -induced mitochondrial impairment was also assessed by comparative measurements of Mitotracker red CMX staining in H_2O_2 -treated N2A, N2AmtGFP, and N2AmtGFPER β cells. CMX is a red fluorescence indicator used to detect oxidation–reduction reactions, which principally occur in active mitochondria of live cells. Thus, CMX stains mitochondria in live cells and its accumulation is dependent upon mitochondrial membrane potential [29–31]. As is shown in Figure 5, treatment of N2A, N2AmtGFP, and N2AmtGFPER β cells with 2 mM H_2O_2 for 2.5 h revealed a statistically significant reduction in mitochondrial staining of H_2O_2 -treated N2A and N2AmtGFP cells compared to that in N2AmtGFPER β cells.




Figure 5. Effect of mtER β on H₂O₂-induced mitochondrial impairment in N2AmtGFP (A), N2Awt (B), and N2AmtGFPER β (C) cells. (A–C) Representative confocal microscopy single section images of GFP, CMX (mitochondrial staining) and Hoechst 33342 (nuclear staining). Bars indicate 10 μ m. (D) Quantification of the relative CMX fluorescence density. Data were analyzed by two-way ANOVA. Relative CMX fluorescence density was expressed as mean \pm S.D. ($n = 30$ – 50), ** $p < 0.01$, *** $p < 0.001$. CMX fluorescence density at vehicle treated N2A cells was set at 1. Vertical asterisks represent statistically significant differences compared to the respective vehicle treated cells of each cells line, while horizontal lines and the respective asterisks correspond to statistically significant differences between cells from different cell lines subjected to H₂O₂ treatment, as indicated.

2.5. Effect of Mitochondrial ER β on ATP Content

To assess whether mitochondrial ER β protective activity is associated with its involvement in the regulation of mitochondrial energy production, assessment of the ATP levels in N2A, N2AmtGFP, and N2AmtGFPER β cells was performed in the presence or absence of 0.2 μ M rotenone, which is an inhibitor of complex I of the mitochondrial respiratory chain. Treatment with rotenone was applied to evaluate the mitochondrial origin of ATP. As is shown in Figure 6, N2AmtGFPER β exhibited approximately a 1.5-fold statistically significant increase in ATP content compared to that in the N2AmtGFP and N2A cells. Preincubation of the cells with 0.2 μ M rotenone for 16 h caused elimination of ATP production in both cell lines, validating the mitochondrial origin of ATP.




Figure 6. Effect of the mitochondrial ER β on ATP levels in N2A, N2AmtGFP, and N2AmtGFPER β cells. Equal numbers of N2A, N2AmtGFP, and N2AmtGFPER β cells were plated on six well plate. At 70–80% confluency, cells were incubated with 0.2 μ M rotenone (diluted in DMSO) for 16 h. Control cells were treated with equal volumes of DMSO. Subsequently, cells were washed in PBS, harvested, and subjected to ATP measurements, applying ENLITEN ATP bioluminescence assay as described in experimental section. Results are expressed as relative luciferase units, normalized against relative protein levels and represent the mean \pm SD of three independent experiments. ** $p < 0.01$, *** $p < 0.001$. ATP levels at vehicle-treated N2A cells were set at 1. Vertical asterisks represent statistically significant differences compared to the respective vehicle-treated cells of each cell line, while horizontal lines and the respective asterisks correspond to statistically significant differences between cells from different cell lines.

3. Discussion

In this study, the direct involvement of the mitochondrial estrogen receptor beta (mtER β) in the activation of the mitochondrial defense antioxidant and anti-apoptotic activities is demonstrated. Accumulated evidence substantiates the regulatory and protective actions of estrogens on the brain, affecting both glial and neuronal physiology [17,21,32–35]. Particularly, the regulation of neuronal mitochondrial functions by estrogen is widely accepted [36,37]. Most of these actions are mediated by the nuclear or the membrane estrogen receptors [6,13]. In addition, considerable amounts of experimental data showed ER-independent estrogen actions [17,35], or suggested the involvement of the mitochondrial estrogen receptors, especially ER β , in these actions [38–43]. Thus, direct actions of estrogens on mitochondria are proposed to be exerted via their mitochondrial-localized estrogen receptors. This hypothesis is based on data showing mitochondrial localization of ER β in a variety of neuronal cells and tissues [10,11,44]. In addition, data showing the interaction of ER β with the mitochondrial DNA [45], the mitochondrial Hydroxyacyl-CoA Dehydrogenase Trifunctional Multienzyme Complex Subunit Beta (HADHB) fatty acid oxidation enzyme, [46,47], the mitochondrial adenosine triphosphatase (ATPase) [47,48], and the pro-apoptotic BAD protein [49,50] indicate possible involvement of the mitochondrial ER β in the regulation of mitochondrial transcription, energy metabolism and apoptosis.

Mitochondrial ATP production by oxidative phosphorylation is critical for neurons to meet their energy demands and survive. In addition, mitochondrial impairment, induced by excessive cytoplasmic Ca²⁺ loading or increased ROS generation, could lead to mitochondrial membrane potential depolarization, mitochondrial swelling, release of apoptotic factors, caspase activation, and eventually the induction of apoptosis and cell death. Thus, mitochondrial dysfunction is closely related to neuronal viability and neurodegeneration [38]. A non-genomic effect of E2 on the regulation of Ca²⁺ transport in brain mitochondria and the involvement of ER β in this action is proposed [51,52]. Thus, the verification and elucidation of the direct actions of estrogens on mitochondria is of great interest. The presence of ER β in mitochondria could establish a direct control of

mitochondrial activities by estrogens and could ensure estrogens' actions on the integration and orchestration of nuclear and mitochondrial functions. Due to the widely accepted neuroprotective actions of estrogens and to the great importance of the preservation of mitochondrial functions in neurodegenerative diseases [17,22,53], the delineation of the putative mitochondrial estrogen receptors' anti-apoptotic and anti-oxidant actions could uncover novel therapeutic targets for those high-risk disorders.

An obstacle to the verification of the direct mitochondrial ER β actions, and to the delineation of the biochemical mechanisms of its actions, is the questionable availability of specific antibodies against ER β [54]. To overcome this impediment, neuroblastoma N2A cell lines stably overexpressing a mitochondrial-targeted GFPER β (N2A mtGFPER β) were generated, and the cells were further analyzed for their resistance to apoptotic and oxidative stimuli in comparative studies. Overexpression of the mtGFPER β protein in the selected G418-resistant colonies was confirmed by Western blot analysis, using antibodies against GFP and antibodies against ER β . Results were verified by real-time PCR using ER β -specific primers. Moreover, the mitochondrial localization of the expressed mtGFPER β protein was validated in colocalization studies of the GFP and ER β proteins with the mitochondrial CMX dye, applying immunofluorescence analysis and confocal microscopy studies. Verification of the expression and the mitochondrial localization of the green fluorescence protein in N2AmtGFP control cells was also achieved. Colonies exhibiting the highest expression of the mtGFPER β and mtGFP proteins were used in this study.

Mitochondrial ROS production and induction of apoptosis is closely related to many degenerative diseases, including neurodegeneration [55–57]. In the current study, we showed mitochondrial ER β -dependent resistance in the induction of apoptosis upon treatment of the cells with either the apoptotic stimulus staurosporine or the oxidative stress stimulus H₂O₂. Thus, assessment of caspase-9 and caspase-3 activation, applying Western blot analysis, revealed decreased production of cleaved caspase-9 and -3, and reduced proteolytic degradation of procaspase-9 and -3 in N2AmtGFPER β cells compared to that in N2AmtGFP cells when treated either with 0.5 to 2 μ M staurosporine or 2 mM H₂O₂. Moreover, upon treatment with relatively low concentrations of apoptotic stimuli, the presence of E2 strengthened the mtER β -dependent resistance in the induction of apoptosis in N2AmtER β cells, whereas the inability of E2 to reverse this effect in control N2AmtGFP cells was observed. This observation indicates that the presence of mtER β is adequate to offer protection against apoptotic stimuli, but its activity is also at least in part estrogen-dependent. The protective antioxidant activities of the mtER β were also verified in fluorescence microscopy studies of the CMX mitochondrial dye, which is incorporated exclusively in functionally active mitochondria [29,30]. Upon H₂O₂-induced oxidative stress, the percentage of sustained functionally active mitochondria was higher in mtER β -overexpressing N2A cells, compared to control N2A and N2AmtGFP cells. This observation is in agreement with previous studies showing anti-apoptotic and antioxidant activities of estrogens in mitochondria, which is proposed to be mediated by nuclear and membranous ERs, as well as by estrogen receptor-independent estrogen action [58–62]. The results from this study provide evidence for the direct involvement of the mitochondrial ER β in this process. Thus, since endogenous ER β levels in N2A cells are limited [25,27], the results from this study indicate that estrogens' protective actions are also mediated, at least in part, via the mitochondrial estrogen receptor beta.

In accordance with results from Western blot analysis and fluorescence microscopy studies, MTT cell viability assay confirmed the protective effect of the mtER β overexpression in cells treated with H₂O₂ at a concentration range of 10–500 μ M, even upon conditions of a longer incubation time (12 h/2.5 h). This effect was not detectable at higher concentrations of H₂O₂, possibly due to severe induction of the oxidative stress, which could not be reversible by the protective activity of mtER β . No statistically significant differences between N2AmtGFPER β and N2AmtGFP cell viability were also observed in cells treated with staurosporine. Nevertheless, results from the Western blot analysis

uncovered the protective actions of mtER β , as was shown by the reduced activation of caspase-9, which corresponds to the initial steps of mitochondrial-induced apoptosis.

Mitochondria are the main energy suppliers of the cell. The maintenance of functional, active mitochondria ensures adequate energy production, which is essential for neuron survival and the prevention of degenerative processes that lead to neurodegenerative diseases and aging [24]. Thus, factors that ensure and reinforce energy production are crucial regulators of neuronal viability and function. In this context, mtER β can be considered such a valuable factor, as indicated by our observations showing increased ATP content in N2AmtGFPER β compared to that in N2AmtGFP cells. The elimination of ATP level by rotenone substantiates its mitochondrial origin, further supporting the involvement of mitochondrial ER β in this action.

Thus, anti-apoptotic and antioxidant activities of the mitochondrial-localized ER β could be attributed to its proposed actions on the mitochondrial transcriptional activation, and thus oxidative phosphorylation enzyme activation [12,42], that ends up in an increase in ATP production and a reduction in ROS production. In addition, anti-apoptotic and antioxidant activities of mtER β can also be mediated via their involvement in the regulation of the activity of mitochondrial apoptosis-associated and antioxidant molecules, such as BAD and MnSOD [4,50,63,64], via mtER β interaction with other mitochondrial components or via its interfering with other estrogen-dependent or estrogen-independent signaling pathways, such as Ca²⁺ modulating effects [11,51,52]. Moreover, since E2 is known to activate the synthesis of many nuclear-encoded antioxidant enzymes, namely MnSOD, thioredoxin, and glutathione peroxidase [65–68], the involvement of mtER β in the orchestration of the synthesis of nuclear-encoded mitochondrial antioxidant- and apoptosis-associated molecules is also an interesting issue that remains to be explored.

To sum up, the results from this study revealed an increased resistance to staurosporine- and H₂O₂-induced apoptotic stimuli in N2A cells overexpressing a mitochondrial-targeted estrogen receptor beta, as indicated by the reduced activation of caspase-9 and -3, the increased cell viability and ATP production, and the increased resistance to mitochondrial impairment in the presence or absence of E2. Thus, the direct involvement of mtER β in the antioxidant and anti-apoptotic E2 actions is documented, rendering ER β a promising therapeutic target for mitochondrial-related degenerative diseases.

4. Materials and Methods

4.1. Chemicals

Dulbecco's modified eagle medium (DMEM), trypsin, fetal bovine serum (FBS), Lipofectamin 2000, and MitoTracker Red CMXRos (CMX) were obtained from Thermo Fisher Scientific (Thermo Fisher Scientific GmbH, Basel, Switzerland). Molecular weight markers, complete protease cocktail inhibitors, and Western blotting luminol reagent were purchased from Thermo Fisher Scientific, Roche (Mannheim, Germany) and Santa Cruz Biotechnology (Santa Cruz Biotechnology, Inc., Dallas, TX, USA), respectively. Geneticin (G418) was provided by Calbiochem (Merck, Kenilworth, NJ, USA). All other chemicals, including Hoechst 33342, staurosporine, estradiol (E2), and H₂O₂, were purchased from Sigma-Aldrich (St. Louis, MO, USA).

4.2. Antibodies

Monoclonal antibodies against human ER β (B-3) and α -tubulin were provided by Santa Cruz Biotechnology. Monoclonal antibodies against β -actin were provided by Sigma Aldrich. Monoclonal antibodies against caspase-9 and -3 and polyclonal antibodies against cleaved caspase-3 were provided by Cell Signaling (Cell Signaling Technology, Danvers, MA, USA). Monoclonal antibodies against GFP were provided by Roche. Horseradish peroxidase (HRP)- and Alexa 568-conjugated secondary antibodies were purchased from Thermo Fisher Scientific.

4.3. Cell Culture—Generation of N2A Cells Stably Expressing a Mitochondrial Targeted GFPER β Fusion Protein

Mouse N2A neuroblastoma cells (obtained from the American type culture collection—ATCC) were maintained in low glucose (1 g/L) DMEM, supplemented with 10% FBS, 2 mM glutamine, and 50 units/mL penicillin/streptomycin (DMEM complete medium). Cells were grown at 37 °C in a humidified atmosphere with 5% CO₂. For stable cell lines generation, at day 1 before transfection, N2A cells were plated in 60 mm culture dishes so that they reached 60–70% confluency at the time of transfection. To obtain stable expression of the human mitochondrial ER β , the human estrogen receptor beta gene (AB006590) [69] was inserted into the pEGPC2 vector (Clontech-Takara, Mountain View, CA, USA), in frame with the enhanced green fluorescence protein (EGFP) gene between Xho1 and BamH1 sites, to produce the pEGFPC2ER β vector. Subsequently, the following sequence: atggctcagc-gacttctctgaggaggttctggcctctgtcatctccaggaagccctctcagggtcagtgccaccctcactccagagcctgcag accccacaatgcagtcctggctggcctgctgaaccccacccagcccgacaatacaccacaggatctcttgaca, which encodes a mitochondrial targeting peptide, was inserted in frame with the EGFP Gene between the NheI and AgeI sites of the pEGFPC2ER β construct to produce the pmtEGFPER β construct. The produced construct was transfected into the N2A cells using Lipofectamin 2000 (Thermo Fisher Scientific GmbH, Europe), according to the manufacturer's instructions. Control cells (N2AmtGFP cells) were prepared by transfection of the pmtEGFPC2 construct [70]. After 36 h incubation at 37 °C, cells were trypsinized and passed at a 1:14 dilution in selective growth medium [DMEM supplemented with 10% (*v/v*) FBS, containing 1.5 mg/mL G418]. On the next day, the medium in all plates was replaced with fresh selective medium and cells were cultured for 2 weeks. G418-resistant colonies were expanded, cloned independently, and analyzed by immunofluorescence and Western blot analysis to confirm the expression of the mitochondrial-GFP (mtGFP) and -GFPER β (mtGFPER β) proteins in the N2AmtGFP and N2AmtGFPER β cells, respectively.

4.4. Western Blot Analysis

Western blot analysis of molecules involved in mitochondria-mediated apoptosis, such as caspase-9 and caspase-3, was performed in extracts from N2AmtGFP and N2AmtGFPER β cells, treated, or not, with 0.5–2 μ M staurosporine or 2 mM H₂O₂, in the presence or absence of estradiol, using specific antibodies. Briefly, cells plated on 6 well plates were cultured for 40–48 h in low glucose DMEM without phenol red, supplemented with 10% dextran-coated charcoal stripped fetal bovine serum. Then, cells were incubated with 0.5–2 μ M staurosporine or 2 mM H₂O₂ at various time points, as indicated in the results section. After the washing of the cells with phosphate buffer saline (PBS), cells were lysed in lysis buffer (20 mM Tris pH: 7.5, 250 mM NaCl, 0.5% Triton, 3 mM EDTA) supplemented with cocktail protease inhibitors. Protein determination, electrophoresis and Western blot analysis were performed as previously described [71]. Quantification of the results was carried out by applying ImageJ (1.52 p) analysis (NIH, Bethesda, MD, USA).

4.5. Immunofluorescence—Confocal Microscopy

Cells grown on coverslips for 48 h in DMEM complete medium were further incubated for 30 min at 37 °C with 200 nM CMX. In the case of H₂O₂ treatment, cells were treated with 1 mM or 2 mM H₂O₂ for 2.5 h at 37 °C in the presence of 200 nM CMX. Subsequently, cells were washed 3 \times 5 min with PBS, fixed for 10 min in ice-cold methanol, and transferred to ice-cold acetone for 2 min. For nuclear staining, specimens were incubated with 1 μ g/mL Hoechst 33,342 in PBS for 30 min at room temperature. Antibodies against ER β (B-3, Santa Cruz Biotechnology, Dallas, TX, USA) were applied for ER β detection. Secondary anti-mouse Alexa 568-conjugated antibodies were used. The specimens were washed in PBS and mounted in anti-fading medium. Images were taken using an inverted laser-scanning confocal microscope (Zeiss laser scanning microscope 800, Zeiss LSM 800, Zeiss, Jena, Germany). Image analysis and quantification of fluorescence staining were performed as previously described [25] using the ImageJ (1.52 p) analysis program. Briefly, the total

corrected fluorescence of area of interest (TCF) = integrated density – (selected area × mean fluorescence of background readings) was calculated. The TCF of treated cells was then normalized against the mean TCF of the relative control cells (Relative Fluorescence Density). Quantitative colocalization analysis of the GFP fusion proteins and the CMX and Hoechst staining, and of the ER β and the CMX staining, was performed by measuring the Pearson's correlation coefficient, applying the ZEN Zeiss LSM 800 software, according to manufacturer's instructions.

4.6. Cell Viability Assay

In order to investigate the role of mtER β on the cell viability under stress conditions, the MTT assay [72] was performed. Cells of the two stable cell lines were plated on 96-well plates and were cultured in low glucose DMEM without phenol red, supplemented with 10% dextran-coated charcoal stripped fetal bovine serum. After 40–48 h, the cells were incubated with various concentrations of staurosporine or H₂O₂ in the presence or absence of E2 for 12 h, as indicated in the results section. Then, the cells were incubated with 0.5 mg/mL MTT for 3 h at 37 °C and 5% CO₂ and the formed formazan crystals were dissolved with isopropanol. The color intensity at 570 nm (reference filter at 690 nm) was measured using a multimode plate reader (EnSpire, PerkinElmer, UK). Relative cell viability corresponds to absorbance values of treated cells normalized against absorbance values of the relative control cells.

4.7. ATP Measurements

ATP contents in N2A, N2AmtGFP, and N2AmtGFPER β was measured as previously described [70]. Briefly, cells grown in DMEM complete medium were trypsinized and plated in a 6-well plate in hormone-free DMEM medium for 48 h. Subsequently, cells were treated in the presence or absence of 0.2 μ M rotenone for 16 h. Cells were washed in PBS, harvested and centrifuged at 800× *g*. ATP from the cell pellet was extracted with 1% trichloroacetic acid and ATP content was measured in a Luminometre (LB 9508, Berthold, Europe) by bioluminescence using the luciferin–luciferase reaction kit (Enliten, Promega Corporation, Madison, WI, USA) following the manufacturer's instructions.

4.8. Statistical Analysis

All results are expressed as mean \pm SD. Data were analyzed by two-way ANOVA followed by Tukeys's post-hoc test using StatPlus LE 7.3.0 software. Differences were considered significant at a two-tailed *p* value < 0.05.

Supplementary Materials: The following are available online at <https://www.mdpi.com/article/10.3390/ijms22147620/s1>.

Author Contributions: A.-M.G.P.; validation, I.T., A.G., M.E.K., F.D.K. and A.G.K.; formal analysis, I.T. and A.G.; investigation, I.T., M.E.K.; resources, A.-M.G.P. and D.D.L.; data curation, I.T. and A.-M.G.P.; writing—original draft preparation, A.-M.G.P.; writing—review and editing, A.-M.G.P., I.T. and D.D.L.; visualization, I.T.; supervision, A.-M.G.P.; project administration, A.-M.G.P.; funding acquisition, A.-M.G.P. and D.D.L. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the project “Synthetic Biology: From omics technologies to genomic engineering” (OMIC-ENGINE) (MIS 5002636) which is implemented under the action “reinforcement of the Research and Innovation Infrastructure” funded by the Operational Programme “Competitiveness, Entrepreneurship and Innovation (NSRF 2014–2020) and co-financed by Greece and the European Union (European Regional Development Fund) (to A.-M.G.P., I.T., A.G. and D.D.L.). The research work was supported by the Hellenic Foundation for Research and Innovation (HFRI) under the HFRI PhD Fellowship grant (Fellowship Number: 6246 to I.T.). This work was also supported by the Postgraduate Programme “Application of Molecular Biology-Molecular Genetics-Molecular Markers”, Department of Biochemistry and Biotechnology, University of Thessaly (to A.-M.G.P.).

Acknowledgments: The authors would like to thank the Bodossaki Foundation for their optiMOS camera donation (Qimaging, Surrey, BC, Canada). The optiMOS camera was used in the initial evaluation of fluorescence staining.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Hess, R.A.; Bunick, D.; Lee, K.-H.; Bahr, J.; Taylor, J.A.; Korach, K.; Lubahn, D.B. A role for oestrogens in the male reproductive system. *Nat. Cell Biol.* **1997**, *390*, 509–512. [[CrossRef](#)]
2. Hofbauer, L.C.; Khosla, S.; Dunstan, C.; Lacey, D.L.; Spelsberg, T.C.; Riggs, B.L. Estrogen Stimulates Gene Expression and Protein Production of Osteoprotegerin in Human Osteoblastic Cells. *Endocrinology* **1999**, *140*, 4367–4370. [[CrossRef](#)]
3. Rosenfeld, C.S.; Ganjam, V.K.; Taylor, J.A.; Yuan, X.; Stiehr, J.R.; Hardy, M.P.; Lubahn, D.B. Transcription and Translation of Estrogen Receptor- α in the Male Reproductive Tract of Estrogen Receptor- α Knock-Out and Wild-Type Mice. *Endocrinology* **1998**, *139*, 2982–2987. [[CrossRef](#)]
4. Klinge, C.M. Estrogens regulate life and death in mitochondria. *J. Bioenerg. Biomembr.* **2017**, *49*, 307–324. [[CrossRef](#)]
5. Kocanova, S.; Mazaheri, M.; Caze-Subra, S.; Bystricky, K. Ligands specify estrogen receptor alpha nuclear localization and degradation. *BMC Cell Biol.* **2010**, *11*, 98. [[CrossRef](#)]
6. Psarra, A.-M.G.; Sekeris, C.E. Steroid and thyroid hormone receptors in mitochondria. *IUBMB Life* **2008**, *60*, 210–223. [[CrossRef](#)] [[PubMed](#)]
7. Heldring, N.; Pike, A.; Andersson, S.; Matthews, J.; Cheng, G.; Hartman, J.; Tujague, M.; Ström, A.; Treuter, E.; Warner, M.; et al. Estrogen Receptors: How Do They Signal and What Are Their Targets. *Physiol. Rev.* **2007**, *87*, 905–931. [[CrossRef](#)]
8. Levin, E.R. Extranuclear Steroid Receptors Are Essential for Steroid Hormone Actions. *Annu. Rev. Med.* **2015**, *66*, 271–280. [[CrossRef](#)] [[PubMed](#)]
9. Fuentes, N.; Silveyra, P. Estrogen receptor signaling mechanisms. *Adv. Protein Chem. Struct. Biol.* **2019**, *116*, 135–170. [[CrossRef](#)] [[PubMed](#)]
10. Psarra, A.-M.; Solakidi, S.; Sekeris, C. The mitochondrion as a primary site of action of steroid and thyroid hormones: Presence and action of steroid and thyroid hormone receptors in mitochondria of animal cells. *Mol. Cell. Endocrinol.* **2006**, *246*, 21–33. [[CrossRef](#)] [[PubMed](#)]
11. Simpkins, J.W.; Yang, S.-H.; Sarkar, S.N.; Pearce, V. Estrogen actions on mitochondria—Physiological and pathological implications. *Mol. Cell. Endocrinol.* **2008**, *290*, 51–59. [[CrossRef](#)] [[PubMed](#)]
12. Chen, J.-Q.; Cammarata, P.R.; Baines, C.; Yager, J.D. Regulation of mitochondrial respiratory chain biogenesis by estrogens/estrogen receptors and physiological, pathological and pharmacological implications. *Biochim. Biophys. Acta (BBA)-Bioenerg.* **2009**, *1793*, 1540–1570. [[CrossRef](#)]
13. Mattingly, K.A.; Ivanova, M.M.; Riggs, K.A.; Wickramasinghe, N.S.; Barch, M.J.; Klinge, C.M. Estradiol Stimulates Transcription of Nuclear Respiratory Factor-1 and Increases Mitochondrial Biogenesis. *Mol. Endocrinol.* **2008**, *22*, 609–622. [[CrossRef](#)] [[PubMed](#)]
14. McEwen, B.S.; Akama, K.T.; Spencer-Segal, J.L.; Milner, T.A.; Waters, E.M. Estrogen effects on the brain: Actions beyond the hypothalamus via novel mechanisms. *Behav. Neurosci.* **2012**, *126*, 4–16. [[CrossRef](#)] [[PubMed](#)]
15. Emmanuelle, N.-E.; Marie-Cécile, V.; Florence, T.; Jean-François, A.; Françoise, L.; Coralie, F.; Alexia, V. Critical Role of Estrogens on Bone Homeostasis in Both Male and Female: From Physiology to Medical Implications. *Int. J. Mol. Sci.* **2021**, *22*, 1568. [[CrossRef](#)] [[PubMed](#)]
16. Ueda, K.; Adachi, Y.; Liu, P.; Fukuma, N.; Takimoto, E. Regulatory Actions of Estrogen Receptor Signaling in the Cardiovascular System. *Front. Endocrinol.* **2020**, *10*, 909. [[CrossRef](#)] [[PubMed](#)]
17. Azcoitia, I.; Barreto, G.E.; Garcia-Segura, L.M. Molecular mechanisms and cellular events involved in the neuroprotective actions of estradiol. Analysis of sex differences. *Front. Neuroendocr.* **2019**, *55*, 100787. [[CrossRef](#)]
18. Jia, M.; Dahlman-Wright, K.; Gustafsson, J.Å. Estrogen receptor alpha and beta in health and disease. *Best Pract. Res. Clin. Endocrinol. Metab.* **2015**, *29*, 557–568. [[CrossRef](#)]
19. Arevalo, M.-A.; Azcoitia, I.; Garcia-Segura, L. The neuroprotective actions of oestradiol and oestrogen receptors. *Nat. Rev. Neurosci.* **2015**, *16*, 17–29. [[CrossRef](#)]
20. Baez-Jurado, E.; Rincón-Benavides, M.; Lanussa, O.A.H.; Guio-Vega, G.; Ashraf, G.M.; Sahebkar, A.; Echeverria, V.; Garcia-Segura, L.; Barreto, G. Molecular mechanisms involved in the protective actions of Selective Estrogen Receptor Modulators in brain cells. *Front. Neuroendocr.* **2019**, *52*, 44–64. [[CrossRef](#)]
21. Saldanha, C.J. Estrogen as a Neuroprotectant in Both Sexes: Stories from the Bird Brain. *Front. Neurol.* **2020**, *11*, 497. [[CrossRef](#)] [[PubMed](#)]
22. Uddin, M.S.; Rahman, M.M.; Jakaria, M.; Rahman, M.S.; Hossain, M.S.; Islam, A.; Ahmed, M.; Mathew, B.; Omar, U.M.; Barreto, G.E.; et al. Estrogen Signaling in Alzheimer’s Disease: Molecular Insights and Therapeutic Targets for Alzheimer’s Dementia. *Mol. Neurobiol.* **2020**, *57*, 2654–2670. [[CrossRef](#)]
23. Kann, O.; Kovács, R. Mitochondria and neuronal activity. *Am. J. Physiol. Physiol.* **2007**, *292*, C641–C657. [[CrossRef](#)]
24. Johnson, J.; Mercado-Ayón, E.; Mercado-Ayón, Y.; Na Dong, Y.; Halawani, S.; Ngaba, L.; Lynch, D.R. Mitochondrial dysfunction in the development and progression of neurodegenerative diseases. *Arch. Biochem. Biophys.* **2020**, *108698*, 108698. [[CrossRef](#)]

25. Tsialtas, I.; Gorgogietas, V.A.; Michalopoulou, M.; Komninou, A.; Liakou, E.; Georgantopoulos, A.; Kalousi, F.D.; Karra, A.G.; Protopapa, E.; Psarra, A.-M.G. Neurotoxic effects of aluminum are associated with its interference with estrogen receptors signaling. *NeuroToxicology* **2020**, *77*, 114–126. [[CrossRef](#)]
26. Méndez, P.; Garcia-Segura, L.M. Phosphatidylinositol 3-Kinase and Glycogen Synthase Kinase 3 Regulate Estrogen Receptor-Mediated Transcription in Neuronal Cells. *Endocrinology* **2006**, *147*, 3027–3039. [[CrossRef](#)]
27. Su, C.; Rybalchenko, N.; Schreihof, D.A.; Singh, M.; Abbassi, B.; Cunningham, R.L. Cell Models for the Study of Sex Steroid Hormone Neurobiology. *J. Steroids Horm. Sci.* **2012**; *3*, (Suppl. 2). [[CrossRef](#)]
28. Li, H.-L.; Wang, H.-H.; Liu, S.-J.; Deng, Y.-Q.; Zhang, Y.-J.; Tian, Q.; Wang, X.-C.; Chen, X.-Q.; Yang, Y.; Zhang, J.-Y.; et al. Phosphorylation of tau antagonizes apoptosis by stabilizing beta-catenin, a mechanism involved in Alzheimer's neurodegeneration. *Proc. Natl. Acad. Sci. USA* **2007**, *104*, 3591–3596. [[CrossRef](#)] [[PubMed](#)]
29. Wang, Y.-F.; Li, C.-C.; Cai, J.-X. Aniracetam attenuates H₂O₂-induced deficiency of neuron viability, mitochondria potential and hippocampal long-term potentiation of mice in vitro. *Neurosci. Bull.* **2006**, *22*, 274–280. [[PubMed](#)]
30. Maharjan, S.; Oku, M.; Tsuda, M.; Hoseki, J.; Sakai, Y. Mitochondrial impairment triggers cytosolic oxidative stress and cell death following proteasome inhibition. *Sci. Rep.* **2014**, *4*, 05896. [[CrossRef](#)] [[PubMed](#)]
31. Chiaradonna, F.; Gaglio, D.; Vanoni, M.; Alberghina, L. Expression of transforming K-Ras oncogene affects mitochondrial function and morphology in mouse fibroblasts. *Biochim. Biophys. Acta (BBA)-Bioenerg.* **2006**, *1757*, 1338–1356. [[CrossRef](#)] [[PubMed](#)]
32. Wang, J.; Sareddy, G.R.; Lu, Y.; Pratap, U.P.; Tang, F.; Greene, K.M.; Meyre, P.L.; Tekmal, R.R.; Vadlamudi, R.K.; Brann, D.W. Astrocyte-Derived Estrogen Regulates Reactive Astroglia and is Neuroprotective following Ischemic Brain Injury. *J. Neurosci.* **2020**, *40*, 9751–9771. [[CrossRef](#)]
33. Wang, J.; Hou, Y.; Zhang, L.; Liu, M.; Zhao, J.; Zhang, Z.; Ma, Y.; Hou, W. Estrogen Attenuates Traumatic Brain Injury by Inhibiting the Activation of Microglia and Astrocyte-Mediated Neuroinflammatory Responses. *Mol. Neurobiol.* **2021**, *58*, 1052–1061. [[CrossRef](#)]
34. Lu, Y.; Sareddy, G.R.; Wang, J.; Zhang, Q.; Tang, F.-L.; Pratap, U.P.; Tekmal, R.R.; Vadlamudi, R.K.; Brann, D.W. Neuron-Derived Estrogen Is Critical for Astrocyte Activation and Neuroprotection of the Ischemic Brain. *J. Neurosci.* **2020**, *40*, 7355–7374. [[CrossRef](#)]
35. Guo, H.; Liu, M.; Zhang, L.; Wang, L.; Hou, W.; Ma, Y.; Ma, Y. The Critical Period for Neuroprotection by Estrogen Replacement Therapy and the Potential Underlying Mechanisms. *Curr. Neuropharmacol.* **2020**, *18*, 485–500. [[CrossRef](#)] [[PubMed](#)]
36. Dykens, J.A.; Moos, W.H.; Howell, N. Development of 17 β -Estradiol as a Neuroprotective Therapeutic Agent: Rationale and Results from a Phase I Clinical Study. *Ann. N. Y. Acad. Sci.* **2005**, *1052*, 116–135. [[CrossRef](#)]
37. Stanojlovic, M.; Guševac, I.; Grković, I.; Mitrović, N.; Zlatković, J.; Horvat, A.; Drakulić, D.; Martinović, J. Repeated Estradiol Treatment Attenuates Chronic Cerebral Hypoperfusion-Induced Neurodegeneration in Rat Hippocampus. *Cell. Mol. Neurobiol.* **2015**, *36*, 989–999. [[CrossRef](#)]
38. Simpkins, J.W.; Yi, K.D.; Yang, S.-H.; Dykens, J.A. Mitochondrial mechanisms of estrogen neuroprotection. *Biochim. Biophys. Acta (BBA)-Gen. Subj.* **2010**, *1800*, 1113–1120. [[CrossRef](#)]
39. Chen, J.-Q.; Yager, J.D.; Russo, J. Regulation of mitochondrial respiratory chain structure and function by estrogens/estrogen receptors and potential physiological/pathophysiological implications. *Biochim. Biophys. Acta (BBA)-Bioenerg.* **2005**, *1746*, 1–17. [[CrossRef](#)]
40. Chen, J.Q.; Eshete, M.; Alworth, W.L.; Yager, J.D. Binding of MCF-7 cell mitochondrial proteins and recombinant human estrogen receptors and to human mitochondrial dna estrogen response elements. *J. Cell. Biochem.* **2004**, *93*, 358–373. [[CrossRef](#)] [[PubMed](#)]
41. Yang, S.-H.; Liu, R.; Perez, E.J.; Wen, Y.; Stevens, S.M.; Valencia, T.; Brun-Zinkernagel, A.-M.; Prokai, L.; Will, Y.; Dykens, J.; et al. Mitochondrial localization of estrogen receptor. *Proc. Natl. Acad. Sci. USA* **2004**, *101*, 4130–4135. [[CrossRef](#)]
42. Psarra, A.-M.G.; Sekeris, C.E. Nuclear receptors and other nuclear transcription factors in mitochondria: Regulatory molecules in a new environment. *Biochim. Biophys. Acta (BBA)-Bioenerg.* **2008**, *1783*, 1–11. [[CrossRef](#)]
43. Borrás, C.; Gambini, J.; Lopez-Gruoso, R.; Pallardó, F.V.; Viña, J. Direct antioxidant and protective effect of estradiol on isolated mitochondria. *Biochim. Biophys. Acta (BBA)-Mol. Basis Dis.* **2010**, *1802*, 205–211. [[CrossRef](#)] [[PubMed](#)]
44. Levin, E.R. Membrane estrogen receptors signal to determine transcription factor function. *Steroids* **2018**, *132*, 1–4. [[CrossRef](#)] [[PubMed](#)]
45. Chen, J.Q.; Delannoy, M.; Cooke, C.; Yager, J.D. Mitochondrial localization of ER α and ER β in human MCF7 cells. *Am. J. Physiol. Metab.* **2004**, *286*, E1011–E1022. [[CrossRef](#)] [[PubMed](#)]
46. Zhou, Z.; Zhou, J.; Du, Y. Estrogen receptor beta interacts and colocalizes with HADHB in mitochondria. *Biochem. Biophys. Res. Commun.* **2012**, *427*, 305–308. [[CrossRef](#)]
47. Burstein, S.R.; Kim, H.J.; Fels, J.A.; Qian, L.; Zhang, S.; Zhou, P.; Starkov, A.A.; Iadecola, C.; Manfredi, G. Estrogen receptor beta modulates permeability transition in brain mitochondria. *Biochim. Biophys. Acta (BBA)-Bioenerg.* **2018**, *1859*, 423–433. [[CrossRef](#)]
48. Álvarez-Delgado, C.; Mendoza-Rodríguez, C.A.; Picazo, O.; Cerbón, M. Different expression of α and β mitochondrial estrogen receptors in the aging rat brain: Interaction with respiratory complex V. *Exp. Gerontol.* **2010**, *45*, 580–585. [[CrossRef](#)] [[PubMed](#)]
49. Zhang, G.; Yanamala, N.; Lathrop, K.L.; Zhang, L.; Klein-Seetharaman, J.; Srinivas, H. Ligand-Independent Antiapoptotic Function of Estrogen Receptor- β in Lung Cancer Cells. *Mol. Endocrinol.* **2010**, *24*, 1737–1747. [[CrossRef](#)]
50. Liang, J.; Xie, Q.; Li, P.; Zhong, X.; Chen, Y. Mitochondrial estrogen receptor β inhibits cell apoptosis via interaction with Bad in a ligand-independent manner. *Mol. Cell. Biochem.* **2014**, *401*, 71–86. [[CrossRef](#)] [[PubMed](#)]

51. Petrovic, S.; Veličković, N.; Stanojević, I.; Milosevic, M.; Drakulic, D.; Stanojlovic, M.; Horvat, A. Inhibition of mitochondrial Na⁺-dependent Ca²⁺ efflux by 17β-estradiol in the rat hippocampus. *Neuroscience* **2011**, *192*, 195–204. [[CrossRef](#)]
52. Petrović, S.; Milosevic, M.; Drakulic, D.; Grkovic, I.; Stanojlovic, M.; Mitrovic, N.; Horvat, A. 17β-estradiol modulates mitochondrial Ca²⁺ flux in rat caudate nucleus and brain stem. *Neuroscience* **2012**, *220*, 32–40. [[CrossRef](#)]
53. Crespo-Castrillo, A.; Arevalo, M.-A. Microglial and Astrocytic Function in Physiological and Pathological Conditions: Estrogenic Modulation. *Int. J. Mol. Sci.* **2020**, *21*, 3219. [[CrossRef](#)]
54. Nelson, A.W.; Groen, A.J.; Miller, J.L.; Warren, A.Y.; Holmes, K.A.; Tarulli, G.A.; Tilley, W.; Katzenellenbogen, B.S.; Hawse, J.R.; Gnanapragasam, V.J.; et al. Comprehensive assessment of estrogen receptor beta antibodies in cancer cell line models and tissue reveals critical limitations in reagent specificity. *Mol. Cell. Endocrinol.* **2017**, *440*, 138–150. [[CrossRef](#)] [[PubMed](#)]
55. Wang, B.; Huang, M.; Shang, D.; Yan, X.; Zhao, B.; Zhang, X. Mitochondrial Behavior in Axon Degeneration and Regeneration. *Front. Aging Neurosci.* **2021**, *13*, 650038. [[CrossRef](#)]
56. Kent, A.C.; El Baradie, K.B.Y.; Hamrick, M.W. Targeting the Mitochondrial Permeability Transition Pore to Prevent Age-Associated Cell Damage and Neurodegeneration. *Oxidative Med. Cell. Longev.* **2021**, *2021*, 1–15. [[CrossRef](#)] [[PubMed](#)]
57. Lee, Y.M.; He, W.; Liou, Y.-C. The redox language in neurodegenerative diseases: Oxidative post-translational modifications by hydrogen peroxide. *Cell Death Dis.* **2021**, *12*, 1–13. [[CrossRef](#)]
58. Chainy, G.B.N.; Sahoo, D.K. Hormones and oxidative stress: An overview. *Free Radic. Res.* **2019**, *54*, 1–26. [[CrossRef](#)]
59. Green, P.S.; Simpkins, J.W. Neuroprotective effects of estrogens: Potential mechanisms of action. *Int. J. Dev. Neurosci.* **2000**, *18*, 347–358. [[CrossRef](#)]
60. Green, P.S.; Simpkins, J.W. Estrogens and estrogen-like non-feminizing compounds. Their role in the prevention and treatment of Alzheimer's disease. *Ann. N. Y. Acad. Sci.* **2006**, *924*, 93–98. [[CrossRef](#)] [[PubMed](#)]
61. Wise, P.M.; Dubal, D.B.; E Wilson, M.; Rau, S.W.; Böttner, M.; Rosewell, K.L. Estradiol is a protective factor in the adult and aging brain: Understanding of mechanisms derived from in vivo and in vitro studies. *Brain Res. Rev.* **2001**, *37*, 313–319. [[CrossRef](#)]
62. McEwen, B.S. Invited review: Estrogens effects on the brain: Multiple sites and molecular mechanisms. *J. Appl. Physiol.* **2001**, *91*, 2785–2801. [[CrossRef](#)] [[PubMed](#)]
63. Lone, M.-U.-D.; Baghel, K.S.; Kanchan, R.K.; Shrivastava, R.; A Malik, S.; Tewari, B.N.; Tripathi, C.; Negi, M.P.S.; Garg, V.K.; Sharma, M.; et al. Physical interaction of estrogen receptor with MnSOD: Implication in mitochondrial O₂ – upregulation and mTORC2 potentiation in estrogen-responsive breast cancer cells. *Oncogene* **2016**, *36*, 1829–1839. [[CrossRef](#)] [[PubMed](#)]
64. Razandi, M.; Pedram, A.; Jordan, V.C.; Fuqua, S.; Levin, E.R. Tamoxifen regulates cell fate through mitochondrial estrogen receptor beta in breast cancer. *Oncogene* **2013**, *32*, 3274–3285. [[CrossRef](#)] [[PubMed](#)]
65. Liu, Z.; Gou, Y.; Zhang, H.; Zuo, H.; Zhang, H.; Liu, Z.; Yao, D. Estradiol improves cardiovascular function through up-regulation of SOD2 on vascular wall. *Redox Biol.* **2014**, *3*, 88–99. [[CrossRef](#)]
66. Strehlow, K.; Rotter, S.; Wassmann, S.; Adam, O.; Grohé, C.; Laufs, K.; Bohm, M.; Nickenig, G. Modulation of Antioxidant Enzyme Expression and Function by Estrogen. *Circ. Res.* **2003**, *93*, 170–177. [[CrossRef](#)]
67. Chiueh, C.C.; Lee, S.Y.; Andoh, T.; Murphy, D.L. Induction of Antioxidative and Antiapoptotic Thioredoxin Supports Neuroprotective Hypothesis of Estrogen. *Endocrine* **2003**, *21*, 27–32. [[CrossRef](#)]
68. Klinge, C.M. Estrogenic control of mitochondrial function and biogenesis. *J. Cell. Biochem.* **2008**, *105*, 1342–1351. [[CrossRef](#)]
69. Ogawaab, S.; Inoueab, S.; Watanabea, T.; Hiroia, H.; Orimoa, A.; Hosoib, T.; Ouchib, Y.; Muramatsu, M. The Complete Primary Structure of Human Estrogen Receptor β (hERβ) and Its Heterodimerization with ER αin Vivoandin Vitro. *Biochem. Biophys. Res. Commun.* **1998**, *243*, 122–126. [[CrossRef](#)]
70. Psarra, A.-M.G.; Sekeris, C.E. Glucocorticoids induce mitochondrial gene transcription in HepG2 cells: Role of the mitochondrial glucocorticoid receptor. *Biochim. Biophys. Acta (BBA)-Bioenerg.* **2011**, *1813*, 1814–1821. [[CrossRef](#)]
71. Psarra, A.-M.; Solakidi, S.; Trougakos, I.P.; Margaritis, L.H.; Spyrou, G.; Sekeris, C.E. Glucocorticoid receptor isoforms in human hepatocarcinoma HepG2 and SaOS-2 osteosarcoma cells: Presence of glucocorticoid receptor alpha in mitochondria and of glucocorticoid receptor beta in nucleoli. *Int. J. Biochem. Cell Biol.* **2005**, *37*, 2544–2558. [[CrossRef](#)] [[PubMed](#)]
72. Mosmann, T. Rapid colorimetric assay for cellular growth and survival: Application to proliferation and cytotoxicity assays. *J. Immunol. Methods* **1983**, *65*, 55–63. [[CrossRef](#)]