

RESEARCH ARTICLE

# Regulation of the Flavonoid Biosynthesis Pathway Genes in Purple and Black Grains of *Hordeum vulgare*

Olesya Yu. Shoeva<sup>1\*</sup>, Hans-Peter Mock<sup>2</sup>, Tatjana V. Kukoeva<sup>1</sup>, Andreas Börner<sup>2</sup>, Elena K. Khlestkina<sup>1,3</sup>

**1** Institute of Cytology and Genetics (ICG), Siberian Branch, Russian Academy of Sciences, Novosibirsk, Russia, **2** Leibniz Institute of Plant Genetics and Crop Plant Research (IPK), Gatersleben, Germany, **3** Food Security Research Center, Novosibirsk State University (NSU), Novosibirsk, Russia

\* [olesya\\_ter@bionet.nsc.ru](mailto:olesya_ter@bionet.nsc.ru)



CrossMark  
click for updates

OPEN ACCESS

**Citation:** Shoeva OY., Mock H-P, Kukoeva TV, Börner A, Khlestkina EK (2016) Regulation of the Flavonoid Biosynthesis Pathway Genes in Purple and Black Grains of *Hordeum vulgare*. PLoS ONE 11(10): e0163782. doi:10.1371/journal.pone.0163782

**Editor:** Dragan Perovic, Institute for Resistance Research and Stress Tolerance, GERMANY

**Received:** May 2, 2016

**Accepted:** September 14, 2016

**Published:** October 5, 2016

**Copyright:** © 2016 Shoeva et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its Supporting Information files.

**Funding:** This work was supported by Russian Foundation for Basic Research (grant no. 16-34-60052, <http://www.rfbr.ru/rffi/eng>, OYuS) and Institute of Cytology and Genetics SB RAS project (No 0324-2015-0005, <http://www.bionet.nsc.ru>, EKK, TVK). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

## Abstract


Barley grain at maturity can have yellow, purple, blue, and black pigmentations which are suggested to play a protective role under stress conditions. The first three types of the colors are caused by phenolic compounds flavonoids; the last one is caused by phytomelanins, oxidized and polymerized phenolic compounds. Although the genetic basis of the flavonoid biosynthesis pathway in barley has been thoroughly studied, there is no data yet on its regulation in purple and black barley grains. In the current study, genetic model of *Hordeum vulgare* ‘Bowman’ near-isogenic lines (NILs) was used to investigate the regulation of the flavonoid biosynthesis in white, purple, and black barley grains. Microsatellite genotyping revealed donor segments in the purple- and black-grained lines on chromosomes 2H (in region of the *Ant2* gene determining purple color of grains) and 1H (in region of the *Blp* gene determining black lemma and pericarp), respectively. The isolated dominant *Ant2* allele of the purple-grained line has high level of sequence similarity with the recessive Bowman’s *ant2* in coding region, whereas an insertion of 179 bp was detected in promoter region of *ant2*. This structural divergence between *Ant2* and *ant2* alleles may underlie their different expression in grain pericarp: Bowman’s *Ant2* is not transcribed, whereas it was up-regulated in the purple-grained line with coordinately co-expressed flavonoid biosynthesis structural genes (*Chs*, *Chi*, *F3h*, *F3’h*, *Dfr*, *Ans*). This led to total anthocyanin content increase in purple-grained line identified by ultra-performance liquid chromatography (HPLC). Collectively, these results proved the regulatory function of the *Ant2* gene in anthocyanin biosynthesis in barley grain pericarp. In the black-grained line, the specific transcriptional regulation of the flavonoid biosynthesis pathway genes was not detected, suggesting that flavonoid pigments are not involved in development of black lemma and pericarp trait.

**Competing Interests:** The authors have declared that no competing interests exist.

## Introduction

Barley (*Hordeum vulgare* L.,  $2n = 2x = 14$ , HH) grain at maturity may have different pigmentations. The most studied ones are yellow, purple, red, and blue caused by different subgroups of flavonoid compounds (Fig 1). The yellow color is due to proanthocyanidins synthesized in seed coat (testa layer) [1]; purple and red pigments are anthocyanins synthesized in pericarp and glumes [2]; blue color is caused by anthocyanins synthesized in aleurone layer of the grain [2]. In white barley grains, the pigments are absent. Scientific interest to the pigments is caused by their protective functions under different environments [3] as well as their undoubted health benefit [4]. In barley, flavonoid biosynthesis metabolic pathway is well studied [5]; structural genes encoding enzymes of the pathway, as well as regulatory genes, predetermining temporal and spatial patterns of the structural gene expressions have been described (S1 Table).

Chalcone synthase (CHS) represents in the barley genome a gene family with at least seven copies [6]. One copy of the gene has been isolated by heterologous hybridization method using a cDNA clone from *Antirrhinum majus* as a probe [7, 8]. Another copy of the gene with unusual substrate preference has been identified in pathogen-induced barley leaves cDNA



**Fig 1. Flavonoid biosynthetic pathway in plants.** The enzymes are: CHS (chalcone synthase); CHI (chalcone-flavanone isomerase); F3H (flavanone 3-hydroxylase); FLS (flavonol synthase); FNS (flavone synthase); F3'H (flavonoid 3'-hydroxylase); F3'5'H (flavonoid 3',5'-hydroxylase); DFR (dihydroflavonol 4-reductase); ANS (anthocyanidin synthase); GT (glycosyltransferase); MT (methyltransferase), RT (rhamnosyltransferase); and LAR (leucoanthocyanidin reductase).

doi:10.1371/journal.pone.0163782.g001

library [6]. One of the *Chs* gene copies has been mapped to the short arm of chromosome 1H using gDNA clone from barley genome as a probe [9]. Three non-overlapping genetic markers for the *Chs* gene have been mapped to chromosomes 1HS, 1HL, and 6HS [10]. The chalcone-flavanone isomerase gene (*Chi*) has been identified and mapped to the long arm of chromosome 5H using Southern blot technic with nucleotide sequence of the maize *Chi* gene as a probe [11]. Flavanone 3-hydroxylase gene (*F3h*) has been identified in cDNA barley library using cDNA probe from *A. majus* [12]. The gene has been mapped to chromosome 2HL [10, 13]. A full-length cDNA copy of the barley dihydroflavanol reductase gene (*Dfr*) has been isolated from kernel-specific cDNA library using cDNA of the maize *Dfr* gene as a probe [14]. The gene has been localized on the long arm of chromosome 3H [10]. Possible nucleotide sequence for the barley leucoanthocyanidin reductase (LAR) has been identified among plant expressed sequence tag (EST) sequences by querying the EST database with the Desmodium protein sequence using the tBLASTn procedure [15]. The *Lar* gene has not been located in barley genome yet. The full-length nucleotide sequence of the gene for the UDP glucose:flavonol 3-O-glucosyltransferase (UFGT) has been isolated from barley genome by Southern blot method with the nucleotide sequence of the maize *Ufgt* gene as a probe. The gene has been mapped to the short arm of chromosome 7H [16]. The full-length nucleotide sequences of the flavonoid 3'-hydroxylase (*F3'h*) and anthocyanidin synthase (*Ans*) genes were identified in the current study (S1 and S2 Files, respectively). The identified contigs of the *F3'h* and *Ans* genes are localized on chromosomes 1H and 5HL, respectively.

Among barley flavonoid biosynthesis regulatory genes full-length nucleotide sequences have been recently isolated for the *Ant1* [17, 18], *Ant2* [19] and *Ant28* [20] genes.

In *ant1* mutants, anthocyanin pigments are not observed in stem, auricles, awns, or lemma [21]. The *Ant1* gene encodes for a R2R3 MYB-type transcription factor that activates transcription of the anthocyanin biosynthesis structural genes (*Chi*, *F3h*, *Dfr*, *Ans*) in leaf sheaths [18]. The gene has been located on chromosome 7HS [17, 22].

The *Pre2* gene controls purple lemma and pericarp trait development in barley [23]. The gene has been mapped to the short arm of chromosome 2H [24], very close to the microsatellite marker linked with the *Ant2* gene, which encodes for a transcription factor with a basic Helix-Loop-Helix domain (bHLH) and is suggested to be a regulatory gene for the anthocyanin biosynthesis pathway [19, 25]. In *ant2* mutants, anthocyanin pigments are not observed in stem, auricles, awns, or lemma [21]. The *Ant2* gene is expressed in auricles, awns, and lemma [19]; however its influence on the structural genes expression has not been shown yet.

In *ant28* mutants, proanthocyanidins are not synthesized in grain, whereas wild-type level of anthocyanin content is retained in the vegetative tissue [5]. This data in addition to the observation of reduced enzymatic activities of DFR and LAR (which are essential for proanthocyanidin synthesis, Fig 1) in the *ant28* mutant are suggested that *Ant28* is a specific regulator of proanthocyanidin synthesis [5]. The gene has been mapped to the long arm of chromosome 3H [20, 26].

Blue aleurone color occurs in the presence of the five complementary dominant genes *Blx1*, *Blx2*, *Blx3*, *Blx4* and *Blx5* [27]. *Blx1*, *Blx3* and *Blx4* are closely linked to each other and mapped to chromosome 4HL. *Blx2* and *Blx5* have been assigned to chromosome 7HL [27]. The *Blx* genes have not been sequenced yet and their certain functions in the synthesis of blue anthocyanin substances have not been studied thus far.

Barley grain may have also black pigmentation which is caused by phytomelanins synthesized in glumes and/or pericarp [2]. The trait is controlled by the *Blp* locus, mapped to chromosome 1HL [23].

In the current study, we used Bowman's NILs [28] to reveal the specific features of the anthocyanin biosynthesis regulation in barley grain and to understand the involvement of the flavonoid biosynthesis pathway in development of black lemma and pericarp trait.

## Materials and Methods

### Plant material, DNA/RNA extraction, cDNA synthesis

A set of the barley cultivar Bowman near-isogenic lines (NILs) were used in the study: PLP (NGB22213) having purple pericarp, BLP (NGB20470) having black lemma and pericarp, and cultivar Bowman (NGB22812) having green lemma and uncolored pericarp (S1 Fig). The set of the lines was provided by the Nordic Gene Bank (NGB, [www.nordgen.org](http://www.nordgen.org)). The plants were grown using resources of ICG Greenhouse Core Facilities (Novosibirsk, Russia) under 12 h of light per day at 20–25°C. DNA was extracted from fresh leaves of plants following [29]. Lemma and pericarp for RNA extraction were peeled by scalpel from grains at early dough stage of maturity simultaneously for all genotypes. RNA was extracted applying a Zymo Research Plant RNA MiniPrep™ kit ([www.zymoresearch.com](http://www.zymoresearch.com)) followed by DNase treatment. A 180 ng aliquot of total RNA was converted to single-stranded cDNA via reverse transcription using a Fermentas RevertAid™ first strand cDNA synthesis kit ([www.thermoscientificbio.com/fermentas/](http://www.thermoscientificbio.com/fermentas/)) primed with (dT)<sub>15</sub> in a 20 μl reaction volume.

### Microsatellite genotyping

Primers to barley microsatellite loci mapped to chromosome 1H (*Xbmac0032*, *Xbmac0063*, *Xbmac0090*, *Xbmac0154*, *Xbmac0211*, *Xbmac0399*, *Xbmac0579*, *Xbmac0872*, *Xgbms0012*, *Xgbms0037*, *Xgbms0054*, *Xgbms0062*, *Xgbms0184* [30, 31]) and 2H (*Hvm23*, *Hvm54*, *Xbmac0125*, *Xbmac0140*, *Xbmac0607* [31]) were kindly provided by Dr Marion Röder (IPK-Gatersleben, Germany). The PCR-conditions were as described in [32]. Amplicons were separated through 5% ACTGene agarose gels (ACTGene, Inc., Piscataway, NJ, USA). Data on marker localization were taken from Barley consensus map, 2007 (chromosome 1H) and Barley consensus map, 2003 (chromosome 2H) deposited at web-site: [wheat.pw.usda.gov/GG2/](http://wheat.pw.usda.gov/GG2/).

### The *Ant2* gene re-sequencing and sequence analysis

The nucleotide sequence of contig 857662 corresponding to the *Ant2* gene of barley cultivar Bowman (S3 File) was found in BARLEX database (<http://apex.ipk-gatersleben.de/apex/f?p=284:10> [33]) using nucleotide sequence of the barley *Ant2* gene encoding protein with bHLH domain (GenBank accession HM370298 [19]) as a query. The full-length *Ant2* sequence of the PLP line was re-constructed from a series of the overlapping amplicons covering the relevant stretch of genomic DNA. The amplicons were generated by primer pairs described by Cockram et al. [19] and by overlapping primer pairs covering 5' regulatory region designed in the current study (S2 Table). The primers were designed using OLISO software [34]. Amplification of gDNA templates was performed in 20 μl PCRs each containing 1 U *Taq* DNA polymerase (Medigen, Novosibirsk, Russia), 1× PCR buffer (Medigen), 1.8 mM MgCl<sub>2</sub>, 0.2 mM dNTP and 0.25 μM of each primer. Amplification program was initiated with a denaturing step of 95°C / 2 min, 45 cycles of 95°C / 45 s, 55°C / 1 min, 72°C / 2 min, and a final extension of 72°C / 10 min. The amplified fragments were purified from a 2% agarose gel, using a DNA Clean kit (Cytokine, St. Petersburg, Russia), and then sequenced in both directions. DNA sequencing was performed by SB RAS Genomics Core Facilities (Novosibirsk, Russia, <http://sequest.niboch.nsc.ru>). Multiple sequence alignments were carried out using Multalin v5.4.1 software [35]. Secondary structures of the transcription factors were predicted by the Chou & Fasman Secondary Structure Prediction Server (CFSSP, <http://cho-fas.sourceforge.net/index.php> [36]). Structural elements of promoters were identified using the PLACE database (<http://www.dna.affrc.go.jp/PLACE/signalscan.html> [37, 38]).

## qRT-PCR

qRT-PCR was performed with the primers amplifying parts of the flavonoid biosynthesis structural genes *Chs*, *Chi*, *F3h*, *Dfr* [18], *F3'h*, *Ans*, and regulatory gene *Ant2* (F-SSR/R-SSR [19]). *F3'h* (Forward 5'-gccaggagttcaaggaca-3', Reverse 5'-ctcgctgatgaatccgtcca-3') primers were designed based on the barley predicted *F3'h* sequence AK363912 [39] found in NCBI database using BLASTp algorithm with *Zea mays F3'h* gene sequence HQ699781 [40] as a query (S1 File). *Ans* (Forward 5'-aagaggagtgaggaggacta-3', Reverse 5'-cgaggagaggatggcga-3') primers were designed based on the contigs identified in the BARLEX database using wheat *Ans* sequence AB247918 as a query (S2 File). A fragment of the *Actin* gene sequence was used for reference purposes [41]. The amplifications were performed in an ABI Prism 7000 Sequence Detection System (Applied Biosystems, <http://www.lifetechnologies.com>) applying a SYNTOL SYBR Green I kit (<http://www.syntol.ru/productmix.htm>). Pre-determined amounts of cloned cDNA were used to generate standard curves. Each sample was run in three replicates. The differences among the lines were tested by Mann-Whitney *U*-test, taking  $p \leq 0.05$  as significant.

## Chromatographic analysis of anthocyanins and related phenylpropanoids

Mature seeds were pulverized in a Retsch mill (Haan, Germany) and aliquots of 180 mg were extracted using 900  $\mu$ l of 70% MeOH with 2% formic acid. After centrifugation (28,000 g at 4°C for 10 min), the pellet was re-extracted with 600  $\mu$ l extraction solvent and centrifuged. Supernatants were combined and reduced under vacuum centrifugation to a final volume of 500  $\mu$ l. Then 200  $\mu$ l of extract were mixed with eluent A (see below) and stored at 4°C until analysis in the dark. Prior to chromatography, extracts were centrifuged again and filtered (Ultrafree MC-HV, PVDF, 0.45  $\mu$ m; Merck-Millipore, Darmstadt, Germany). The supernatant was filled into vials and 5  $\mu$ l were injected. Two independent extraction procedures were performed for each genotype studied.


Profiles of phenylpropanoids were obtained by using ultra-performance liquid chromatography (UPLC) in combination with photodiode array (PDA) as well as MS detection of separated compounds essentially with the system described by Petridis et al. [42] without recording fluorescent traces. Diode array data were extracted at 280 nm for phenolic compounds in general and at 515 nm for anthocyanins. The gradient was modified as follows: starting concentration was 98% eluent A (0.1% v/v formic acid in MS grade water) and 2% eluent B (0.1% v/v formic acid in acetonitrile, MS grade). 10% B were reached after 3 min, 20% B after 7 min and 98% B after 10 min in a linear mode. The column was flushed for further 3 min at this concentration and then equilibrated at the initial conditions prior to injection of the next sample. Anthocyanins were quantified using a calibration curve obtained by injecting cyanidin 3-*O*-glucoside as a standard at different concentrations with the help of the diode array data. A mass range from 50 to 1300 m/z was recorded.

## Results

### Microsatellite genotyping of the barley NILs

Microsatellite genotyping of the barley NILs showed, that the PLP line has donor fragment located on chromosome 2HL, in the region of the microsatellite locus *Xbmag140*, near the *Ant2* locus, predetermining purple lemma and pericarp (Fig 2A).

In the BLP line, donor segment was located between microsatellites *Xgbms0012* and *Xgbms0184*, near the *Blp* gene, controlling black lemma and pericarp trait (Fig 2B). Thus, the NILs carry donor fragments with the genes, predetermining different colors of the barley lemma and pericarp.



**Fig 2. Microsatellite genotyping of the PLP (A) and BLP (B) lines.** Purple and black colors mark introgressed fragments in the PLP and BLP lines, respectively.

doi:10.1371/journal.pone.0163782.g002

### Structural and functional comparison of the different *Ant2* alleles

The nucleotide sequences of the *Ant2* genes of Bowman and the PLP line were determined and compared. The full-length nucleotide sequence 6,419 bp in length for the Bowman *Ant2* gene was found in the BARLEX database (S3 File). The PLP *Ant2* nucleotide sequence 5,066 bp in


length was obtained by sequencing overlapping PCR products (the sequence was submitted to NCBI GenBank under accession number KX035100). The coding sequences of Bowman and PLP *Ant2* genes are split into eight exons with total length 1,680 bp (S2 Fig).

The *Ant2* nucleotide sequence identity between Bowman and PLP was 98% (the full-length exonic-intronic sequences were compared). The observed polymorphism between Bowman and PLP *Ant2* alleles (7 synonymous and 6 non-synonymous substitutions in the coding region and 45 SNPs and 2 indels 4, 5 bp in length within the intronic region) suggested the PLP to be a carrier of the *Ant2* allele from a donor parent (S2 Fig).

The predicted polypeptide sequence is 559 amino acid residues. Comparison with the known transcription factor LC, regulating anthocyanin biosynthesis in maize [43] revealed the conservative basic helix-loop-helix (bHLH) domain important for DNA binding and protein-protein interaction (S3 Fig). The observed non-synonymous substitutions between Bowman and PLP lay outside of the bHLH domain and are assumed to have no essential effect on the functionality of ANT2 since these substitutions are very similar by the level of hydrophobicity (S3 Table) and not changed the predicted secondary structure of the proteins (S4 Fig).

The *Ant2* genes of Bowman and PLP differ significantly in the 5' regulatory region. In Bowman, a 179 bp insertion was identified in comparison with the PLP *Ant2* gene (S5 Fig). The insertion disturbs the order and spatial localization of the *cis*-acting regulatory elements identified in promotor regions of the genes. Among the identified elements those related to different stimuli such as light, biotic and abiotic stress factors, phytohormones and elements recognized by other regulatory factors participating in physiological functions have been identified (S4 Table).

The expression assay of the *Ant2* gene in grain of Bowman and the PLP line shows that level of the *Ant2* gene transcription was almost 17-fold abundant in the PLP line in comparison with Bowman and the BLP NIL (Fig 3). The data suggested that the rearrangements in



**Fig 3. Expression of the *Ant2* gene in lemma and pericarp of NILs differing by the coloration.** The data are presented as mean value  $\pm$  standard error. \*—differences are statistically significant between NILs and Bowman at  $p \leq 0.05$  (*U*-test).

doi:10.1371/journal.pone.0163782.g003

promotor region of the *Ant2* gene lead to reduction of transcription level of *Ant2* in Bowman, resulting in the uncolored grain phenotype.

## Expression of the flavonoid biosynthesis structural genes in lemma and pericarp of differently colored barley grain

To show regulatory functions of the *Ant2* gene and to investigate the involvement of the flavonoid pathway in black lemma and pericarp trait development the expression analysis of the flavonoid biosynthesis structural genes was performed in NILs having green, purple and black color of lemma and pericarp. In the purple-grained PLP line, expression of the structural genes *Chs*, *Chi*, *F3h*, *F3'h*, *Dfr*, and *Ans* was up-regulated in comparison with the green and black grains (Fig 4). All the genes were co-regulated as one regulatory unit. The data proved the regulatory functions of the *Ant2* gene in anthocyanin biosynthesis in barley grain pericarp.

In the black-grained BLP line, no differentially expressed genes (with the exception of *Chi*) in comparison with Bowman were found (Fig 4). The data demonstrated that there is no specific transcriptional regulation of the flavonoid biosynthesis in the black-grained barley line, suggesting that flavonoid pigments are not involved in development of black lemma and pericarp trait. Thus, distinct genetic networks underlay black and purple pigmentation of barley grain.

## Profiles of anthocyanins and related phenylpropanoids

In order to correlate expression of the flavonoid biosynthesis pathway genes with metabolic profiles, methanolic whole seed extracts were separated by UPLC and compounds were detected by absorption as well as by mass spectrometry. Six peaks differing in elution time and mass could be detected in anthocyanin profile for the PLP genotype (Fig 5). In seed extracts of Bowman and BLP, only one compound corresponding in mass and elution time to peak two of PLP was found. Calculation of anthocyanin contents based on total peak areas showed similar low amounts in Bowman and BLP and about nine times higher value for anthocyanin content in the PLP line (S5 Table), consistent with the gene expression data (Figs 3 and 4).


Analysis of the PDA data at 280 nm revealed complex metabolic profiles of the three genotypes, representing phenolic compounds in general, but not excluding other cellular compounds with similar polarity (S6 Fig). Although the metabolic patterns of all three genotypes appeared similar, genotype BLP showed higher abundance in many of the individual compounds detected at 280 nm.

## Discussion

### New polymorphism underlies different grain color phenotype

In the current study, new polymorphism underlying different grain coloration in barley was revealed. The polymorphism is suggested to be associated with the rearrangement in the 5' regulatory region of the *Ant2* gene in white-grained cultivar Bowman in comparison with its purple-grained NIL PLP. The rearrangement leads to reduction of the transcription level of the *Ant2* genes in Bowman (Fig 3) resulting in a decline of the anthocyanin biosynthesis structural genes expressions (Fig 4), absence of anthocyanins (Fig 5) and hence conferring a white-grained phenotype (S1 Fig).


Earlier the diagnostic marker distinguishing dominant and recessive alleles of the *Ant2* genes (*Ant2* / *ant2*) has been developed. The marker has been designed to a region carrying a 16-bp deletion within exon 6 that results in truncation of the predicted protein upstream of the bHLH domain [19]. Although the deletion has been identified in all 21 white-grained genotypes studied by Cockram et al. [19] and perfectly co-segregated with *ant2* alleles in biparental



**Fig 4. Expression of the flavonoid biosynthesis structural genes in grains of the barley NILs having different coloration of lemma and pericarp.** The data are presented as mean value  $\pm$  standard error. \*—differences are statistically significant between NILs and Bowman at  $p \leq 0.05$  (*U*-test).

doi:10.1371/journal.pone.0163782.g004

mapping population, it appeared to be not common for further white-grained barley cultivars, such as Bowman (S2 Fig). Also the *Ant2* marker was monomorphic between purple-grained Yuyaohongdamai and white-grained ACCA [24]. The authors [24] have been focused on the fine mapping of the *Pre2* gene which controls purple lemma and pericarp trait [23]. They have found close linkage (0.1 cM) of *Pre2* with the *HvOs04g47170* marker linked to *Ant2* allowed suggesting that *Pre2* is synonymous to *Ant2*, but because the *Ant2* marker has revealed no differences between the cultivars having different colors of grain the authors have stated that the gene controlling purple color in grain of Yuyaohongdamai is not *Ant2*. The transcriptional activity assay of the *Ant2* gene in purple-grained variety Retriever and white-grained variety Saffron has shown that the gene is transcribed in auricles, awns, and lemma of both cultivars; moreover transcriptional level of the *Ant2* gene did not differ between the cultivars [19].



**Fig 5. Anthocyanin profiles of Bowman (A), PLP (B) and BLP (C) genotypes.** Seed extracts were prepared using acidified aqueous methanol as described in the materials and methods section. Extracts were separated by UPLC and compound elution was monitored by photodiode array (PDA) detection followed by MS analysis. The chromatograms were obtained by extracting the PDA data at 515 nm. X-axis represents time (min) and Y-axis represents absorbance in milliabsorbance units (mAU).

doi:10.1371/journal.pone.0163782.g005

Such different results on transcriptional activity of the *Ant2* gene in white-grained vs purple-grained lines/cultivars obtained in the current study (Fig 3) and by Cockram et al. [19] can be explained by different mutations in the *Ant2* gene leading to the white-grained phenotype: Cockram et al. [19] have revealed mutation causing structural damage of the ANT2 protein, but it did not influence the transcription of the gene; our data demonstrate another mutation which caused damage in transcriptional regulation of the *Ant2* gene but did not have influence on the structure of the protein.

Overall, the data on co-localization of the *Ant2* and *Pre2* genes, the absence of another bHLH-type gene in the region of localization of the *Ant2* gene [19, 24] and the differences in functional activity of the *Ant2* gene in colored and uncolored barley grains (Fig 3) suggest together that purple color of barley grain is controlled by *Ant2* that is synonymous to the *Pre2* gene.

### Co-regulation of the anthocyanin biosynthesis pathway genes in barley grain

Although the molecular-genetic basis for anthocyanin biosynthesis is common for the plant species: structural genes encoding for enzymes of the pathway (Fig 1) are regulated by transcriptional factors of MYB-, bHLH- (MYC-) and WD40-types [44], some species-specific differences have been noted in the regulation of the structural genes [45, 46]. Depending on co-expression of the structural genes in the presence of the dominant alleles of the regulatory genes following patterns of the anthocyanin biosynthesis regulation could be distinguished: (1) the set of the genes is co-regulated separately from the others genes (the subsets of the 'early' and 'late' genes); (2) anthocyanin biosynthesis is regulated at a certain stage of the pathway; (3) whole set of the genes is co-regulated as one regulatory unit.

The first regulatory pattern, for example, has been observed in petals of *Petunia hybrida* where the 'late' genes *Dfr*, *Ans*, *Mt*, and *Rt* were co-regulated by transcriptional factors AN1, AN2, and AN11 (bHLH, MYB, and WD40, respectively) separately from the 'early' genes *Chs*, *Chi*, and *F3h* [45]. In flowers of *Antirrhinum majus*, the 'late' genes *F3h*, *Dfr*, *Ans*, and *Ufgt* where co-expressed in the presence of DELILA (bHLH type) regulatory factor separately from the 'early' genes *Chs*, *Chi* [47]. In seedlings of *Oryza sativa*, *F3h*, *Dfr*, and *Ans* were co-regulated by the OSB2 (bHLH-type) transcriptional regulator separately from *Chs*, *Chi*, *F3'h* [48, 49].

The second pattern of the anthocyanin pathway regulation is similar to the first one, but characterized by specific regulation of one of the structural genes in dependence of the dominant allele of the regulatory genes. Such regulation pattern was observed in berries of *Vitis vinifera* [50, 51] and pericarp of *Litchi chinensis* fruits [52, 53], where anthocyanin biosynthesis is regulated at a stage of the *Ufgt* gene expression. In grain pericarp of *Triticum aestivum*, the pathway is regulated at a step of *F3h* gene expression [54].

The third regulatory pattern has been revealed in aleurone layer of kernel and husk of *Zea mays*, where *Chs*, *Dfr*, and *Ufgt* were co-regulated in the presence of the dominant allele *R(S)/C1* and *B/Pl* (MYB and bHLH, respectively) [55, 56]. As shown in the current study, the anthocyanin biosynthesis structural genes in barley grain are co-regulated as one regulatory unit in the presence of the ANT2 (bHLH) regulatory factor. Similar phenomenon has been observed by Meldgaard M. [12], who has studied transcription of the *Chs*, *F3h*, and *Dfr* genes in testa pericarp tissues in the proanthocyanidin-free barley *ant13* mutant line. This regulatory pattern is characteristic not only to such monocot plant species as maize and barley. The whole set of the genes is co-expressed in dicot plants having parts colored by anthocyanins such as flash of tuberous roots in *Ipomoea batatas* [57], skin of apple fruits [58], or taproots in *Daucus carota* [59, 60].

As seen the regulation pattern of anthocyanin biosynthesis in barley grain is not unique in comparison with the other plant species but it is quite different from such cereal relatives as wheat and rice.

### 'Pure' melanin-like pigmentation of the black grains of barley

Black color of barley lemma and pericarp is caused by melanin-like pigments, called phytomelanins [2]. Plant 'Melanins' is a general name for groups of high-molecular black/brown pigmented irregular polymers, arising in the course of oxidation and polymerization of phenolic

compounds. Although black pigmentation is considered to be an important agronomic trait because of its protective functions against severe environments and infections [61–63] chemical nature of the black pigments and the biosynthesis pathway leading to them is not clear [61, 64].

Some studies of phytochemical composition of barley grains with black color have revealed that the real melanin-like black pigments can be mixed with anthocyanins and other related copigments that contribute to the total content of phenols [65, 66] and prevent determination of 'pure' melanin-like compounds and their precursors in barley.

In the current study, the comparative chromatographic analysis of whole seed extracts and transcriptional assay of the flavonoid biosynthesis pathway genes was performed for the NILs having different colors of pericarp and lemma (S1 Fig). Although the minor peak corresponding to the anthocyanins was detected in whole seed extracts of black-grained line as well as of uncolored Bowman (Fig 5), none of the studied genes involved in flavonoid biosynthesis is expressed significantly in lemma and pericarp of these genotypes (Figs 3 and 4). This allows suggesting that anthocyanins and the other flavonoids unlikely participate in black pigmentation of barley lemma and pericarp and identified peak may belong to compound from the other parts of grain such as seed coat, aleurone layer or endosperm. The lines studied have similar phenylpropanoid profiles with higher abundance in many of the individual compounds in BLP (S6 Fig). Further studies are needed to address the question about the chemical composition of black barley grains and metabolic pathway leading to them. The characterized NILs are proper model for this goal.

## Supporting Information

**S1 Fig. Spikes and grains of Bowman (A), PLP (B) and BLP (C) lines used in the current study.**

(DOCX)

**S2 Fig. Alignment of the *Ant2* gene nucleotide sequences of Bowman and PLP line.** Exonic sequences are marked by green color.

(DOCX)

**S3 Fig. Alignment of the ANT2 proteins of Bowman and PLP and the related LC protein, regulating anthocyanin biosynthesis in maize.** Conservative basic helix-loop-helix (bHLH) domain is marked.

(DOCX)

**S4 Fig. Secondary structure of the ANT2 protein of Bowman and PLP predicted by CFSSP Server.** The residues differing in Bowman and PLP are marked by pink color.

(DOCX)

**S5 Fig. Alignment of the nucleotide sequences of the 5' regulatory region of the *Ant2* gene of Bowman and PLP line.** Exonic sequences are marked by green color.

(DOCX)

**S6 Fig. Phenylpropanoid profiles of Bowman, PLP and BLP seed extracts.** The overlay of chromatograms obtained by extracting the PDA data at 280 from extract 1 of all genotypes is shown. Chromatograms are depicted in black (Bowman), blue (PLP) and red (BLP).

(DOCX)

**S1 Table. Data on the structural and regulatory genes for flavonoid biosynthesis in barley identified up to date.**

(DOCX)

**S2 Table. Primer pairs amplifying *Ant2* gene 5' regulatory region designed in the current study.**

(DOCX)

**S3 Table. Amino acid substitutions identified between ANT2 of Bowman and PLP and its classification based on the polarity of the side chain group.**

(DOCX)

**S4 Table. Putative *cis*-acting regulatory elements identified in promoter regions of Bowman (936 bp before ATG codon) and PLP (754 bp before ATG codon) *Ant2* genes using the PLACE database.**

(DOCX)

**S5 Table. Quantification of total anthocyanins in grains of Bowman, PLP and BLP.** Peaks areas of PDA data at 515 nm were integrated. The sum of all peak areas was taken for calculating total anthocyanin contents. A calibration curve using authentic cyanidin 3-*O*-glucoside was used.

(DOCX)

**S1 File. The nucleotide sequences of Morex, Barke, and Bowman contigs corresponding to the *F3'h* gene found in BARLEX database.**

(DOCX)

**S2 File. The nucleotide sequences of Bowman contigs corresponding to the *Ans* gene found in BARLEX database. Reconstructed full-length nucleotide sequence of *Ans* is also present. Yellow color marks coding sequence of the gene.**

(DOCX)

**S3 File. The nucleotide sequence of Bowman contig corresponding to the *Ant2* gene found in BARLEX database. Exonic sequences are marked by green color.**

(DOCX)

## Acknowledgments

We thank Ms Galina Generalova (ICG SB RAS) and Mrs Elena Barthel (IPK Gatersleben) for technical assistance.

## Author Contributions

**Conceptualization:** EKK AB.

**Formal analysis:** OYuS HPM.

**Investigation:** OYuS TVK.

**Resources:** AB HPM.

**Writing – original draft:** OYuS.

## References

1. Aastrup S, Outtrup H, Erdal K. Location of the proanthocyanidins in the barley grain. *Carlsberg Res Commun.* 1984; 49(1):105–109. doi: [10.1007/BF02913969](https://doi.org/10.1007/BF02913969)
2. Harlan HV. Some distinctions in our cultivated barleys with reference to their use in plant breeding. *U S Dept Agric Bul.* 1914; 137. doi: [10.5962/bhl.title.109258](https://doi.org/10.5962/bhl.title.109258)

3. Khlestkina E. The adaptive role of flavonoids: emphasis on cereals. *Cereal Res Commun.* 2013; 41(2):185–198. doi: [10.1556/CRC.2013.0004](https://doi.org/10.1556/CRC.2013.0004)
4. Tarahovsky YS, Kim YA, Abdrasilov BS, Muzafarov EN. Flavonoids: biochemistry, biophysics, medicine. Pushchino: Synchronobook; 2013.
5. Jende-Strid B. Genetic control of flavonoid biosynthesis in barley. *Hereditas.* 1993; 119(2):187–204. doi: [10.1111/j.1601-5223.1993.00187.x](https://doi.org/10.1111/j.1601-5223.1993.00187.x)
6. Christensen AB, Gregersen PL, Schröder J, Collinge DB. A chalcone synthase with an unusual substrate preference is expressed in barley leaves in response to UV light and pathogen attack. *Plant Mol Biol.* 1998; 37(5):849–857. doi: [10.1023/A:1006031822141](https://doi.org/10.1023/A:1006031822141) PMID: [9678580](https://pubmed.ncbi.nlm.nih.gov/9678580/)
7. Rohde W, Barzen E, Marocco A, Schwarz-Sommer Z, Saedler H, Salamini F. Isolation of genes that could serve as traps for transposable elements in *Hordeum vulgare*. In: Yasuda, S, Konishi, T, editors. Barley genetics V: proceedings of the fifth international barley genetics symposium. Okayama; 1987. p. 553–541.
8. Rohde W, Dörr S, Salamini F, Becker D. Structure of a chalcone synthase gene from *Hordeum vulgare*. *Plant Mol Biol.* 1991; 16(6):1103–1106. doi: [10.1007/BF00016087](https://doi.org/10.1007/BF00016087) PMID: [1863766](https://pubmed.ncbi.nlm.nih.gov/1863766/)
9. Pecchioni N, Vale G, Toubia-Rahme H, Faccioli P, Terzi V, Delogu G. Barley—*Pyrenophora graminea* interaction: QTL analysis and gene mapping. *Plant Breed.* 1999; 118(1):29–35. doi: [10.1046/j.1439-0523.1999.118001029.x](https://doi.org/10.1046/j.1439-0523.1999.118001029.x)
10. Peukert M, Weise S, Röder MS, Matthies IE. Development of SNP markers for genes of the phenylpropanoid pathway and their association to kernel and malting traits in barley. *BMC Genet.* 2013; 14:97. doi: [10.1186/1471-2156-14-97](https://doi.org/10.1186/1471-2156-14-97) PMID: [24088365](https://pubmed.ncbi.nlm.nih.gov/24088365/)
11. Druka A, Kudrna D, Rostoks N, Brueggeman R, von Wettstein D, Kleinbartsch A. Chalcone isomerase gene from rice (*Oryza sativa*) and barley (*Hordeum vulgare*): physical, genetic and mutation mapping. *Gene.* 2003; 302(1–2):171–178. doi: [10.1016/S0378-1119\(02\)01105-8](https://doi.org/10.1016/S0378-1119(02)01105-8) PMID: [12527208](https://pubmed.ncbi.nlm.nih.gov/12527208/)
12. Meldgaard M. Expression of chalcone synthase, dihydroflavonol reductase, and flavanone-3-hydroxylase in mutants of barley deficient in anthocyanin and proanthocyanidin biosynthesis. *Theor Appl Genet.* 1992; 83(6–7):695–706. doi: [10.1007/BF00226687](https://doi.org/10.1007/BF00226687) PMID: [24202743](https://pubmed.ncbi.nlm.nih.gov/24202743/)
13. Khlestkina E, Salina E, Matthies I, Leonova I, Börner A, Röder M. Comparative molecular marker-based genetic mapping of flavanone 3-hydroxylase genes in wheat, rye and barley. *Euphytica.* 2011; 179(2):333–341. doi: [10.1007/s10681-010-0337-2](https://doi.org/10.1007/s10681-010-0337-2)
14. Kristiansen KN, Rohde W. Structure of the *Hordeum vulgare* gene encoding dihydroflavonol-4-reductase and molecular analysis of *ant18* mutants blocked in flavonoid synthesis. *Mol Gen Genet.* 1991; 230(1):49–59. doi: [10.1007/BF00290650](https://doi.org/10.1007/BF00290650) PMID: [1720864](https://pubmed.ncbi.nlm.nih.gov/1720864/)
15. Bogs J, Downey MO, Harvey JS, Ashton AR, Tanner GJ, Robinson SP. Proanthocyanidin synthesis and expression of genes encoding leucoanthocyanidin reductase and anthocyanidin reductase in developing grape berries and grapevine leaves. *Plant Physiol.* 2005; 139(2):652–663. doi: [10.1104/pp.105.064238](https://doi.org/10.1104/pp.105.064238) PMID: [16169968](https://pubmed.ncbi.nlm.nih.gov/16169968/)
16. Wise RP, Rohde W, Salamini F. Nucleotide sequence of the *Bronze-1* homologous gene from *Hordeum vulgare*. *Plant Mol Biol.* 1990; 14(2):277–279. doi: [10.1007/BF00018568](https://doi.org/10.1007/BF00018568) PMID: [2151660](https://pubmed.ncbi.nlm.nih.gov/2151660/)
17. Himi E, Taketa S. Isolation of candidate genes for the barley *Ant1* and wheat *Rc* genes controlling anthocyanin pigmentation in different vegetative tissues. *Mol Genet Genomics.* 2015; 290(4):1287–1298. doi: [10.1007/s00438-015-0991-0](https://doi.org/10.1007/s00438-015-0991-0) PMID: [25585663](https://pubmed.ncbi.nlm.nih.gov/25585663/)
18. Shoeva OY, Kukoeva TV, Börner A, Khlestkina EK. Barley *Ant1* is a homolog of maize *C1* and its product is part of the regulatory machinery governing anthocyanin synthesis in the leaf sheath. *Plant Breed.* 2015; 134(4):400–405. doi: [10.1111/pbr.12277](https://doi.org/10.1111/pbr.12277)
19. Cockram J, White J, Zuluaga DL, Smith D, Comadran J, Macaulay M, et al. Genome-wide association mapping to candidate polymorphism resolution in the unsequenced barley genome. *Proc Natl Acad Sci U S A.* 2010; 107(50):21611–21616. doi: [10.1073/pnas.1010179107](https://doi.org/10.1073/pnas.1010179107) PMID: [21115826](https://pubmed.ncbi.nlm.nih.gov/21115826/)
20. Himi E, Yamashita Y, Haruyama N, Yanagisawa T, Maekawa M, Taketa S. *Ant28* gene for proanthocyanidin synthesis encoding the R2R3 MYB domain protein (Hvmyb10) highly affects grain dormancy in barley. *Euphytica.* 2012; 188(1):141–151. doi: [10.1007/s10681-011-0552-5](https://doi.org/10.1007/s10681-011-0552-5)
21. Jende-Strid B, Lundqvist U. Diallelic tests of anthocyanin-deficient mutants. *Barley Genet Newsl.* 1978; 8:57–59.
22. Jende-Strid B. Coordinator's report: anthocyanin genes. *Barley Genet Newsl.* 1984; 14:76–79.
23. Franckowiak JD, Lundqvist U, Konishi T. New and revised descriptions of barley genes. *Barley Genet Newsl.* 1997; 26:22–516.
24. Jia Q, Zhu J, Wang J, Yang J, Zhang G. Genetic mapping and molecular marker development for the gene *Pre2* controlling purple grains in barley. *Euphytica.* 2016; 208(2):215–223. doi: [10.1007/s10681-015-1593-y](https://doi.org/10.1007/s10681-015-1593-y)

25. Lahaye T, Hartmann S, Töpsch S, Freialdenhoven A, Yano M, Schulze-Lefert P. High-resolution genetic and physical mapping of the *Rar1* locus in barley. *Theor Appl Genet.* 1998; 97(4):526–534. doi: [10.1007/s001220050927](https://doi.org/10.1007/s001220050927)
26. Garvin DF, Miller-Garvin JE, Viccars EA, Jacobsen JV, Brown AHD. Identification of molecular markers linked to *ant28-484*, a mutation that eliminates proanthocyanidin production in barley seeds. *Crop Sci.* 1998; 38(5):1250–1255. doi: [10.2135/cropsci1998.0011183X003800050023x](https://doi.org/10.2135/cropsci1998.0011183X003800050023x)
27. Finch RA, Simpson E. New colours and complementary colour genes in barley. *Z Pflanzenzüchtg.* 1978; 81(1):40–53.
28. Druka A, Franckowiak J, Lundqvist U, Bonar N, Alexander J, Houston K, et al. Genetic dissection of barley morphology and development. *Plant Physiol.* 2011; 155(2):617–627. doi: [10.1104/pp.110.166249](https://doi.org/10.1104/pp.110.166249) PMID: [21088227](https://pubmed.ncbi.nlm.nih.gov/21088227/)
29. Plaschke J, Ganai MW, Röder MS. Detection of genetic diversity in closely related bread wheat using microsatellite markers. *Theor Appl Genet.* 1995; 91(6–7):1001–1007. doi: [10.1007/BF00223912](https://doi.org/10.1007/BF00223912) PMID: [24169989](https://pubmed.ncbi.nlm.nih.gov/24169989/)
30. Li JZ, Sjakste TG, Röder MS, Ganai MW. Development and genetic mapping of 127 new microsatellite markers in barley. *Theor Appl Genet.* 2003; 107(6):1021–1027. doi: [10.1007/s00122-003-1345-6](https://doi.org/10.1007/s00122-003-1345-6) PMID: [12879255](https://pubmed.ncbi.nlm.nih.gov/12879255/)
31. Ramsay L, Macaulay M, degli Ivanissivich S, MacLean K, Cardle L, Fuller J, et al. A simple sequence repeat-based linkage map of barley. *Genet.* 2000; 156(4):1997–2005. PMID: [11102390](https://pubmed.ncbi.nlm.nih.gov/11102390/)
32. Röder MS, Korzun V, Wendehake K, Plaschke J, Tixier M-H, Leroy P, et al. A microsatellite map of wheat. *Genet.* 1998; 149(4):2007–2023.
33. Colmsee C, Beier S, Himmelbach A, Schmutzer T, Stein N, Scholz U, et al. BARLEX—the barley draft genome explorer. *Mol Plant.* 2015; 8(6):964–966. doi: [10.1016/j.molp.2015.03.009](https://doi.org/10.1016/j.molp.2015.03.009) PMID: [25804976](https://pubmed.ncbi.nlm.nih.gov/25804976/)
34. Offerman JD, Rychlik W. Oligo primer analysis software. In: Krawetz SA, Womble DD, editors. *Introduction to bioinformatics: a theoretical and practical approach.* New York: Humana Press; 2003. p. 345–361. doi: [10.1007/978-1-59259-335-4](https://doi.org/10.1007/978-1-59259-335-4)
35. Corpet F. Multiple sequence alignment with hierarchical clustering. *Nucleic Acids Res.* 1988; 16(22):10881–10890. doi: [10.1093/nar/16.22.10881](https://doi.org/10.1093/nar/16.22.10881) PMID: [2849754](https://pubmed.ncbi.nlm.nih.gov/2849754/)
36. Chou PY, Fasman GD. Prediction of protein conformation. *Biochem.* 1974; 13(2):222–245. doi: [10.1021/bi00699a002](https://doi.org/10.1021/bi00699a002)
37. Higo K, Ugawa Y, Iwamoto M, Korenaga T. Plant *cis*-acting regulatory DNA elements (PLACE) database. *Nucleic Acids Res.* 1999; 27(1):297–300. doi: [10.1093/nar/27.1.297](https://doi.org/10.1093/nar/27.1.297) PMID: [9847208](https://pubmed.ncbi.nlm.nih.gov/9847208/)
38. Prestridge DS. SIGNAL SCAN: a computer program that scans DNA sequences for eukaryotic transcriptional elements. *Comput Appl Biosci.* 1991; 7(2):203–206. doi: [10.1093/bioinformatics/7.2.203](https://doi.org/10.1093/bioinformatics/7.2.203) PMID: [2059845](https://pubmed.ncbi.nlm.nih.gov/2059845/)
39. Matsumoto T, Tanaka T, Sakai H, Amano N, Kanamori H, Kurita K, et al. Comprehensive sequence analysis of 24,783 barley full-length cDNAs. *Plant Physiol.* 2011; 156(1):20–28. doi: [10.1104/pp.110.171579](https://doi.org/10.1104/pp.110.171579) PMID: [21415278](https://pubmed.ncbi.nlm.nih.gov/21415278/)
40. Sharma M, Cortes-Cruz M, Ahern KR, McMullen M, Brutnell TP, Chopra S. Identification of the *pr1* gene product completes the anthocyanin biosynthesis pathway of maize. *Genet.* 2011; 188(1):69–79. doi: [10.1534/genetics.110.126136](https://doi.org/10.1534/genetics.110.126136) PMID: [21385724](https://pubmed.ncbi.nlm.nih.gov/21385724/)
41. von Zitzewitz J, Szucs P, Dubcovsky J, Yan L, Francia E, Pecchioni N, et al. Molecular and structural characterization of barley vernalization genes. *Plant Mol Biol.* 2005; 59(3):449–467. doi: [10.1007/s11103-005-0351-2](https://doi.org/10.1007/s11103-005-0351-2) PMID: [16235110](https://pubmed.ncbi.nlm.nih.gov/16235110/)
42. Petridis A, Döll S, Nichelmann L, Bilger W, Mock H-P. *Arabidopsis thaliana* G2-LIKE FLAVONOID REGULATOR and BRASSINOSTEROID ENHANCED EXPRESSION1 are low-temperature regulators of flavonoid accumulation. *New Phytol.* 2016; 211(3):912–925. doi: [10.1111/nph.13986](https://doi.org/10.1111/nph.13986) PMID: [27125220](https://pubmed.ncbi.nlm.nih.gov/27125220/)
43. Ludwig SR, Habera LF, Dellaporta SL, Wessler SR. *Lc*, a member of the maize *R* gene family responsible for tissue-specific anthocyanin production, encodes a protein similar to transcription activators and contains the *myc*-homology region. *Proc Natl Acad Sci U S A.* 1989; 86(18):7092–7096. doi: [10.1073/pnas.86.18.7092](https://doi.org/10.1073/pnas.86.18.7092) PMID: [2674946](https://pubmed.ncbi.nlm.nih.gov/2674946/)
44. Mol J, Grotewold E, Koes R. How genes paint flowers and seeds. *Trends Plant Sci.* 1998; 3(6):212–217. doi: [10.1016/S1360-1385\(98\)01242-4](https://doi.org/10.1016/S1360-1385(98)01242-4)
45. Quattrocchio F, Wing JF, Leppen HTC, Mol JNM, Koes RE. Regulatory genes controlling anthocyanin pigmentation are functionally conserved among plant species and have distinct sets of target genes. *Plant Cell.* 1993; 5(11):1497–1512. doi: [10.1105/tpc.5.11.1497](https://doi.org/10.1105/tpc.5.11.1497) PMID: [12271045](https://pubmed.ncbi.nlm.nih.gov/12271045/)

46. Adzhieva VF, Babak OG, Shoeva OY, Kilchevsky AV, Khlestkina EK. Molecular-genetic mechanisms underlying fruit and seed coloration in plants. *Vavilovskii Zhurnal Genetiki i Seleksii* (Vavilov Journal of Genetics and Breeding). 2015; 19(5):561–573. doi: [10.18699/VJ15.073](https://doi.org/10.18699/VJ15.073)
47. Martin C, Prescott A, MacKay S, Bartlett J, Vrijlandt E. Control of anthocyanin biosynthesis in flowers of *Antirrhinum majus*. *Plant J*. 1991; 1(1):37–49. PMID: [1844879](https://pubmed.ncbi.nlm.nih.gov/1844879/)
48. Sakulsingharoj C, Inta P, Sukkasem R, Pongjaroenkit S, Chowpongpan S, Sangtong V. Overexpression of *OSB2* gene in transgenic rice up-regulated expression of structural genes in anthocyanin biosynthesis pathway. *Thai J Genet*. 2014; 7(3):173–182.
49. Shih CH, Chu H, Tang LK, Sakamoto W, Maekawa M, Chu IK, et al. Functional characterization of key structural genes in rice flavonoid biosynthesis. *Planta*. 2008; 228(6):1043–1054. doi: [10.1007/s00425-008-0806-1](https://doi.org/10.1007/s00425-008-0806-1) PMID: [18726614](https://pubmed.ncbi.nlm.nih.gov/18726614/)
50. Boss PK, Davies C, Robinson SP. Expression of anthocyanin biosynthesis genes in red and white grapes. *Plant Mol Biol*. 1996; 32(3):565–569. doi: [10.1007/BF00019111](https://doi.org/10.1007/BF00019111) PMID: [8980508](https://pubmed.ncbi.nlm.nih.gov/8980508/)
51. Ramazzotti S, Filippetti I, Intrieri C. Expression of genes associated with anthocyanin synthesis in red-purple, pink, pinkish-green and green grape berries from mutated 'Sangiovese' biotypes: a case study. *Vitis*. 2008; 47(3):147–151.
52. Wei Y-Z, Hu F-C, Hu G-B, Li X-J, Huang X-M, Wang H-C. Differential expression of anthocyanin biosynthetic genes in relation to anthocyanin accumulation in the pericarp of *Litchi chinensis* Sonn. *PLoS ONE*. 2011; 6(4):e19455. doi: [10.1371/journal.pone.0019455](https://doi.org/10.1371/journal.pone.0019455) PMID: [21559331](https://pubmed.ncbi.nlm.nih.gov/21559331/)
53. Zhao ZC, Hu GB, Hu FC, Wang HC, Yang ZY, Lai B. The UDP glucose:flavonoid-3-O-glucosyltransferase (*UGFT*) gene regulates anthocyanin biosynthesis in litchi (*Litchi chinensis* Sonn.) during fruit coloration. *Mol Biol Rep*. 2012; 39(6):6409–6415. doi: [10.1007/s11033-011-1303-3](https://doi.org/10.1007/s11033-011-1303-3) PMID: [22447536](https://pubmed.ncbi.nlm.nih.gov/22447536/)
54. Tereshchenko OY, Arbuzova VS, Khlestkina EK. Allelic state of the genes conferring purple pigmentation in different wheat organs predetermines transcriptional activity of the anthocyanin biosynthesis structural genes. *J Cereal Sci*. 2013; 57(1):10–13. doi: [10.1016/j.jcs.2012.09.010](https://doi.org/10.1016/j.jcs.2012.09.010)
55. Dooner HK. Coordinate genetic regulation of flavonoid biosynthetic enzymes in maize. *Mol Gen Genet*. 1983; 189(1):136–141. doi: [10.1007/BF00326066](https://doi.org/10.1007/BF00326066)
56. Cone KC, Cocciolone SM, Burr FA, Burr B. Maize anthocyanin regulatory gene *pl* is a duplicate of *c1* that functions in the plant. *Plant Cell*. 1993; 5(12):1795–1805. doi: [10.1105/tpc.5.12.1795](https://doi.org/10.1105/tpc.5.12.1795) PMID: [8305872](https://pubmed.ncbi.nlm.nih.gov/8305872/)
57. Mano H, Ogasawara F, Sato K, Higo H, Minobe Y. Isolation of a regulatory gene of anthocyanin biosynthesis in tuberous roots of purple-fleshed sweet potato. *Plant Physiol*. 2007; 143(3):1252–1268. doi: [10.1104/pp.106.094425](https://doi.org/10.1104/pp.106.094425) PMID: [17208956](https://pubmed.ncbi.nlm.nih.gov/17208956/)
58. Honda C, Kotoda N, Wada M, Kondo S, Kobayashi S, Soejima J, et al. Anthocyanin biosynthetic genes are coordinately expressed during red coloration in apple skin. *Plant Physiol Biochem*. 2002; 40(11):955–962. doi: [10.1016/S0981-9428\(02\)01454-7](https://doi.org/10.1016/S0981-9428(02)01454-7)
59. Xu ZS, Huang Y, Wang F, Song X, Wang GL, Xiong AS. Transcript profiling of structural genes involved in cyanidin-based anthocyanin biosynthesis between purple and non-purple carrot (*Daucus carota* L.) cultivars reveals distinct patterns. *BMC Plant Biol*. 2014; 14:262. doi: [10.1186/s12870-014-0262-y](https://doi.org/10.1186/s12870-014-0262-y) PMID: [25269413](https://pubmed.ncbi.nlm.nih.gov/25269413/)
60. Yildiz M, Willis DK, Cavagnaro PF, Iorizzo M, Abak K, Simon PW. Expression and mapping of anthocyanin biosynthesis genes in carrot. *Theor Appl Genet*. 2013; 126(7):1689–1702. doi: [10.1007/s00122-013-2084-y](https://doi.org/10.1007/s00122-013-2084-y) PMID: [23525633](https://pubmed.ncbi.nlm.nih.gov/23525633/)
61. Pandey AK, Dhakal MR. Phytomelanin in Compositae. *Curr Sci*. 2001; 80(8):933–940.
62. Choo TM, Vigier B, Ho KM, Ceccarelli S, Grando S, Franckowiak JD. Comparison of black, purple, and yellow barleys. *Genet Resour Crop Evol*. 2005; 52(2):121–126. doi: [10.1007/s10722-003-3086-4](https://doi.org/10.1007/s10722-003-3086-4)
63. Choo T-M, Vigier BJ, Savard ME, Blackwell BA, Martin RA, Wang JM, et al. Black barley as a means of mitigating deoxynivalenol contamination. *Crop Sci*. 2015; 55(3):1096–1103. doi: [10.2135/cropsci2014.05.0405](https://doi.org/10.2135/cropsci2014.05.0405)
64. Jana BK, Mukherjee SK. Notes on the distribution of phytomelanin layer in higher plants—a short communication. *J Pharm Biol*. 2014; 4(3):131–132.
65. Kim MJ, Hyun JN, Kim JA, Park JC, Kim MY, Kim JG, et al. Relationship between phenolic compounds, anthocyanins content and antioxidant activity in colored barley germplasm. *J Agric Food Chem*. 2007; 55(12):4802–4809. doi: [10.1021/jf0701943](https://doi.org/10.1021/jf0701943) PMID: [17516656](https://pubmed.ncbi.nlm.nih.gov/17516656/)
66. Siebenhandl S, Grausgruber H, Pellegrini N, Del Rio D, Fogliano V, Pernice R, et al. Phytochemical profile of main antioxidants in different fractions of purple and blue wheat, and black barley. *J Agric Food Chem*. 2007; 55(21):8541–8547. doi: [10.1021/jf072021j](https://doi.org/10.1021/jf072021j) PMID: [17894457](https://pubmed.ncbi.nlm.nih.gov/17894457/)