



Predictors of Blood Transfusion in Patients Undergoing Coronary Artery Bypass Grafting Surgery

Saleh Sandoughdaran¹, Mahmood Reza Sarzaem^{1*}, Jamshid Bagheri¹, Mohammad Jebelli¹,
Mohammad Hossein Mandegar¹

¹Cardiac Surgery and Transplantation Research Center, Shariati Hospital, Tehran University of Medical Sciences, Tehran, IR Iran

ARTICLE INFO

Article Type:
Brief Report

Article History:
Received: 16 Feb 2013
Revised: 23 Feb 2013
Accepted: 2 Mar 2013

Keywords:
Blood Transfusion
Coronary Artery Bypass
Blood Cells

ABSTRACT

Objectives: The aim of this retrospective study is to identify intraoperative patient's characteristics predicting the need for blood transfusion during CABG in our local surgical service.

Methods: This study included 1835 consecutive patients, 1311 males and 524 females with mean age 58.8 ± 9.9 years, undergoing coronary artery bypass grafting. Risk factors detected by univariate study were entered in a multivariate logistic regression model of the relationship between preoperative variables and blood transfusion.

Results: Blood transfusion was used in 435 patients (29.9%). Univariate analysis identified hemoglobin, smoking, hypertension, sex, diabetes, BMI and use of cardiopulmonary bypass (CPB) as significant predictors. Multivariate analysis revealed hemoglobin (OR: 0.8; CI: 0.74-0.86; $P < 0.001$), CPB use (OR: 12.2; CI: 8.2-18.1; $P < 0.001$) and female gender (OR: 2.29; CI: 1.72-3.04; $P < 0.001$) as independent risk factors for blood transfusion.

Conclusions: The predictors of RBC transfusion after isolated CABG were performing CPB, preoperative hemoglobin and female gender. These factors can be used as a clinical tool to preserve blood bank resources without increasing patient's risk.

► Implication for health policy/practice/research/medical education:

Ability to identify patients at risk of blood transfusion would save blood bank efforts and resources and allow the employment of a targeted blood conservation policy in CABG patients.

► Please cite this paper as:

Sandoughdaran S, Sarzaem MR, Bagheri J, Jebelli M, Mandegar MH. Predictors of Blood Transfusion in Patients Undergoing Coronary Artery Bypass Grafting Surgery. *Int Cardiovasc Res J*.2013;7(1):25-8.

1. Introduction

Coronary artery bypass grafting (CABG) surgery is a surgical procedure associated with one of the highest rates of transfusion(1). Nearly 20% of all cardiac surgeries in the United States require blood transfusions (2).

The transfusion of red blood cells (RBCs) in cardiac surgery is associated with increased morbidity and mortality (3,4). It has been suggested that blood conservation techniques, such as autologous blood donation, red cell salvage, hemostatic agents

and normovolemic hemodilution, either alone or in combination, in patients undergoing cardiac surgery, could result in an estimated 75% reduction of unnecessary transfusions (2).

The identification of pre-operative variables associated with the need for transfusion will reveal the transfusion risk and thus allows for the use of cost-effective blood conservation methods. The aim of this study was determine the clinical and demographic variables associated with blood product transfusion in patients undergoing elective CABG.

2. Methods

We reviewed the records of patients undergoing CABG surgery at the Division of Cardiovascular Surgery Shariati Hospital from May 2007 to

*Corresponding author: Mahmood Reza Sarzaem, Cardiac Surgery and Transplantation Research Center, Dr. Shariati Hospital, North Karegar Ave., Tehran, IR Iran PO: 1411713137.
Tel: +98-9125268001, Fax: +9821-44453449,
E-mail: sarzaem_mr@yahoo.com

Table 1. The Baseline of Patients' Characteristics Stratified by Blood Transfusion

	Without Transfusion (n=1395)	Transfusion (n=435)	P value
Age (yr)	58.8±10.1	59.4±9.1	0.21
Sex			
Female	332 (23.8)	187 (42.8)	<0.001
Male	1062 (76.2)	250 (57.2)	
BMI (m/Kg ²)	27.0	26.9	0.35
Preoperative Hb (g/dL)	14.0	13.1	<0.001
On-pump	776 (55.5)	402 (91.8)	<0.001
Diabetes	449 (35.7)	184 (42.0)	0.03
COPD	35 (2.5)	6 (1.4)	0.20
Smoking	259 (18.5)	62 (14.2)	0.03
HTN	695 (49.7)	254 (58.0)	0.02
Previous MI	512 (36.6)	167 (37.4)	0.77
Renal Failure	29 (2.1)	11 (2.5)	0.57
Ejection Fraction	46.85	46.74	0.82

Data are presented as mean ± SD or No. (%).

Abbreviations: Hb, Hemoglobin; COPD, Chronic Obstructive Pulmonary Disease; BMI, Body Mass Index; HTN, Hypertension; MI, Myocardial Infarction.

October 2012. Standard demographic and clinical characteristics were obtained from institutional cardiac surgery database. These included history of diabetes mellitus (DM), hypertension (HTN), preoperative hemoglobin, operative transfusion data, cardiopulmonary bypass, ejection fraction (EF), chronic obstructive pulmonary disease (COPD), and preoperative creatinine. Preoperative hemoglobin data were measured closest to the time of surgery. Apheresis products were not used. Indications for allogeneic transfusion were based on routine laboratory measurements of International Normalized Ratio (INR) platelet counts and hemoglobin values, in addition to measurements of hemodynamic data, the rate of blood loss, and existing concomitant diseases. Transfusions of allogeneic packed red cells were given at the discretion of the attending surgeon. Student's test was used to compare differences in pre-operative hemoglobin, ejection fraction and BMI between patients with and without transfusion. The relationship between transfusion requirements and categorical variables was examined using the chi-squared test. The relationship between transfusion requirements and all variables adjusted to one another was examined using backward stepwise multiple logistic regression.

3. Results

Between May 2007 to October 2012, 1992 patients underwent CABG surgery. We excluded 157 patients who underwent concomitant cardiac surgical procedures, leaving 1835 patients who underwent isolated primary CABG surgery for the study. The study population had a mean age of 58.8 ± 9.9 SD years. Baseline patients' characteristics stratified by transfusion of RBCs are shown in Table 1.

Blood transfusions (mean quantity 1.1 units) were given to 435 of 1835 patients (24%); 187 of 524 women (35.6%) required transfusion, compared with 250 of 1311 men (19.0%) ($P<0.001$). Of those who received blood, 380 (87.2%) received 1 unit; 47 (10.8%), 2 units, 8 (1.8%), 3 units, and 1 (0.2%), 4 units.

The results of the univariable analysis are given in Table 1. Sex, BMI, diabetes, on-pump surgery, preoperative hemoglobin, smoking, and hypertension were significantly associated with intraoperative blood transfusion. The multivariable stepwise logistic regression analysis determined that in females, use of cardiopulmonary bypass (CPB) and preoperative hemoglobin level were independently associated with an increased risk of blood transfusion. The regression coefficients, odds ratios, and P -values are summarized in Table 2.

Table 2. Multivariate Analyses of Preoperative Risk Factors

	Odds Ratio	95% Confidence Interval		P value
		Lower	Upper	
Diabetes	0.925	0.712	1.201	0.558
Hypertension	1.171	0.905	1.516	0.229
On-Pump Surgery	12.215	8.218	18.157	<0.001
Female Gender	2.291	1.718	3.055	<0.001
Smoking	0.815	0.574	1.155	0.250
Preoperative Hemoglobin	0.801	0.745	0.860	<0.001

4. Discussion

This study was undertaken to determine variables indicating allogeneic RBC transfusion in patients undergoing CABG, which is associated with a high risk of blood transfusion. Despite Society of Thoracic Surgeons guidelines and other reports, blood transfusion rate varies significantly between institutions (5-7). Allogeneic RBC transfusion rates between 8–100% have been reported during CABG in different studies (7-9). This wide variability may be explained by a variety of facts including differences in patient population among the study centers, preoperative medication with anti-platelet agents and anticoagulants and several surgical and procedure-related factors (9-11). Therefore, it is essential for individual centers to pay due attention to their current transfusion practice.

Although several variables showed association with the need for Allogeneic RBC transfusion during CABG in the univariate analysis in our study, three of these including preoperative hemoglobin, on-pump surgery and female gender proved to be independent predictors of intraoperative blood transfusion.

Several studies have reported anemia as a major predictor for need of transfusion. However it is still unclear what hemoglobin level vindicates the need for transfusion. Latest Published guideline for blood transfusion and conservation in cardiac surgery by The Society of Thoracic Surgeons, recommends transfusion for a hemoglobin level of <7 g/dl. However, the level of evidence is 'C' with a recommendation of 'Class 2a', which means that supporting evidence is still insufficient (5). Various strategies exist for dealing with preoperative anemia in patient undergoing CABG. In patients with delayed surgery, erythropoiesis-stimulating agents should begin before surgery. The use of off-pump CABG should be considered for some anemic patients, while bypass pumps that require minimal priming volumes, resulting in lesser dilution of the patient's blood volume, should be used in patients in need of cardiac surgery with cardiopulmonary bypass (12).

In agreement with other studies, the present authors found that female sex is a predictive factor for the transfusion of RBCs (13-15). The reason for such a gender difference in requiring blood transfusion is not clear. Some studies suggested low Hematocrit, as a major reason for the greater need for blood transfusion in females (16). However, difference in blood transfusion was present even when patients with similar preoperative Hct levels were compared in other studies (13,14,16,17). Our findings are consistent with those of the aforementioned studies where blood volume was included in the analysis, and sex was independently associated with the need for transfusion.

Several studies have shown differences in blood loss and allogeneic transfusion requirements between on-pump and off-pump CABG. Nuttall et al.(18)

showed that intraoperative transfusion of allogeneic red blood cells and platelets was greater in patients undergoing CPB than Off-pump coronary artery bypass (OPCAB). In another study Ascione et al. (19) reported that avoiding CPB decreases perioperative bleeding and consequently reduces the use of blood products after CABG. Several factors may contribute to greater need for transfusion in patients undergoing CPB including hypothermia, hemodilution, activation of coagulation, endothelial cell and tissue injury, foreign surface contact, platelet dysfunction and hyperfibrinolysis (20). The present results show that on-pump surgery has a detrimental effect on postoperative blood transfusion.

There are a number of potential limitations to this study. First, this study was performed at a large tertiary-care teaching hospital that followed current blood-conservation and transfusion guidelines; thus, the practice pattern should be comparable to similar institutions. Second, the effects of unknown or unmeasured confounders on the observed association cannot be ruled out. Taking into account the limitations of the present study, we conclude that the best predictor for transfusion risk is the preoperative hemoglobin level, female gender and on-pump surgery.

Acknowledgements

There is no Acknowledgement.

Financial Disclosure

None declared.

Funding/ Support

None declared.

References

- Chiavetta JA, Herst R, Freedman J, Axcell TJ, Wall AJ, van Rooy SC. A survey of red cell use in 45 hospitals in central Ontario, Canada. *Transfusion*. 1996;**36**(8):699-706.
- Shander A, Rijhwani TS. Clinical outcomes in cardiac surgery: conventional surgery versus bloodless surgery. *Anesthesiol Clin North America*. 2005;**23**(2):327-45, vii.
- Engoren MC, Habib RH, Zacharias A, Schwann TA, Riordan CJ, Durham SJ. Effect of blood transfusion on long-term survival after cardiac operation. *Ann Thorac Surg*. 2002;**74**(4):1180-6.
- Murphy GJ, Reeves BC, Rogers CA, Rizvi SI, Culliford L, Angelini GD. Increased mortality, postoperative morbidity, and cost after red blood cell transfusion in patients having cardiac surgery. *Circulation*. 2007;**116**(22):2544-52.
- Ferraris VA, Brown JR, Despotis GJ, Hammon JW, Reece TB, Saha SP, et al. 2011 update to the Society of Thoracic Surgeons and the Society of Cardiovascular Anesthesiologists blood conservation clinical practice guidelines. *Ann Thorac Surg*. 2011;**91**(3):944-82.
- Karkouti K, Wijjeyundera DN, Beattie WS, Callum JL, Cheng D, Dupuis JY, et al. Variability and predictability of large-volume red blood cell transfusion in cardiac surgery: a multicenter study. *Transfusion*. 2007;**47**(11):2081-8.
- Legare JF, Buth KJ, King S, Wood J, Sullivan JA, Hancock Friesen C, et al. Coronary bypass surgery performed off pump does not result in lower in-hospital morbidity than coronary artery bypass grafting performed on pump. *Circulation*. 2004;**109**(7):887-92.
- Rogers MA, Blumberg N, Saint S, Langa KM, Nallamothu BK. Hospital variation in transfusion and infection after cardiac surgery: a cohort study. *BMC Med*. 2009;**7**:37.
- Snyder-Ramos SA, Mohnle P, Weng YS, Bottiger BW, Kulier A, Levin J, et al. The ongoing variability in blood transfusion practices

- in cardiac surgery. *Transfusion*. 2008;**48**(7):1284-99.
10. Ferraris VA, Ferraris SP, Saha SP, Hessel EA, 2nd, Haan CK, Royston BD, et al. Perioperative blood transfusion and blood conservation in cardiac surgery: the Society of Thoracic Surgeons and The Society of Cardiovascular Anesthesiologists clinical practice guideline. *Ann Thorac Surg*. 2007;**83**(5 Suppl):S27-86.
 11. Slight RD, Bappu NJ, Nzewi OC, Lee RJ, McClelland DB, Mankad PS. Factors predicting loss and gain of red cell volume in cardiac surgery patients. *Transfus Med*. 2006;**16**(3):169-75.
 12. Gardner TJ. To transfuse or not to transfuse. *Circulation*. 2007;**116**(5):458-60.
 13. Karkouti K, Cohen MM, McCluskey SA, Sher GD. A multivariable model for predicting the need for blood transfusion in patients undergoing first-time elective coronary bypass graft surgery. *Transfusion*. 2001;**41**(10):1193-203.
 14. Shevde K, Pagala M, Kashikar A, Tyagaraj C, Shahbaz N, Iqbal M, et al. Gender is an essential determinant of blood transfusion in patients undergoing coronary artery bypass graft procedure. *J Clin Anesth*. 2000;**12**(2):109-16.
 15. van Straten AH, Kats S, Bekker MW, Verstappen F, ter Woorst JF, van Zundert AJ, et al. Risk factors for red blood cell transfusion after coronary artery bypass graft surgery. *J Cardiothorac Vasc Anesth*. 2010;**24**(3):413-7.
 16. Utley JR, Wallace DJ, Thomason ME, Mutch DW, Staton L, Brown V, et al. Correlates of preoperative hematocrit value in patients undergoing coronary artery bypass. *J Thorac Cardiovasc Surg*. 1989;**98**(3):451-3.
 17. Shevde K, Pagala M, Tyagaraj C, Udeh C, Punjala M, Arora S, et al. Preoperative blood volume deficit influences blood transfusion requirements in females and males undergoing coronary bypass graft surgery. *J Clin Anesth*. 2002;**14**(7):512-7.
 18. Nuttall GA, Erchul DT, Haight TJ, Ringhofer SN, Miller TL, Oliver WC, Jr, et al. A comparison of bleeding and transfusion in patients who undergo coronary artery bypass grafting via sternotomy with and without cardiopulmonary bypass. *J Cardiothorac Vasc Anesth*. 2003;**17**(4):447-51.
 19. Ascione R, Williams S, Lloyd CT, Sundaramoorthi T, Pitsis AA, Angelini GD. Reduced postoperative blood loss and transfusion requirement after beating-heart coronary operations: a prospective randomized study. *J Thorac Cardiovasc Surg*. 2001;**121**(4):689-96.
 20. Green JA, Spiess BD. Current status of antifibrinolytics in cardiopulmonary bypass and elective deep hypothermic circulatory arrest. *Anesthesiol Clin North America*. 2003;**21**(3):527-51 viii.