



Expression of DISC1-Interactome Members Correlates with Cognitive Phenotypes Related to Schizophrenia

Antonio Rampino^{1,2}, Rosie May Walker¹, Helen Scott Torrance¹, Susan Maguire Anderson¹, Leonardo Fazio², Annabella Di Giorgio^{2,3}, Paolo Taurisano², Barbara Gelao², Raffaella Romano², Rita Masellis², Gianluca Ursini², Grazia Caforio², Giuseppe Blasi², J. Kirsty Millar¹, David John Porteous^{1,4}, Pippa Ann Thomson^{1,4}, Alessandro Bertolino^{2,5*}, Kathryn Louise Evans^{1,4*}

1 Medical Genetics Section, Centre for Genomic and Experimental Medicine, Medical Research Council Institute of Genetics and Molecular Medicine, The University of Edinburgh, Edinburgh, United Kingdom, **2** Department of Neuroscience and Sense Organs, University of Bari, Bari, Italy, **3** Istituti di Ricovero e Cura a Carattere Scientifico Casa Sollievo della Sofferenza, Section of Liaison Psychiatry, San Giovanni Rotondo (FG), Italy, **4** Centre for Cognitive Ageing and Cognitive Epidemiology, Medical Genetics Section, Centre for Genomic and Experimental Medicine, Medical Research Council Institute of Genetics and Molecular Medicine, University of Edinburgh, Edinburgh, United Kingdom, **5** Imaging, Pharma Research and Early Development, Neuroscience Discovery and Translational Area, Hoffman-La Roche, Basel, Switzerland

Abstract

Cognitive dysfunction is central to the schizophrenia phenotype. Genetic and functional studies have implicated *Disrupted-in-Schizophrenia 1 (DISC1)*, a leading candidate gene for schizophrenia and related psychiatric conditions, in cognitive function. Altered expression of *DISC1* and *DISC1*-interactors has been identified in schizophrenia. Dysregulated expression of *DISC1*-interactome genes might, therefore, contribute to schizophrenia susceptibility via disruption of molecular systems required for normal cognitive function. Here, the blood RNA expression levels of *DISC1* and *DISC1*-interacting proteins were measured in 63 control subjects. Cognitive function was assessed using neuropsychiatric tests and functional magnetic resonance imaging was used to assess the activity of prefrontal cortical regions during the N-back working memory task, which is abnormal in schizophrenia. Pairwise correlations between gene expression levels and the relationship between gene expression levels and cognitive function and N-back-elicited brain activity were assessed. Finally, the expression levels of *DISC1*, *AKAP9*, *FEZ1*, *NDEL1* and *PCMI* were compared between 63 controls and 69 schizophrenic subjects. We found that *DISC1*-interactome genes showed correlated expression in the blood of healthy individuals. The expression levels of several interactome members were correlated with cognitive performance and N-back-elicited activity in the prefrontal cortex. In addition, *DISC1* and *NDEL1* showed decreased expression in schizophrenic subjects compared to healthy controls. Our findings highlight the importance of the coordinated expression of *DISC1*-interactome genes for normal cognitive function and suggest that dysregulated *DISC1* and *NDEL1* expression might, in part, contribute to susceptibility for schizophrenia via disruption of prefrontal cortex-dependent cognitive functions.

Citation: Rampino A, Walker RM, Torrance HS, Anderson SM, Fazio L, et al. (2014) Expression of *DISC1*-Interactome Members Correlates with Cognitive Phenotypes Related to Schizophrenia. PLoS ONE 9(6): e99892. doi:10.1371/journal.pone.0099892

Editor: Ya-Ping Tang, Louisiana State University Health Sciences Center, United States of America

Received: January 7, 2014; **Accepted:** May 19, 2014; **Published:** June 18, 2014

Copyright: © 2014 Rampino et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by the Chief Scientist Office (<http://www.cso.scot.nhs.uk/>) [grant number CZB/4/782]; the Scottish Funding Council (<http://www.sfc.ac.uk/>) Strategic Research Development Grant “Biomarkers for Battling Chronic Disease”; and Research Council UK (<http://www.rcuk.ac.uk/Pages/Home.aspx>) fellowships to PAT and JKM. KLE, PAT and DJP are members of The University of Edinburgh Centre for Cognitive Ageing and Cognitive Epidemiology (CCACE), part of the cross-council Lifelong Health and Wellbeing Initiative [grant number G0700704/84698]. CCACE is funded by the Biotechnology and Biosciences Research Council (<http://www.bbsrc.ac.uk/home/home.aspx>); Engineering and Physical Sciences Research Council (<http://www.epsrc.ac.uk/Pages/default.aspx>); Economic and Social Research Council (<http://www.esrc.ac.uk/>); and Medical Research Council (<http://www.mrc.ac.uk/index.htm>). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: Alessandro Bertolino is a full-time employee of Hoffman-La Roche Ltd. This does not alter the authors' adherence to PLOS ONE policies on sharing data and materials. All other authors declare that they have no conflicts of interest.

* E-mail: kathy.evans@ed.ac.uk (KLE); a.bertolino@psichiat.uniba.it (AB)

Introduction

Abnormal cognitive function is recognized as a core component of the schizophrenia phenotype [1,2]. Cognitive deficits are often apparent before the onset of positive symptoms, are heritable [3], and are seen in unaffected relatives of patients [4]. Moreover, while deficits in working memory are a good predictor of functional outcome [5], cognitive deficits are often refractory to current drug treatments [6].

A network of brain regions, including the prefrontal cortex (PFC), contributes to cognition [7]. Several of these regions demonstrate abnormal activity in schizophrenia [8]. Schizophrenic

patients are impaired in the performance of PFC-dependent tasks such as the N-back task, a test of working memory [9]. This task elicits a heritable pattern of activity in the PFC [3], meaning it can be used to assess whether “schizophrenia-risk” polymorphisms exert their pathogenic effect by altering PFC function.

Variation in *Disrupted in Schizophrenia 1 (DISC1)* is a risk-factor for schizophrenia [10,11] and is implicated in cognitive function in both healthy individuals and schizophrenic patients [12].

The *DISC1* protein interacts with several proteins to link multiple cellular functions and molecular pathways, including neuronal migration (*LIS1*, *NDE1*, *NDEL1*, *Dixdc1*), axonal bundling and elongation (*FEZ1*), neural progenitor proliferation

(GSK3 β), cell cycle regulation (PCM1), neuronal signalling (Girdin, GSK3 β , PDE4), and signal transduction (Kal7, TNK1, AKAP9) [10,13]. Polymorphisms in several genes encoding DISC1-interactors have been associated with schizophrenia [10,14], and altered hippocampal expression of *NDE1*, *FEZ1*, and *LIS1* has been detected in post-mortem brains of schizophrenic patients [15]. Thus, alterations in brain function that confer risk for psychiatric illness might arise from dysfunction at several loci in the DISC1-interactome. In support of this assertion, Hennah and Porteous [16] have shown genetic variation in *DISC1* and DISC1-interacting proteins to modulate RNA expression of genes involved in schizophrenia-relevant processes, including cytoskeletal functions, synaptogenesis, neurodevelopment and sensory perception. This finding is particularly pertinent in light of the evidence for the involvement of perturbed gene expression in schizophrenia [17].

In this study, we test the hypotheses that: (i) genes encoding DISC1-interactome members show coordinated expression; (ii) expression levels of these genes are associated with cognitive function; (iii) frontal cortical activity elicited by the N-back task is related to the expression of genes whose expression level is associated with cognitive function; and (iv) the expression of genes whose expression correlates with cognitive function is abnormal in schizophrenic patients. We demonstrate associations between: (i) the expression levels of several DISC1-interactome members; (ii) DISC1-interactome members' expression levels and cognitive performance; and (iii) genes whose expression is related to cognitive performance and frontal cortical activity. Moreover, we find the expression of *DISC1* and *NDEL*, whose expression levels correlate with cognitive performance, to be reduced in the blood of schizophrenic patients.

Materials and Methods

Ethics Statement

Procedures were carried out in accordance with The Code of Ethics of the World Medical Association (Declaration of Helsinki) and approval was given by the local ethics committee ("Comitato Etico Indipendente Locale-Azienda Ospedaliera Ospedale Policlinico Consorziato di Bari"), who examined the study protocol and the suitability of the environment and investigators involved. Participants gave written informed consent. The capacity of participants to consent was assessed by a psychiatrist. If participants were not deemed capable of giving informed consent, a legal representative gave consent on their behalf. In accordance with Articles 414 and 415 of the Civil Code, the Legal Representative was a member of the patient's family, a clinician or a lawyer nominated by the Local Court in order to represent the patient's civil rights.

Subjects

We examined 63 control subjects (age: mean = 26.2, standard deviation (SD) = 5.4; 36 females) and 69 subjects with schizophrenia (age: mean = 37.3, SD = 10.8, 16 females). Control subjects were screened using the Non-Patient Structured Clinical Interview for DSM-IV to ensure they were unaffected by any psychiatric condition. In addition, control subjects were screened using the Family Interview for Genetic Studies [18] to ensure absence of psychotic disorders in their first-degree relatives. Diagnosis of schizophrenia was made using the Structured Clinical Interview for the DSM-IV Axis I disorders, which was administered by psychiatrists. Both patients and control subjects were excluded if they had: a significant history of drug or alcohol abuse; active drug abuse in the previous year; experienced a head trauma with a loss

of consciousness; or if they suffered from any other significant medical condition.

Gene Expression Levels

Blood for RNA extraction was collected using the Tempus Blood Collection System (Applied Biosystems). RNA was extracted from 3 ml whole blood using Tempus Blood RNA Isolation Kits and subsequently reverse transcribed using Transcriptor First Strand cDNA Synthesis Kits (Roche Applied Science). mRNA expression was measured by quantitative reverse-transcription polymerase chain reaction (qRT-PCR), using TaqMan probe-based assays (Applied Biosystems; see Table S1 in File S1 for probe IDs). Reactions were performed using an ABI Prism 7900 Sequence Detection System (Applied Biosystems), at the Wellcome Trust Clinical Research Facility, Edinburgh. Assays spanned an exon-exon boundary and amplified the maximum possible number of isoforms. Expression levels were calculated using the relative standard curve method and gene-of-interest expression levels (*DISC1*, *AKAP9*, *FEZ1*, *GSK3 β* , *LIS1*, *NDE1*, *NDEL1*, *PCM1*, *PDE4B*, *PDE4D*) were normalised to the geometric mean of two reference genes (*RPLPO* and *HNRMPD*). These reference genes were selected on the basis of geNorm analysis [19], which showed these genes to be stably expressed in a representative sub-set of the blood samples assessed in this study (N = 15 schizophrenic patients and 15 controls). Each sample was measured in technical triplicate. Between-plate differences were accounted for by normalisation to a calibrator. Outliers were identified and excluded following Burns et al. [20].

Cognitive Assessment

All subjects were assessed with a battery of cognitive tests comprising the: (i) N-Back paradigm (2-back condition); (ii) Continuous Performance Test, assessing sustained attention; (iii) Wechsler Memory scale, giving a memory quotient; (iv) Wisconsin Card Sorting Test, evaluating executive functions; (v) Trail Making A and B, assessing cognitive flexibility; and (vi) Controlled Oral Word Association Test (Benton, 1967), measuring Semantic Fluency and Phonological Fluency. For further details, see File S1 Materials and Methods.

Gene-by-gene Expression Interactions

Pearson pairwise correlation coefficients were calculated for each pair of gene expression levels using SPSSv17.0 software (www.spss.com). To control for the effects of multiple testing, permutation analysis (10^5 permutations) was performed using Resampling (<http://core.ecu.edu/psyc/wuenschk/StatHelp/Resampling.htm>, East Carolina University) which adopts a classical "randomization without replacement" algorithm. The same software was used to derive 95% bootstrap confidence intervals for correlation coefficients. Bootstrapping was conducted by running a total of 10^5 re-samples drawn pairwise from our original sample. Corrected $p \leq 0.05$ indicated statistical significance.

Association between Gene Expression and Cognitive Performance

Principal component analysis. Principal component analysis (PCA) on a correlation matrix followed by varimax rotation, retaining factors with an eigenvalue ≥ 1 was used to group cognitive functions. In addition to providing an insight into the architecture of cognitive function, this approach reduces the number of dependent variables, thus reducing the type 1 error rate [21,22].

Correlation of gene expression with cognition. Multiple regression analysis with backward stepwise removal was conducted for the expression levels of ten genes against the scores obtained for each of the four principal components (PCs). The variable with the smallest partial correlation with the dependent variable was considered for removal first, and removed if the significance level of its F-value was $p > 0.05$. For all PCs, a higher score indicated better performance. For each analysis, the model with the highest R^2 and the lowest p -value was taken to best explain the variance within the relevant PC. All gene expression levels and their pairwise interactions were then regressed within each model against the relevant PC. The threshold for statistical significance was $p \leq 0.05$. Analyses were conducted using SPSSv.17.0.

fMRI Paradigm

The fMRI experimental paradigm consisted of the N-back task, as described previously [23]. The stimuli consisted of numbers (1–4) shown in random sequence and displayed at the points of a diamond-shaped box. A non-memory-guided control condition (0-back) presented the same stimuli but simply required subjects to identify the current stimulus. In the working memory condition, the task required recollection of a stimulus seen two stimuli previously (2-back). Performance data were recorded as the percentage of correct responses and reaction time (msecs).

Echo planar imaging blood oxygenation level dependent fMRI data were acquired as described previously [23] (TE = 30 m secs, TR = 2 secs, 20 contiguous slices, voxel size = $3.75 \times 3.75 \times 5$) on a GE 3T scanner with a standard head coil. A simple block design was used in which each block consisted of 8 alternating 0-back and 2-back conditions (each 30 secs), obtained in 4 min and 8 secs, 120 whole-brain fMRI volumes. The first four scans of the time series were discarded to allow for signal equilibration effects.

fMRI Data Analysis

Images were pre-processed using Statistical Parametric Mapping 8 software (SPM8; <http://www.fil.ion.ucl.ac.uk/spm>). Images were realigned to the first volume in the time series to correct for head motion (< 2 voxels translation, $< 1^\circ$ rotation), re-sampled to a 2 mm isotropic voxel size, spatially normalized into a standard stereotactic space (Montreal Neurological Institute template) using a 12 parameter affine model and smoothed to minimize noise and residual differences in gyral anatomy with a Gaussian filter, set at 6 mm full-width at half-maximum. Voxel-wise signal intensities were ratio normalized to the whole-brain global mean. For each experimental condition, a box-car model convolved with the haemodynamic response function at each voxel was modelled. Pre-determined condition effects at each voxel were calculated using a t-statistic, producing a statistical image for the contrast of 2-back versus 0-back. Individual contrast images were then used in second-level random effects models to determine task-specific regional responses at the group level for the entire sample.

To assess the association between gene expression and fMRI activation, contrast images of all subjects were used as dependent variables in a general linear model using SPM8. A priori evidence for involvement of the frontal cortex in the N-back task [3,9] meant this region was of particular interest. Regions of interest (ROIs) were specified by dividing the frontal cortex into its constituent Brodmann Areas (BAs), as defined by the WFU_Pick-Atlas [24]. Within each ROI, a family wise error (FWE) small volume correction with a $p \leq 0.05$ was applied to control for type I errors.

Comparison of Gene Expression Levels in Patients with Schizophrenia and Healthy Controls

Case-control comparisons were carried out for *AKAP9*, *DISC1*, *FEZ1*, *NDELI* and *PCMI* expression levels. To assess the influence of potentially confounding variables, the effects of age and gender on expression were assessed for each gene using one-way analysis of variance (ANOVA) and linear regression analysis, respectively. Comparisons of gene expression levels between schizophrenic patients and controls were made using either ANOVA or analysis of covariance (ANCOVA) with the independent variable of diagnostic status and the dependent variable of expression level. Covariates were included where appropriate.

Results

Gene-by-gene Expression Interactions

In the control subjects, we assessed correlations between RNA expression levels of *DISC1* and nine DISC1-interactors by calculating Pearson correlation coefficients for each pair-wise combination. Following correction for multiple testing by permutation and bootstrap analyses, *DISC1* expression showed significant correlation with *AKAP9*, *GSK3 β* , and *NDELI*. Several other significant pair-wise correlations were detected (Table 1; Figure S1 in File S1).

Principal Component Analysis of Cognitive Test Data

As we had carried out multiple related cognitive tests we defined a set of linearly independent cognitive variables using principal component analysis (PCA). PCA revealed four principal components (PCs) that together explained 66.5% of the variance in cognitive performance ($p = 0.003$; Figure 1; Table S2 in File S1).

Correlation between Gene Expression and Cognitive Performance

To explore the correlation between gene expression and cognition we carried out backward removal multiple regression analysis of gene expression levels against the four PCs. For each PC, variation in the expression of either a single gene or a combination of genes was found to explain a significant proportion of the variance (Table 2). We next examined the effect of gene-by-gene interactions within PCs 2, 3 and 4 (PC1 was best explained by a single-gene model). Backward removal multiple regression analysis revealed that inclusion of interaction terms improved the fit and significance of each model (Table 3; Figure S2 in File S1). The combination of genes and gene \times gene interactions that best explained variation in each PC will henceforth be referred to as the “expression model”.

Correlation between Gene Expression and Frontal Cortical Function

We assessed the relationship between the four expression models and N-back-elicited frontal cortical activity, as measured by fMRI, in the control subjects. Several findings at the fMRI level were consistent with findings at the behavioural level (Table 4; Table S3 in File S1). The PC 2, 3, and 4 expression models showed significant relationships with the activity of frontal regions (Figures S3 and S4 in File S1). The PC1 expression model showed a nominally significant relationship with activity in BA 9 (uncorrected $p < 0.0001$); however, this relationship did not withstand multiple testing correction (corrected $p = 0.093$).

Table 1. Gene co-expression within the DISC1-interactome.

Co-expressed gene pair	Pearson correlation coefficient (<i>r</i>)	Uncorrected p-value (2-tailed)	Corrected p-value ^a	95% bootstrap Confidence interval
DISC1 × AKAP9	+0.27	0.046	0.048	0.012; 0.53
DISC1 × GSK3β	−0.35	0.0082	0.012	−0.55; −0.34
DISC1 × NDEL1	+0.29	0.036	0.036	0.038; 0.51
AKAP9 × FEZ1	+0.42	0.0033	0.0052	0.14; 0.42
AKAP9 × PCMI	+0.39	0.0032	0.0052	0.036; 0.39
FEZ1 × NDE1	−0.45	0.0012	0.0012	−0.68; −0.18
NDE1 × GSK3β	+0.57	0.000033	0.000055	0.37; 0.72
NDE1 × NDEL1	+0.32	0.033	0.033	0.003; 0.56
GSK3β × PDE4B	+0.32	0.023	0.041	0.11; 0.32

^aCorrected by permutation analysis (10⁵ permutations).
doi:10.1371/journal.pone.0099892.t001

Comparison of the Expression Levels of DISC1-Interactome Members in Schizophrenic Patients and Healthy Controls

The expression levels of five members of the DISC1-interactome whose expression was correlated with cognitive performance in control subjects (*AKAP9*, *DISC1*, *FEZ1*, *NDEL1*, and *PCMI*) were measured in the peripheral blood of schizophrenic patients. As age significantly affected *DISC1* expression ($p = 0.012$), it was included as a covariate in the DISC1 analysis. Comparing schizophrenic patients to controls revealed significant differences in *DISC1* ($p = 0.0015$, ANCOVA) and *NDEL1* ($p = 0.028$, ANOVA) expression. Both genes showed reduced RNA expression in schizophrenic patients (*DISC1*: mean = 1.33, SD = 0.810; *NDEL1*: mean = 1.88, SD = 0.294) relative to controls (*DISC1*: mean = 2.10, SD = 1.20; *NDEL1*: mean = 2.15, SD = 0.936).

Discussion

Multiple lines of evidence support a role for variation in *DISC1* and DISC1-interactors in conferring risk for psychiatric illness [10,11], potentially, in part, via an effect on cognitive function [12]. *DISC1*'s role as a molecular scaffold [10] suggests that the

coordinated expression of DISC1-interactome members is essential for normal brain development and function, and that alteration may result in cognitive deficits and susceptibility to psychiatric illness.

Here, in healthy individuals, we demonstrated significant correlations between the expression levels of several members of the DISC1-interactome. Moreover, RNA expression of five of these genes (*AKAP9*, *DISC1*, *FEZ1*, *NDEL1*, and *PCMI*) correlated with cognitive performance. Cognitive function, as assessed by several measures, was reduced to four PCs. The expression level of at least one DISC1-interactome gene was significantly correlated with each PC. Interestingly, where multiple genes best explained performance, the majority of these genes showed significant between-gene correlations in their expression levels. It is noteworthy that genes showing correlated expression often conferred opposite effects on cognitive function. Furthermore, the inclusion of interaction terms improved the fit and the significance of the models. These findings are suggestive of a complex scenario whereby members of gene pairs showing correlated expression contribute to cognitive function via both independent and overlapping mechanisms. Our findings suggest that it is unlikely that genes showing correlated expression contribute to the same cognitive PC via a mechanism that simply involves one gene regulating the expression of the other and, thus, conferring an effect on cognition. Nevertheless, it would be of interest in the future to perform experiments *in vitro* to establish whether manipulating the expression level of one member of a correlated gene pair alters the expression of the other member of the gene pair. Our results indicate the inter-dependency of DISC1-interactome members and highlight the possibility for variation in their expression levels to confer risk for schizophrenia via a deleterious effect on cognition.

We report that N-back task-elicited activity in BAs 9, 45, and 6 is correlated with the PC1, PC2, PC3, and PC4 expression models. Our observation of N-back task-elicited activity in these regions is consistent with previous studies [25]. The expression models for PCs 1, 2, and 4 correlated with activity in BAs 9 and 45, regions that have been implicated in working memory function [25,26]. BA6 has traditionally been viewed as motor cortex; however, the observation that this area is active in non-motor mental operation tasks that require the updating of spatial and verbal representations in memory suggests it may subserve both cognitive and motor functions [27,28]. PC3, the cognitive model for which expression correlated with BA6 activity, related to phonological




Figure 1. Graphical depiction of the principal components derived from performance on the neurocognitive tests. Four principal components were defined by principal components analysis. Each principal component was named according to the specific cognitive functions assessed by the tests that loaded onto it.
doi:10.1371/journal.pone.0099892.g001

Table 2. Relationship between the expression levels of DISC1-interactome genes and cognitive performance.

PC ^a	Cognitive functions assessed	Model ^b	β coefficient	p-value	Whole model p-value	Whole model R ²
1	Working Memory –Attention	NDEL1	+0.34	0.02	0.023	0.12
2	Memory -Perseverative Errors	AKAP9	+0.48	0.002	0.0014	0.35
		FEZ1	−0.47	0.002		
		PCM1	−0.38	0.01		
3	Phonological Fluency - Speed of Processing	DISC1	+0.28	0.072	0.029	0.16
		NDEL1	−0.38	0.015		
4	Semantic Fluency	DISC1	−0.26	0.067	0.0052	0.23
		AKAP9	+0.41	0.002		

^aPC = principal component.

^bThe model explaining the most variation within each PC was determined by backward removal multiple regression analysis including main effects only.

doi:10.1371/journal.pone.0099892.t002

fluency and speed of processing. Interestingly, activity in BA6 has previously been found to correlate with response rate on a word fluency task [29].

Within each expression model, the effect of each gene on N-back task-elicited activity depended on the expression of the other DISC1-interactome genes in the model and the specific brain region considered. Furthermore, significant correlations were observed between several gene-by-gene interaction terms and frontal cortical activity. For genes contributing to PCs 2, 3 and 4, the relationship between gene expression and N-back task-elicited activity was in the opposite direction for the gene-by-gene interaction terms and the related main effects. This pattern of results is compatible with a scenario whereby genes contributing to a given PC act via both mutually dependent and independent pathways, which confer opposite effects on the activity of a given brain region in response to changes in gene expression. These findings support a functional interaction between DISC1-interactome members, adding to our understanding of their known cellular interactions, and highlight the need to consider genetic variables in the context of their appropriate biological networks. Moreover, we are the first to report a correlation between the expression of a gene in the blood and frontal cortical activity.

Of the five genes whose expression correlated with cognitive function in healthy controls, *DISC1* and *NDEL1*, showed reduced expression in schizophrenic patients. This finding is in keeping

with the positive correlation between *DISC1* and *NDEL1* expression observed in control subjects. In light of this result, one possibility is that the reduced expression of both genes in schizophrenic patients reflects the inter-dependency of the expression levels of these genes. This possibility could be assessed *in vitro* by manipulating the expression level of each gene individually and measuring the expression of the other gene.

Our study is the first to compare the peripheral blood expression of *NDEL1* between patients and controls and is consistent with a previous study, which demonstrated reduced *NDEL1* plasma activity in schizophrenia [30]. In contrast to our findings, a recent study found *DISC1* RNA expression to be increased in peripheral blood mononuclear cells (PBMCs) from treatment-naïve patients with schizophrenia and to remain increased following six to eight weeks of antipsychotic treatment [31]. The discrepancy with our findings is, therefore, unlikely to be attributable to treatment effects, with the caveat that antipsychotic treatment may take longer than eight weeks to alter *DISC1* expression. Another potential explanation for this discrepancy is the difference in the cellular composition (whole blood vs. PMBCs) of the RNA source. Our finding of reduced *DISC1* expression is consistent with our previous work [32], where we observed ~50% reduction in *DISC1* protein expression in lymphoblastoid cell lines derived from carriers of at (1;11) translocation that disrupts *DISC1* and is a strong genetic risk factor for psychiatric illness [33].

Table 3. Relationship between the expression levels of DISC1-interactome genes and their pairwise interactions and cognitive performance.

PC ^a	Model ^b	β coefficient	p-value	Whole model p-value	Whole model R ²
2	AKAP9	+0.41	0.0028	0.00038	0.38
	FEZ1	−0.45	0.0021		
	PCM1	−0.34	0.015		
	FEZ1 × PCM1	−0.65	0.000019		
3	NDEL1	−0.39	0.004	0.0052	0.21
	DISC1	+0.33	0.014		
	NDEL1 × DISC1	−0.65	0.0058		
4	AKAP9	+0.47	0.00046	0.0013	0.23
	DISC1	−0.26	0.038		
	AKAP9 × DISC1	−0.28	0.0003		

^aPC = principal component.

^bThe model explaining the most variation within each PC was determined by backward removal multiple regression analysis including main effects and gene-by-gene interaction terms.

doi:10.1371/journal.pone.0099892.t003

Table 4. Relationship of gene expression levels to activity in the prefrontal cortex elicited by the N-back task.

PC ^a	Regression term (gene/gene × gene interaction)	Brodmann area showing significantly correlated activity	Direction of correlation	Corrected <i>p</i> -value ^b	Whole model <i>p</i> -value
1	NDEL1	9	POSITIVE	0.093	0.093
2	AKAP9	9	NEGATIVE	0.035	0.01
	PCM1	45	NEGATIVE	0.029	
	FEZ1	45	NEGATIVE	0.040	
	FEZ1 × PCM1	45	POSITIVE	0.026	
3	NDEL1	6	NEGATIVE	0.025	0.00083
	DISC1	6	NEGATIVE	0.023	
	NDEL1 × DISC1	6	POSITIVE	0.022	
4	AKAP9	9	POSITIVE	0.019	0.00078
	DISC1	9	POSITIVE	0.007	
	AKAP9 × DISC1	9	NEGATIVE	0.009	

^aPC = principal component.

^bCorrected by permutation analysis (10⁵ permutations).

doi:10.1371/journal.pone.0099892.t004

Further studies are required to characterise the relationship between blood expression of *DISC1* and schizophrenia.

Assaying expression in peripheral blood rather than post-mortem brains avoids the confounds of agonal conditions and post-mortem delay, which can affect RNA integrity and thus gene expression [34,35]. Cognisant of these caveats, it should be noted that Lipska et al. [15] found the hippocampal mRNA expression of *NDEL1* but not *DISC1* to be reduced in schizophrenic patients. In contrast, ~75 kDa *DISC1* hippocampal protein expression was found to be increased in schizophrenic patients [15]. A recent study identified decreased 91 kDa *DISC1* immunoreactivity in the dorsolateral prefrontal cortex (DLPFC) in schizophrenic patients [36], while studies of *DISC1* mRNA expression in this region have found no differences between schizophrenic patients and controls [15,37]. There are several possible explanations for the apparent discrepancies between these studies: schizophrenia-associated changes in gene expression might be isoform, tissue- and/or brain region- specific; and/or aetiological heterogeneity might have resulted in systematic between-study sample differences. It is clear that further studies are required to distinguish between these possibilities.

While the relevance of blood expression to brain expression is a subject of debate, the use of blood RNA is a practical necessity when conducting within-subjects studies of gene expression and cognition. Additional studies in which blood and brain expression are compared within-samples would aid in the interpretation of blood-based expression findings; however, regardless of the relationship between blood and brain expression levels, our findings provide encouragement for the possibility raised by previous studies that blood-based gene expression profiling may eventually aid in diagnosis [38,39].

In summary, this study highlights the contribution of the coordinated expression of *DISC1*-interactome genes to cognitive functions relevant to schizophrenia. Future studies should attempt to clarify the specificity of the relationship between the expression levels of *DISC1*-interactome members and sub-domains of cognitive function in independent samples of healthy controls, as well as attempting to replicate our findings of aberrant expression in schizophrenic patients. In addition it would be of interest to (i) assess the relationship between the expression of *DISC1* and *NDEL1* and cognitive function in schizophrenic patients and (ii) identify the molecular processes affected by the altered expression of these genes. Such studies have the potential to both identify new

targets for therapeutic action, thus permitting the development of medications with greater efficacy in ameliorating the cognitive deficits of schizophrenia, and identify blood-based biomarkers, thus improving diagnosis.

Supporting Information

File S1 Table S1, Details of TaqMan probes used for qRT-PCR. List of TaqMan (Applied Biosystems) probes used to explore expression levels in ten genes of the *DISC1* interactome. Probes were chosen to span an intron-exon boundary and to amplify the maximum possible number of isoforms possible. Table S2, Factor loadings of cognitive tasks. Factors with eigenvalues of at least 1 were identified by principal components analysis followed by varimax rotation. For each cognitive test, the function explored is detailed together with the loading of the test on each of the four identified factors. Table S3, Assessment of the effect of gene expression levels on prefrontal cortical activity elicited by the N-back task. Correlations were assessed between gene expression and activity in prefrontal cortex during the N-back task (2-back condition). For each correlation, the Brodmann's area (BA) involved, Talairach coordinates, *k* values, *z*-score, and familywise error (FWE) rate corrected *p*-value are indicated. Where a *p*-value did not survive multiple testing correction, this is indicated and the uncorrected *p*-value is presented. Figure S1, Patterns of co-expression amongst *DISC1* pathway genes. Graphical representation of the co-expression patterns amongst *DISC1* and nine *DISC1*-interactor genes. Lines between pairs of genes represent significant positive (continuous lines) and negative (dashed lines) Pearson correlations. Line width represents a visual proxy of correlation strength, as indicated. Figure S2, Heat map representations of the relationship between the expression of *DISC1* pathway genes and cognitive performance. Depicted are heat maps representing the relationship between A. changes in PC3 (Phonological Fluency-Speed of Processing) in relation to changes in expression levels of *DISC1* and *NDEL1*, and B. changes in PC4 (Semantic Fluency) in relation to changes in expression levels of *DISC1* and *AKAP9*. Figure S3, Relationship between *DISC1* and *NDEL1* expression levels and frontal cortex function. The relationship between frontal cortical activity elicited by the N-back task and the expression levels of *NDEL1*, *DISC1*, and the interaction between *NDEL1* and *DISC1* expression was assessed using the general linear model. A. Heat map graphical

representation of changes in fMRI BOLD signal in BA 6 during the N-back task in relation to changes in expression levels of DISC1 and NDEL1. B. Negative main effect of NDEL1 expression level on BA 6 activity during the N-back task when including NDEL1 expression level, DISC1 expression level and the DISC1-by-NDEL1 interaction term in the model. C. Negative main effect of DISC1 expression level on BA 6 activity during the N-back task when including NDEL1 expression level, DISC1 expression level and the NDEL1-by-DISC1 interaction term in the model. D. Positive effect of the interaction term, DISC1-by-NDEL1 expression level, on BA 6 activity during the N-back task when including NDEL1, DISC1, and the DISC1-by-NDEL1 interaction term in the model. Figure S4, Relationship between AKAP9 and DISC1 expression levels and prefrontal cortex function. The relationship between frontal cortical activity elicited by the N-back task and the expression levels of AKAP9, DISC1, and the interaction between AKAP9 and DISC1 expression was assessed using the general linear model. The model included AKAP9 expression level, DISC1 expression level and the AKAP9-by-DISC1 interaction. A. Heat map graphical representation of changes in fMRI BOLD signal in BA 6 during the N-back task in relation to changes in expression levels of DISC1 and AKAP9. B. Positive main effect of DISC1 expression level on BA 9 activity during the N-back task when including AKAP9 expression level,

DISC1 expression level and the DISC1-by-AKAP9 interaction term in the model. C. Positive main effect of AKAP9 expression level on BA 9 activity during the N-back task when including AKAP9 expression level, DISC1 expression level and the DISC1-by-AKAP9 interaction term in the model. D. Negative effect of the AKAP9-by-DISC1 interaction on BA 9 activity during the N-back task when including AKAP9, DISC1 and the DISC1-by-AKAP9 interaction term in the model. (DOC)

Acknowledgments

We thank Riccarda Lomuscio, Rosemary Tomczynski and Heather Davidson for management of the collaboration between Bari and Edinburgh. We thank Andrew McIntosh and Heather Whalley for helpful comments on a draft manuscript. TaqMan qRT-PCR was carried out at the Wellcome Trust CRF Genetics Core, Western General Hospital, Edinburgh.

Author Contributions

Conceived and designed the experiments: AR JKM DJP AB KLE. Performed the experiments: AR HST SMA LF ADG PT BG RR RM GU GC. Analyzed the data: AR RMW LF PAT. Wrote the paper: AR RMW HST SMA LF ADG PT BG RR RM GU GC GB JKM DJP PAT AB KLE.

References

- Barch DM, Ceaser A (2012) Cognition in schizophrenia: core psychological and neural mechanisms. *Trends Cogn Sci* 16: 27–34.
- Barch DM, Csernansky JG, Conturo T, Snyder AZ (2002) Working and long-term memory deficits in schizophrenia: is there a common prefrontal mechanism? *J Abnorm Psychol* 111: 478–494.
- Blokland GA, McMahon KL, Thompson PM, Martin NG, de Zubicaray GI, et al. (2011) Heritability of working memory brain activation. *J Neurosci* 31: 10882–10890.
- Snitz BE, Macdonald AW 3rd, Carter CS (2006) Cognitive deficits in unaffected first-degree relatives of schizophrenia patients: a meta-analytic review of putative endophenotypes. *Schizophr Bull* 32: 179–194.
- Green MF, Kern RS, Braff DL, Mintz J (2000) Neurocognitive deficits and functional outcome in schizophrenia: are we measuring the “right stuff”? *Schizophr Bull* 26: 119–136.
- Money TT, Scarr E, Udawela M, Gibbons AS, Jeon WJ, et al. (2010) Treating schizophrenia: novel targets for the cholinergic system. *CNS Neurol Disord Drug Targets* 9: 241–256.
- Bressler SL, Menon V (2010) Large-scale brain networks in cognition: emerging methods and principles. *Trends Cogn Sci* 14: 277–290.
- Libby LA, Ragland JD (2011) fMRI as a Measure of Cognition Related Brain Circuitry in Schizophrenia. *Curr Top Behav Neurosci*.
- Callicott JH, Bertolino A, Mattay VS, Langheim FJ, Duyn J, et al. (2000) Physiological dysfunction of the dorsolateral prefrontal cortex in schizophrenia revisited. *Cereb Cortex* 10: 1078–1092.
- Bradshaw NJ, Porteous DJ (2012) DISC1-binding proteins in neural development, signalling and schizophrenia. *Neuropharmacology* 62: 1230–1241.
- Chubb JE, Bradshaw NJ, Soares DC, Porteous DJ, Millar JK (2008) The DISC1 locus in psychiatric illness. *Mol Psychiatry* 13: 36–64.
- Thomson PA, Malavasi EL, Grunewald E, Soares DC, Borkowska M, et al. (2013) DISC1 genetics, biology and psychiatric illness. *Front Biol (Beijing)* 8: 1–31.
- Camargo LM, Collura V, Rain JC, Mizuguchi K, Hermyjakob H, et al. (2007) Disrupted in Schizophrenia 1 Interactome: evidence for the close connectivity of risk genes and a potential synaptic basis for schizophrenia. *Mol Psychiatry* 12: 74–86.
- Tomppo L, Hennah W, Lahermo P, Loukola A, Tuulio-Henriksson A, et al. (2009) Association between genes of Disrupted in schizophrenia 1 (DISC1) interactors and schizophrenia supports the role of the DISC1 pathway in the etiology of major mental illnesses. *Biol Psychiatry* 65: 1055–1062.
- Lipska BK, Peters T, Hyde TM, Halim N, Horowitz C, et al. (2006) Expression of DISC1 binding partners is reduced in schizophrenia and associated with DISC1 SNPs. *Hum Mol Genet* 15: 1245–1258.
- Hennah W, Porteous D (2009) The DISC1 pathway modulates expression of neurodevelopmental, synaptogenic and sensory perception genes. *PLoS One* 4: e4906.
- Richards AL, Jones L, Moskvina V, Kirov G, Gejman PV, et al. (2012) Schizophrenia susceptibility alleles are enriched for alleles that affect gene expression in adult human brain. *Mol Psychiatry* 17: 193–201.
- Maxwell ME (1992) Family Interview for Genetic Studies (FIGS): A Manual for FIGS. Bethesda, MD: NIMH: Clinical Neurogenetics Branch.
- Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, et al. (2002) Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol* 3: RESEARCH0034.
- Burns MJ, Nixon GJ, Foy CA, Harris N (2005) Standardisation of data from real-time quantitative PCR methods - evaluation of outliers and comparison of calibration curves. *BMC Biotechnol* 5: 31.
- Dickinson D, Goldberg TE, Gold JM, Elvevag B, Weinberger DR (2010) Cognitive factor structure and invariance in people with schizophrenia, their unaffected siblings, and controls. *Schizophr Bull* 37: 1157–1167.
- Genderson MR, Dickinson D, Diaz-Asper CM, Egan MF, Weinberger DR, et al. (2007) Factor analysis of neurocognitive tests in a large sample of schizophrenic probands, their siblings, and healthy controls. *Schizophr Res* 94: 231–239.
- Bertolino A, Caforio G, Blasi G, De Candia M, Latorre V, et al. (2004) Interaction of COMT (Val108/158)Met genotype and olanzapine treatment on prefrontal cortical function in patients with schizophrenia. *Am J Psychiatry* 161: 1798–1805.
- Maldjian JA, Laurienti PJ, Kraft RA, Burdette JH (2003) An automated method for neuroanatomic and cytoarchitectonic atlas-based interrogation of fMRI data sets. *Neuroimage* 19: 1233–1239.
- Owen AM, McMillan KM, Laird AR, Bullmore E (2005). “N-back working memory paradigm: a meta-analysis of normative functional neuroimaging studies.” *Hum Brain Mapp* 25(1): 46–59.
- Mottaghy FM, Gangitano M, Sparing R, Krause BJ, Pascual-Leone A (2002). “Segregation of areas related to visual working memory in the prefrontal cortex revealed by rTMS.” *Cereb Cortex* 12(4): 369–375.
- Hanakawa T, Honda M, Sawamoto N, Okada T, Yonekura Y, et al. (2002). “The role of rostral Brodmann area 6 in mental-operation tasks: an integrative neuroimaging approach.” *Cereb Cortex* 12(11): 1157–1170.
- Tanaka S, Honda M, Sadato N (2005). “Modality-specific cognitive function of medial and lateral human Brodmann area 6.” *J Neurosci* 25(2): 496–501.
- Condon B, Montaldi D, Wilson JT, Hadley D (1997). “The relation between MRI neuroactivation changes and response rate on a word-fluency task.” *Appl Neuropsychol* 4(4): 201–207.
- Gadelha A, Machado MF, Yonamine CM, Sato JR, Juliano MA, et al. (2013) Plasma Ndel1 enzyme activity is reduced in patients with schizophrenia—a potential biomarker? *J Psychiatr Res* 47: 657–663.
- Kumarasinghe N, Beveridge NJ, Gardiner E, Scott RJ, Yasawardene S, et al. (2013) Gene expression profiling in treatment-naive schizophrenia patients identifies abnormalities in biological pathways involving AKT1 that are corrected by antipsychotic medication. *Int J Neuropsychopharmacol* 16: 1483–1503.
- Millar JK, Pickard BS, Mackie S, James R, Christie S, et al. (2005) DISC1 and PDE4B are interacting genetic factors in schizophrenia that regulate cAMP signaling. *Science* 310: 1187–1191.

33. Millar JK, Wilson-Annan JC, Anderson S, Christie S, Taylor MS, et al. (2000) Disruption of two novel genes by a translocation co-segregating with schizophrenia. *Hum Mol Genet* 9: 1415–1423.
34. Ferrer I, Martinez A, Boluda S, Parchi P, Barrachina M (2008) Brain banks: benefits, limitations and cautions concerning the use of post-mortem brain tissue for molecular studies. *Cell Tissue Bank* 9: 181–194.
35. Tomita H, Vawter MP, Walsh DM, Evans SJ, Choudary PV, et al. (2004) Effect of agonal and postmortem factors on gene expression profile: quality control in microarray analyses of postmortem human brain. *Biol Psychiatry* 55: 346–352.
36. Ratta-Apha W, Hishimoto A, Mouri K, Shiroyiwa K, Sasada T, et al. (2013). “Association analysis of the DISC1 gene with schizophrenia in the Japanese population and DISC1 immunoreactivity in the postmortem brain.” *Neurosci Res* 77(4): 222–227.
37. Rastogi A, Zai C, Likhodi O, Kennedy JL, Wong AH (2009) Genetic association and post-mortem brain mRNA analysis of DISC1 and related genes in schizophrenia. *Schizophr Res* 114: 39–49.
38. Lai CY, Yu SL, Hsieh MH, Chen CH, Chen HY, et al. (2011) MicroRNA expression aberration as potential peripheral blood biomarkers for schizophrenia. *PLoS One* 6: e21635.
39. Takahashi M, Hayashi H, Watanabe Y, Sawamura K, Fukui N, et al. (2010) Diagnostic classification of schizophrenia by neural network analysis of blood-based gene expression signatures. *Schizophr Res* 119: 210–218.