

Culicoides hildebrandoi, a new species of the *reticulatus* species group from the Brazilian Amazon Region (Diptera, Ceratopogonidae)

Emanuelle de Sousa Farias¹, Antonio Marques Pereira Júnior^{2,3},
Maria Luiza Felipe-Bauer⁴, Felipe Arley Costa Pessoa¹,
Jansen Fernandes Medeiros³, Maria Clara Alves Santarém⁴

1 Laboratório de Ecologia de Doenças Transmissíveis da Amazônia (EDTA), Centro de Pesquisa Leônidas e Maria Deane, Fiocruz Amazônia, Rua Terezina, 476, Adrianópolis, CEP 69057-070, Manaus, Amazonas, Brazil **2** Fundação Universidade Federal de Rondônia, UNIR, Campus BR 364, Km 9.5, CEP 76801-059, Porto Velho, Rondônia, Brazil **3** Laboratório de Entomologia, Fundação Oswaldo Cruz/Rondônia, Rua da Beira, 7671, Lagoa, CEP 76812-245, Porto Velho, Rondônia, Brazil **4** Coleção de Ceratopogonidae, Laboratório de Diptera, Instituto Oswaldo Cruz-Fiocruz, Av. Brasil, 4365, 21040-900 Rio de Janeiro, RJ, Brazil

Corresponding author: Jansen Fernandes Medeiros (jmdeiro@gmail.com)

Academic editor: Art Borkent | Received 28 November 2015 | Accepted 5 February 2016 | Published 7 March 2016

<http://zoobank.org/8B94BD74-A373-4722-9D9A-5C928C5B7277>

Citation: Farias ES, Pereira Júnior AM, Felipe-Bauer ML, Pessoa FAC, Medeiros JF, Santarém MCA (2016) *Culicoides hildebrandoi*, a new species of the *reticulatus* species group from the Brazilian Amazon Region (Diptera, Ceratopogonidae). ZooKeys 571: 105–111. doi: 10.3897/zookeys.571.7341

Abstract

A new species of biting midge (Diptera: Ceratopogonidae), *Culicoides hildebrandoi* **sp. n.**, is described and illustrated based on female and male specimens from the states of Amazonas and Rondônia, Brazil. This new species belongs to the *reticulatus* species group and differs from the 24 other species of this group by the elongate slightly swollen 3rd palpal segment with scattered capitate sensilla but lacking a sensory pit.

Keywords

Amazonas, biting midges, Brazil, Rondônia

Introduction

The biting midges genus *Culicoides* Latreille (Diptera, Ceratopogonidae), presently includes 1355 extant worldwide species (Borkent 2015a), some of which can transmit pathogenic viruses and filarial nematodes to humans and other vertebrates. Due to their epidemiological importance, *Culicoides* are the best studied genus of Ceratopogonidae. Most species of *Culicoides* have wings with unique patterns of light and dark spots. Species with similar wing patterns have been included in subgenera or species groups with other similar morphological characters that are important for species identification (Felippe-Bauer 2003; Borkent 2015b).

The Neotropical *reticulatus* species group includes 24 species (Santarém et al. 2015). Santarém et al. (2014) redescribed *Culicoides reticulatus* Lutz and described seven new species, five of which were from three states in the Amazon Region of Brazil (Amazonas, Pará and Roraima). Santarém et al. (2014) noted an apparently undescribed, poorly mounted female from the state of Amazonas that they declined to describe. During our study of material from the state of Rondônia, we discovered an additional 12 female and four male specimens which we describe and illustrate herein.

Materials and methods

Specimens were collected with CDC light traps in the forest on Rancho Colorado farm, Porto Velho municipality, state of Rondônia, Brazil. The specimens were preserved in 70% ethanol and subsequently slide-mounted in phenol-balsam with the methods described by Wirth and Marston (1968). The female specimen from Balbina, Amazonas studied by Santarém et al. (2014) was also examined. Diagnostic features were microphotographed using a Digital System (Synoptics™, Cambridge, UK) adapted to an optical microscope (Leica DMTM1000, Frankfurt, Germany). Images were taken with a digital camera (JVC 3CCDTM, Wayne, USA), and Auto Montage 4.0 used to obtain the final images.

Morphological terms are from the chapter on Ceratopogonidae by Borkent et al. (2009), in the recent the Manual of Central American Diptera. Terms of structures specific to *Culicoides* follow those described by Felippe-Bauer (2003). Measurements of spermathecae are in micrometers, whereas those of wings are in millimeters. Meristic information is presented as ranges of values, followed by mean and sample size. The holotype, allotype and some paratypes were deposited in the Ceratopogonidae Collection of Oswaldo Cruz Foundation (CCER), Rio de Janeiro, RJ, Brazil; other paratypes were deposited in the Laboratório de Ecologia de Doenças Transmissíveis da Amazônia, Leônidas and Maria Deane Institute (ILMD), Manaus, Amazonas, Brazil.

Results

Culicoides hildebrandoi sp. n.

<http://zoobank.org/78B8970A-218C-4CBC-8388-AA4B248A11A2>

Figs 1–3

Diagnosis. Female: only species of *Culicoides* in the Neotropical Region with the following combination of features: 2nd radial cell in dark spot, r_3 with four sparsely distributed pale spots, r-m crossvein pale; hind femur with subapical pale band; scutum with two anterior submedian clover-leaf shaped spots; third palpal segment elongate, slightly swollen, cylindrical, without a sensory pit but with capitate sensilla scattered on the surface cuticle. Male: only species in the Neotropical Region with the following combination of features: tergite 9 with a posteromedial notch, parameres with slightly sinuous stem, swollen on mid-portion and without a ventral lobe and the basal arch of aedeagus extending 2/3 of total length.

Description. Female. *Head* (Fig. 2d). Brown. Eyes bare, separated by distance equal to diameter of nearly one ommatidium (Fig. 2a). Antennal pedicel brown; flagellum pale brown, flagellomeres 1–8 pale on proximal ½; AR 0.90–1.00 (0.95, n = 8); sensilla coeloconica on flagellomeres 1, 6–8, two on 1, three on 6, three or four on 7 and four on 8. *Palpus* (Fig. 2b) brown; third segment elongate cylindrical, slightly swollen, without sensory pit, with sensilla scattered on surface; PR 3.30–3.90 (3.54, n = 12). Proboscis moderately long; P/H ratio 0.87–1.00 (0.95, n = 13); mandible (Fig. 2c) with 19–25 (22, n = 12) teeth.

Thorax (Fig. 2e). Dark brown, with prominent pattern of well-defined yellowish patches, humeral depression pale. Scutum with two anterior submedian clover-leaf shaped spots, two posterior submedian pale areas; lateral portion with anterior, posterior pale areas; prescutellar depressions pale; scutellum with yellowish lateral margins; postscutellum brown with pale median area. *Wing* (Fig. 1a) with contrasting pattern of pale and dark spots; distal ½ of 1st, all of 2nd radial cell in dark spot; pale spot over r-m extending from M_1 to just below radius; another pale spot from dorsal portion of radius to margin of costa; r_3 with four small separated pale spots: 1st rounded, between 2nd radial cell and M_1 ; 2nd extending posterior to 2nd radial cell, abutting wing margin; 3rd ovoid, in mid portion of cell, larger than 2nd spot; 4th distal pale spot close to 3rd spot, extending to wing margin; m_1 with two pale spots, 1st small, ovoid, beyond fork of M_1 and M_2 , 2nd larger than 1st, not close to wing margin; m_2 with four pale spots: 1st proximal to CuA, 2nd and 3rd between medial and mediocubital forks, 4th larger, not reaching wing margin; cua_1 with a rounded pale spot in middle of cell; anal cell with faint basal sinuous pale area and one distal pale spot near mediocubital fork that is nearly subdivided, abutting wing margin; wing base with faint pale spot on M ; apices of M_1 , M_2 and CuA_1 broadly pale; macrotrichia sparsely distributed on distal half of wing; wing length 1.10–1.30 (1.25, n = 13) mm, breadth 0.50–0.58 (0.56, n = 12) mm; CR 0.60–0.68 (0.65, n = 13). Halter stem pale, knob brown. *Legs* (Fig. 2g)




Figure 1. *Culicoides hildebrandoi* sp. n. Wing. **a** Female **b** Male.




Figure 2. a–g *Culicoides hildebrandoi*, female sp. n. **a** Eye separation, anterior view **b** Palpal segment 3 **c** Mandibular teeth **d** Head, anterior view **e** Thorax, dorsal view **f** Spermathecae **g** Legs (right to left) fore-, mid- and hind.




Figure 3. *Culicoides hildebrandoi*, male terminalia sp. n. **a** Terminalia and aedeagus (parameres removed), ventral view **b** Parameres, ventral view.

brown; femora with subapical, tibiae with subbasal pale bands; apex of hind tibia pale; hind tibial comb with four spines, that nearest spur longest.

Abdomen. Brown. Two subequal-size ovoid spermathecae (Fig. 2f), measuring 37.5–47.5 × 32.5–35.0 (n = 2) μm and 37.5–40.0 × 30.0–32.5 (n = 2) μm, slen-

der sclerotized necks with 7.5 μm ; third slender, elongate rudimentary spermatheca, length 27.5–30.0 ($n = 2$) μm .

Male. Similar to female with usual sexual differences. Sensilla coeloconica on flagellomeres 1, 6–10, one on 1, 6–8, two on 9, three on 10; AR 0.80–0.87 (0.85, $n = 4$). PR 1.6–2.0 (1.78, $n = 4$). Wing with pattern of pale spots as in Fig. 1b, wing length 0.95–1.08 (1.00, $n = 4$), breadth 0.35–0.40 (0.38, $n = 4$); CR 0.62–0.64 (0.63, $n = 4$). Terminalia (Fig. 3a): Tergite 9 long, tapered slightly at mid length, distal portion broader with short conical apicolateral processes, with distinct posteromedial notch; sternite 9 with rounded posteromedian excavation. Gonocoxite twice as long as broad, ventral, dorsal roots slender, elongated, sclerotized; gonostylus tapering distally, distal portion curved, apex broader with beak-like tip. Parameres (Fig. 3b) separate, each one with heavily sclerotized basal knob; stem long, curved near base, slightly sinuous, swollen on mid portion without ventral lobe; apical portion tapered, elongate, abruptly bent without lateral fringe of spicules. Aedeagus (Fig. 3a) Y-shaped; basal arms heavily sclerotized; basal arch triangular, extending 2/3 of total length; distal portion moderately slender, apex rounded.

Type material. Holotype female, labeled “Brazil, Rondônia, Porto Velho, Rancho Colorado Farm, 08°42'3.7"S; 63°59'3.8"W, 20.VIII.2014, CDC light trap, forest, Jansen F Medeiros col.” (CCER). Allotype male labeled as for female (CCER). Paratypes 12 females and three males: 11 females and three males same data as holotype (7 females, 2 males ILMD; 4 females, 1 male CCER); 1 female, Brazil, Amazonas, Balbina, 08.V.1985, CDC light trap, E. Castellón & S. Gomes col. (ILMD).

Distribution and bionomics. This is a forest species of the Amazon region of Brazil in the states of Amazonas and Rondônia.

Etymology. This species is named in memory of the late Dr. Luiz Hildebrando Pereira da Silva, a parasitologist who dedicated many years to studying Tropical Diseases such as Malaria and Chagas Disease. During his long career, Dr. Hildebrando was director of the Pasteur Institute, France. In Brazil he was a Medical School Professor at the University of São Paulo and Federal University of Rondônia. He also created the Institute for Research of Tropical Pathologies in Rondônia and was pivotal for the implementation of Oswaldo Cruz Foundation in Rondônia.

Taxonomic discussion

Culicoides hildebrandoi sp. n. is very similar to *C. reticulatus* and seven other closely related species described by Santarém et al. (2014) based on the number and distribution of pale spots on the wing and the clover-leaf pattern of yellowish patches on the anterio-submedian portion of the scutum. *Culicoides hildebrandoi* sp. n. can be distinguished from these eight congeners by its elongate, cylindrical third palpal segment with scattered capitate sensilla on its surface (third palpal segment is swollen and the capitate sensilla are in sensory pits in other species). Females of *C. hildebrandoi* sp. n. are larger than related species with a wing length of 1.10–1.30 mm (wing length < 1.0 mm in *C. amazonicus* Santarém, Felipe-Bauer & Trindade, *C. diplus* Santarém

& Felipe-Bauer, *C. profundus* Santarém, Felipe-Bauer & Trindade, *C. pseudoreticulatus* Santarém, Felipe-Bauer & Castellón and *C. rhombus* Santarém, Felipe-Bauer & Castellón), they have a more slender 3rd palpal segment, PR 3.3–3.9 (PR < 3.2 in the other species) and, a moderately long proboscis, P/H ratio 0.87–1.00 (P/H ratio ≥ 1.00 in *C. amazonicus*, *C. diplus*, *C. fluminensis* Santarém & Felipe-Bauer, *C. pseudoreticulatus*, *C. reticulatus* and *C. rhombus*).

This new species along with *C. amazonicus*, *C. irregularis* Santarém, Felipe-Bauer & Castellón, *C. profundus*, *C. pseudoreticulatus* and *C. rhombus* are associated with forested environments in the Brazilian Amazon Region, while the other three congeners are associated with mangrove swamps in the coastal regions of Colombia and Panama (*C. diplus*), Rio de Janeiro, Brazil (*C. fluminensis*), and Bahia and Pernambuco, Brazil (*C. reticulatus*).

With the description of *C. hildebrandoi* sp. n. here the “*reticulatus* species group” now has 25 species distributed throughout the Neotropics.

Acknowledgements

We would like to thank Bruno Mathieu, Gustavo R. Spinelli and William L. Grogan for their comments and for revision of the English. We further thank Tiago do Nascimento da Silva for assistance with the production of the Figure 3.

References

- Borkent A (2015a) World species of Biting Midges (Diptera: Ceratopogonidae). Illinois Natural History Museum. <http://www.inhs.illinois.edu/research/FLYTREE/Borkent.html>
- Borkent A (2015b) The subgeneric classification of species of *Culicoides*. Illinois Natural History Museum. <http://www.inhs.uiuc.edu/research/FLYTREE/CulicoidesSubgenera.pdf>
- Borkent A, Spinelli GR, Grogan WL (2009) Ceratopogonidae (biting midges, purrujas). Chapter 29. In: Brown BV, Borkent A, Cumming JM, Wood DM, Woodley NE, Zumbado MA (Eds) Manual of Central American Diptera. Volume 1. NRC Research Press, Ottawa, Ontario, Canada, 407–435.
- Felipe-Bauer ML (2003) A importância do padrão das manchas das asas em *Culicoides* (Latreille, 1809). *Entomologia y Vectores* 10: 595–600.
- Santarém MCA, Farias ES, Felipe-Bauer ML (2015) *Culicoides castelloni* sp. n. from the Brazilian Amazon Region with a revision of the *reticulatus* species group (Diptera, Ceratopogonidae). *Anais da Academia Brasileira de Ciências* 87: 955–972. doi: 10.1590/0001-3765201520140517
- Santarém MCA, Trindade RL, Silva TN, Castellón EG, Patiú CAM, Felipe-Bauer ML (2014) New Neotropical *Culicoides* and redescription of *Culicoides reticulatus* Lutz (Diptera: Ceratopogonidae). *Zootaxa* 3795: 255–274. doi: 10.11646/zootaxa.3795.3.2
- Wirth WW, Marston N (1968) A method for mounting small insects on microscope slides in Canada Balsam. *Annals of the Entomological Society of America* 61: 783–784. doi: 10.1093/aesa/61.3.783