



Redox Modulation at Work: Natural Phytoprotective Polysulfanes From *Alliums* Based on Redox-Active Sulfur

Awais Anwar^{1,2} · Emma Gould^{1,2} · Ryan Tinson^{1,2} · Javaid Iqbal^{1,2} · Chris Hamilton^{1,2}

Published online: 21 September 2018
© The Author(s) 2018

Abstract

Purpose of review This article provides a brief overview of natural phytoprotective products of *allium* with a special focus on the therapeutic potential of diallyl polysulfanes from garlic, their molecular targets and their fate in the living organisms. A comprehensive overview of antimicrobial and anticancer properties of published literature is presented for the reader to understand the effective concentrations of polysulfanes and their sensitivity towards different human pathogenic microbes, fungi, and cancer cell lines.

Recent findings The article finds polysulfanes potentials as new generation novel antibiotics and chemo preventive agent. The effective dose rates of polysulfanes for antimicrobial properties are in the range of 0.5–40 mg/L and for anticancer 20–100 μ M. The molecular targets for these redox modulators are mainly cellular thiols as well as inhibition and/or activation of certain cellular proteins in cancer cell lines.

Summary Antimicrobial and anticancer activities of polysulfanes published in the literature indicate that with further development, they could be promising candidates for cancer prevention due to their selectivity towards abnormal cells.

Keywords Garlic · *Allium sativum* · Polysulfanes · Reactive sulfur species · Glutathione · Redox modulators · Chemo-prevention

Introduction

Sulfur plays a major role in biology and is found in numerous peptides, proteins, and low molecular weight metabolites. Among the sulfur compounds found in plants, bacteria, fungi, and animals are many agents with unique chemical and biochemical properties linked to redox processes, metal binding, and catalytic reactions to name a few.

Nature provides a range of sulfur redox modulators from plants, fungi, bacteria, and animals that have been investigated to determine their therapeutic potential. Of which the genus *allium* presents a range of sulfur-based natural products with many benefits to human health from antimicrobial to anticancer effects. These natural products are particularly prominent

in garlic (*Allium sativum*) and onion (*Allium cepa*) mainly consisting of thiosulfates and polysulfanes. The biological activity of such compounds is often associated with a broad spectrum of (bio)chemical properties. Their modes of action are often associated with redox activity, catalysis, metal binding, enzyme inhibition, and/or radical generation allowing these reactive sulfur species (RSS) to interact with oxidative stressors, to affect the function of redox-sensitive cysteine proteins, and to disrupt the integrity of DNA and cellular membranes. This has been discussed in various reviews previously [1, 2]. In some cases, the biological activity of sulfur-containing plant products depends on initial enzymatic activation, which allows thiosulfates to be generated with high target selectivity. The antibiotic and anticancer activities of RSS make them interesting from a pharmacological perspective. Not surprisingly, research into the biochemical and pharmacological properties of these sulfur chemotypes is advancing rapidly especially as anticancer agents [3].

In order to understand how these sulfur compounds develop their biological activities, we need to consider the rather complicated chemistry of various sulfur chemotypes and their biochemical transformations. Sulfur redox networks provide a glimpse of sulfur-centered formation and transformation pathways in vivo [4]. Although such networks are continuously

This article is part of the Topical Collection on *Redox Modulators*

✉ Awais Anwar
awais@ecospray.com

¹ Ecospray Limited, Grange Farm, Hilborough, Thetford, Norfolk IP26 5BT, UK

² School of Pharmacy, University of East Anglia, Norwich Research Park, Norwich NR4 7TJ, UK

expanding, they serve as a snapshot of the sulfur redox chemistry known to date and illustrate the complexity of redox-active sulfur species. For full details on sulfur redox mechanisms and pathways, a previous review article by Jacob is worth reading to understand this chemistry [2].

Redox Sulfanes of *Allium*

Sulfur metabolism in plants provides a treasure of reactive sulfur species (RSS) that includes several chemically unusual substances, such as thiosulfonates and polysulfanes from *Alliums*.

Polysulfanes are the most abundant organosulfur metabolites produced by garlic (Fig. 1) and are the result of an enzymatic reaction involving the non-protein amino acid alliin (*S*-allyl-L-cysteine sulfoxide), stored in large amounts in the cytosol of the plant cells (5–14 mg/g fresh dry weight, 1.4% of fresh weight) and an enzyme, allinase, present in the vacuole [6]. Upon crushing the enzyme substrate reaction produces and intermediate thiosulfonates, allicin which is not very stable at room temperature (half-life 3.1 h at 20 °C) [7].

Upon heating, allicin undergoes a cascade of further chemical rearrangements leading to other organosulfur molecules such as ajoene, dithiines, and predominantly diallyl polysulfanes (Fig. 1). The diallyl polysulfanes with DAS1–DAS4 constitute a major part of garlic extract and oils with others like methylated polysulfanes [8]. In this article, we will limit discussion to the diallyl polysulfanes mentioned in Fig. 1.

These diallyl polysulfanes (DAS1–DAS6) depicted in Fig. 1 exhibit distinct redox properties, which provide an interesting spectrum of biological activities in vivo, such as antibiotic, fungicidal, pesticidal, or anticancer activity. The last decade has provided an insight into the molecular basis for such activity and has achieved a better knowledge of the in

vitro properties of diallyl polysulfanes. This has led to an improved understanding of their impact on intracellular redox signaling and control pathways in living cells.


The full impact of sulfur in living systems becomes apparent by considering the diversity of sulfur species and their reactions. The biological activity of RSS can be attributed to sulfur and its uniqueness to exist in various oxidation states that exist naturally and able to transform in in vivo environments to plethora of different chemotypes. The oxidation states and various chemotypes have been discussed in a detailed review previously [2]. The redox-active sulfur species are able to oxidize thiols to generate oxidative stressors (e.g., peroxides, hydroxyl radicals, hydroperulfides, persulfides) which in turn results in cocktail of RSS that can adopt various pathways in the cell.

Medicinal/Pharmacological Properties of Sulfanes

Antimicrobial Activity

The key studies accredited allicin as the main contributor for the antimicrobial activity of garlic [9]. Allicin was found to inhibit bacterial growth as a vapor of lung pathogenic bacteria from the genera *Pseudomonas*, *Streptococcus*, and *Staphylococcus*, including multi-drug-resistant (MDR) strains, suggesting that it could be used to combat bacterial lung infections via direct inhalation; currently, there are no volatile antibiotics available to treat pulmonary infections [10]. Growth inhibition of *Escherichia coli* during allicin exposure coincides with a depletion of the glutathione pool and *S*-allylmercapto-modification of proteins, resulting in an overall decreased total sulfhydryl levels, which is accompanied by the induction of the oxidative and heat stress response. The mode of action of allicin is a combination of a decrease of

Fig. 1 A selection of the redox-sulfur chemistry found in garlic. Garlic produces various other chemicals which are not part of this review. The polysulfanes produced by garlic on crushing vastly depend on methods of extraction and temperature [5]. In figure DAS1 (diallyl sulfane), DAS2 (diallyl disulfane), DAS3 (diallyl trisulfane), DAS4 (diallyl tetrasulfane), DAS5 (diallyl pentasulfane)



glutathione levels, unfolding stress, and inactivation of crucial metabolic enzymes through *S*-allylmercapto-modification of cysteines [11]. It is suggested that allicin's ability to permeabilize cell membranes may contribute to its antimicrobial activity independently of its activity as a thiol reagent [12].

Garlic oil and its major diallyl polysulfanes constituents, as well as garlic extract, and allicin possess significant activity against *H. pylori* [13]. DAS2 and DAS3 were found to be the most abundant components in garlic oil. Interestingly, DAS4 also inhibits *H. pylori*, an activity which might be relevant in the context of garlic consumption and stomach ulcers. The minimum inhibitory concentrations (MICs) and minimum bactericidal concentrations (MBCs) of DAS4 against the NCTC 11637, and 107018 B strains of *H. pylori* are between 3 and 6 µg/mL, i.e., even lower than the values for allicin (6–12 µg/mL). DAS1, DAS2, and DAS3 were less active, with MIC values of 2074–4148, 100–200, and 13–25 µg/mL, respectively [14].

There have been many studies using garlic to combat clinically important bacteria and fungi. Garlic extract was tested against three major antibiotic resistant pathogens: *C. albicans*, MRSA, and *P. aeruginosa*. Interestingly, a synergism effect was reported of garlic extract improving the effectiveness of the antibiotics [15]. A separate study on 30 clinical isolates of MRSA found that allicin, extracted from garlic, caused 88% of the strains tested to have MICs of 16 µg/mL and all strains were inhibited at 32 µg/mL [16]. Garlic extract was used in a study on the drug resistant pathogens *E. coli*, *P. aeruginosa*, *B. subtilis*, *S. aureus*, *K. pneumoniae*, *S. sonnei*, *S. epidermidis*, and *S. typhi*. All pathogens showed high susceptibility to garlic extract where the lowest MIC was 0.05 mg/mL [17]. Studies using synthetic allicin, which included multi-drug resistant (MDR) strains, showed that the growth of the majority of *Pseudomonas*, *Streptococcus*, and *Staphylococcus* isolates was completely inhibited by 64 µg/mL allicin. *S. pyogenes* (SNo 67467), *S. pneumoniae* (SNo 68668), and *S. aureus* (ATCC 43300) were completely inhibited by 32 µg/mL allicin and all *A. baumannii* isolates were completely inhibited by 16 µg/mL. However, *K. pneumoniae* isolates were slightly more resistant, with a MIC of 128 µg/mL. *P. aeruginosa* (DSM2659) showed high resistance to allicin (MIC 512 µg/mL) compared to *P. aeruginosa* (PAO1 SBUG8 and PAO25), MIC 64 µg/mL. MDR and non-MDR *S. pneumoniae* strains tested were equally susceptible to allicin and showed MICs from 32 to 64 µg/mL allicin and MBCs from 64 to 128 µg/mL allicin, respectively. This study shows that different strains have different susceptibilities to garlic and its constituents [10].

Antifungal studies found that garlic oil can penetrate the cell membrane of *C. albicans* as well as organelle membranes, such as mitochondria, which would result in destruction of the organelle and ultimately cell death. Due to their lipophilic nature, it is likely that many of the diallylpolysulfanes can pass through membranes of various organisms [18] and they

have been shown to interact with membrane lipids to modify membrane fluidity [19].

Garlic oil has been shown to induce differential expression of important genes such as those involved in pathogenesis, oxidation-reduction, and cellular response to drugs and starvation [20]. Allicin and aged garlic extracts exhibit antimicrobial properties against the *Burkholderia cepacia* complex (Bcc), an intrinsically multi-resistant and life-threatening human pathogen showing the modification of cysteine residue which suggest allicin ability as a general electrophilic reagent targeting protein thiols [21, 22].

An important observation, which is demonstrated in Tables 1 and 2, is the different susceptibilities of pathogens to the different garlic constituents. DAS3 and DAS4 from garlic exhibit a wide spectrum of antimicrobial, antibacterial and antifungal activities. For example, DAS3 and DAS4 both inhibit *S. aureus* (MIC 2.0 and 0.5 µg/mL, respectively), *S. aureus* (MRSA) (MIC 8.0 and 2.0 µg/mL, respectively), *C. albicans* (MIC 1.0 and 0.5 µg/mL, respectively), *C. krusei* (MIC 8.0 and 4.0 µg/mL, respectively), *C. glabrata* (MIC 4.0 and 2.0 µg/mL, respectively), *A. niger* (MIC 2.0 and 1.0 µg/mL, respectively), *A. fumigatus* (MIC 8.0 and 4.0 µg/mL, respectively), and *A. flavus* (MIC 4.0 and 2.0 µg/mL, respectively). While the DAS3 is consistently less active, DAS4 possesses an antibiotic activity comparable to that of allicin. For example, against *H. pylori*, DAS4 has a MIC of 3–6 mg/L and allicin has a MIC of 6–12 mg/L [23].

Garlic was tested for synergistic effects with antibiotics (levofloxacin, gentamicin, azithromycin, and doxycycline) against *Pseudomonas* and *Acinetobacter* genera. This results in a decrease in the antibiotic MIC of 4->32, 4->2048, 2->2048 and 2->128 fold, respectively. The garlic increased the rate of lethality of the antibiotics against the bacteria. While these results show a potential for the synergistic use of garlic with antibiotics, a notable weakness of this study is it does not provide any details of the garlic preparation that is used [37, 38]. This unfortunately limits the meaningfulness of this piece of work.

Anticancer Activities

Polysulfanes have also been studied as a potential anticancer agent. Most studies have been conducted on DAS3 which shows this molecule as promising chemopreventive therapy for cancer. Studies with polysulfanes on different cancer lines and effective dose rates are summarized in Table 3. This research area has developed in the last decade enormously and various research groups have identified different targets in different types of cancer cell lines. The details of molecular targets identified by polysulfanes are highlighted in next section. This was noticed that some cancer cell lines are more sensitive to polysulfanes than others (Table 3). For example, DAS3 was effective at very low concentration (~2–9 mg/mL) in colon and breast cancer compared to gastric and skin (~29 mg/mL).

Table 1 MIC values (mg/L) of polysulfanes against different human pathogenic bacteria. MIC values have been converted to mg/L where other concentration units were reported in the literature

Garlic component/ preparation	Organism	MIC (mg/L)	Reference
Allicin	<i>H. pylori</i>	6–12	[23]
	<i>Pseudomonas</i> spp.	64	[10]
	<i>Streptococcus</i> spp.	64	
	<i>Staphylococcus</i> spp.	64	
	<i>P. aeruginosa</i>	64	
	<i>S. pneumonia</i>	32	
	<i>S. pyogenes</i>	32	
	<i>S. aureus</i>	32	
	<i>A. baumannii</i>	16	
	<i>E. coli</i>	23	[11]
DAS1	<i>H. pylori</i>	2074–4148	[23]
	<i>B. cereus</i>	64	[24]
	<i>C. jejuni</i>	56	
	<i>C. botulinium</i>	64	
	<i>E. coli</i>	72	
	<i>L. monocytogenes</i>	48	
	<i>S. enteric</i>	54	
	<i>S. aureus</i>	64	
	<i>V. cholerae</i>	72	
	<i>S. aureus</i>	2	[23]
DAS2	<i>H. pylori</i>	100	
	<i>B. cereus</i>	14	[24]
	<i>C. jejuni</i>	12	
	<i>C. botulinium</i>	20	
	<i>E. coli</i>	20	
	<i>L. monocytogenes</i>	8	
	<i>S. enteric</i>	12	
	<i>S. aureus</i>	16	
	<i>V. cholerae</i>	24	
	<i>S. aureus</i>	0.5	[23]
DAS3	<i>H. pylori</i>	13–25	
	<i>B. cereus</i>	4	[24]
	<i>C. jejuni</i>	2	
	<i>C. botulinium</i>	4	
	<i>E. coli</i>	12	
	<i>L. monocytogenes</i>	2	
	<i>S. enteric</i>	2	
	<i>S. aureus</i>	8	
	<i>V. cholera</i>	12	
	<i>M. tuberculosis</i>	2.5	[25]
DAS4	<i>H. pylori</i>	3–6	[23]
	<i>B. cereus</i>	1	[24]
	<i>C. jejuni</i>	1	
	<i>C. botulinium</i>	1	
	<i>E. coli</i>	4	
	<i>L. monocytogenes</i>	0.5	
	<i>S. enteric</i>	0.5	
	<i>S. aureus</i>	2	
	<i>V. cholerae</i>	4	

Table 1 (continued)

Garlic component/ preparation	Organism	MIC (mg/L)	Reference
Garlic oil ^a	<i>H. pylori</i>	8–32	[23]
	<i>B. cereus</i>	40	[24]
	<i>C. jejuni</i>	36	
	<i>C. botulinium</i>	32	
	<i>E. coli</i>	48	
	<i>L. monocytogenes</i>	20	
	<i>S. enteric</i>	32	
	<i>S. aureus</i>	36	
	<i>V. cholera</i>	40	
	<i>S. boydii</i>	2.75	[26]
	<i>S. flexnar</i>	2.75	
	<i>S. fluvialis</i>	2.75–5.5	
	<i>V. metschnikovii</i>	0.34	
	<i>V. parahaemolyticus</i>	0.08	
	<i>V. enterocolitica</i>	0.68	
	<i>C. coli</i>	0.49	
	<i>C. lari</i>	0.49	
	<i>B. fragilis</i>	0.04	
	<i>B. subtilis</i>	0.17–0.68	
	<i>E. aerogenes</i>	0.68	
	<i>E. faecalis</i>	0.34	
	<i>K. aerogenes</i>	0.17	
	<i>P. vulgaris</i>	2.74	
<i>L. acidophilus</i>	0.34–2.75		
<i>S. faecalis</i>	0.34		
<i>S. mutans</i>	0.08		
<i>S. pyogenes</i>	0.04		
Garlic extract ^a	<i>P. gingivalis</i>	16.6	[27]
	<i>P. aeruginosa</i>	6	
	<i>A. actinomycetemcomitans</i>	62.5	
	<i>S. aureus</i>	4	[28]
	<i>E. coli</i>	7	
	<i>B. subtilis</i>	0.1	[29]
	<i>K. pneumonia</i>	0.2	
	<i>S. epidermidis</i>	0.9	
	<i>S. typhi</i>	0.02	
	<i>Proteus</i> spp.	7–21	[30]
	<i>H. pylori</i>	2–5	[31]
	<i>S. epidermidis</i>	22.9	[32]
	<i>S. pneumoniae</i>	30.3	
	<i>S. pyogenes</i>	33	
	<i>H. influenzae</i>	30.5	
<i>Shigella</i> spp.	15.6		
<i>P. aeruginosa</i>	3.5		
<i>S. mutans</i>	4–32	[33]	

^a Activity depends on how garlic oil and garlic extract are manufactured. Various papers depicting the biological activity of polysulfanes did not mention the characterization of such preparations. The concentrations of individual polysulfanes in an extract or oil widely depend on method of extraction or distillation

Table 2 MIC (mg/L) of polysulfanes against different pathogenic fungal species. MIC values have been converted to mg/L where other concentration units were reported in the literature

Garlic component/ preparation	Organism	MIC (mg/L)	Percentage inhibition at MIC (%)	Reference
Allicin	<i>C. albicans</i>	0.8	100	[34]
	<i>C. neoformans</i>	0.3		
	<i>C. parapsilosis</i>	0.15		
	<i>C. tropicalis</i>	0.3		
	<i>C. krusei</i>	0.3		
	<i>T. glabrata</i>	0.3		
	<i>Aspergillus</i> spp.	8–32		
DAS2	<i>T. rubrum</i>	12.5	90	[36]
	<i>C. krusei</i>	8	100	[14]
	<i>C. albicans</i>	1		
	<i>C. krusei</i>	8		
	<i>C. glabrata</i>	4		
	<i>A. niger</i>	2		
	<i>A. fumigates</i>	8		
DAS3	<i>A. flavus</i>	4		
	<i>C. krusei</i>	2	100	[14]
	<i>C. albicans</i>	0.5		
	<i>C. krusei</i>	4		
	<i>C. glabrata</i>	2		
	<i>A. niger</i>	1		
	<i>A. fumigates</i>	4		
<i>A. flavus</i>	2			
Garlic oil ^a	<i>C. albicans</i>	0.35	100	[20]
	<i>P. funiculosum</i>	0.69		
Garlic extract ^a	<i>Candida</i> sp.	14.9	100	[32]

^a Activity depends on how garlic oil and garlic extract are manufactured. Various papers depicting the biological activity of polysulfanes did not mention the characterization of such preparations. The concentrations of polysulfanes in an extract or oil widely depend on method of extraction or distillation

General reviews have been published on polysulfanes as chemopreventives with a special focus on DAS3 for pediatric cancer treatment. A recent review indicated that DAS3 is of importance not only for its remarkable antitumor and cancer preventive effects, as suggested by many in vitro and in vivo studies, but also because of its many health benefits, like improvements to immune-system function, radioprotection, and protection against microbial infections. These features make it a potential candidate for the treatment of pediatric cancers. Herman-Antosiewicz and Yi concluded the molecular targets of polysulfanes in cellular environment showing that it selectively targets the cancer cells [52, 53].

DAS3 is more potent than mono- and disulfides against skin cancer. DAS3 inhibits cell growth of human melanoma A375 cells and basal cell carcinoma (BCC) cells by increasing the levels of intracellular reactive oxygen species (ROS) and DNA damage and by inducing G2/M arrest, endoplasmic reticulum (ER) stress. A recent review focuses on the molecular mechanisms of garlic-derived allyl sulfides on skin cancer prevention [54]. Similar findings were observed in human

colon cancer cells HCT-15 and DLD-1. The growth of the cells was significantly suppressed by DAS3, but neither DAS nor DAS2 showed such an effect [55]. An interesting paper depicts that allyl group in polysulfanes are responsible for the disruption of microtubule network formation in human colon cancer cell line HT-29 cells [43]. The effective dose rate in different cancer cell lines and their molecular targets are summarized in Table 3 below.

Molecular Targets and Metabolism of Sulfanes

Low molecular weight (LMW) thiols such as glutathione (GSH) serve as intracellular redox buffers in most aerobic organisms. They play a central role in the essential maintenance of an intracellular reducing environment and neutralize the damaging effects of toxic oxidants. When the cellular concentrations of these LMW thiols are dramatically reduced, the ability to defend against, and survive, oxidative stress is severely impaired [56].

Table 3 Effect of polysulfanes on different human cancer cell lines and their molecular targets. The concentration units reported in the literature have all been converted to mg/mL so they can be compared with the antimicrobial activities in Tables 1 and 2

Redox modulator	Cancer type/cell line	Dose (mg/mL)	Target	Effect	Reference
Ajoene	Leukemia HL-60, U937, HEL and OCIM-1	9.36	Bcl-2 Caspase-3	Inhibition of proliferation and induction of apoptosis	[39]
DAS2	Breast cancer MDA-MB-231	2.6	Estrogen receptor (ER)-positive (KPL-1 and MCF-7) and -negative (MDA-MB-231 and MKL-F)	Growth inhibition of cancer cells by inducing apoptosis	[40]
DAS2	Breast MCF-7	29.2	Kinase protein Caspase-3	Inhibited proliferation of MCF-7 cells and increased apoptotic ratio	[41]
DAS2	Colon Caco-2, HT-29	29.2	Histone H3 and H4	Inhibition of cancer proliferation by interaction with HDAC pathway	[42]
DAS3	Liver HepG2	35	Caspase-3	Increased H ₂ O ₂ levels, lowered thiol levels and inhibited cell proliferation	[3]
DAS3	Colon HCT-15 DLD-1	2.0 2.3	Tubulin	Suppression of cell growth	[43]
DAS3	Prostate PC-3	7.1	CDK1	Inhibition of cells by dose dependent manner.	[44]
DAS3	Gastric BGC-823	20.5	Bcl-2	Inhibited viability of BGC-823 in vitro and modulated Bcl-2.	[45]
DAS3	Breast MDA-MB-231 HS 578t	1.78	MMP2/9	Suppressed metastasis	[46]
DAS3	Colon HT29	8.91	Focal adhesion kinase (FAK)	Inhibition of angiogenesis	[47]
DAS3	A375 Skin	17.8	Mitochondrial caspase pathway	Increase in ROS	[48]
DAS3	Lymphoma BC2, BC3, BCBL1, HBL6	3.56	NF- κ B	Apoptosis in primary effusion lymphoma [PEL]	[49]
DAS3	Prostate LNCaP, C4-2, TRAMP-C1	3.56	Androgen receptor (AR)	Decrease in AR levels	[50]
DAS3	Glioblastoma U87MG Neuroblastoma SH-SY5Y	17.8	Bcl-2	Inhibition and proliferation	[51]

Upon entering the target (i.e., microbes and cancer cells) polysulfanes undergo rapid thiol-polysulfide exchange reactions with these LMW thiols. The implications of this process are twofold: (a) cellular LMW thiol concentrations decrease making them more susceptible to oxidative stresses and (b) in parallel, RSS such as allyl persulfide, hydropolysulfide species are formed, which can enhance the production of toxic oxidants (e.g., hydrogen peroxide, superoxide) thereby increasing oxidative stress [2].

In addition, polysulfanes can also react with exposed cysteine thiols on proteins to form *S*-allyl modified proteins. Such cysteinyl-*S*-allylations processes can result in altered/impaired protein function [57]. In addition to the redox activity of these molecules, their lipophilicity may contribute to their biological activity (e.g., by interrupting membrane integrity, binding

to hydrophobic sites on proteins) [18, 58]. These different modes of action are summarized in Fig. 2.

In pathological conditions, it has been shown that organosulfur compounds from garlic have protective effects, which are mostly associated with antioxidant properties. Because of the ability of polysulfanes to target multiple biochemical pathways, many researchers have studied their roles. The major metabolic pathways of sulfanes in mammals include methylation, oxidation, glutathione, and *N*-acetyl conjugations, and in certain instances, after ingestion of raw garlic by a human, allyl methyl sulfide, allyl methyl disulfide, DAS2, and DAS3 were discovered in the breath of tested volunteers [63], therefore providing an early example of their biochemical transformations within the body.

mediated by H₂S-dependent mechanisms [86]. Computational and experimental studies have revealed that glutathione and cysteine are capable of releasing H₂S from diallyl trisulfide (DAS3), via thiol-polysulfide exchange pathways, but diallyl disulfide (DAS2) is a much poorer H₂S donor via an α -carbon nucleophilic substitution pathway [87, 88].

Intraperitoneal (*ip*) administration of DAS1, DAS2, and DAS3 in mice increased the activity of rhodanese. Moreover, DAS2 and DAS3 increased the total sulfane sulfur level and γ -cystathionase activity in the normal mouse kidney. Aldehyde dehydrogenase activity was inhibited in the kidney after DAS3 administration. The results indicated that none of the studied polysulfanes affected the level of bound sulfur or H₂S. Thus, it can be concluded that garlic-derived DAS2 and DAS3 can be a source of sulfane sulfur for renal cells but they are not connected with persulfide formation [89]. DAS1, DAS2, and DAS3 dissolved in corn oil were given intraperitoneally to mice for 10 days. It showed that polysulfanes had a beneficial effect in the mouse liver, decreasing reactive oxygen species and malondialdehyde levels, and increasing glutathione S-transferase activity and non-protein sulfhydryl group level. Moreover, DAS2 and DAS3 elevated the total sulfane sulfur pool and activity suggesting its antioxidant and regulatory capacities [90].

Some studies of DAS4 have shown that it induces reactive oxygen species (ROS) in normal cells similar to cancer cells in a time (0 to 60 min) and dose-dependent manner (0 to 50 μ M) [91]; it also activates both the eIF2 α and Nrf2/HO-1 pathways [92].

Conclusion

This review summarizes the numerous biological activities garlic diallyl polysulfanes are involved in. While there is extensive evidence that show the various medicinal benefits of garlic polysulfanes, there is a critical need for controlled clinical studies to strictly evaluate the safety and efficacy of these compounds for establishing sufficient application methods before medical use. Furthermore, polysulfanes are emerging as promising, environmentally benign pesticides which are not only safe for humans, but also various non-target species endangered by synthetic chemical pesticides [8, 93]. With further work, diallyl polysulfanes could provide the basis for the innovative development of novel antibiotics, fungicides, pesticides, and anticancer agents.

Compliance with Ethical Standards

Human and Animal Rights and Informed Consent This article does not contain any studies with human or animal subjects performed by any of the authors.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Münchberg U, Anwar A, Mecklenburg S, Jacob C. Polysulfides as biologically active ingredients of garlic. *Org Biomol Chem*. 2007;5: 1505–18. <https://doi.org/10.1039/b703832a>.
- Jacob C. A scent of therapy: pharmacological implications of natural products containing redox-active sulfur atoms. *Nat Prod Rep*. 2006;23:851–63.
- Iciek M, Kwiecien I, Wlodek L. Biological properties of garlic and garlic-derived organosulfur compounds. *Environ Mol Mutagen*. 2009;50(3):247–65. <https://doi.org/10.1002/em.20474>.
- Filomeni G, Rotilio G, Ciriolo MR. Molecular transduction mechanisms of the redox network underlying the antiproliferative effects of allyl compounds from garlic. *J Nutr*. 2008;138(11):2053–7.
- Block E, editor. *Garlic and other Alliums, the Lore and the Science*. Cambridge: RSC Publishing; 2010.
- Lawson LD. Garlic: a review of its medicinal effects and indicated active compounds. *Phytomedicines of Europe*. ACS Symposium Series, vol 691: American Chemical Society; 1998. p. 176–209.
- Fujisawa H, Suma K, Origuchi K, Seki T, Ariga T. Thermostability of allicin determined by chemical and biological assays. *Biosci Biotechnol Biochem*. 2008;72(11):2877–83. <https://doi.org/10.1271/bbb.80381>.
- Anwar A, Gould E, Tinson R, Groom M, Hamilton C. Think yellow and keep green—role of sulfanes from garlic in agriculture. *Antioxidants*. 2017;6(1):3.
- Cavallito CJ, Bailey JH. Allicin, the antibacterial principle of *Allium sativum*. I. Isolation, physical properties and antibacterial action. *J Am Chem Soc*. 1944;66(11):1950–1. <https://doi.org/10.1021/ja01239a048>.
- Reiter J, Levina N, van der Linden M, Gruhlke M, Martin C, Slusarenko AJ. Diallylthiosulfinate (allicin), a volatile antimicrobial from garlic (*Allium sativum*), kills human lung pathogenic bacteria, including MDR strains, as a vapor. *Molecules*. 2017;22(10). <https://doi.org/10.3390/molecules22101711>.
- Muller A, Eller J, Albrecht F, Prochnow P, Kuhlmann K, Bandow JE, et al. Allicin induces thiol stress in bacteria through S-allylmercapto modification of protein cysteines. *J Biol Chem*. 2016;291(22):11477–90. <https://doi.org/10.1074/jbc.M115.702308>.
- Gruhlke MC, Hemmis B, Noll U, Wagner R, Lühring H, Slusarenko AJ. The defense substance allicin from garlic permeabilizes membranes of *Beta vulgaris*, *Rhoeo discolor*, *Chara corallina* and artificial lipid bilayers. *Biochim Biophys Acta*. 2015;1850(4):602–11. <https://doi.org/10.1016/j.bbagen.2014.11.020>.
- Zardast M, Namakin K, Esmaelian Kaho J, Hashemi SS. Assessment of antibacterial effect of garlic in patients infected with *Helicobacter pylori* using urease breath test. *Avicenna J Phytomed*. 2016;6(5):495–501.
- O’Gara EA, Hill DJ, Maslin DJ. Activities of garlic oil, garlic powder, and their diallyl constituents against *Helicobacter pylori*. *Appl Environ Microbiol*. 2000;66(5):2269–73.
- Li G, Ma X, Deng L, Zhao X, Wei Y, Gao Z, et al. Fresh garlic extract enhances the antimicrobial activities of antibiotics on

- resistant strains in vitro. *Jundishapur J Microbiol.* 2015;8(5): e14814. <https://doi.org/10.5812/jjm.14814>.
16. Cutler RR, Wilson P. Antibacterial activity of a new, stable, aqueous extract of allicin against methicillin-resistant *Staphylococcus aureus*. *Br J Biomed Sci.* 2004;61(2):71–4.
 17. Gull I, Saeed M, Shaukat H, Aslam SM, Samra ZQ, Athar AM. Inhibitory effect of *Allium sativum* and *Zingiber officinale* extracts on clinically important drug resistant pathogenic bacteria. *Ann Clin Microbiol Antimicrob.* 2012;11:8. <https://doi.org/10.1186/1476-0711-11-8>.
 18. Anwar A, Burkholz T, Scherer C, Abbas M, Lehr C-M, Diederich M, et al. Naturally occurring reactive sulfur species, their activity against Caco-2 cells, and possible modes of biochemical action (vol 28, pg 251, 2008). *J Sulfur Chem.* 2008;29(5):573. <https://doi.org/10.1080/17415990802519103>.
 19. Tsuchiya H, Nagayama M. Garlic allyl derivatives interact with membrane lipids to modify the membrane fluidity. *J Biomed Sci.* 2008;15(5):653–60. <https://doi.org/10.1007/s11373-008-9257-8>.
 20. Li WR, Shi QS, Dai HQ, Liang Q, Xie XB, Huang XM, et al. Antifungal activity, kinetics and molecular mechanism of action of garlic oil against *Candida albicans*. *Sci Rep.* 2016;6:22805. <https://doi.org/10.1038/srep22805>.
 21. Wallock-Richards D, Doherty CJ, Doherty L, Clarke DJ, Place M, Govan JRW, et al. Garlic revisited: antimicrobial activity of allicin-containing garlic extracts against *Burkholderia cepacia* complex. *PLoS One.* 2014;9(12):e112726. <https://doi.org/10.1371/journal.pone.0112726>.
 22. Liu Q, Meng X, Li Y, Zhao C, Tang G, Li H. Antibacterial and antifungal activities of spices. *Int J Mol Sci.* 2017;18(6). <https://doi.org/10.3390/ijms18061283>.
 23. O'Gara EA, Hill DJ, Maslin DJ. Activities of garlic oil, garlic powder, and their diallyl constituents against *Helicobacter pylori*. *Appl Environ Microbiol.* 2000;66(5):2269–73.
 24. Rattanachaikunsopon P, Phumkhachorn P. Diallyl sulfide content and antimicrobial activity against food-borne pathogenic bacteria of chives (*Allium schoenoprasum*). *Biosci Biotechnol Biochem.* 2008;72(11):2987–91. <https://doi.org/10.1271/bbb.80482>.
 25. Oosthuizen C, Arbach M, Meyer D, Hamilton C, Lall N. Diallyl polysulfides from *Allium sativum* as immunomodulators, hepatoprotectors, and antimycobacterial agents. *J Med Food.* 2017;20(7):685–90. <https://doi.org/10.1089/jmf.2016.0137>.
 26. Ross ZM, O'Gara EA, Hill DJ, Sleightholme HV, Maslin DJ. Antimicrobial properties of garlic oil against human enteric bacteria: evaluation of methodologies and comparisons with garlic oil sulfides and garlic powder. *Appl Environ Microbiol.* 2001;67(1): 475–80. <https://doi.org/10.1128/AEM.67.1.475-480.2001>.
 27. Shetty S, Thomas B, Shetty V, Bhandary R, Shetty RM. An in-vitro evaluation of the efficacy of garlic extract as an antimicrobial agent on periodontal pathogens: a microbiological study. *Ayu.* 2013;34(4):445–51. <https://doi.org/10.4103/0974-8520.127732>.
 28. Saha SK, Saha S, Akhter SM, Khatun S, Islam MM, Roy P. In vitro determination of minimum inhibitory concentration of aqueous garlic extract and imipenem against *Staphylococcus aureus* and *Escherichia coli*. *Mymensingh Med J.* 2016;25(3):477–84.
 29. Gull I, Saeed M, Shaukat H, Aslam SM, Samra ZQ, Athar AM. Inhibitory effect of *Allium sativum* and *Zingiber officinale* extracts on clinically important drug resistant pathogenic bacteria. *Ann Clin Microbiol Antimicrob.* 2012;11:8. <https://doi.org/10.1186/1476-0711-11-8>.
 30. Durairaj, Srinivasan, Lakshmanaperumalsamy. In vitro antibacterial activity and stability of garlic extract at different pH and temperature. *Electron J Biol.* 2009;5(1).
 31. Cellini L, Di Campi E, Masulli M, Di Bartolomeo S, Allocati N. Inhibition of *Helicobacter pylori* by garlic extract (*Allium sativum*). *FEMS Immunol Med Microbiol.* 1996;13(4):273–7.
 32. Iwalokun BA, Ogunledun A, Ogbolu DO, Bamiro SB, Jimi-Omojola J. In vitro antimicrobial properties of aqueous garlic extract against multidrug-resistant bacteria and *Candida* species from Nigeria. *J Med Food.* 2004;7(3):327–33. <https://doi.org/10.1089/jmf.2004.7.327>.
 33. Fani MM, Kohanteb J, Dayaghi M. Inhibitory activity of garlic (*Allium sativum*) extract on multidrug-resistant *Streptococcus mutans*. *J Indian Soc Pedod Prev Dent.* 2007;25(4):164–8.
 34. Ankri S, Mirelman D. Antimicrobial properties of allicin from garlic. *Microbes Infect.* 1999;1(2):125–9.
 35. Shadkchan Y, Shemesh E, Mirelman D, Miron T, Rabinkov A, Wilchek M, et al. Efficacy of allicin, the reactive molecule of garlic, in inhibiting *Aspergillus* spp. in vitro, and in a murine model of disseminated aspergillosis. *J Antimicrob Chemother.* 2004;53(5): 832–6. <https://doi.org/10.1093/jac/dkh174>.
 36. Aala F, Yusuf UK, Jamal F, Rezaie S. Antimicrobial effects of allicin and ketoconazole on *Trichophyton rubrum* under in vitro condition. *Braz J Microbiol.* 2012;43(2):786–92. <https://doi.org/10.1590/s1517-83822012000200044>.
 37. Abouelfetouh M. Enhancement of antimicrobial activity of four classes of antibiotics combined with garlic. *Asian J Plant Sci.* 2012;11:148–52.
 38. Li G, Ma X, Deng L, Zhao X, Wei Y, Gao Z, et al. Fresh garlic extract enhances the antimicrobial activities of antibiotics on resistant strains in vitro. *Jundishapur J Microbiol.* 2015;8(5):e14814. <https://doi.org/10.5812/jjm.14814>.
 39. Hassan HT. Ajoene (natural garlic compound): a new anti-leukaemia agent for AML therapy. *Leuk Res.* 2004;28(7):667–71. <https://doi.org/10.1016/j.leukres.2003.10.008>.
 40. Nakagawa H, Kiyozuka Y, Uemura Y, Senzaki H, Shikata N, Hioki K, et al. Resveratrol inhibits human breast cancer cell growth and may mitigate the effect of linoleic acid, a potent breast cancer cell stimulator. *J Cancer Res Clin Oncol.* 2001;127(4):258–64.
 41. Lei X-y, Yao S-q, Zu X-y, Huang Z-x, Liu L-j, Zhong M, et al. Apoptosis induced by diallyl disulfide in human breast cancer cell line MCF-7. *Acta Pharmacol Sin.* 2008;29:1233. <https://doi.org/10.1111/j.1745-7254.2008.00851.x>.
 42. Druesne N, Pagniez A, Mayeur C, Thomas M, Cherbuy C, Duée P-H, et al. Diallyl disulfide (DADS) increases histone acetylation and p21 waf1/cip1 expression in human colon tumor cell lines. *Carcinogenesis.* 2004;25(7):1227–36. <https://doi.org/10.1093/carcin/bgh123>.
 43. Hosono T, Hosono-Fukao T, Inada K, Tanaka R, Yamada H, Iitsuka Y, et al. Alkenyl group is responsible for the disruption of microtubule network formation in human colon cancer cell line HT-29 cells. *Carcinogenesis.* 2008;29(7):1400–6. <https://doi.org/10.1093/carcin/bgn124>.
 44. Arunkumar A, Vijayababu MR, Kanagaraj P, Balasubramanian K, Aruldas MM, Arunakaran J. Growth suppressing effect of garlic compound diallyl disulfide on prostate cancer cell line (PC-3) *in vitro* *in vivo*. *Biol Pharm Bull.* 2005;28(4):740–3. <https://doi.org/10.1248/bpb.28.740>.
 45. Jiang XY, Zhu XS, Xu HY, Zhao ZX, Li SY, Li SZ, et al. Diallyl trisulfide suppresses tumor growth through the attenuation of Nrf2/Akt and activation of p38/JNK and potentiates cisplatin efficacy in gastric cancer treatment. *Acta Pharmacol Sin.* 2017;38(7):1048–58. <https://doi.org/10.1038/aps.2016.176>.
 46. Liu Y, Zhu P, Wang Y, Wei Z, Tao L, Zhu Z, et al. Antimetastatic therapies of the polysulfide diallyl trisulfide against triple-negative breast cancer (TNBC) via suppressing MMP2/9 by blocking NF-kappaB and ERK/MAPK signaling pathways. *PLoS One.* 2015;10(4):e0123781. <https://doi.org/10.1371/journal.pone.0123781>.
 47. Lai KC, Hsu SC, Yang JS, Yu CC, Lein JC, Chung JG. Diallyl trisulfide inhibits migration, invasion and angiogenesis of human colon cancer HT-29 cells and umbilical vein endothelial cells, and

- suppresses murine xenograft tumour growth. *J Cell Mol Med.* 2015;19(2):474–84. <https://doi.org/10.1111/jcmm.12486>.
48. Wang HC, Yang JH, Hsieh SC, Sheen LY. Allyl sulfides inhibit cell growth of skin cancer cells through induction of DNA damage mediated G2/M arrest and apoptosis. *J Agric Food Chem.* 2010;58(11):7096–103. <https://doi.org/10.1021/jf100613x>.
 49. Shigemi Z, Furukawa Y, Hosokawa K, Minami S, Matsuhiro J, Nakata S, et al. Diallyl trisulfide induces apoptosis by suppressing NF-kappaB signaling through destabilization of TRAF6 in primary effusion lymphoma. *Int J Oncol.* 2016;48(1):293–304. <https://doi.org/10.3892/ijo.2015.3247>.
 50. Stan SD, Singh SV. Transcriptional repression and inhibition of nuclear translocation of androgen receptor by diallyl trisulfide in human prostate cancer cells. *Clin Cancer Res.* 2009;15(15):4895–903. <https://doi.org/10.1158/1078-0432.Ccr-09-0512>.
 51. Jurkowska H, Wróbel M, Kaczor-Kamińska M, Jasek-Gajda E. A possible mechanism of inhibition of U87MG and SH-SY5Y cancer cell proliferation by diallyl trisulfide and other aspects of its activity. *Amino Acids.* 2017;49(11):1855–66. <https://doi.org/10.1007/s00726-017-2484-4>.
 52. Antosiewicz J, Ziolkowski W, Kar S, Powolny AA, Singh SV. Role of reactive oxygen intermediates in cellular responses to dietary cancer chemopreventive agents. *Planta Med.* 2008;74(13):1570–9. <https://doi.org/10.1055/s-2008-1081307>.
 53. Yi L, Su Q. Molecular mechanisms for the anti-cancer effects of diallyl disulfide. *Food Chem Toxicol.* 2013;57:362–70. <https://doi.org/10.1016/j.fct.2013.04.001>.
 54. Wang Z, Xia Q, Cui J, Diao Y, Li J. Reversion of P-glycoprotein-mediated multidrug resistance by diallyl trisulfide in a human osteosarcoma cell line. *Oncol Rep.* 2014;31(6):2720–6. <https://doi.org/10.3892/or.2014.3154>.
 55. Seki T, Hosono T, Hosono-Fukao T, Inada K, Tanaka R, Ogihara J, et al. Anticancer effects of diallyl trisulfide derived from garlic. *Asia Pac J Clin Nutr.* 2008;17(Suppl 1):249–52.
 56. Masip L, Veeravalli K, Georgiou G. The many faces of glutathione in bacteria. *Antioxid Redox Signal.* 2006;8(5–6):753–62. <https://doi.org/10.1089/ars.2006.8.753>.
 57. Ariga T, Seki T. Antithrombotic and anticancer effects of garlic-derived sulfur compounds: a review. *BioFactors.* 2006. p. 93–103.
 58. Schneider T, Ba LA, Khairan K, Zwergel C, Bach ND, Bernhardt I, et al. Interactions of polysulfanes with components of red blood cells. *Med Chem Commun.* 2011;2(3):196–200. <https://doi.org/10.1039/c0md00203h>.
 59. Davenport DM, Wargovich MJ. Modulation of cytochrome P450 enzymes by organosulfur compounds from garlic. *Food Chem Toxicol.* 2005;43(12):1753–62. <https://doi.org/10.1016/j.fct.2005.05.018>.
 60. Nian H, Delage B, Pinto JT, Dashwood RH. Allyl mercaptan, a garlic-derived organosulfur compound, inhibits histone deacetylase and enhances Sp3 binding on the P21WAF1 promoter. *Carcinogenesis.* 2008;29(9):1816–24. <https://doi.org/10.1093/carcin/bgn165>.
 61. Lawson LD, Wang ZJ. Allicin and allicin-derived garlic compounds increase breath acetone through allyl methyl sulfide: use in measuring allicin bioavailability. *J Agric Food Chem.* 2005;53(6):1974–83. <https://doi.org/10.1021/jf048323s>.
 62. Germain E, Auger J, Ginies C, Siess MH, Teyssier C. In vivo metabolism of diallyl disulphide in the rat: identification of two new metabolites. *Xenobiotica.* 2002;32(12):1127–38.
 63. Taucher J, Hansel A, Jordan A, Lindinger W. Analysis of compounds in human breath after ingestion of garlic using proton-transfer-reaction mass spectrometry. *J Agric Food Chem.* 1996;44(12):3778–82. <https://doi.org/10.1021/jf960640e>.
 64. Xiao D, Herman-Antosiewicz A, Antosiewicz J, Xiao H, Brisson M, Lazo JS, et al. Diallyl trisulfide-induced G(2)-M phase cell cycle arrest in human prostate cancer cells is caused by reactive oxygen species-dependent destruction and hyperphosphorylation of Cdc25C. *Oncogene.* 2005;24(41):6256–68.
 65. Ha MW, Ma R, Shun LP, Gong YH, Yuan Y. Effects of allitridi on cell cycle arrest of human gastric cancer cells. *World J Gastroenterol.* 2005;11(35):5433–7.
 66. Liu KL, Chen HW, Wang RY, Lei YP, Sheen LY, Lii CK. DATS reduces LPS-induced iNOS expression, NO production, oxidative stress, and NF-kappaB activation in RAW 264.7 macrophages. *J Agric Food Chem.* 2006;54(9):3472–8.
 67. Powolny AA, Singh SV. Multitargeted prevention and therapy of cancer by diallyl trisulfide and related Allium vegetable-derived organosulfur compounds. *Cancer Lett.* 2008;269(2):305–14. <https://doi.org/10.1016/j.canlet.2008.05.027>.
 68. Puccinelli MT, Stan SD. Dietary bioactive diallyl trisulfide in cancer prevention and treatment. *Int J Mol Sci.* 2017;18(8). doi:<https://doi.org/10.3390/ijms18081645>.
 69. Kiesel VA, Stan SD. Diallyl trisulfide, a chemopreventive agent from Allium vegetables, inhibits alpha-secretases in breast cancer cells. *Biochem Biophys Res Commun.* 2017;484(4):833–8. <https://doi.org/10.1016/j.bbrc.2017.01.184>.
 70. Mikaili P, Maadirad S, Moloudizargari M, Aghajanshakeri S, Sarahroodi S. Therapeutic uses and pharmacological properties of garlic, shallot, and their biologically active compounds. *Iranian J Basic Med Sci.* 2013;16(10):1031–48.
 71. Chandra-Kuntal K, Lee J, Singh SV. Critical role for reactive oxygen species in apoptosis induction and cell migration inhibition by diallyl trisulfide, a cancer chemopreventive component of garlic. *Breast Cancer Res Treat.* 2013;138(1):69–79. <https://doi.org/10.1007/s10549-013-2440-2>.
 72. Wallace GC, Haar CP, Vandergrift WA 3rd, Giglio P, Dixon-Mah YN, Varma AK, et al. Multi-targeted DATS prevents tumor progression and promotes apoptosis in ectopic glioblastoma xenografts in SCID mice via HDAC inhibition. *J Neuro-Oncol.* 2013;114(1):43–50. <https://doi.org/10.1007/s11060-013-1165-8>.
 73. Yang CS, Chhabra SK, Hong JY, Smith TJ. Mechanisms of inhibition of chemical toxicity and carcinogenesis by diallyl sulfide (DAS) and related compounds from garlic. *J Nutr.* 2001;131(3s):1041S–5S.
 74. Gao, Jiang, Wang, Zhao, Wang. Drug metabolism and pharmacokinetics of organosulfur compounds from garlic. *J Drug Metab Toxicol* 2013;4(159).
 75. Chen C, Pung D, Leong V, Hebbar V, Shen G, Nair S, et al. Induction of detoxifying enzymes by garlic organosulfur compounds through transcription factor Nrf2: effect of chemical structure and stress signals. *Free Radic Biol Med.* 2004;37(10):1578–90. <https://doi.org/10.1016/j.freeradbiomed.2004.07.021>.
 76. Gong P, Hu B, Cederbaum AI. Diallyl sulfide induces heme oxygenase-1 through MAPK pathway. *Arch Biochem Biophys.* 2004;432(2):252–60. <https://doi.org/10.1016/j.abb.2004.09.024>.
 77. Kim S, Lee HG, Park SA, Kundu JK, Keum YS, Cha YN, et al. Keap1 cysteine 288 as a potential target for diallyl trisulfide-induced Nrf2 activation. *PLoS One.* 2014;9(1):e85984. <https://doi.org/10.1371/journal.pone.0085984>.
 78. Fisher CD, Augustine LM, Maher JM, Nelson DM, Slitt AL, Klaassen CD, et al. Induction of drug-metabolizing enzymes by garlic and allyl sulfide compounds via activation of constitutive androstane receptor and nuclear factor E2-related factor 2. *Drug Metab Dispos.* 2007;35(6):995–1000. <https://doi.org/10.1124/dmd.106.014340>.
 79. Lee BC, Park BH, Kim SY, Lee YJ. Role of Bim in diallyl trisulfide-induced cytotoxicity in human cancer cells. *J Cell Biochem.* 2011;112(1):118–27. <https://doi.org/10.1002/jcb.22896>.
 80. Das A, Banik NL, Ray SK. Garlic compounds generate reactive oxygen species leading to activation of stress kinases and cysteine proteases for apoptosis in human glioblastoma T98G and U87MG

- cells. *Cancer*. 2007;110(5):1083–95. <https://doi.org/10.1002/cncr.22888>.
81. Kim YA, Xiao D, Xiao H, Powolny AA, Lew KL, Reilly ML, et al. Mitochondria-mediated apoptosis by diallyl trisulfide in human prostate cancer cells is associated with generation of reactive oxygen species and regulated by Bax/Bak. *Mol Cancer Ther*. 2007;6(5):1599–609.
 82. Ferrer-Sueta G, Manta B, Botti H, Radi R, Trujillo M, Denicola A. Factors affecting protein thiol reactivity and specificity in peroxide reduction. *Chem Res Toxicol*. 2011;24(4):434–50. <https://doi.org/10.1021/tx100413v>.
 83. Allah DR, Schwind L, Asali IA, Nasim J, Jacob C, Gotz C, et al. A scent of therapy: synthetic polysulfanes with improved physico-chemical properties induce apoptosis in human cancer cells. *Int J Oncol*. 2015;47(3):991–1000. <https://doi.org/10.3892/ijo.2015.3093>.
 84. Benavides GA, Squadrito GL, Mills RW, Patel HD, Isbell TS, Patel RP, et al. Hydrogen sulfide mediates the vasoactivity of garlic. *Proc Natl Acad Sci U S A*. 2007;104(46):17977–82.
 85. Zhang F, Jin H, Wu L, Shao J, Zhu X, Chen A, et al. Diallyl trisulfide suppresses oxidative stress-induced activation of hepatic stellate cells through production of hydrogen sulfide. *Oxidative Med Cell Longev*. 2017;2017:1406726–13. <https://doi.org/10.1155/2017/1406726>.
 86. Bradley JM, Organ CL, Lefer DJ. Garlic-derived organic polysulfides and myocardial protection. *J Nutr*. 2016;146(2):403s–9s. <https://doi.org/10.3945/jn.114.208066>.
 87. Cai YR, Hu CH. Computational study of H₂S release in reactions of diallyl polysulfides with thiols. *J Phys Chem B*. 2017;121(26):6359–66. <https://doi.org/10.1021/acs.jpcc.7b03683>.
 88. Liang D, Wu H, Wong MW, Huang D. Diallyl trisulfide is a fast H₂S donor, but diallyl disulfide is a slow one: the reaction pathways and intermediates of glutathione with polysulfides. *Org Lett*. 2015;17(17):4196–9. <https://doi.org/10.1021/acs.orglett.5b01962>.
 89. Iciek M, Bilska-Wilkosz A, Gorny M, Sokolowska-Jezewicz M, Kowalczyk-Pachel D. The effects of different garlic-derived allyl sulfides on anaerobic sulfur metabolism in the mouse kidney. *Antioxidants* (Basel, Switzerland). 2016;5(4). <https://doi.org/10.3390/antiox5040046>.
 90. Iciek MB, Kowalczyk-Pachel D, Kwiecien I, Dudek MB. Effects of different garlic-derived allyl sulfides on peroxidative processes and anaerobic sulfur metabolism in mouse liver. *Phytother Res*. 2012;26(3):425–31. <https://doi.org/10.1002/ptr.3572>.
 91. Saidu NEB, Abu Asali I, Czepukojc B, Seitz B, Jacob C, Montenarh M. Comparison between the effects of diallyl tetrasulfide on human retina pigment epithelial cells (ARPE-19) and HCT116 cells. *Biochim Biophys Acta Gen Subj*. 2013;1830(11):5267–76.
 92. Saidu NEB, Touma R, Asali IA, Jacob C, Montenarh M. Diallyl tetrasulfane activates both the eIF2 α and Nrf2/HO-1 pathways. *Biochim Biophys Acta Gen Subj*. 2013;1830(1):2214–25.
 93. Anwar A, Groom M, Arbach M, Hamilton CJ. How to turn the chemistry of garlic into a ‘botanical’ pesticide. In: Jacob C, Kirsch G, Slusarenko A, Winyard PG, Burkholz T, editors. *Recent advances in redox active plant and microbial products: from basic chemistry to widespread applications in medicine and agriculture*. Dordrecht: Springer Netherlands; 2014. p. 323–41.