

Research



Clinical and biochemical correlates of hypogonadism in men with type 2 diabetes mellitus

Ezekiel Musa, Jibril Mohammed El-Bashir, Fatima Sani-Bello, Adamu Girei Bakari

Corresponding author: Ezekiel Musa, Department of Internal Medicine, Kaduna State University, Kaduna, Kaduna State, Nigeria. ezemusa2000@gmail.com

Received: 23 Aug 2020 - **Accepted:** 03 Mar 2021 - **Published:** 19 Mar 2021

Keywords: Type 2 diabetes mellitus, hypogonadism, total testosterone, testicular volume

Copyright: Ezekiel Musa et al. Pan African Medical Journal (ISSN: 1937-8688). This is an Open Access article distributed under the terms of the Creative Commons Attribution International 4.0 License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Cite this article: Ezekiel Musa et al. Clinical and biochemical correlates of hypogonadism in men with type 2 diabetes mellitus. Pan African Medical Journal. 2021;38(292). 10.11604/pamj.2021.38.292.25719

Available online at: <https://www.panafrican-med-journal.com//content/article/38/292/full>

Clinical and biochemical correlates of hypogonadism in men with type 2 diabetes mellitus

Ezekiel Musa^{1,2,&}, Jibril Mohammed El-Bashir³, Fatima Sani-Bello⁴, Adamu Girei Bakari⁴

¹Department of Internal Medicine, Kaduna State University, Kaduna, Kaduna State, Nigeria, ²Division of Endocrinology, Department of Medicine, University of Cape Town, Cape Town, South Africa, ³Department of Chemical Pathology, Ahmadu Bello University, Zaria, Kaduna State, Nigeria, ⁴Department of Medicine, Ahmadu Bello University, Zaria, Kaduna State, Nigeria

[&]Corresponding author

Ezekiel Musa, Department of Internal Medicine, Kaduna State University, Kaduna, Kaduna State, Nigeria

Abstract

Introduction: there is an association between hypogonadism and obesity, chronic hyperglycaemia, and ageing in men with type 2 diabetes mellitus (T2DM). T2DM is known to be associated with low testosterone. There is a paucity of data on the risk factors of hypogonadism in Nigerian men with T2DM. The objective of this study was to determine the clinical and biochemical correlates of hypogonadism and clinical predictors of low total testosterone levels in men with T2DM.

Methods: this was a cross-sectional study consisting of 358 men with T2DM and 179 non-diabetic men (controls). Structured Androgen Deficiency in the Ageing Male questionnaire was administered. Clinical and biochemical parameters were measured. Free testosterone was calculated from albumin, SHBG and total testosterone using Vermeulen's method. Hypogonadism was defined as fasting TT as < 8 nmol/L with or without symptoms or TT of 8-12 nmol/L with symptoms of androgen deficiency. Low testosterone was defined as serum total testosterone levels ≤ 12 nmol/L.

Results: the mean (\pm SD) total testosterone of men with T2DM and controls were 8.79 ± 3.35 nmol/L and 15.41 ± 3.79 nmol/L respectively ($p < 0.001$). The risk of hypogonadism was associated with central obesity (Odds ratio [OR] 2.24, 95% confidence interval [CI] 0.38-13.07), systolic hypertension (OR 3.93, 95% CI 0.67-23.10), hyperglycaemia (OR 2.48, 95% CI 0.37-16.46) and hypercholesterolaemia (OR 2.50, 95% CI 0.43-14.61). In a multivariable regression analysis, there was a significant negative correlation between total testosterone and triglycerides ($r -1.85$, 95% CI -3.58 - 0.12, $P = 0.04$) and HDL cholesterol ($r -1.25$, 95% CI -5.95-3.45, $P = 0.02$). **Conclusion:** this study shows that in men with T2DM, triglycerides and HDL cholesterol are independent correlates of hypogonadism but not central adiposity, systolic blood pressure and glycaemia. Further large prospective studies are recommended.

Introduction

Hypogonadism is a clinical syndrome comprising both symptoms, with or without signs, combined with biochemical evidence of testosterone deficiency [1,2]. In cross-sectional studies, between 20-80.4% of men with T2DM have hypogonadism [3-5]. Type 2 diabetes mellitus has been reported to be associated with low testosterone [3, 6-9]. Low testosterone levels are reportedly associated with a high risk of developing T2DM in men, a sequel to its insulin-resistance effect [10, 11]. Some of the clinical features of symptomatic hypogonadism are erectile dysfunction, loss of libido, depression, irritability, fatigue, anaemia, decreased intellectual activity, sleep disturbance, increased abdominal fat, decreased body hair and bone mineral density and lean body mass [2]. Erectile dysfunction, a commonly documented hypogonadism feature, was three times more common in men with T2DM than those without DM [12]. Furthermore, another study found that loss of libido and erectile dysfunction were the prominently reported symptoms by most type 2 diabetic men [9].

Cross-sectional studies have reported a significant relationship between body mass index (BMI), waist circumference and testosterone levels [7, 9, 13]. In South Africa, Paruk *et al.* found a significant relationship between ageing and BMI as well as an inverse relationship between total testosterone and BMI, waist circumference and metabolic syndrome [9]. Similarly, Kapoor *et al.* reported a significant negative correlation between testosterone levels and BMI and waist circumference in men with T2DM [7]. Meanwhile, another study found BMI was inversely correlated with both free and total testosterone and a strong correlation between equilibrium dialysis-measured free testosterone and calculated free testosterone but not with SHBG [3]. Furthermore, Kemp *et al.* reported waist circumference and known cardiovascular disease as the significant factors associated with low total testosterone [13]. Ugwu *et al.* reported older age and central obesity as the

predictors of hypogonadism among type 2 diabetic men [14].

There is currently a paucity of data on the risk factors of hypogonadism in Nigerian men with T2DM. Availability of relevant clinical and biochemical markers that will signal the early screening and diagnosis of hypogonadism in men with T2DM is timely. This will perhaps herald the appropriate care and improvement of the quality of life of men with T2DM. This study aimed to determine the clinical and biochemical correlates of hypogonadism in men with T2DM as well as the clinical predictors of low testosterone in men with T2DM.

Methods

Study design and setting

This was a cross-sectional study undertaken at the endocrine clinic of the Ahmadu Bello University Teaching Hospital (ABUTH), Zaria, from October 2015 to June 2016.

Study population

The study consisted of 358 male patients with T2DM and 179 non-diabetic men (controls). Men with T2DM (age 21 years and above) were recruited consecutively by convenience sampling as they fulfilled all the inclusion criteria. One hundred and seventy-nine healthy non-diabetic men were recruited consecutively. Those with chronic liver disease, chronic kidney disease, panhypopituitarism, HIV, type 1 diabetes mellitus, suspected prostate or testicular cancer, history of previous or present treatment of hypogonadism with testosterone and anti-androgen or related drugs, hospital admission, age less than 21 years, and refusal to participate in the study were excluded. Additionally, 43 men each with T2DM and controls were randomly selected in a case-control fashion using a computer from the cohorts of 358 men with T2DM and 179 controls, respectively, for sub-analysis.

Data collection

A structured Androgen Deficiency in the Ageing Male (ADAM) questionnaire was administered to obtain low testosterone symptoms. Anthropometric measurements were obtained. Body mass index (BMI) was calculated as weight in kilogram (kg) divided by height in meters squared (m^2). Weight was measured in kg with the subjects standing steadily on a scale, and height was measured in meters using a stadiometer. According to the World Health Organization (WHO) protocol, Waist circumference was measured at the midpoint between the iliac crest and the costal margin (lower rib). Meanwhile, hip circumference was measured at the widest portion of the buttocks with the tape parallel to the floor [15, 16]. The BMI classification was based on the WHO definition for obesity, while waist circumference cut-off of ≥ 94 cm defined abdominal obesity based on the International Diabetes Federation (IDF) criteria [17, 18]. Blood pressure (BP) was measured by a sphygmomanometer, and participants' BP was considered uncontrolled if their target BP was $\geq 140/90$ mmHg, according to the American Diabetes Association (ADA) recommendations [19]. A Prader orchidometer was used to measure the testicular volume of participants [20].

The laboratory biochemical tests performed on all participants included glycated haemoglobin (HbA1c), fasting plasma glucose (FPG), fasting serum total testosterone (TT), sex hormone-binding globulin (SHBG), albumin and lipid profile. Free testosterone was calculated from albumin, SHBG and total testosterone using Vermeulen's method [21]. The Glycated haemoglobin and FPG targets were based on ADA recommendations [22]. Fasting venous blood samples (10 ml) were collected in the morning between 8:00 AM and 10:00 AM into plain tubes and allowed to clot within 30 minutes. Samples were centrifuged for 10 minutes, and sera were collected into sample bottles and frozen at $-20^{\circ}C$ for future analysis. The sera samples were used for the analysis of total testosterone, SHBG, albumin and lipid profile. Separate 4 ml of blood was collected into an

ethylenediaminetetraacetic acid (EDTA) bottle for HbA1c assay.

Glycated haemoglobin was measured using the Clover A1C (Infopia Co. Ltd, Gyeonggi-do, South Korea), plasma glucose was measured using the oxidase-peroxidase method, serum albumin was measured by modified bromocresol green colorimetric method using kit acquired from Randox laboratory limited, Crumlin, United Kingdom. Lipid profile was measured using reagents acquired from Randox laboratory limited, Crumlin, United Kingdom. SHBG was measured by enzyme immunoassay technique using kits acquired from Diagnostic Automation Inc, California, USA, and serum total testosterone was measured by the enzyme immunoassay technique using Testosterone AccuBind ELISA kits from Monobind Inc, California, USA.

Definitions

Hypogonadism was defined as total testosterone levels < 8 nmol/L with or without symptoms or total testosterone levels of 8-12 nmol/L with the presence of symptoms. Low testosterone was defined as serum total testosterone levels \leq 12 nmol/L.

Statistical analysis

Data were analysed using Statistical Package for Social Sciences (SPSS) version 21 after validation. Quantitative variables such as BMI, lipid profile and testosterone levels were presented as means and standard deviations. The Student's t-test was used to determine the difference in means of clinical and biochemical parameters between type 2 diabetic men and controls. Chi-Square and regression analysis were used to analyse associations between hypogonadism and clinical and biochemical parameters, and univariable and multivariable correlations between clinical and biochemical parameters and low testosterone, respectively. Logistic regression analysis was used to determine the clinical predictors of low testosterone levels

among men with T2DM. P values < 0.05 were considered statistically significant.

Ethical considerations

All participants gave written informed consent. Ethical clearance for the study was obtained from ABUTH ethical committee (ABUTHZ/HREC/NP18/2015), and confidentiality was observed throughout the study.

Results

General characteristics of the study population

A total of 358 men with T2DM and 179 controls participated in the study. The mean (\pm SD) age for the men with T2DM was 46.34 ± 5.66 years, while that of the controls was 44.09 ± 12.39 years ($p = 0.57$). In our study, 177 (49.4%) of the men with T2DM had a duration of diabetes ranging 0-5 years, while 91 (25.4%), 55 (15.4%), 13 (3.6%), and 22 (6.2%) had diabetes for a duration of 6-10, 11-15, 16-20 and more than 20 years respectively (Table 1). Among the six categories of medications men with T2DM were taken, majority (34.6%) were on a metformin-glimepiride combination. Those who received metformin alone and metformin and insulin combinations were 55 (15.4%) and 52 (14.5%), respectively (Table 1). Two hundred and sixty-nine (75%) of the men with T2DM were hypertensive, with the majority (73.2%) on bendrofluazide, while 43.3% were taking lisinopril. The mean (\pm SD) systolic blood pressure (SBP) for men with T2DM and controls was 143.36 ± 25.64 mmHg and 128.70 ± 22.59 mmHg, respectively. There was a statistically significant difference between men's mean SBP with T2DM and the controls ($p < 0.001$, Table 1). Also, the mean (\pm SD) diastolic blood pressure (DBP) of men with T2DM was statistically higher than the controls (82.54 ± 13.05 mmHg versus 79.82 ± 14.05 mmHg, $p = 0.03$, Table 1). Abdominal obesity was found in 43.6% of men with T2DM, and the mean waist circumference of type 2 diabetic men was higher than the controls (92.40 ± 11.96 cm versus 79.49 ± 9.84 cm, $p < 0.001$) (Table 1). The mean (\pm SD) testicular volume of men

with T2DM and controls was 20.01 ± 4.15 ml and 21.37 ± 3.54 ml, respectively. The mean testicular volume of the controls was higher than that of the men with T2DM, which was statistically significant ($p < 0.001$, Table 1). Three hundred and seven (85.8%) of the men with T2DM did not achieve target metabolic control using glycated haemoglobin. The mean (\pm SD) glycated haemoglobin was significantly higher in men with T2DM compared to controls ($p < 0.001$) (Table 1). There was a significant difference between type 2 diabetic men and controls for their mean total cholesterol ($p < 0.001$), LDL cholesterol (< 0.001), and triglycerides (< 0.001) (Table 1).

Association of clinical and biochemical parameters with hypogonadism in men with T2DM

A sub-group analysis of 43 men with T2DM was performed to assess the risk factors of hypogonadism. All obese type 2 diabetic men had hypogonadism, with grade 1 obesity accounting for all the cases. None of the hypogonadal men with T2DM had either grade 2 or morbid obesity. Among the type 2 diabetic men who had abdominal obesity, 89.5% had hypogonadism, and 79.2% of the non-obese men with T2DM had hypogonadism. The odds of developing hypogonadism among type 2 diabetic men with abdominal obesity was twice that of the non-obese men, but this did not reach statistical significance ($X^2 = 0.83$, $p = 0.36$, $OR = 2.24$, $CI = 0.38 - 13.07$, Table 2).

Twenty-two (91.7%) of type 2 diabetic men with above-target systolic blood pressure (SBP) had hypogonadism, while 73.7% of those with SBP within target had hypogonadism as well. There was about a four-fold likelihood of developing hypogonadism among type 2 diabetic men with elevated SBP compared to normal SBP ($X^2 = 2.52$, $p = 0.11$, $OR = 3.93$, $CI = 0.67 - 23.10$, Table 2). All men with T2DM who had high diastolic blood pressure (DBP) were hypogonadal compared to 77.4% of those with normal DBP ($X^2 = 0.01$, $p = 0.92$, $OR = 1.10$, $CI = 0.18 - 6.56$, Table 2). There was a higher frequency of hypogonadism (90%) among type 2 diabetic men with poor glycaemic control

compared to those who achieved target fasting plasma glucose (FPG) using ADA criteria. Thirty-one (86.1%) of those who did not achieve target glycated haemoglobin had a higher prevalence of hypogonadism than those whose glycated haemoglobin was well-controlled (71.4%). The risk of having hypogonadism among men with T2DM with poor glycated haemoglobin was about 2.5 times more than with well-controlled glycated haemoglobin, but a statistically significant association between glycated haemoglobin and hypogonadism was not achieved ($X^2 = 0.93$, $p = 0.34$, $OR = 2.48$, $CI = 0.37 - 16.46$, Table 2). Among the men with T2DM who had elevated total cholesterol, 75% had hypogonadism. Meanwhile, 85.7% of those who had normal total cholesterol were hypogonadal. There was no statistically significant association between total cholesterol level and hypogonadism ($X^2 = 0.55$, $p = 0.46$, $OR = 2.00$, $CI = 0.31 - 12.84$, Table 2). The odds of having hypogonadism were 2.5 times higher in men with T2DM who had elevated LDL cholesterol as opposed to having normal LDL cholesterol ($X^2 = 1.08$, $p = 0.30$, $OR = 2.50$, $CI = 0.43 - 14.61$, Table 2).

Univariable and multivariable regression analyses of correlates of hypogonadism in men with T2DM

In a multivariable regression, there was a significant negative correlation between total testosterone and triglycerides ($r = -1.85$, 95% $CI = -3.58 - 0.12$, $p = 0.04$) and HDL cholesterol ($r = -1.25$, 95% $CI = -5.95 - 3.45$, $p = 0.02$) (Table 3). However, a univariable regression showed a significant positive correlation between total testosterone and LDL cholesterol ($r = 0.38$, 95% $CI = 0.38 - 2.92$, $p = 0.01$).

Logistic regression analysis of the clinical predictors of low total testosterone in men with T2DM with hypogonadism

In a logistic regression analysis using total testosterone as the dependent variable and average testicular volume, BMI, erectile dysfunction, decreased libido, duration of diabetes and waist circumference as independent variables, the clinical variables that had a higher likelihood of

predicting low total testosterone levels in men with T2DM were erectile dysfunction (OR 6.00, 95% CI 0.97-37.12, $p = 0.05$) and a decreased libido (OR 4.67, 95% CI 0.86-25.31, $p = 0.07$) (Table 4).

Discussion

This study found a significantly lower total and calculated free testosterone in men with T2DM than non-diabetic controls. Asare-Anane *et al.* and Onung *et al.* found similar findings in Accra, Ghana and Lagos, Nigeria, respectively [6, 23]. Furthermore, Paruk *et al.* in South Africa also found lower total and free testosterone levels in type 2 diabetic men than in the controls. Two Meta-analyses by Ding *et al.* and Corona *et al.* which included 20 cross-sectional studies (850 diabetic men and 2000 non-diabetic controls) and 28 cross-sectional studies (1,822 men with diabetes and 10,009 non-diabetic controls), respectively, found consistent lower total testosterone levels in men with diabetes compared with non-diabetic controls [24, 25]. The association between low total testosterone and diabetes has been previously attributed to low SHBG observed commonly in T2DM [26]. However, in our study, calculated free testosterone was significantly lower in men with T2DM than non-diabetic controls, suggesting that other factors may play a role in lowering testosterone levels and not SHBG alone.

Despite the higher risk of hypogonadism among type 2 diabetic men with poor glycaemic control, the association between poor glycaemic control and hypogonadism did not reach statistical significance but was clinically significant. This finding is consistent with a report from a Nigerian study [8]. Some studies reported a significant association between hypogonadism and FPG and glycated haemoglobin [6, 7]. Prolonged hyperglycaemia impairs the synthesis and secretion of testosterone, resulting in low testosterone levels. The high proportion of type 2 diabetic men with uncontrolled and well-controlled glycated haemoglobin may explain the apparent insignificant statistical association between glycaemia and hypogonadism. This study did not

show a statistically significant association between total cholesterol and hypogonadism. It was similar to findings from Nigerian and United Kingdom studies [4, 8]. A study reported a significant association with low total testosterone [27].

Furthermore, this study showed no statistically significant association between LDL cholesterol and hypogonadism, despite the high mean LDL cholesterol in hypogonadal subjects. This finding is supported by Mirzaei *et al.*'s study, which found no correlation between total testosterone and lipid profile reported [28]. Conversely, Wickramatilake and colleagues showed a significant association between LDL cholesterol and hypogonadism [27]. The association observed between hypogonadism and both systolic and diastolic blood pressure was not significant in this study. These findings are consistent with those reported in other studies [4, 29]. In contrast, some studies reported an observed association between hypogonadism and blood pressure [7, 30]. There was an observed clinically but not a statistically significant association between hypogonadism and abdominal obesity using waist circumference despite a higher frequency of abdominal obesity in hypogonadal type 2 diabetic men. Zheng *et al.* also reported a lack of significant association between hypogonadism and abdominal obesity but the opposite finding was reported in a study by Laaksonen *et al.* [31, 32]. This study found no association between hypogonadism and global obesity similar to prior studies [8, 33]. Observations by some researchers showed a significant association between obesity (using BMI) and hypogonadism [13, 34]. The lack of a statistically significant association between obesity and hypogonadism despite the higher odds of developing hypogonadism in obese type 2 diabetic men compared to non-obese could be attributed to a considerably high proportion of hypogonadism in the non-obese men.

There was a significant negative correlation between total testosterone and triglycerides and HDL cholesterol in multivariable analysis with a linear decrease in the total testosterone levels as

these variables increased. These findings were consistent with other researchers reports [34, 35] but contradicted the findings by Asare-Anane *et al.* and Chang *et al.* [6, 36]. These indicate that male patients with T2DM who have dyslipidaemia may often exhibit hypogonadism. Low testosterone is associated with increased adiposity, T2DM, increased inflammation, hyperlipidaemia [37]. The findings in this study show no significant association between total testosterone and anthropometric measurements and FPG as opposed to the findings by Kapoor *et al.* and Asare-Anane *et al.* [6, 7].

Erectile dysfunction and decreased libido showed a higher likelihood of predicting low total testosterone levels, with the odds approaching statistical significance. Meanwhile, waist circumference, testicular volume, duration of diabetes and BMI were not found to increase the likelihood of predicting low total testosterone. The ADAM questionnaire, which has a sensitivity of 88%, established that erectile dysfunction and loss of libido are predictors of low testosterone [38]. In contrast to the findings in this study, Mahmoud *et al.* found testicular volume was a predictor of low testosterone [39] while Travison *et al.* reported that obesity using WC and BMI and duration of diabetes as predictors of low total testosterone [40]. The much lower cut-off for defining abdominal obesity in sub-Saharan African men, according to IDF, may account for the seeming lack of statistical significance despite the observed higher odds of having low testosterone with abdominal obesity. Additionally, the exceptionally high proportion of both hypogonadal and eugonadal men who gave positive responses to erectile dysfunction and loss of libido could have attenuated any inherent significance.

This study, being hospital-based, might not have given an actual reflection of hypogonadism in men with T2DM. Also, considering its cross-sectional nature, findings are mainly associations, therefore, a well-designed prospective study to investigate the clinical and biochemical correlates of

hypogonadism in male patients with T2DM is recommended.

Conclusion

This study shows that triglycerides and HDL cholesterol are independent correlates of hypogonadism in men with T2DM. There was no association between hypogonadism and abdominal obesity, systolic hypertension, and hyperglycaemia in men with T2DM. Erectile dysfunction and decreased libido are the likely clinically important predictors of low testosterone levels in type 2 diabetic men with hypogonadism.

What is known about this topic

- *An association between hypogonadism and obesity and chronic hyperglycaemia in men with T2DM exists;*
- *T2DM is known to be associated with low testosterone in male patients.*

What this study adds

- *This study found a lower testicular volume in men with T2DM than non-diabetic men; to the best of our knowledge, this is the first study in Africa to utilise a Prader orchidometer in assessing testicular volume in T2DM;*
- *This is also the first study to determine calculated free testosterone using Vermeulen's equation with approximate precision to equilibrium dialysis (goal standard) for measuring serum free testosterone; and our research found lower levels of free testosterone in men with T2DM compared to non-diabetics;*
- *Our findings suggest that erectile dysfunction and decreased libido may be clinically important in assessing hypogonadism in men with T2DM.*

Competing interests

The authors declare no competing interests.

Authors' contributions

All contributing authors (EM, JME, FS and AGB) were fully involved in this study's design. EM and AGB conceptualized the research question. EM wrote the final draft of this manuscript. All authors have read and agreed to the final version of this manuscript.

Tables

Table 1: clinical and biochemical features of men with T2DM and controls

Table 2: association of clinical and biochemical parameters with hypogonadism in men with T2DM

Table 3: univariable and multivariable correlates of hypogonadism in men with T2DM

Table 4: logistic regression analysis of the clinical correlates of low total testosterone in men with T2DM with hypogonadism

References

1. Bhasin S, Cunningham GR, Hayes FJ, Matsumoto AM, Snyder PJ, Swerdloff RS *et al.* Testosterone therapy in men with androgen deficiency syndromes: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab.* 2010 Jun;95(6): 2536-59. **PubMed | Google Scholar**
2. Dhatriya K, Nagi D, Jones T. ABCD position statement on the management of hypogonadal males with type 2 diabetes. *Pract Diabetes.* 2010;27(9): 408-12. **Google Scholar**
3. Dhindsa S, Prabhakar S, Sethi M, Bandyopadhyay A, Chaudhuri A, Dandona P. Frequent occurrence of hypogonadotropic hypogonadism in type 2 diabetes. *J Clin Endocrinol Metab.* 2004 Nov;89(11): 5462-8. **PubMed | Google Scholar**
4. Hackett GI, Cole NS, Deshpande AA, Popple MD, Kennedy D, Wilkinson P. Biochemical hypogonadism in men with type 2 diabetes in primary care practice. *Br J Diabetes Vasc Dis.* 2009;9(5): 226-31. **Google Scholar**
5. Musa E, El-Bashir JM, Sani-Bello F, Bakari AG. Hypergonadotropic hypogonadism in Nigerian men with type 2 diabetes mellitus. *Clin Diabetol.* 2021;10(1): 129-37. **Google Scholar**
6. Asare-Anane H, Ofori E, Yeboah F, Tagoe E, Bani S, Bawah A *et al.* Primary hypogonadism in Ghanaian men with type 2 diabetes mellitus. *IJSTR.* 2013;2(5): 310-5. **Google Scholar**
7. Kapoor D, Aldred H, Clark S, Channer KS, Jones TH. Clinical and biochemical assessment of hypogonadism in men with type 2 diabetes: correlations with bioavailable testosterone and visceral adiposity. *Diabetes care.* 2007 Apr;30(4): 911-7. **PubMed | Google Scholar**
8. Ogbera OA, Sonny C, Olufemi F, Wale A. Hypogonadism and subnormal total testosterone levels in men with type 2 diabetes mellitus. *J Coll Physicians Surg Pak.* 2011 Sep;21(9): 517-21. **PubMed | Google Scholar**
9. Paruk IM, Pirie FJ, Nkwanyana NM, Motala AA. Prevalence of low serum testosterone levels among men with type 2 diabetes mellitus attending two outpatient diabetes clinics in KwaZulu-Natal Province, South Africa. *S Afr Med J.* 2019 Nov 27;109(12): 963-970. **PubMed | Google Scholar**
10. Svartberg J. Epidemiology: testosterone and the metabolic syndrome. *Int J Impot Res.* Mar-Apr 2007;19(2): 124-8. **PubMed | Google Scholar**
11. Zitzmann M, Faber S, Nieschlag E. Association of specific symptoms and metabolic risks with serum testosterone in older men. *J Clin Endocrinol Metab.* 2006 Nov;91(11): 4335-43. **PubMed | Google Scholar**
12. De Berardis G, Franciosi M, Belfiglio M, Di Nardo B, Greenfield S, Kaplan SH *et al.* Erectile dysfunction and quality of life in type 2 diabetic patients: a serious problem too often overlooked. *Diabetes care.* 2002 Feb;25(2): 284-91. **PubMed | Google Scholar**
13. Kemp T, Rheeder P. The prevalence and association of low testosterone levels in a South African male, diabetic, urban population. *JEMDSA.* 2015;20(2): 92-7. **Google Scholar**

14. Ugwu TE, Ikem RT, Kolawole BA, Ezeani IU. Clinicopathologic assessment of hypogonadism in men with type 2 diabetes mellitus. *Indian J Endocrinol Metab.* Sep-Oct 2016;20(5): 667-673. **PubMed | Google Scholar**
15. Jimoh KA, Adediran OS, Agboola SM, Olugbodi D, Idowu AA, Adebisi SA. A study of correlation between derived and basic anthropometric indices in type 2 diabetes mellitus. *Eur J Sci Res.* 2009;36(3): 437-44. **Google Scholar**
16. Consultation WE. Waist circumference and waist-hip ratio: report of a WHO expert consultation, Geneva, 8-11 December 2008. 2014. **Google Scholar**
17. Alberti KG, Zimmet P, Shaw J. Metabolic syndrome--a new world-wide definition. A Consensus Statement from the International Diabetes Federation. *Diabet Med.* 2006 May;23(5): 469-80. **PubMed | Google Scholar**
18. Organization WH. Obesity: preventing and managing the global epidemic. Report of a WHO consultation. *World Health Organ Tech Rep Ser.* 2000;894: i-xii, 1-253. **PubMed | Google Scholar**
19. de Boer IH, Bangalore S, Benetos A, Davis AM, Michos ED, Muntner P *et al.* Diabetes and Hypertension: A Position Statement by the American Diabetes Association. *Diabetes care.* 2017 Sep;40(9): 1273-1284. **PubMed | Google Scholar**
20. Prader A. Testicular size: assessment and clinical importance. *Triangle.* 1966;7(6): 240-3. **PubMed | Google Scholar**
21. Vermeulen A, Verdonck L, Kaufman JM. A critical evaluation of simple methods for the estimation of free testosterone in serum. *J Clin Endocrinol Metab.* 1999 Oct;84(10): 3666-72. **PubMed | Google Scholar**
22. Association AD. 5. Glycemic targets. *Diabetes care.* 2016 Jan;39 Suppl 1: S39-46. **PubMed | Google Scholar**
23. Onung S, Young E, Ugwu T, Fasanmade O. Hypogonadism in Nigerian men with type 2 diabetes mellitus. *Int J Diabetes Dev Ctries.* 2017;37(3): 254-61. **Google Scholar**
24. Ding EL, Song Y, Malik VS, Liu S. Sex differences of endogenous sex hormones and risk of type 2 diabetes: a systematic review and meta-analysis. *Jama.* 2006 Mar 15;295(11): 1288-99. **PubMed | Google Scholar**
25. Corona G, Monami M, Rastrelli G, Aversa A, Sforza A, Lenzi A *et al.* Type 2 diabetes mellitus and testosterone: A meta-analysis study. *Int J Androl.* 2011 Dec;34(6 Pt 1): 528-40. **PubMed | Google Scholar**
26. Lakshman KM, Bhasin S, Araujo AB. Sex hormone-binding globulin as an independent predictor of incident type 2 diabetes mellitus in men. *J Gerontol A Biol Sci Med Sci.* 2010 May;65(5): 503-9. **PubMed | Google Scholar**
27. Wickramatilake CM, Mohideen MR, Pathirana C. Association of serum testosterone with lipid abnormalities in patients with angiographically proven coronary artery disease. *Indian J Endocrinol Metab.* 2013 Nov;17(6): 1061-5. **PubMed | Google Scholar**
28. Mirzaei MR, Amini M, Aminorroaya A. The prevalence of hypogonadism in diabetic men in Isfahan Endocrine and Metabolism Research Center, Isfahan, Iran. *J Res Med Sci.* 2012 Jul;17(7): 602-6. **PubMed | Google Scholar**
29. Alhayek AA, Khader YS, Jafal S, Khawaja N, Ajlouni K. Hypogonadism among Jordanian men with type 2 diabetes: Prevalence and associated factor. *Int J Diabetes Mellit.* 2015;3(1): 31-6. **Google Scholar**
30. Phillips GB, Jing T, Heymsfield SB. Relationships in men of sex hormones, insulin, adiposity, and risk factors for myocardial infarction. *Metabolism.* 2003 Jun;52(6): 784-90. **PubMed | Google Scholar**
31. Zheng R, Cao L, Cao W, Chu X, Hu Y, Zhang H *et al.* Risk Factors for Hypogonadism in Male Patients with Type 2 Diabetes. *J Diabetes Res.* 2016;2016: 5162167. **PubMed | Google Scholar**
32. Laaksonen DE, Niskanen L, Punnonen K, Nyyssonen K, Tuomainen TP, Salonen R *et al.* Sex hormones, inflammation and the metabolic syndrome: a population-based study. *Eur J Endocrinol.* 2003 Dec;149(6): 601-8. **PubMed | Google Scholar**

33. Tripathy D, Dhindsa S, Garg R, Khaishagi A, Syed T, Dandona P. Hypogonadotropic hypogonadism in erectile dysfunction associated with type 2 diabetes mellitus: a common defect? *Metab Syndr Relat Disord*. 2003 Mar;1(1): 75-80. **Google Scholar**
34. Diaz-Arjonilla M, Schwarcz M, Swerdloff RS, Wang C. Obesity, low testosterone levels and erectile dysfunction. *Int J Impot Res*. Mar-Apr 2009;21(2): 89-98. **PubMed | Google Scholar**
35. Fukui M, Soh J, Tanaka M, Kitagawa Y, Hasegawa G, Yoshikawa T *et al*. Low serum testosterone concentration in middle-aged men with type 2 diabetes. *Endocr J*. 2007 Dec;54(6): 871-7. **PubMed | Google Scholar**
36. Chang TC, Tung CC, Hsiao YL. Hormonal changes in elderly men with non-insulin-dependent diabetes mellitus and the hormonal relationships to abdominal adiposity. *Gerontology*. 1994;40(5): 260-7. **PubMed | Google Scholar**
37. Wang C, Jackson G, Jones TH, Matsumoto AM, Nehra A, Perelman MA *et al*. Low testosterone associated with obesity and the metabolic syndrome contributes to sexual dysfunction and cardiovascular disease risk in men with type 2 diabetes. *Diabetes care*. 2011 Jul;34(7): 1669-75. **PubMed | Google Scholar**
38. Morley JE, Charlton E, Patrick P, Kaiser FE, Cadeau P, McCready D *et al*. Validation of a screening questionnaire for androgen deficiency in aging males. *Metabolism*. 2000 Sep;49(9): 1239-42. **PubMed | Google Scholar**
39. Mahmoud AM, Goemaere S, El-Garem Y, Van Pottelbergh I, Comhaire FH, Kaufman JM. Testicular volume in relation to hormonal indices of gonadal function in community-dwelling elderly men. *J Clin Endocrinol Metab*. 2003 Jan;88(1): 179-84. **PubMed | Google Scholar**
40. Travison TG, Araujo AB, Kupelian V, O'Donnell AB, McKinlay JB. The relative contributions of aging, health, and lifestyle factors to serum testosterone decline in men. *J Clin Endocrinol Metab*. 2007 Feb;92(2): 549-55. **PubMed | Google Scholar**

Table 1: clinical and biochemical features of men with T2DM and controls

	T2DM	Controls	
	n = 358	n = 179	Significance
	Mean±SD	Mean±SD	P-value
Age (years)	46.34±5.66	44.09 ± 12.39	0.57
Clinical parameters:			
SBP (mmHg)	143.36±25.64	128.70 ± 22.59	<0.001
DBP (mmHg)	82.54±13.05	79.82 ± 14.05	0.03
WC (cm)	92.40 ± 11.96	79.49 ± 9.84	<0.001
BMI (m/kg ²)	25.26 ± 4.45	24.31 ± 3.79	0.01
Testicular volume (ml)	20.01 ± 4.15	21.37 ± 3.54	<0.001
Biochemical parameters:			
FPG (mmol/L)	8.30 ± 4.77	4.43 ± 1.79	<0.001
HbA1c (%)	9.13 ± 2.12	5.07 ± 0.74	<0.001
Total cholesterol (mmol/L)	4.28 ± 0.92	3.98 ± 0.70	<0.001
LDL cholesterol (mmol/L)	2.63 ± 0.74	2.37 ± 0.56	<0.001
HDL cholesterol (mmol/L)	1.04 ± 0.33	1.10 ± 0.34	0.06
Triglycerides (mmol/L)	1.32 ± 0.49	1.11 ± 0.37	<0.001
Total testosterone (nmol/L)	8.79 ± 3.35	15.41 ± 3.79	<0.001
SHBG (nmol/L)	85.59 ± 63.96	95.51 ± 76.29	0.14
Albumin (mmol/L)	39.68 ± 3.26	41.53 ± 3.42	<0.001
Calculated free testosterone (nmol/L)	0.11 ± 0.07	0.22 ± 0.16	<0.001
Duration of T2DM (years) n (%):			
0-5	177 (49.4)		
6-10	91 (25.4%)		
11-15	55 (15.4%)		
16-20	13 (3.6%)		
> 20	22 (6.2%)		
Medications n (%):			
Metformin alone	55 (15.4%)		
Metformin + glimepiride	124 (34.6%)		
Metformin + glibenclamide	78 (21.8%)		
Metformin + glimepiride + insulin	8 (2.2%)		
Metformin + glibenclamide + insulin	6 (1.7%)		
Metformin + insulin	52 (14.5%)		
Others	35 (9.8%)		

P <0.05 is statistically significant. BMI: body mass index, DBP: diastolic blood pressure, FBG: fasting plasma glucose, HbA1c: glycated haemoglobin, HDL: high density lipoprotein, LDL: low density lipoprotein, SHBG: sex hormone-binding globulin, SBP: systolic blood pressure, WC: waist circumference. Please note that BMI, FPG, HbA1c, total testosterone and WC of men with T2DM and controls have been published already (Musa et al 2021) [5].

Table 2: association of clinical and biochemical parameters with hypogonadism in men with T2DM

Characteristics	Hypogonadism	Eugonadism	Significance
	Frequency (%)	Frequency (%)	P-Values
BMI			
≥ 30 kg/m ²	5 (100)	0 (0)	0.64
< 30 kg/m ²	31 (81.6)	7 (18.4)	
Obesity by WC			
≥ 94 cm	17 (89.5)	2 (10.5)	0.36
< 94 cm	19 (79.2)	5 (20.8)	
Obesity by WHR			
≥ 0.90	35 (83.3)	7 (16.7)	0.66
< 0.90	1 (100)	0 (0)	
SBP			
≥ 140 mmHg	22 (91.7)	2 (8.3)	0.11
< 140 mmHg	14 (73.7)	5 (26.3)	
DBP			
≥ 90 mmHg	12(100)	0 (0)	0.92
< 90 mmHg	24(77.4)	7 (22.6)	
FPG			
>7.2 mmol/L	18 (90)	2 (10)	0.58
4.4 - 7.2 mmol/L	14 (77.8)	4 (22.2)	
< 4.4 mmol/L	4 (80)	1 (20)	
HbA1c			
≥7%	31(86.1)	5 (13.9)	0.34
< 7 %	5(71.4)	2 (28.6)	
T C			
≥ 5.2 mmol/L	6 (75)	2(25)	0.46
< 5.2 mmol/L	30 (85.7)	5(14.3)	
LDL cholesterol			
≥ 2.6 mmol/L	18 (78.3)	5 (21.7)	0.30
< 2.6 mmol/L	18 (90)	2(10)	
HDL cholesterol			
< 1.1 mmol/L	14 (87.5)	2(12.5)	0.61
≥ 1.1 mmol/L	22 (81.5)	5(18.5)	
Triglycerides			
≥ 1.7 mmol/L	8 (100)	0(0)	0.17
< 1.7 mmol/L	28 (80)	7 (20)	

BMI: body mass index, DBP: diastolic blood pressure, FPG: fasting plasma glucose, HbA1c: glycated haemoglobin, HDL: high density lipoprotein, LDL: low density lipoprotein, SBP: systolic blood pressure, TC: total cholesterol, WC: waist circumference, WHR: waist-hip ratio.

Table 3: univariable and multivariable correlates of hypogonadism in men with T2DM

	Univariable analysis		Multivariable analysis	
	Coefficient of Correlation (r) (95% CI)	P-value	Coefficient of Correlation (r) (95% CI)	P-value
FPG	0.07 (-0.30 - 0.18)	0.64	-0.08 (-0.53 - 0.37)	0.72
HbA1C	0.01 (-0.44 - 0.43)	0.97	0.33 (-0.50 - 1.15)	0.43
TC	0.28 (-0.09 - 1.95)	0.07	1.79 (0.01 - 3.56)	0.83
HDL	0.18 (-1.14 - 4.14)	0.26	-1.25 (-5.95 - 3.45)	0.02*
LDL	0.38 (0.38 - 2.92)	0.01*	1.62 (-0.10 - 3.34)	0.06
TG	0.25 (-2.45 - 0.27)	0.11	-1.85 (-3.58 - 0.12)	0.04*
SBP	0.05 (-0.03 - 0.04)	0.77	-0.07 (-0.07 - 0.05)	0.82
DBP	0.03 (-0.06 - 0.07)	0.84	0.04 (-0.08 - 0.15)	0.52
BMI	0.09 (-0.31 - 0.17)	0.57	-0.26 (-0.70 - 0.19)	0.25
WC	0.01 (-0.09 - 0.09)	0.96	0.10 (-0.10 - 0.29)	0.33
WHR	0.12 (-24.14 - 11.03)	0.46	-4.67 (-30.47 - 21.14)	0.72

Note: *p<0.05 is statistically significant. BMI: body mass index, DBP: diastolic blood pressure, FPG: fasting plasma glucose, HbA1c: glycated haemoglobin, HDL: high density lipoprotein, LDL: low density lipoprotein, SBP: systolic blood pressure, WC: waist circumference, WHR: waist-hip ratio.

Table 4: logistic regression analysis of the clinical correlates of low total testosterone in men with T2DM with hypogonadism

	P-values	Odd Ratio	95% CI	
			Lowerbound	Upperbound
Average testicular volume	0.10	0.00	0.00	0.00
BMI	0.37	0.61	0.20	1.80
WC	0.37	2.24	0.38	13.07
ED	0.05	6.00	0.97	37.12
Decreased libido	0.07	4.67	0.86	25.31
Duration of diabetes mellitus	0.20	0.45	0.13	1.52

BMI: body mass index, ED: erectile dysfunction, WC: waist circumference.