

BMJ Open Keloid risk in patients with atopic dermatitis: a nationwide retrospective cohort study in Taiwan

Ying-Yi Lu,^{1,2,3} Chun-Ching Lu,⁴ Wei-Wen Yu,¹ Li Zhang,⁵ Qing-Rui Wang,⁶ Cong-Liang Zhang,⁷ Chieh-Hsin Wu^{3,8,9}

To cite: Lu Y-Y, Lu C-C, Yu W-W, *et al.* Keloid risk in patients with atopic dermatitis: a nationwide retrospective cohort study in Taiwan. *BMJ Open* 2018;**8**:e022865. doi:10.1136/bmjopen-2018-022865

► Prepublication history for this paper is available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2018-022865>).

Received 10 March 2018
Revised 9 May 2018
Accepted 11 June 2018

ABSTRACT

Objective The pathogenesis of keloid is largely unknown. Because keloid and atopic dermatitis have overlapping pathophysiological mechanisms, we aimed to evaluate keloid risk in patients with atopic dermatitis.

Study design Population-based retrospective cohort study.

Setting The Taiwan National Health Insurance Research Database was used to analyse data for people who had been diagnosed with atopic dermatitis.

Participants We identified 8371 patients with newly diagnosed atopic dermatitis during 1996–2010. An additional 33 484 controls without atopic dermatitis were randomly identified and frequency matched at a one-to-four ratio.

Primary and secondary outcome measure The association between atopic dermatitis and keloid risk was estimated using Cox proportional hazard regression models.

Results After adjustment for covariates, the atopic dermatitis patients have a 3.19-fold greater risk of developing keloid compared with the non-atopic dermatitis group (3.19vs1.07 per 1000 person-years, respectively). During the study period, 163 patients with atopic dermatitis and 532 patients without atopic dermatitis developed keloid. Notably, keloid risk increased with severity of atopic dermatitis, particularly in patients with moderate to severe atopic dermatitis.

Conclusions Our results indicate that patients with atopic dermatitis had a higher than normal risk of developing keloid and suggest that atopic dermatitis may be an independent risk factor for keloid.

INTRODUCTION

Keloid is a pathological disorder as a result of dermal injury and excess production in the extracellular matrix (ECM) (mainly type I collagen and proteoglycans). Keloids are characterised by concomitant annoying symptoms such as pain and pruritus. The recurrence rate after surgical excision is high. The pathogenesis of keloids is largely unknown,^{1 2} and no standard treatment has emerged. Although the cause of this disorder is not well understood, known risk factors for keloids are age 10–30 years, female gender^{3 4}

Strengths and limitations of this study

- The cohort study evaluates the risk of developing keloids in patients with atopic dermatitis by using multivariable Cox proportional hazard regression models.
- This study was based on secondary dataset, which encrypted patient information to keep privacy.
- The strengths of this study are the large population size, the same ethnic group and long follow-up period.
- Detailed personal information including risk factors was not found in the database, which compromised our findings and hindered further analyses.
- All insurance reimbursement was peer reviewed and checked by well-trained specialists.

and dark skin pigmentation (eg, in Asian, African-American and Hispanic people). Notably, the risk of keloid development is highest in populations with the darkest complexions.^{5 6}

In a recent case-control study performed in Northeast China, comparisons of blood test results for 283 subjects with keloids and 290 controls suggested that keloid development may be associated with the single-nucleotide polymorphisms, a disintegrin and metalloprotease 33 (ADAM33).¹ Earlier studies have also shown that ADAM33 is associated with immune-mediated disorders such as atopic dermatitis (AD).^{7 8} In the first epidemiological study of the relationship between ADAM33 gene polymorphisms and AD risk, Matsusue *et al* reported a significant association between ADAM33 gene polymorphisms and AD in comparisons of 140 children with AD and 258 healthy controls recruited from June 2006 to January 2007. We hypothesised that AD and keloids may develop through common pathways, particularly in the ADAM33-related gene. Since no population-based data were available for investigating the association between AD and keloids, this study retrospectively analysed data from the Taiwan National Health



© Author(s) (or their employer(s)) 2018. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to
Professor Chieh-Hsin Wu;
wujoey@gmail.com

Insurance Research Database (NHIRD) to investigate this relationship.

MATERIALS AND METHODS

Source of data

The data used in this population-based cohort study were obtained from the NHIRD established by the Taiwan national healthcare system. This encrypted secondary database, which was implemented in March 1995 contains medical reports for almost 99% of the 23.74 million residents of Taiwan. The NHIRD contains records of medical services and enrolment data for all beneficiaries. The database is publicly available to researchers, but its use is strictly regulated by Taiwan Personal Electronic Data Protection Law. The NHIRD has been used in many studies published in scientific journals.^{9–11} The current study used the Longitudinal Health Insurance Database 2010, in which diseases are coded according to the International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM).

The study was proceeded according to Declaration of Helsinki guidelines. Because all NHIRD data had been deidentified and personal information traced was anonymised before data analysis, informed consent was not needed.

Study design

We examined medical claims and identified 8371 patients who had at least one AD diagnosis (ICD-9-CM code 691.8) during 1996–2010.¹² To ensure accurate data, the analysis was limited to patients who had received ≥ 3 AD diagnoses during outpatient visits or ≥ 1 AD diagnoses during inpatient medical service. The date of the first clinical

visit for AD was defined as the index date. Patients with moderate to severe AD were defined as patients who had received systemic corticosteroids.^{13–16} The exclusion criterion was any diagnosis of a keloid (ICD-9-CM code 701.4) before the index date.¹⁷ During the follow-up period, the occurrence of keloid was defined as ≥ 3 keloid diagnoses in outpatient visits or ≥ 1 keloid diagnoses in inpatient medical service. The two cohorts were limited to patients who had an ICD-9 code for keloid assigned by a dermatologist. Another 33 484 controls without AD were randomly identified after eliminating study case cohort and frequency matched at a one-to-four ratio for age, gender and index year (year of AD diagnosis).¹⁸ Both cohorts were followed up until 31 December 2010, or until the occurrence of keloids. Figure 1 shows a flow chart of the study procedure.

Ethics approval

The study was conducted in accordance with the Declaration of Helsinki guidelines.

Measures and definition

As in previous NHIRD studies, the patients were classified into three urbanisation levels according to their residence addresses recorded in the databases: urban (those living in the ‘city’), suburban (those living in ‘Jeng’ areas) and rural (those living in ‘Xiang’ areas).^{19–21}

In addition to demographic risk factors (urbanisation level, gender and age), concurrent allergic diseases identified by ICD-9-CM codes in the claims records before the index date included asthma (ICD-9-CM code 493), allergic rhinitis (ICD-9-CM code 477) and allergic conjunctivitis (ICD-9-CM code 372.05, 372.10 and 372.14). Baseline comorbidities were also identified by ICD-9-CM codes,




Figure 1 Flow diagram summarising the process of the enrolment. LHID, Longitudinal Health Insurance Database.

including hypertension (ICD-9-CM codes 401–405), chronic obstructive pulmonary disease (ICD-9-CM code 491, 492, 494 and 496), hyperlipidaemia (ICD-9-CM code 272), chronic kidney disease (ICD-9-CM code 582, 583, 585, 586 and 588), diabetes mellitus (ICD-9-CM code 250), chronic liver disease (ICD-9-CM code 571), stroke (ICD-9-CM code 430–438), mental disorders (ICD-9-CM code 290–319), coronary artery disease (ICD-9-CM code 410–414) and alcohol-attributed disease (ICD-9-CM codes 291.0–9, 303, 305.0, 357.5, 425.5, 535.3, 571.0–3980.0 and V11.3). Concurrent allergic diseases and comorbidities were ascertained at least more than two outpatient claims during study periods.

Statistical analysis

We used χ^2 test to compare distributions of categorical demographic and clinical characteristics between the AD and non-AD control cohorts. Student's t-test and Wilcoxon rank-sum test were used to compare mean age and follow-up time (y) between the two cohorts, as appropriately. In the patients with AD, survival time was calculated until hospitalisation, an ambulatory visit for keloids or the end of the study period (31 December 2010), whichever came first. Keloid incidence rates per 1000 person-years were compared between the two cohorts. Kaplan-Meier analysis was used to estimate cumulative incidence of keloids, and the differences between the curves were compared by two-tailed log-rank test. Univariable and multivariable Cox proportional hazard regression models were used to calculate HRs and 95% CIs for keloids in patients with AD when the proportional hazards assumption was satisfied. Multivariate Cox models were adjusted for age, gender, urbanisation level, concurrent allergic diseases and relevant comorbidities. Statistical Analysis Software, V.9.4 was used for data analyses; a p value less than 0.05 in a two-tailed test was considered statistically significant.

Patient and public involvement

There were neither patients nor the public involved in this study.

Results

Table 1 shows the demographic characteristics and comorbidities of patients in the AD and non-AD cohorts. In the AD cohort, 50.71% patients were younger than 40 years old, and 64.81% patients were female. Compared with the non-AD cohort, the AD cohort had significantly higher percentages of patients with asthma (25.41 vs 12.91, $p<0.001$), allergic rhinitis (54.78 vs 35.92, $p<0.001$), allergic conjunctivitis (57.22 vs 42.08, $p<0.001$), hypertension (34.09 vs 21.60, $p<0.001$), diabetes mellitus (23.51 vs 13.46, $p<0.001$), hyperlipidaemia (34.56 vs 21.08, $p<0.001$), chronic kidney disease (12.03 vs 6.20, $p<0.001$), chronic liver disease (35.83 vs 21.98, $p<0.001$), chronic obstructive pulmonary disease (26.74 vs 15.08, $p<0.001$), stroke (5.85 vs 2.97, $p<0.001$), coronary artery disease (5.02 vs 2.71, $p<0.001$), mental disorders (18.05 vs

11.33, $p<0.001$) and alcohol-attributable disease (3.72 vs 2.28, $p<0.001$). During a median observation time of 1.9 years, 1.95% (163) of the patients with AD had keloids (IQR=0.8–4.3). The incidence of keloids was significantly ($p=0.022$) higher in the AD cohort compared with the non-AD cohort (532 patients with keloids out of 33 484 controls matched for age, and gender (1.59%)) during a median observation time of 9.0 years (IQR=6.2–11.8). The median duration of follow-up for keloids was significantly shorter in the AD group (1.9 years) compared with the non-AD group (9.0 years).

Table 2 stratifies the keloid incidence rates and HRs by age, gender and concurrent allergic disease or comorbidity. During the follow-up period, keloids developed in 1.95% (163) of the patients with AD and in 1.59% (532) of the patients without AD. The overall keloid risk was 3.19 times greater in the AD group compared with the non-AD group (3.19 vs 1.07 per 1000 person-years, respectively) after adjusting for age, gender, urbanisation level, concurrent allergic diseases (asthma, allergic rhinitis and allergic conjunctivitis) and related comorbidities (hypertension, hyperlipidaemia, diabetes mellitus, chronic liver disease, chronic obstructive pulmonary disease, chronic kidney diseases, stroke, mental disorders, coronary artery disease and alcohol attributed disease).

In both the AD and non-AD cohorts, gender-specific analyses showed that the incidence of keloids was higher in women than in men (3.52 and 2.61 per 1000 person-years, respectively, in the AD cohort; 1.27 vs 0.69 per 1000 person-years, respectively, in the non-AD cohort). However, men in the AD group had a significantly higher keloid risk compared with men in the non-AD group (adjusted HR=3.67, 95% CI 2.47 to 5.44) and women in the AD group had a significantly higher keloid risk compared with women in the non-AD group (adjusted HR=3.03, 95% CI 2.39 to 3.83).

In all age groups, the AD cohort also had a significantly higher incidence of keloids compared with the non-AD cohort. Age-specific risk comparisons showed that compared with the non-AD cohort, the AD cohort had a significantly higher keloid risk in all age groups, and the risk was highest in patients under 40 years old. Keloid risk was higher in patients with AD than in patients without AD, regardless of concurrent allergic disease or comorbidity.

Table 3 shows that keloid risk increased with severity of AD, particularly in patients with moderate to severe AD.

Figure 2 compares the Kaplan-Meier curves for the cumulative incidence of keloids between the AD and non-AD groups at the 15-year follow-up. The baseline risk of keloid development was significantly higher in the AD group (2.83%, $p<0.001$) compared with the non-AD group (1.59%). The Kaplan-Meier curves also showed a significantly higher cumulative incidence of keloids in the AD group compared with the non-AD group (log-rank test, $p<0.001$).

Table 1 Baseline characteristics of patients with and without atopic dermatitis

Variables	Atopic dermatitis		P values
	Yes (n=8371)	No (n=33484)	
Keloid patients, n (%)	163 (1.95)	532 (1.59)	0.022
Period of developing keloid, median (IQR), years	1.9 (0.8–4.3)	9.0 (6.2–11.8)	<0.001
Mean age at diagnosis of keloid, years	33.1 (14.8)	36.8 (14.4)	<0.001
Age group, n (%)			
<40	4245 (50.71)	16980 (50.71)	
40–59	2601 (31.07)	10404 (31.07)	
≥60	1525 (18.22)	6100 (18.22)	1.000
Gender, n (%)			
Men	2946 (35.19)	11784 (35.19)	
Women	5425 (64.81)	21700 (64.81)	1.000
Degree of urbanisation, n (%)			
Urban	5267 (62.92)	20463 (61.11)	
Suburban	1332 (15.91)	5665 (16.92)	
Rural	1772 (21.17)	7356 (21.97)	0.008
Concurrent allergic disease, n (%)			
Asthma	2127 (25.41)	4323 (12.91)	<0.001
Allergic rhinitis	4586 (54.78)	12028 (35.92)	<0.001
Allergic conjunctivitis	4790 (57.22)	14089 (42.08)	<0.001
Comorbidity, n (%)			
Hypertension	2854 (34.09)	7232 (21.60)	<0.001
Diabetes mellitus	1968 (23.51)	4508 (13.46)	<0.001
Hyperlipidaemia	2893 (34.56)	7059 (21.08)	<0.001
Chronic kidney disease	1007 (12.03)	2076 (6.20)	<0.001
Chronic liver disease	2999 (35.83)	7361 (21.98)	<0.001
Chronic obstructive pulmonary disease	2238 (26.74)	5050 (15.08)	<0.001
Stroke	490 (5.85)	994 (2.97)	<0.001
Coronary artery disease	420 (5.02)	909 (2.71)	<0.001
Mental disorders	1511 (18.05)	3794 (11.33)	<0.001
Alcohol attributed disease	311 (3.72)	765 (2.28)	<0.001

DISCUSSION

The above comparisons supported our study hypothesis that patients with AD have a higher than normal risk of subsequently developing keloids. This study is the first to demonstrate an association between AD and keloids in an Asian population. The keloid risk was 3.19-fold higher in the AD group compared with the non-AD group. Notably, the females in the AD group had a significantly higher incidence rate of keloids compared with males. Specifically, compared with patients in the non-AD group, the AD group had a significantly higher keloid risk in all age groups, especially in those less than 40 years old. In the AD group, keloid risk was significantly associated with the presence of comorbidities and concurrent allergic diseases and keloid risk increased with the severity of AD.

Previous studies of the association between AD and keloids have obtained inconsistent results. In

Hajdarbegovic *et al.*⁷ for example, a case–control study of 131 patients with keloids and 258 controls recruited during 2000–2012 revealed no association between AD and keloids in the adjusted model (adjusted OR 1.27; 95% CI 0.76 to 2.13). However, the questionnaire-based reporting method used in that study may have caused information bias resulting in under-diagnosis of AD. Another limitation is that the authors did not report AD-compatible itchy flexural rash with AD, which may have also contributed to underdiagnosis of AD. Underdiagnosis of keloids is also common because patients with keloids are often immigrants, who may be unaccustomed to seeking prompt medical attention for non-critical medical problems. Additionally, the diverse ancestry of the population analysed in Hajdarbegovic *et al.*⁷ may have had a confounding effect on the association between AD and keloids. A final limitation of Hajdarbegovic *et al.* is its small sample size.

Table 2 Incidence and HRs of keloid by demographic characteristics and concurrent allergic disease or comorbidity among patients with or without atopic dermatitis

Variables	Patients with atopic dermatitis			Patients without atopic dermatitis			Compared with non-atopic dermatitis		
	Keloid	PYs	Rate	Keloid	PYs	Rate	Crude HR (95% CI)	Adjusted HR* (95% CI)	Adjusted HR† (95% CI)
Overall	163	50990.85	3.19	532	499087.75	1.07	3.84 (3.17 to 4.66)‡	3.80 (3.14 to 4.61)‡	3.19 (2.60 to 3.89)‡
Gender									
Men	47	17992.05	2.61	123	176032.98	0.69	4.81 (3.31 to 6.99)‡	4.72 (3.24 to 6.85)‡	3.67 (2.47 to 5.44)‡
Women	116	32998.80	3.52	409	323054.76	1.27	3.56 (2.84 to 4.45)‡	3.53 (2.82 to 4.42)‡	3.03 (2.39 to 3.83)‡
Stratify by age (years)									
<40	122	26311.81	4.64	75	63234.61	1.19	4.07 (3.27 to 5.07)‡	4.07 (3.27 to 5.08)‡	3.44 (2.74 to 4.31)‡
40–59	30	15707.99	1.91	29	38829.76	0.75	3.31 (2.19 to 4.99)‡	3.32 (2.20 to 4.99)‡	2.74 (1.79 to 4.17)‡
≥60	11	8971.06	1.23	11	22791.71	0.48	2.87 (1.49–5.56)§	2.82 (1.46–5.44)§	2.36 (1.22–4.59)¶
Concurrent allergic disease**									
No	22	9661.48	2.28	139	194784.11	0.71	5.41 (4.02 to 7.27)‡	4.03 (2.55 to 6.37)‡	4.41 (3.27 to 5.94)‡
Yes	141	41329.37	3.41	393	304303.64	1.29	3.19 (2.52 to 4.05)‡	3.34 (2.72 to 4.12)‡	2.62 (2.05 to 3.34)‡
Comorbidity††									
No	59	14141.34	4.17	247	246626.44	1.00	4.13 (2.62 to 6.52)‡	4.85 (3.61 to 6.52)‡	3.78 (2.38 to 5.98)‡
Yes	104	36849.51	2.82	285	252461.31	1.12	3.37 (2.74 to 4.15)‡	2.82 (2.22 to 3.58)‡	3.08 (2.48 to 3.82)‡

*Model adjusted for age and gender.

†Model adjusted for age, gender, urbanisation, concurrent allergic diseases (asthma, allergic rhinitis and allergic conjunctivitis) and relevant comorbidities (hypertension, diabetes mellitus, hyperlipidaemia, chronic kidney diseases, chronic liver disease, chronic obstructive pulmonary disease, stroke, coronary artery disease, mental disorders and alcohol attributed disease).

‡P<0.001.

§P=0.002.

¶P=0.011.

**Patients with any concurrent allergic disease, including asthma, allergic rhinitis and allergic conjunctivitis, were classified as the allergic disease group.

††Patients with any examined comorbidities, including hypertension, diabetes mellitus, hyperlipidaemia, chronic kidney diseases, chronic liver disease, chronic obstructive pulmonary disease, stroke, coronary artery disease, mental disorders and alcohol attributed disease, were classified as the comorbidity group.

PYs, person-years; rate, incidence rate in per 1000 person-years.

Table 3 Incidence rates and HRs of keloid risk in patients with different severity of atopic dermatitis

Variables	N	keloid	Rate	Crude HR (95% CI)	Crude HR (95% CI)	AdjustedHR* (95% CI)	Adjusted HR* (95% CI)
Without atopic dermatitis	33484	532	1.07	1.00 (reference)		1.00 (reference)	
With atopic dermatitis	8371	163	3.19	3.84 (3.17 to 4.66)†		3.19 (2.60 to 3.89)†	
Mild	5193	81	2.58	3.10 (2.42 to 3.97)†	1.00 (reference)	2.54 (1.98 to 3.27)†	1.00 (reference)
Moderate to severe	3178	82	4.19	5.02 (3.93 to 6.42)†	1.62 (1.19-2.21)‡	4.39 (3.38 to 5.70)†	1.73 (1.26 to 2.36)†

*Model adjusted for age, gender, urbanisation, concurrent allergic diseases (asthma, allergic rhinitis and allergic conjunctivitis) and relevant comorbidities (hypertension, diabetes mellitus, hyperlipidaemia, chronic kidney diseases, chronic liver disease, chronic obstructive pulmonary disease, stroke, coronary artery disease, mental disorders and alcohol attributed disease).

†P<0.001.

‡P=0.002.

IRR, incidence rate ratio; PYs, person-years; rate, incidence rate in per 1000 person-years.

Although the exact mechanisms underlying the relationship between AD and keloids are unclear, several possible links have been hypothesised. First, thymic stromal lymphopoietin (TSLP), an interleukin (IL)-7-like cytokine, is believed to cause Th2-type inflammatory responses.^{22 23} Upregulation of Th2 cytokines may be linked to atopic diseases.²⁴ The TSLP is also reportedly linked to AD. In 2010, Lee *et al* collected sera from 232 children, including 75 with AD. Comparisons showed higher serum TSLP in children with AD compared with normal controls. Studies of patients with AD also indicate that skin lesions are associated with increased TSLP gene expression.²⁵ In another recent study, Shin *et al*² showed that TSLP expression was higher in keloidal tissue compared with normal tissue. Their data showed that by inducing production of collagen and tissue growth factor beta (TGF- β), TSLP contributes to

excessive deposition of ECM. Since TSLP expression is apparently associated with keloids in patients with AD, keloids and AD may be induced through similar inflammatory pathways.

Periostin, which is in the fasciclin family of proteins, has important functions in the ECM and is also a matricellular protein associated with allergic inflammation.²⁶ Kou *et al* reported that periostin expression in serum or in inflamed sites of patients with AD was associated with the clinical severity of AD, which suggests that periostin mediates AD.^{27 28} Furthermore, periostin can activate TGF- β and NF- κ B to induce IL-6 in eosinophils and lead keratinocytes to produce TSLP. Periostin in fibroblasts can then be induced by activated eosinophils through TGF- β reciprocally. Besides, Zhou *et al*²⁹ reported that increased expression of periostin was found in keloid scars. Since periostin participates in the epithelial-mesenchymal




Figure 2 Cumulative incidence of keloid among patients with atopic dermatitis and the control cohort. The incidence of keloid in atopic dermatitis patients and controls was compared using the log-rank test by a Kaplan-Meier analysis.

transition, periostin is important in forming both AD and keloids.

Sirtuins, which are class III histone deacetylases, are implicated in fibrotic disease, inflammation and cellular proliferation, apoptosis and differentiation.³⁰ Ming *et al* demonstrated that loss of sirtuin can reduce the epidermal formation of filaggrin, resulting in disruption of the epidermal barrier and a predisposition to occurrence of AD.^{31 32} Bai *et al*³³ showed that expression of sirtuin was suppressed in fibrotic scarring and that sirtuin deletion enhanced fibrotic wound healing. Sirtuin over-expression can attenuate the expression of ECM induced by TGF- β . Loss of sirtuin leads to both AD development and excessive fibrotic scarring. Finally, we hypothesised that individuals with AD may be predisposed to subsequent development of keloids.

TGF- β is a main fibrotic growth factor involved in wound healing and tissue repair.³⁴ Upregulation of TGF- β gene with proinflammation cytokines was found in keloids,^{35 36} resulting in keloidal fibroblasts formation together with collagen synthesis.³⁷ Previously mentioned literatures showed periostin, TSLP and loss of sirtuins can activate TGF- β in AD and keloids. Therefore, patients with AD are prone to develop keloids.

Psychological stress can trigger various physiological responses in the endocrine, nervous and immune systems.^{38–45} Psychoemotional stress is associated with a higher than normal incidence of several skin diseases including AD,^{38 46–53} and it is also known to aggravate AD. Anxiety and depression have been reported to correlate with AD.^{54–56} In Chen *et al*,⁵⁵ the patients with AD had a higher risk of developing major depression, any depressive disorder and anxiety disorders in later life. Other studies have reported an association between anxiety and keloids.⁵⁷ In Furtado *et al*,⁵⁷ for example, psychological stress was evaluated in 25 patients with keloids before undergoing surgical resection and postoperative radiotherapy. They found that stress had a direct correlation with development of keloids. Notably, the hypothalamic–pituitary–adrenal axis has a well-known mediating role in the eventual contributing effects of psychological stress on development of both AD and keloids.^{48 58} Psychological stress also increases concentrations of nerve growth factor (NGF).⁵⁹ Ewa and colleagues demonstrated that comparing to healthy ones, patients with AD had significant increased plasma levels of substance P and NGF.⁶⁰ The NGF is made by keratinocytes and contributes to neurogenic inflammation.⁶¹ Akaishi *et al* found that neurogenic inflammation has an important role in keloid formation.⁶² Therefore, we suggest that AD should be considered a risk factor for keloids.

A notable strength of this study is the use of a large representative population-based dataset, which enabled analysis of all cases of AD and keloids during the study period. Moreover, the large sample size also provided sufficient statistical power to compare the complex relationships between these two diseases in the AD and non-AD groups. Notably, since most of the Taiwan population

is ethnic Chinese, analysis of this population may yield valuable insights into the roles of genetic ancestry in AD and keloids. However, several limitations of this study are noted. First, the data analysed in this study were originally collected for insurance purposes rather than for research purposes. Second, data for diagnoses of AD and keloids in the study population were based on ICD-9-CM codes entered by different doctors, which may have introduced coding errors. The Taiwan Bureau of National Health Insurance has attempted to minimise this problem by performing regular audits and by imposing large fines on hospitals that make coding errors. Nevertheless, the NHIRD has been rigorously examined and extensively used in many studies in recent years.^{63–65} Third, despite the relatively low healthcare costs in Taiwan, some patients attempt to eliminate AD and keloids through folk remedies or through private treatment by local pharmacists rather than by medical professionals. Moreover, the NHIRD does not record all information regarding risk factors such as body mass index, blood groups, smoking, alcohol consumption and operation sites, which potentially compromises our results. Lacking information about trauma history, anatomical sites, keloid size and number, acne and the different methods of surgery is also a flaw in the report. Finally, the retrospective design of this study only revealed statistically significant relationships. Further studies are needed to validate actual causal relationships. Further clinical trials are needed to clarify the underlying mechanisms of this association.

In conclusion, this study showed that AD is associated with development of keloids and suggested that AD could be an important risk factor in the future development of keloids. Nevertheless, further studies are needed to clarify this association and to explore pathophysiological mechanisms that are common to AD and keloids.

Author affiliations

¹Department of Dermatology, Kaohsiung Veterans General Hospital, Kaohsiung, Taiwan

²Cosmetic Applications and Management Department, Yuh-Ing Junior College of Health Care & Management, Kaohsiung, Taiwan

³Graduate Institute of Medicine, College of Medicine, Kaohsiung Medical University, Kaohsiung, Taiwan

⁴Department of Orthopedics, Taipei Veterans General Hospital, Taipei, Taiwan

⁵Department of Neurosurgery, Zhangjiagang First People's Hospital, Zhangjiagang, China

⁶Department of Neurosurgery, Qinghe County Central Hospital, Qinghe, China

⁷Department of Cardiology, Hebei Quyang Renji Hospital, Quyang, China

⁸Division of Neurosurgery, Department of Surgery, Kaohsiung Medical University Hospital, Kaohsiung, Taiwan

⁹Department of Surgery, School of Medicine, College of Medicine, Kaohsiung Medical University, Kaohsiung, Taiwan

Contributors Substantial contributions to conception and design: Y-YL, C-CL, W-WY, Q-RW, LZ and C-HW. Drafting and revising the article critically for important intellectual content: Y-YL, Q-RW, C-LZ, C-CL and C-HW. Final approval of the version to be published: Y-YL and C-HW.

Funding The study was supported by Kaohsiung Veterans General Hospital (VGHKS-106-68) (VGHKS-107-70).

Disclaimer The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- Han J, Han J, Yu D, *et al.* Association of ADAM33 gene polymorphisms with keloid scars in a northeastern Chinese population. *Cell Physiol Biochem* 2014;34:981–7.
- Shin JU, Kim SH, Kim H, *et al.* TSLP is a Potential initiator of collagen synthesis and an activator of CXCR4/SDF-1 axis in keloid pathogenesis. *J Invest Dermatol* 2016;136:507–15.
- Viera MH, Caperton CV, Berman B. Advances in the treatment of keloids. *J Drugs Dermatol* 2011;10:468–80.
- Jones ME, Hardy C, Ridgway J. Keloid management: a retrospective case review on a new approach using surgical excision, platelet-rich plasma, and in-office superficial photon X-ray radiation therapy. *Adv Skin Wound Care* 2016;29:303–7.
- Bissek AC, Tabah EN, Kouotou E, *et al.* The spectrum of skin diseases in a rural setting in Cameroon (sub-Saharan Africa). *BMC Dermatol* 2012;12:7.
- Viera MH, Vivas AC, Berman B. Update on keloid management: clinical and basic science advances. *Adv Wound Care* 2012;1:200–6.
- Hajdarbegovic E, Bloem A, Balak D, *et al.* The association between atopic disorders and keloids: a case-control study. *Indian J Dermatol* 2015;60:635.
- Matsusue A, Kiyohara C, Tanaka K, *et al.* ADAM33 genetic polymorphisms and risk of atopic dermatitis among Japanese children. *Clin Biochem* 2009;42:477–83.
- Wu CH, Tung YC, Chai CY, *et al.* Increased risk of osteoporosis in patients with peptic ulcer disease: a nationwide population-based study. *Medicine* 2016;95:e3309.
- Wu CH, Chai CY, Tung YC, *et al.* Herpes zoster as a risk factor for osteoporosis: a 15-year nationwide population-based study. *Medicine* 2016;95:e3943.
- Wu CH, Tung YC, Lin TK, *et al.* Hip fracture in people with erectile dysfunction: a nationwide population-based cohort study. *PLoS One* 2016;11:e0153467.
- Wu LC, Hwang CY, Chung PI, *et al.* Autoimmune disease comorbidities in patients with atopic dermatitis: a nationwide case-control study in Taiwan. *Pediatr Allergy Immunol* 2014;25:586–92.
- Shreberk-Hassidim R, Hassidim A, Gronovich Y, *et al.* Atopic dermatitis in Israeli Adolescents from 1998 to 2013: trends in time and association with migraine. *Pediatr Dermatol* 2017;34:247–52.
- Katayama I, Aihara M, Ohya Y, *et al.* Japanese guidelines for atopic dermatitis 2017. *Allergol Int* 2017;66:230–47.
- Werfel T, Schwerk N, Hansen G, *et al.* The diagnosis and graded therapy of atopic dermatitis. *Dtsch Arztebl Int* 2014;111:509–20.
- Ricci G, Dondi A, Patrizi A, *et al.* Systemic therapy of atopic dermatitis in children. *Drugs* 2009;69:297–306.
- Sun LM, Wang KH, Lee YC. Keloid incidence in Asian people and its comorbidity with other fibrosis-related diseases: a nationwide population-based study. *Arch Dermatol Res* 2014;306:803–8.
- Flohr C, Mann J. New insights into the epidemiology of childhood atopic dermatitis. *Allergy* 2014;69:3–16.
- Shyu CS, Lin HK, Lin CH, *et al.* Prevalence of attention-deficit/hyperactivity disorder in patients with pediatric allergic disorders: a nationwide, population-based study. *J Microbiol Immunol Infect* 2012;45:237–42.
- Chien IC, Chang KC, Lin CH, *et al.* Prevalence of diabetes in patients with bipolar disorder in Taiwan: a population-based national health insurance study. *Gen Hosp Psychiatry* 2010;32:577–82.
- Chen CY, Liu CY, Su WC, Wc S, *et al.* Factors associated with the diagnosis of neurodevelopmental disorders: a population-based longitudinal study. *Pediatrics* 2007;119:e435–e443.
- Liu YJ. Thymic stromal lymphopoietin: master switch for allergic inflammation. *J Exp Med* 2006;203:269–73.
- Sano Y, Masuda K, Tamagawa-Mineoka R, *et al.* Thymic stromal lymphopoietin expression is increased in the horny layer of patients with atopic dermatitis. *Clin Exp Immunol* 2013;171:330–7.
- Irvine AD, McLean WH, Leung DY. Filaggrin mutations associated with skin and allergic diseases. *N Engl J Med* 2011;365:1315–27.
- Lee EB, Kim KW, Hong JY, *et al.* Increased serum thymic stromal lymphopoietin in children with atopic dermatitis. *Pediatr Allergy Immunol* 2010;21:e457–e460.
- Izuhara K, Nunomura S, Nanri Y, *et al.* Periostin in inflammation and allergy. *Cell Mol Life Sci* 2017;74:4293–303.
- Masuoka M, Shiraishi H, Ohta S, *et al.* Periostin promotes chronic allergic inflammation in response to Th2 cytokines. *J Clin Invest* 2012;122:2590–600.
- Kou K, Okawa T, Yamaguchi Y, *et al.* Periostin levels correlate with disease severity and chronicity in patients with atopic dermatitis. *Br J Dermatol* 2014;171:283–91.
- Zhou HM, Wang J, Elliott C, *et al.* Spatiotemporal expression of periostin during skin development and incisional wound healing: lessons for human fibrotic scar formation. *J Cell Commun Signal* 2010;4:99–107.
- Serravallo M, Jagdeo J, Glick SA, *et al.* Sirtuins in dermatology: applications for future research and therapeutics. *Arch Dermatol Res* 2013;305:269–82.
- Ming M, Zhao B, Shea CR, *et al.* Loss of sirtuin 1 (SIRT1) disrupts skin barrier integrity and sensitizes mice to epicutaneous allergen challenge. *J Allergy Clin Immunol* 2015;135:936–45.
- Nomura T, Kabashima K. Advances in atopic dermatitis in 2015. *J Allergy Clin Immunol* 2016;138:1548–55.
- Bai XZ, Liu JQ, Yang LL, *et al.* Identification of sirtuin 1 as a promising therapeutic target for hypertrophic scars. *Br J Pharmacol* 2016;173:1589–601.
- Fujiwara M, Muragaki Y, Ooshima A. Upregulation of transforming growth factor-beta1 and vascular endothelial growth factor in cultured keloid fibroblasts: relevance to angiogenic activity. *Arch Dermatol Res* 2005;297:161–9.
- Liu W, Wang DR, Cao YL. TGF-beta: a fibrotic factor in wound scarring and a potential target for anti-scarring gene therapy. *Curr Gene Ther* 2004;4:123–36.
- Chen W, Fu X, Sun X, *et al.* Analysis of differentially expressed genes in keloids and normal skin with cDNA microarray. *J Surg Res* 2003;113:208–16.
- Andrews JP, Marttala J, Macarak E, *et al.* Keloids: the paradigm of skin fibrosis - Pathomechanisms and treatment. *Matrix Biol* 2016;51:37–46.
- Hall JM, Cruser D, Podawiltz A, *et al.* Psychological stress and the cutaneous immune response: roles of the HPA axis and the sympathetic nervous system in atopic dermatitis and psoriasis. *Dermatol Res Pract* 2012;2012:1–11.
- Dhabhar FS, Miller AH, McEwen BS, *et al.* Effects of stress on immune cell distribution. Dynamics and hormonal mechanisms. *J Immunol* 1995;154:5511–27.
- Bowers SL, Bilbo SD, Dhabhar FS, *et al.* Stressor-specific alterations in corticosterone and immune responses in mice. *Brain Behav Immun* 2008;22:105–13.
- Dhabhar FS, McEwen BS. Acute stress enhances while chronic stress suppresses cell-mediated immunity in vivo: a potential role for leukocyte trafficking. *Brain Behav Immun* 1997;11:286–306.
- Altemus M, Rao B, Dhabhar FS, *et al.* Stress-induced changes in skin barrier function in healthy women. *J Invest Dermatol* 2001;117:309–17.
- Dhabhar FS. Acute stress enhances while chronic stress suppresses skin immunity. The role of stress hormones and leukocyte trafficking. *Ann N Y Acad Sci* 2000;917:876–93.
- Dhabhar FS, McEwen BS. Enhancing versus suppressive effects of stress hormones on skin immune function. *Proc Natl Acad Sci U S A* 1999;96:1059–64.
- Dhabhar FS, Miller AH, Stein M, *et al.* Diurnal and acute stress-induced changes in distribution of peripheral blood leukocyte subpopulations. *Brain Behav Immun* 1994;8:66–79.
- Hochman B, Isoldi FC, Furtado F, *et al.* New approach to the understanding of keloid: psychoneuroimmune-endocrine aspects. *Clin Cosmet Investig Dermatol* 2015;8:67–73.
- Misery L, Touboul S, Vinçot C, *et al.* [Stress and seborrheic dermatitis]. *Ann Dermatol Venerol* 2007;134:833–7.
- Suárez AL, Feramisco JD, Koo J, *et al.* Psychoneuroimmunology of psychological stress and atopic dermatitis: pathophysiological and therapeutic updates. *Acta Derm Venerol* 2012;92:7–15.
- Yu R, Huang Y, Zhang X, *et al.* Potential role of neurogenic inflammatory factors in the pathogenesis of vitiligo. *J Cutan Med Surg* 2012;16:230–44.
- Hadshiew IM, Foitzik K, Arck PC, *et al.* Burden of hair loss: stress and the underestimated psychosocial impact of telogen effluvium and androgenetic alopecia. *J Invest Dermatol* 2004;123:455–7.

51. Chida Y, Mao X. Does psychosocial stress predict symptomatic herpes simplex virus recurrence? A meta-analytic investigation on prospective studies. *Brain Behav Immun* 2009;23:917–25.
52. Yosipovitch G, Tang M, Dawn AG, et al. Study of psychological stress, sebum production and acne vulgaris in adolescents. *Acta Derm Venereol* 2007;87:135–9.
53. Manolache L, Benea V. Stress in patients with alopecia areata and vitiligo. *J Eur Acad Dermatol Venereol* 2007;21:921–8.
54. Hashizume H, Takigawa M. Anxiety in allergy and atopic dermatitis. *Curr Opin Allergy Clin Immunol* 2006;6:335–9.
55. Cheng CM, Hsu JW, Huang KL, et al. Risk of developing major depressive disorder and anxiety disorders among adolescents and adults with atopic dermatitis: a nationwide longitudinal study. *J Affect Disord* 2015;178:60–5.
56. Hashiro M, Okumura M. Anxiety, depression and psychosomatic symptoms in patients with atopic dermatitis: comparison with normal controls and among groups of different degrees of severity. *J Dermatol Sci* 1997;14:63–7.
57. Furtado F, Hochman B, Farber PL, et al. Psychological stress as a risk factor for postoperative keloid recurrence. *J Psychosom Res* 2012;72:282–7.
58. Sheridan JF, Padgett DA, Avitsur R, et al. Experimental models of stress and wound healing. *World J Surg* 2004;28:327–30.
59. Joachim RA, Kuhlmei A, Dinh QT, et al. Neuronal plasticity of the "brain-skin connection": stress-triggered up-regulation of neuropeptides in dorsal root ganglia and skin via nerve growth factor-dependent pathways. *J Mol Med* 2007;85:1369–78.
60. Teresiak-Mikołajczak E, Czarnecka-Operacz M, Jenerowicz D, et al. Neurogenic markers of the inflammatory process in atopic dermatitis: relation to the severity and pruritus. *Postepy Dermatol Alergol* 2013;30:286–92.
61. Tominaga M, Takamori K. Itch and nerve fibers with special reference to atopic dermatitis: therapeutic implications. *J Dermatol* 2014;41:205–12.
62. Akaishi S, Ogawa R, Hyakusoku H. Keloid and hypertrophic scar: neurogenic inflammation hypotheses. *Med Hypotheses* 2008;71:32–8.
63. Wu CH, Zhang ZH, Wu MK, et al. Increased migraine risk in osteoporosis patients: a nationwide population-based study. *Springerplus* 2016;5:1378.
64. Wu CH, Lu YY, Chai CY, et al. Increased risk of osteoporosis in patients with erectile dysfunction: A nationwide population-based cohort study. *Medicine* 2016;95:e4024.
65. Wu CH, Tsai TH, Su YF, et al. Traumatic Brain Injury and Substance Related Disorder: A 10-Year Nationwide Cohort Study in Taiwan. *Neural Plast* 2016;2016:8030676.