

Linoleic Acid-Induced Mitochondrial Ca²⁺ Efflux Causes Peroxynitrite Generation and Protein Nitrotyrosylation

Hong-Mei Zhang¹, Howard Dang², Chih-Ko Yeh^{3,4}, Bin-Xian Zhang^{1,4*}

1 Department of Medicine, University of Texas Health Science Center at San Antonio, San Antonio, Texas, United States of America, **2** Department of Community Dentistry, University of Texas Health Science Center at San Antonio, San Antonio, Texas, United States of America, **3** Department of Dental Diagnostic Science, University of Texas Health Science Center at San Antonio, San Antonio, Texas, United States of America, **4** Geriatric Research, Education and Clinical Center, South Texas Veterans Health Care System, Audie L. Murphy Division, San Antonio, Texas, United States of America

Abstract

It is well known that excessive non-esterified fatty acids in diabetes contribute to the pathogenesis of renal complications although the mechanism remains elusive. Enhanced oxidative stress has been hypothesized as a unified factor contributing to diabetic complications and increased protein nitrotyrosylation has been reported in the kidneys of diabetic patients. In the current manuscript we described that linoleic acid (LA) caused mitochondrial Ca²⁺ efflux and peroxynitrite production, along with increased nitrotyrosine levels of cellular proteins in primary human mesangial cells. The peroxynitrite production by LA was found to depend on mitochondrial Ca²⁺ efflux. Downregulation of hsp90β1, which has been previously shown to be essential for polyunsaturated fatty acid-induced mitochondrial Ca²⁺ efflux, significantly diminished LA-responsive mitochondrial Ca²⁺ efflux and the coupled peroxynitrite generation, implicating a critical role of hsp90β1 in the LA responses. Our results further demonstrated that mitochondrial complexes I and III were directly involved in the LA-induced peroxynitrite generation. Using the well established type 2 diabetic animal model db/db mice, we observed a dramatically enhanced LA responsive mitochondrial Ca²⁺ efflux and protein nitrotyrosylation in the kidney. Our study thus demonstrates a cause-effect relationship between LA and peroxynitrite or protein nitrotyrosylation and provides a novel mechanism for lipid-induced nephropathy in diabetes.

Citation: Zhang H-M, Dang H, Yeh C-K, Zhang B-X (2009) Linoleic Acid-Induced Mitochondrial Ca²⁺ Efflux Causes Peroxynitrite Generation and Protein Nitrotyrosylation. PLoS ONE 4(6): e6048. doi:10.1371/journal.pone.0006048

Editor: Carmine Zoccali, L' Istituto di Biomedicina ed Immunologia Molecolare, Consiglio Nazionale delle Ricerche, Italy

Received: March 20, 2009; **Accepted:** May 27, 2009; **Published:** June 26, 2009

This is an open-access article distributed under the terms of the Creative Commons Public Domain declaration which stipulates that, once placed in the public domain, this work may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose.

Funding: This work was partially supported by grants from the American Heart Association (0235065N to BXZ), the National Institute of Health (HL075011 to BXZ) and the Department of Veterans Affairs (to BXZ and CKY). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: zhangb2@uthscsa.edu

Introduction

Elevated non-esterified fatty acids (NEFA) have been reported in patients and animal models of type 2 diabetes and shown to be associated with increased reactive oxygen species (ROS) production from mitochondria [1,2] and nonmitochondrial sources [3]. Theoretically, excessive NEFA may influence mitochondrial ROS production by affecting the regeneration of reduced glutathione, proton gradients and electron transport [1]. Ionic Ca²⁺ in the mitochondria ([Ca²⁺]_m) regulates not only substrate oxidation and ATP production [4,5], but also ROS generation [6], which has been demonstrated to be a major contributing factor to the pathogenesis of diabetic complications including diabetic nephropathy [7]. Under normal physiological conditions, Ca²⁺ diminishes ROS production from both complexes I and III [6]. Whether excessive NEFA contribute to the massive ROS generation in diabetic nephropathy by altering [Ca²⁺]_m is currently unknown.

Superoxide interacts with nitric oxide (NO) to form peroxynitrite, a strong oxidant that attacks various biomolecules and causes functional defects in cells and tissues [8]. Peroxynitrite interacts with and modifies multiple proteins and enzymes at tyrosine residues to form nitrotyrosine. Protein nitrotyrosylation may either up- or down-regulate activities of cellular pathways in a target-dependent manner [8–10]. Increased formation of nitrotyrosine

has been demonstrated in diabetic kidneys [11] and may be associated with the pathogenesis of nephropathy.

In advanced diabetic nephropathy, mesangial cells are major contributors to glomerular mesangial matrix expansion and capillary basement membrane thickening with increased expression of the extracellular matrix components collagen IV, fibronectin, and laminin [12]. Human mesangial (HM) cells grown in high glucose show largely increased mitochondrial membrane potential and ROS production [13]. Elevated ROS generation may activate NF-κB and contribute to increased extracellular matrix accumulation.

NEFA may influence mitochondrial function by altering gene expression [14], metabolism [15], and/or mitochondrial Ca²⁺ homeostasis [4,16]. We have recently reported that polyunsaturated fatty acids (PUFA) induce Ca²⁺ efflux from mitochondria [16,17], an action that may disrupt mitochondrial function and influence ROS generation. It has yet to be determined whether PUFA-induced mitochondrial Ca²⁺ efflux (PIMCE) links to ROS generation. In the current study, we demonstrated that linoleic acid (LA, an 18:2 n-6 PUFA)-responsive mitochondrial Ca²⁺ efflux caused peroxynitrite generation in HM cells. Our results also showed increased nitrotyrosylation of proteins in LA-treated HM cells and in the kidney of db/db diabetic mice, which may in turn impair kidney function and contribute to the pathogenesis of diabetic renal complications.




Figure 1. LA-induced [Ca²⁺]_i mobilization and ROS generation. Human mesangial cells were labeled with fura-2 (1 μM) alone or together with DCF (1 μM) (37°C, 30 min). The fluorescent intensity in the figures was presented as arbitrary unit (AU). (A): Traces represent the [Ca²⁺]_i responses induced by different concentrations of LA. (B): The curve shows the relationship between the amplitude of [Ca²⁺]_i responses and the LA concentrations. (C): Traces represent LA (30 μM)-induced [Ca²⁺]_i mobilization (black trace) and the associated ROS generation (red trace). doi:10.1371/journal.pone.0006048.g001

Results

LA caused [Ca²⁺]_i mobilization, peroxynitrite generation, and protein nitrotyrosylation in HM cells

PIMCE and the indispensable role of hsp90β1 in the process have previously been described in the human teratocarcinoma NT2 cells [17]. To show that PIMCE is a contributing factor to diabetic nephropathy, we determined whether PIMCE occurs in primary cultured HM cells. As demonstrated in Figure 1, LA caused [Ca²⁺]_i rise in HM cells and the amplitude of the LA responses was dose dependent (Figure 1A and B) with maximal response observed at 30–60 μM. Thus 30 μM of LA was used in all subsequent experiments of this study. Furthermore, the [Ca²⁺]_i response of LA (30 μM) was associated with ROS generation in HM cells (Figure 1C).

Because nitrosative injury as indexed by protein nitrotyrosine was dramatically increased in diabetic nephropathy [11], we wanted to know whether the ROS generated along with LA-induced [Ca²⁺]_i mobilization in HM cells was peroxynitrite. Among different types of ROS, 2',7'-dichlorodihydrofluorescein diacetate (DCF) preferentially reacts with peroxynitrite [18]. To determine the ROS signal observed in Figure 1C (as indexed by the DCF fluorescent intensity) was peroxynitrite, HM cells were treated with 5,10,15,20-Tetrakis(4-sulfonatophenyl) porphyrinato iron (III) (FeTPPS, 0.5 μM), a specific peroxynitrite decomposition catalyst [9,19,20]. Compared to control, pretreatment with FeTPPS did not affect the LA-induced [Ca²⁺]_i rise, but significantly attenuated the ROS production (Figure 2A and B; FeTPPS reduced the LA responsive DCF signal to 14.5±9.2% of control, *P*<0.001, *n*=6), suggesting that the generated ROS were predominantly peroxynitrite. The data also indicated that the LA responsive peroxynitrite generation was either unrelated or occurred downstream of the [Ca²⁺]_i rise.

Since peroxynitrite is formed by the interaction of superoxide and NO, we next measured the generation of the latter two substances along with [Ca²⁺]_i mobilization by measuring dihydroethidium (DHE, a fluorescent indicator of superoxide), and diaminofluorescein-2 diacetate (DAF-2, an indicator of NO) fluorescence, respectively. As shown in Figure 2C and D, LA-induced [Ca²⁺]_i mobilization was associated with the generation of superoxide and NO. The specificity of DHE, DAF-2, and DCF was further demonstrated by using the exogenous NO donor (S-nitroso-N-acetylpenicillamine, 50 μM). The NO donor had no

effect on DHE fluorescence (Figure 2E) but rapidly increased the fluorescent intensity of DAF-2 (Figure 2F). The NO donor also gradually increased DCF fluorescence, indicating the formation of peroxynitrite (Figure 2G).

Peroxynitrite is a strongly reactive oxidant that may interact with multiple proteins and cause nitrosative modification of tyrosine residues to form nitrotyrosine. We next measured the level of nitrotyrosine in cellular proteins of HM cells following LA treatment (30 μM, 1 h at 37°C). As shown in Figure 2H and I, LA treatment caused significantly increased nitrotyrosine levels in a spectrum of proteins in HM cells. The MW of these proteins ranged from 18 to 54 kDa (Figure 2H). These results further supported that the LA-induced [Ca²⁺]_i mobilization was associated with peroxynitrite generation and the latter leading to cellular protein nitrotyrosylation in HM cells.

LA-induced mitochondrial Ca²⁺ efflux caused peroxynitrite production

Bradykinin (BK) and thapsigargin (TG) are known to induce [Ca²⁺]_i mobilization by activation of Ca²⁺ release from the ER and store-operated Ca²⁺ influx through inositol-1, 4, 5-trisphosphate dependent and independent mechanisms, respectively. However, as shown in Figure 3A and B, the [Ca²⁺]_i mobilization induced by both BK and TG was not associated with any detectable peroxynitrite production. These results indicated that the rise of [Ca²⁺]_i from the endoplasmic reticulum (ER) Ca²⁺ release and store-operated Ca²⁺ influx did not lead to peroxynitrite generation.

Even though LA-induced [Ca²⁺]_i mobilization was observed in the absence of extracellular Ca²⁺, chelation of extracellular Ca²⁺ significantly diminished LA-induced [Ca²⁺]_i signal (Figure 3C and D), demonstrating that both Ca²⁺ release from the internal Ca²⁺ stores and Ca²⁺ influx were involved in LA responsive [Ca²⁺]_i mobilization. The peroxynitrite responses to LA were unaffected by extracellular Ca²⁺ (Figure 3C–E). Moreover, clamping the [Ca²⁺]_i with 1,2-Bis(2-aminophenoxy)ethane-N,N,N',N'-tetraacetic acid (BAPTA, a membrane permeable chelator of cytosolic Ca²⁺) in the absence of extracellular Ca²⁺, abolished LA-induced [Ca²⁺]_i signal but had no significant effect on peroxynitrite generation (Figure 3F–H; *P*>0.05, *n*=6), indicating that LA responsive peroxynitrite generation occurred independently of [Ca²⁺]_i rise. Because BAPTA does not prevent Ca²⁺ release from the internal Ca²⁺ stores, including ER and mitochondria, these experiments did not exclude the role of Ca²⁺ release from the internal Ca²⁺ stores in peroxynitrite generation.




Figure 2. The LA-induced ROS signal was identified as peroxynitrite. (A): HM cells were labeled with fura-2 and DCF and traces represent LA-induced $[Ca^{2+}]_i$ mobilization (black trace) and ROS generation (red trace); (B): same measurements performed in cells treated with FeTPPS (a specific peroxynitrite decomposer; 0.5 μ M, 30 min) prior to LA addition and the treatment inhibited the ROS signal significantly; (C) and (E): HM cells were labeled with fura-2 and dihydroethidium (DHE, 1 μ M) and traces represent LA (C) or S-nitroso-N-acetylpenicillamine (exogenous NO donor, ExNO, 50 μ M; E) responsive $[Ca^{2+}]_i$ mobilization (black trace) and superoxide generation (red trace); (D) and (F): HM cells were labeled with fura-2 and 4,5-diaminofluorescein diacetate (DAF-2, 0.5 nM) and traces represent LA (D) or ExNO (F) responsive $[Ca^{2+}]_i$ mobilization (black trace) and NO generation (red trace); (G) HM cells were labeled with fura-2 and DCF and traces represent ExNO responses in $[Ca^{2+}]_i$ (black trace) and peroxynitrite (red trace). The nitrotyrosine level in the cellular proteins was determined by western blot analysis in vehicle-(control) or LA-treated HM cells using α -tubulin as the loading control. The image (H) shows a representative experiment. The graphs (I) represent the mean ratio value \pm SEM of the density of protein band over α -tubulin. * P <0.05, n =4, LA vs control. doi:10.1371/journal.pone.0006048.g002




Figure 3. Mitochondria played an essential role in LA-induced [Ca²⁺]_i mobilization and peroxynitrite generation. HM cells were labeled with fura-2 and DCF to measure [Ca²⁺]_i mobilization (black traces) and peroxynitrite generation (red traces). (A): Bradykinin (BK, 100 nM) responses; (B): thapsigargin (TG, 2 μM) responses; (C) and (F): LA responses measured in the presence of extracellular Ca²⁺ (control); (D): LA responses measured without extracellular Ca²⁺; (G) LA responses measured in cells pretreated with BAPTA (50 μM, 30 min) without extracellular Ca²⁺. In the absence of extracellular Ca²⁺, (I): LA responses (control); (J): LA responses in cells pretreated with TG (2 μM, 5 min); (K) LA responses in cells pretreated with FCCP (a mitochondrial uncoupler; 4 μM, 5 min). The values in graphs (E, H, and L) represent the mean ± SEM of the relative amplitudes of LA-induced [Ca²⁺]_i and peroxynitrite responses, **P* < 0.05, ***P* < 0.01, *n* = 6, treated cells vs control. doi:10.1371/journal.pone.0006048.g003

Pretreatment of the cells with TG (2 μM) in the absence of extracellular Ca²⁺ to completely deplete the ER Ca²⁺ stores (as indicated by a complete inhibition of the BK-induced [Ca²⁺]_i signal following TG treatment, data not shown) only partially reduced the LA-induced [Ca²⁺]_i mobilization and had no significant effect on peroxynitrite generation (Figure 3I, J, L). These results indicated that additional Ca²⁺ stores other than ER, such as mitochondria, may be involved in LA-induced [Ca²⁺]_i response.

Interestingly, in the absence of extracellular Ca²⁺, pretreatment of the cells with carbonyl cyanide-p-trifluoromethoxyph-

nylhydrazone (FCCP, 4 μM), a well-known mitochondrial uncoupler, completely inhibited LA-induced peroxynitrite production and also attenuated the [Ca²⁺]_i response (Figure 3I, K, L). FCCP may alleviate LA-induced [Ca²⁺]_i by collapse of the mitochondrial proton gradient, which prevents Ca²⁺ uptake to the mitochondria and thus reduces the subsequent LA-responsive [Ca²⁺]_m efflux. These data indicated that at least part of the [Ca²⁺]_i response to LA resulted from [Ca²⁺]_m efflux, which might be essential to the peroxynitrite generation in HM cells.




Figure 4. LA-induced mitochondrial Ca²⁺ efflux was responsible for peroxynitrite generation. Mitochondria were prepared from HM cells and labeled with X-rhod-1 (2 μM) and DCF (1 μM) with (A) or without Ca²⁺ (B). LA-induced mitochondrial Ca²⁺ efflux (black traces) and peroxynitrite generation (red traces) were indexed by the gradual decrease in X-rhod-1 and increase in DCF fluorescence intensity, respectively. doi:10.1371/journal.pone.0006048.g004

LA had been shown to increase [Ca²⁺]_i by causing [Ca²⁺]_m efflux in human teratocarcinoma NT2 cells [17]. We thus evaluated the role of LA-induced [Ca²⁺]_m efflux in peroxynitrite generation in isolated mitochondria from HM cells. Isolated mitochondria were labeled with X-rhod-1 and DCF to measure Ca²⁺ efflux and peroxynitrite generation, respectively. As shown in Figure 4A, LA caused Ca²⁺ efflux from the mitochondria as indexed by the reduced X-rhod-1 fluorescence intensity (black trace, Figure 4A). Interestingly, a robust increase in peroxynitrite generation was observed along with the [Ca²⁺]_m efflux, as indexed by the enhanced DCF fluorescence intensity (red trace, Figure 4A). Furthermore, chelation of Ca²⁺ to prevent mitochondria from Ca²⁺ refilling during X-rhod-1 and DCF labeling not only blocked LA-induced Ca²⁺ efflux from mitochondria (black trace, Figure 4B) but also significantly inhibited LA-induced peroxynitrite generation (red trace, Figure 4B), indicating that peroxynitrite generation was dependent on LA-induced [Ca²⁺]_m efflux.

Hsp90β1 played a critical role in LA-induced mitochondrial Ca²⁺ efflux and peroxynitrite generation

Based on the indispensable role of hsp90β1 in PIMCE [17], it is expected that hsp90β1 would be essential for LA-induced peroxynitrite generation if the peroxynitrite response of LA indeed depends on [Ca²⁺]_m efflux in HM cells. We tested the role of hsp90β1 in the LA responses by downregulation of hsp90β1 expression with a well-known hsp90 chaperone inhibitor, 17-(dimethylaminoethylamino)-17-demethoxygeldanamycin (17-DMAG), and hsp90β1 RNAi. As shown in Figure 5A, treatment of HM cells with 17-DMAG (90 nM) or the hsp90β1 RNAi for 48 h effectively downregulated hsp90β1 expression. Of particular note, both treatments not only significantly inhibited LA-induced [Ca²⁺]_m efflux (as indicated by the alleviated [Ca²⁺]_i response) but also attenuated peroxynitrite generation (Figure. 5B). These results indicate that hsp90β1 is critical to LA-responsive [Ca²⁺]_m efflux and the coupled peroxynitrite responses.

Mitochondrial complexes I and III were responsible for LA-induced peroxynitrite generation

It has been known that in the mitochondria ROS are produced predominantly by complexes I and III [6]. We thus evaluated whether mitochondrial complexes I and III were responsible for the LA-induced peroxynitrite generation in HM cells. Pretreatment of HM cells with the complex I inhibitor rotenone (10 μM, 20 min) completely diminished peroxynitrite generation (Figure 6A and B; *P*<0.001, *n*=6) without significant effect on LA-induced [Ca²⁺]_i mobilization (*P*>0.05, *n*=6). Blocking complex III with either antimycin A (1 μM, 20 min) or myxothiazol (0.5 μM,

60 min) also did not alter the [Ca²⁺]_i responses (*P*>0.05, *n*=6) to LA but inhibited peroxynitrite generation (Figure 6A, C and D; *P*<0.001, *n*=6). These results indicate that the mitochondrial complexes I and III are responsible for LA-induced peroxynitrite generation.

LA-induced mitochondrial Ca²⁺ efflux and protein nitrotyrosylation were enhanced in diabetic mouse kidney

Functional and structural abnormalities were reported in the kidney of db/db mice, a type 2 diabetes model [21]. In mitochondria prepared from the kidneys of 12–16 wk old db/+ (littermate control) and db/db mice, LA caused Ca²⁺ efflux (Figure 7A and B). Compared to db/+ mice, the rate of LA-induced mitochondrial Ca²⁺ efflux was significantly enhanced




Figure 5. The effect of hsp90β1 on LA-induced [Ca²⁺]_i mobilization and peroxynitrite generation. HM cells were treated with vehicle (control), hsp90β1 RNAi, or 17-DMAG for 48 h. (A): Cell lysates were prepared and the alterations of hsp90β1 expression were analyzed by western blot analysis using α-tubulin as loading control. (B): The alteration of LA-induced [Ca²⁺]_i mobilization and peroxynitrite generation in hsp90β1 RNAi- and 17-DMAG-treated HM cells relative to control. The values represent the mean ± SEM. **P*<0.05, ***P*<0.01, *n*=6. doi:10.1371/journal.pone.0006048.g005




Figure 6. Mitochondrial complexes I and III were required for LA-induced peroxynitrite generation. HM cells were labeled with fura-2 and DCF to measure LA responsive [Ca²⁺]_m mobilization (black traces) and peroxynitrite generation (red traces). (A): LA responses in untreated cells (control); (B) LA responses in cells pretreated with rotenone (a mitochondrial complex I inhibitor, 10 μM, 20 min); (C) and (D): LA responses in cells pretreated with the mitochondrial complex III blockers, antimycin A (1 μM, 20 min); C) or myxothiazol (0.5 μM, 60 min); D). doi:10.1371/journal.pone.0006048.g006

more than two folds in db/db mice (Figure 7B, $P < 0.01$, $n = 3$). Western blot analysis indicated that nitrotyrosylation in a protein with a MW about 22 kDa was significantly enhanced in db/db mice (Figure 7C and D, $P < 0.005$, $n = 4$). Analysis of the nitrotyrosine level in kidney sections with immunohistochemistry further confirmed that protein nitrotyrosylation was enhanced in db/db mice kidney (Figure 7E and F).

Discussion

A major role of [Ca²⁺]_m under physiological conditions is to stimulate oxidative phosphorylation by activation of multiple dehydrogenases [4] and ATP synthase [22] as well as regulation of other metabolic processes [6]. [Ca²⁺]_m may inhibit the generation of ROS from complexes I and III under normal conditions whereas its overload may promote ROS generation and apoptosis [6]. In our current work, we have demonstrated that LA (an 18:2 n-6 PUFA) induced Ca²⁺ efflux from mitochondria and caused peroxynitrite production in HM cells. The LA responsive mitochondrial Ca²⁺ efflux may diminish the inhibitory effect of Ca²⁺ on ROS production from complexes I or III and thus leads to enhanced peroxynitrite generation (Figures 1–4, 6 and 8). These results revealed a cause and effect relationship between LA-induced [Ca²⁺]_m efflux and peroxynitrite production.

Our observation uncovered a novel mechanism by which increased NEFA may contribute to the pathogenesis of diabetic nephropathy by increasing peroxynitrite generation. It has been shown that LA is a major NEFA component, and its concentration is significantly elevated in type 2 diabetic patients [23]. Elevated NEFA concentration has also been reported in the db/db mice and other diabetic animal models [24]. In the current work, we observed increased nitrotyrosylation in multiple proteins of LA-treated HM cells (Figure 2), demonstrating that the LA responsive peroxynitrite generation enhanced nitrosative stress. Increased nitrotyrosine levels were also observed in db/db mice along with enhanced LA responsive [Ca²⁺]_m efflux (Figure 7). Protein nitrotyrosylation may disrupt the normal functions of mesangial cells and the kidney, leading to abnormalities in the structure and/or function of diabetic kidney. Indeed, increased nitrotyrosine levels have been reported in the plasma and kidney of diabetic patients [25,26].

Peroxynitrite is a highly reactive oxidant and is formed by the reaction between superoxide and NO. It has been reported that the mitochondrial complexes I and III cause superoxide production [7]. Previous studies described the existence of mitochondrial nitric oxide synthase (NOS), which produced NO in an L-arginine dependent manner [27,28]. We have previously

reported the expression of endothelial NOS (eNOS) and inducible NOS (iNOS), but not neuronal NOS, in rat mesangial cells [29]. The eNOS and iNOS expressed in mesangial cells may be responsible for NO productions induced by G protein coupled receptor agonists and inflammatory stimulators, respectively [29,30]. The expression of iNOS in mesangial cells in response to inflammatory stimulators has been shown to be inhibited by conjugated-LA [30]. All three isoforms of NOS were detected in the kidney homogenates and the expression of NOS proteins was downregulated in aged ZSF1 fatty rats, which was associated with enhanced renal peroxynitrite and other structural and functional abnormalities [31]. The importance of NOS in the pathogenesis of diabetic nephropathy has clearly evidenced by studies from animal models and diabetic patients. However, in HM cells the LA-induced peroxynitrite generation was not inhibited by N^G-nitro-L-arginine (0.1–1 mM) or L-N^G-monomethyl arginine (0.5 mM) (data not shown), indicating that the LA responsive NO, a required substrate for peroxynitrite generation, was produced in an L-arginine independent manner. Mitochondrial complex III has been shown to possess nitrite reductase activity and is responsible for NO generation in the mitochondria by an L-arginine independent manner [32–34]. Thus the NO involved in LA responsive peroxynitrite generation was more likely to be produced from the nitrite conversion by mitochondrial complex III in HM cells (as depicted in Figure 8). The inhibition of LA responsive peroxynitrite generation by the complex I blocker (rotenone) or complex III blockers (antimycin A or myxothiazol) suggested that complexes I and III may be responsible for the generation of either superoxide or NO, which are the two primary substrates required to form peroxynitrite.

Our previous works had indicated that hsp90β1 played an important role in PIMCE [17]. In the current study, we showed that LA-induced [Ca²⁺]_m efflux and peroxynitrite generation were inhibited by 17-DMAG and, more specifically, hsp90β1 RNAi, indicating the hsp90β1 involvement (Figure 5). Based on the observations on the essential role of hsp90β1 in PIMCE, it is likely that hsp90β1 is involved in LA-induced peroxynitrite generation via regulation of [Ca²⁺]_m efflux.

In addition to diabetes, other pathological conditions have been shown to cause cellular NEFA accumulations. For example, excessive NEFA have been shown to accumulate in the kidney proximal tubules and cardiovascular tissues during ischemia-reperfusion [35,36] and was responsible for sustained mitochondrial damage. The responses demonstrated in the current work (i.e. activation of [Ca²⁺]_m efflux and generation of peroxynitrite, Figure 1–4, and 6) may partially explain the deleterious effects of NEFA under a variety of disease conditions.




Figure 7. LA-induced mitochondrial Ca²⁺ efflux and protein nitrotyrosylation were enhanced in the kidney of db/db mice. Mitochondria were prepared from fresh kidney tissues of 12–16 wk old db/+ and db/db mice and labeled with X-rhod-1 (2 μ M). (A): LA-induced mitochondrial Ca²⁺ efflux in the kidney of db/+ and db/db mice. (B): The ratio of the initial rate of LA-induced mitochondrial Ca²⁺ efflux (V_{LA}; measured in the first 60 s following LA) to the basal rate (V₀; measured prior to LA). The values represent mean \pm SEM of V_{LA}/V₀, **P<0.01, n=3. (C) and (D): Nitrotyrosine levels in the kidney homogenates of four pairs of db/+ and db/db mice were analyzed by western blot using α -tubulin as loading control. The arrowhead highlights a protein band (~22 kDa) with enhanced nitrotyrosine levels in db/db mice, **P<0.01, n=4. (E) and (F): Representatives of the nitrotyrosine levels in the kidney sections of db/+ and db/db mice as determined by immunohistochemistry. doi:10.1371/journal.pone.0006048.g007

In summary, our work demonstrated that LA-induced [Ca²⁺]_m efflux caused peroxynitrite generation and led to increased nitrotyrosine in proteins. The enhanced nitrosative injury in response to LA may disrupt mesangial cell function in the kidney and thus contribute to the pathogenesis of diabetic nephropathy.

Materials and Methods

Cell culture

Primary human mesangial (HM) cells (a gift of Dr. H. E. Abboud, Department of Medicine, University of Texas Health Science Center at San Antonio) were prepared and cultured as previously described [37].

Measurement of [Ca²⁺]_i and superoxide, NO, or peroxynitrite in HM cells

HM cells grown in 100 mm dishes were labeled with fura-2 (1 μ M) alone to measure [Ca²⁺]_i or together with DHE (1 μ M), DAF-2 (0.5 nM), or DCF (1 μ M) as needed at 37°C for 30 min in RPMI 1640 medium to measure [Ca²⁺]_i and superoxide, NO, or

peroxynitrite simultaneously. Loaded cells were harvested and resuspended in PBS1Ca buffer (containing 1 mM KH₂PO₄, 3 mM Na₂HPO₄, 154 mM NaCl, 1 mM CaCl₂, 1 mM MgSO₄, pH 7.4). [Ca²⁺]_i mobilization and superoxide, NO or peroxynitrite generation were monitored in a fluorometer (QM-6, Photon Technology International, NJ) using a cuvette with the temperature stabilized at room temperature. The ratio of fluorescence excited at 340 and 380 nm with emission of 510 nm was recorded and used to index the [Ca²⁺]_i change as previously reported [17]. With the multi dye mode of the QM-6 fluorometer, fluorescence excited at 473, 488, or 506 nm and emitted at 595, 515, or 529 nm was measured to index the generation of superoxide, NO, or peroxynitrite in HM cells. The cell density was 1–2 \times 10⁶ cells/ml in these experiments. The interference of LA to DHE, DAF-2, and DCF fluorescence was calibrated by subtracting the signal resulted from palmitic acid, which does not cause [Ca²⁺]_i mobilization in HM cells.

In experiments in which [Ca²⁺]_i mobilization was measured in the absence of extracellular Ca²⁺, the loaded cells were washed and measured in PBS0Ca buffer (containing 200 μ M EGTA instead of 1 mM CaCl₂ as in PBS1Ca).




Figure 8. Mechanism of excessive NEFA contributing to pathogenesis of diabetic nephropathy. NEFA, such as linoleic acid (LA, an 18:2 n-6 polyunsaturated fatty acid) induces Ca²⁺ efflux from mitochondria by activating a hsp90β1-dependent pathway (PIMCE). The LA responsive mitochondrial Ca²⁺ efflux diminishes the inhibitory effect of Ca²⁺ on superoxide production from complexes I or III and enhances nitrite conversion to NO by complex III, resulting in increased peroxynitrite formation and protein nitrotyrosylation. Protein nitrotyrosylation may disrupt the normal functions of mesangial cells and the kidney, leading to abnormalities in the structure and/or function of diabetic kidney. doi:10.1371/journal.pone.0006048.g008

Measurement of [Ca²⁺]_m and peroxynitrite in mitochondria

Mitochondria were prepared from HM cells or the kidney of 12–16 wk old db/+ or db/db mice as previously described [17] and were suspended in MB1 buffer (containing 250 mM mannitol, 75 mM succinic acid, 0.1 mM EDTA, 0.5 mM EGTA, 10 mM HEPES, pH 7.4). The samples were diluted with nine volumes of HKG buffer (20 mM NaCl, 100 mM KCl, 1 mM CaCl₂, 1 mM MgCl₂, 10 mM pyruvate, 20 mM HEPES, pH 7.4) [17], and labeled with X-rhod-1 (2 μM) and DCF (1 μM) at 37°C for 30 min. Measurement with loaded mitochondria were performed in PBS0Ca buffer. Alteration of [Ca²⁺]_m and peroxynitrite generation were measured with fluorescence excited at 578 or 506 nm and emitted at 602 or 529 nm in a PTI QM-6 fluorometer, respectively. In these experiments, mitochondrial preparations equal to 40 μg/ml proteins were used.

In experiments in which LA-induced [Ca²⁺]_m efflux and peroxynitrite generation were determined in the absence of Ca²⁺ refilling, the mitochondrial preparations were suspended in MB1 buffer and diluted with nine volumes of HKG-EGTA buffer (containing 200 μM EGTA instead of 1 mM CaCl₂ as in HKG), then labeled with X-rhod-1 and DCF and measured as described above.

Downregulation of hsp90β1

HM cells in 100 mm dishes were grown for 24 h in RPMI 1640 supplemented with 14% FBS prior to treatment with hsp90β1 RNAi or 17-DMAG. The three hsp90β1 RNAi sequences used were: 1). UUAGCAAGACGUGUUCGAUUCGAGU; 2). UUCUGCUGGACCCAGCCAUG AAGUA; 3). UCAAACU-GAGGCCAAGCAUUCUUUC (Invitrogen). Cells were treated with vehicle (Lipofectamine RNAi Max alone), a mixture of hsp90β1 RNAi (20 μM) and Lipofectamine-RNAi-Max, or 17-

DMAG (90 nM) as described previously [17]. The cells were cultured for an additional 48 h and the effects of these treatments on hsp90 β 1 expression were verified by western blot analysis as described [17].

The effects of 17-DMAG and hsp90 β 1 RNAi on LA-induced [Ca²⁺]_i mobilization and peroxynitrite generation in treated HM cells were assessed as described above.

Western blot analysis of nitrotyrosine

HM cells in 100 mm dishes were treated with LA (30 μ M, 60 min at 37°C), lysed in ice-cold RIPA buffer (containing Tris 50 mM, NaCl 150 mM, NP-40 1%, Na-deoxycholate 0.25%, EDTA 1 mM, PMSF 1 mM, aprotinin 1 μ g/ml, leupeptin 1 μ g/ml, pepstatin 1 μ g/ml, Na₃VO₄ 1 mM, NaF 1 mM, pH 7.4), and kept on ice for 60 min. The mixture was centrifuged at 12,000 \times g for 15 min at 4°C. Western blot analysis of the proteins in the supernatant was performed with mouse mAb to nitrotyrosine (1:1000, Upstate, Lake Placid, NY) and α -tubulin was used to control protein loading.

The kidneys of 12–16 wk old db/+ and db/db mice were collected, and cleaned slices (10–25 mg) of kidney tissues were homogenized in 1 ml RIPA buffer with a glass-glass homogenizer manually for 30 strokes. The homogenates were centrifuged at 150 \times g for 15 min at 4°C and the supernatants were collected. The nitrotyrosine levels were determined by western blot analysis.

Immunohistochemistry analysis of nitrotyrosine in kidney sections

The kidneys of 12–16 wk old db/+ and db/db mice were fixed in formalin, embedded in paraffin and sectioned (4 μ m). Paraffin

sections were deparaffinized with xylene and hydrated with a series of graded ethanol washes. Endogenous peroxidase was inactivated by incubation for 30 min in 0.3% H₂O₂. The mouse mAb to nitrotyrosine was added for 1 h at room temperature. Tissue sections were treated with VECTASTAIN[®] ABC-Peroxidase Kit (Vector Laboratories, Burlingame, CA). The DAB Substrate Kit (Vector Laboratories) was used as the enzyme substrate. Sections were counterstained with CAT Hematoxylin (Biocare Medical, Concord, CA). Procedures involving the use of mice were approved by the Institutional Animal Care and Use Committee, University of Texas Health Science Center at San Antonio (0707-001).

Statistics

Results are mean \pm SEM. Experimental data were analyzed using a two-tailed Student's t test for comparisons between two groups. A significant difference was defined at $P < 0.05$.

Acknowledgments

We thank Dr. Michael S. Katz for critical review of the manuscript. The permanent address for Dr. Hong-Mei Zhang: Department of Clinical Oncology, Xijing Hospital, the Fourth Military Medical University, Xi'an, China.

Author Contributions

Conceived and designed the experiments: BXZ. Performed the experiments: HMZ HD BXZ. Analyzed the data: HMZ HD CKY BXZ. Wrote the paper: HMZ HD CKY BXZ.

References

- Schönfeld P, Wojtczak L (2008) Fatty acids as modulators of the cellular production of reactive oxygen species. *Free Radic Biol Med* 45: 231–241.
- Chinen I, Shimabukuro M, Yamakawa K, Higa N, Matsuzaki T, Noguchi K, Ueda S, Sakanashi M, Takasu N (2007) Vascular lipotoxicity: endothelial dysfunction via fatty-acid-induced reactive oxygen species overproduction in obese Zucker diabetic fatty rats. *Endocrinology* 148: 160–165.
- Lassègue B, Clemens RE (2003) Vascular NAD(P)H oxidases: specific features, expression, and regulation. *Am J Physiol Regul Integr Comp Physiol* 285: R277–R297.
- Hansford R, Zorov D (1998) Role of mitochondrial calcium transport in the control of substrate oxidation. *Mol Cell Biochem* 184: 359–369.
- Jouaville LS, Pinton P, Bastianutto C, Rutter GA, Rizzuto R (1999) Regulation of mitochondrial ATP synthesis by calcium: evidence for a long-term metabolic priming. *Proc Natl Acad Sci USA* 96: 13807–13812.
- Brookes PS, Yoon Y, Robotham JL, Anders MW, Sheu SS (2004) Calcium, ATP, and ROS: a mitochondrial love-hate triangle. *Am J Physiol Cell Physiol* 287: C817–C833.
- Brownlee M (2005) The pathobiology of diabetic complications: a unifying mechanism. *Diabetes* 54: 1615–1625.
- Pacher P, Obrosova IG, Mabley JG, Szabó C (2005) Role of nitrosative stress and peroxynitrite in the pathogenesis of diabetic complications. Emerging new therapeutic strategies. *Curr Med Chem* 12: 267–275.
- Ali TK, Matragoon S, Pillai BA, Liou GI, El-Remessy AB (2008) Peroxynitrite mediates retinal neurodegeneration by inhibiting nerve growth factor survival signaling in experimental and human diabetes. *Diabetes* 57: 889–898.
- Kanski J, Hong SJ, Schoneich C (2005) Proteomic analysis of protein nitration in aging skeletal muscle and identification of nitrotyrosine-containing sequences in vivo by nanoelectrospray ionization tandem mass spectrometry. *J Biol Chem* 280: 24261–24266.
- Drel VR, Pacher P, Stevens MJ, Obrosova IG (2006) Aldose reductase inhibition counteracts nitrosative stress and poly(ADP-ribose) polymerase activation in diabetic rat kidney and high-glucose-exposed human mesangial cells. *Free Radic Biol Med* 40: 1454–1465.
- Kanwar YS, Wada J, Sun L, Xie P, Wallner EI, Chen S, Chugh S, Danesh FR (2008) Diabetic nephropathy: mechanisms of renal disease progression. *Exp Biol Med* (Maywood) 233: 4–11.
- Manabe E, Handa O, Naito Y, Mizushima K, Akagiri S, Adachi S, Takagi T, Kokura S, Maoka T, Yoshikawa T (2008) Astaxanthin protects mesangial cells from hyperglycemia-induced oxidative signaling. *J Cell Biochem* 103: 1925–1937.
- Biden TJ, Robinson D, Cordery D, Hughes WE, Busch AK (2004) Chronic effects of fatty acids on pancreatic beta-cell function: new insights from functional genomics. *Diabetes* 53: S159–S165.
- Befroy DE, Petersen KF, Dufour S, Mason GF, de Graaf RA, Rothman DL, Shulman GI (2007) Impaired mitochondrial substrate oxidation in muscle of insulin-resistant offspring of type 2 diabetic patients. *Diabetes* 56: 1376–1381.
- Zhang BX, Ma X, Zhang W, Yeh CK, Lin A, Luo J, Sprague EA, Swerdlow RH, Katz MS (2006) Polyunsaturated fatty acids mobilize intracellular Ca²⁺ in NT2 human teratocarcinoma cells by causing release of Ca²⁺ from mitochondria. *Am J Physiol Cell Physiol* 290: C1321–C1333.
- Zhang H, Li ZH, Zhang MQ, Katz MS, Zhang BX (2008) Heat shock protein 90 β 1 is essential for polyunsaturated fatty acid-induced mitochondrial Ca²⁺ efflux. *J Biol Chem* 283: 7580–7589.
- Possel H, Noack H, Augustin W, Keilhoff G, Wolf G (1997) 2, 7-Dihydrodichlorofluorescein diacetate as a fluorescent marker for peroxynitrite formation. *FEBS Lett* 416: 175–178.
- Duplain H, Sartori C, Dessen P, Jayet PY, Schwab M, Bloch J, Nicod P, Scherrer U (2008) Stimulation of peroxynitrite catalysis improves insulin sensitivity in high fat diet-fed mice. *J Physiol* 586: 4011–4016.
- Misko TP, Highkin MK, Veenhuizen AW, Manning PT, Stern MK, Currie MG, Salvemini D (1998) Characterization of the cytoprotective action of peroxynitrite decomposition catalysts. *J Biol Chem* 273: 15646–15653.
- HA TS, Barnes JL, Stewart JL, Ko CW, Miner JH, Abrahamson DR, Sanes JR, Kasinath BS (1999) Regulation of Renal Laminin in Mice with Type II Diabetes. *J Am Soc Nephrol* 10: 1931–1939.
- Das AM, Harris DA (1990) Control of mitochondrial ATP synthase in heart cells: inactive to active transitions caused by beating or positive inotropic agents. *Cardiovasc Res* 24: 411–417.
- Yi L, He J, Liang Y, Yuan D, Gao H, Zhou H (2007) Simultaneously quantitative measurement of comprehensive profiles of esterified and non-esterified fatty acid in plasma of type 2 diabetic patients. *Chem Phys Lipids* 150: 204–216.
- Zhou YP, Berggren P-O, Grill V (1996) A fatty-acid decrease in pyruvate dehydrogenase activity is an important determinant of beta-cell dysfunction in the obese diabetic db/db mouse. *Diabetes* 45: 580–586.
- Ceriello A, Mercuri F, Quagliaro L, Assaloni R, Motz E, Tonutti L, Taboga C (2001) Detection of nitrotyrosine in the diabetic plasma: evidence of oxidative stress. *Diabetologia* 44: 834–838.
- Thuraisingham RC, Nott CA, Dodd SM, Yaqoob MM (2000) Increased nitrotyrosine staining in kidneys from patients with diabetic nephropathy. *Kidney Int* 57: 1968–1972.

27. Ghafourifar P, Asbury ML, Joshi SS, Kincaid ED (2005) Determination of mitochondrial nitric oxide synthase activity. *Methods Enzymol* 396: 424–444.
28. Parihar MS, Nazarewicz RR, Kincaid E, Bringold U, Ghafourifar P (2008) Association of mitochondrial nitric oxide synthase activity with respiratory chain complex I. *Biochem Biophys Res Commun* 366: 23–28.
29. Zhang W, Meng H, Li ZH, Shu Z, Ma X, Zhang BX (2007) Regulation of STIM1, store-operated Ca²⁺ influx, and nitric oxide generation by retinoic acid in rat mesangial cells. *Am J Physiol Renal Physiol* 292: F1054–F1064.
30. Sheu JN, Lin TH, Lii CK, Chen CC, Chen HW, Liu KL (2006) Contribution of conjugated linoleic acid to the suppression of inducible nitric oxide synthase expression and transcription factor activation in stimulated mouse mesangial cells. *Food Chem Toxicol* 44: 409–416.
31. Prabhakar S, Starnes J, Shi S, Lonis B, Tran R (2007) Diabetic nephropathy is associated with oxidative stress and decreased renal nitric oxide production. *J Am Soc Nephrol* 18: 2945–2952.
32. Kozlov AV, Staniek K, Nohl H (1999) Nitrite reductase activity is a novel function of mammalian mitochondria. *FEBS Lett* 454: 127–130.
33. Castello PR, David PS, McClure T, Crook Z, Poyton RO (2006) Mitochondrial cytochrome oxidase produces nitric oxide under hypoxic conditions: implications for oxygen sensing and hypoxic signaling in eukaryotes. *Cell Metab* 3: 277–287.
34. Lacza Z, Pankotai E, Csordás A, Gero D, Kiss L, Horváth EM, Kollai M, Busija DW, Szabó C (2006) Mitochondrial NO and reactive nitrogen species production: does mtNOS exist? *Nitric Oxide* 14: 162–168.
35. Feldkamp T, Kribben A, Roeser NF, Senter RA, Weinberg JM (2006) Accumulation of nonesterified fatty acids causes the sustained energetic deficit in kidney proximal tubules after hypoxia-reoxygenation. *Am J Physiol Renal Physiol* 290: F465–F477.
36. Hendrickson SC, St Louis JD, Lowe JE, Abdel-aleem S (1997) Free fatty acid metabolism during myocardial ischemia and reperfusion. *Mol Cell Biochem* 166: 85–94.
37. Silver BJ, Jaffer FE, Abboud HE (1989) Platelet-derived growth factor synthesis in mesangial cells: Induction by multiple peptide mitogens. *Proc Natl Acad Sci USA* 86: 1056–1060.