



POSTER PRESENTATION

Open Access

HIV-1 capture and antigen presentation by dendritic cells: enhanced viral capture does not correlate with better T-Cell activation

M Rodriguez-Plata^{1*}, A Urrutia², S Cardinaud², M Buzon¹, N Izquierdo-Useros¹, JG Prado¹, M Puertas¹, I Erkizia¹, P Coulon², S Cedeño¹, B Clotet¹, A Moris³, J Martinez-Picado¹

From AIDS Vaccine 2012
Boston, MA, USA. 9-12 September 2012

Background

During HIV-1 infection, dendritic cells (DC) facilitate dissemination of HIV-1 while trying to trigger adaptive anti-viral immune responses. We examined whether increased HIV-1 capture in DC matured with lipopolysaccharide (LPS) results in more efficient antigen presentation to HIV-1-specific CD4⁺ and CD8⁺ T cells. In order to block the DC-mediated trans-infection of HIV-1 and maximize antigen loading, we also evaluated a non-infectious integrase-deficient HIV-1 isolate, the HIV_{NL4-3ΔIN}.

Methods

Immature DC (iDC), mature DC (mDC) activated with IL-1 β , TNF- α , IL-6, and PGE2 (ITIP) or LPS during viral uptake, and fully mDC matured with ITIP or with LPS for 48 h before viral loading were tested. Antigen presentation to HIV-1-specific CD4⁺ and CD8⁺ T cell clones was quantified by IFN- γ ELISPOT. DC-associated p24^{Gag} HIV-1 and DC-mediated HIV-1 trans-infection were also evaluated in parallel.

Results

We showed that higher viral capture of DC did not guarantee better antigen presentation or T-cell activation. Greater HIV_{NL4-3} uptake by fully LPS-matured DC resulted in higher viral transmission to target cells but poorer stimulation of HIV-1-specific CD4⁺ and CD8⁺ T cells. Conversely, maturation of DC with LPS during—but not before—viral loading enhanced both HLA-I and HLA-II HIV-1-derived antigen presentation. On the other hand, DC maturation with ITIP during viral uptake only

stimulated HIV-1-specific CD8⁺ T cells. Integrase-deficient HIV_{NL4-3ΔIN} was also efficiently captured and presented by DC through HLA-I and HLA-II pathways, but in absence of viral dissemination.

Conclusion

Hence, DC maturation state, activation stimulus, and time lag between DC maturation and antigen loading impact HIV-1 capture and virus antigen presentation. Our results demonstrate a dissociation between the capacity to capture HIV-1 and to present viral antigens. HIV_{NL4-3ΔIN} seems to be an attractive candidate to be explored. These results provide new insights into DC biology and have implications in the optimization of DC-based immunotherapy against HIV-1 infection.

Author details

¹AIDS Research Institute IrsiCaixa, Badalona, Spain. ²INSERM, UMRS-945, Infection and Immunity, Université Pierre et Marie Curie, Paris, France. ³INSERM, UMRS-945, Infection and Immunity, Univ. Pierre et Marie Curie, Paris, France.

Published: 13 September 2012

doi:10.1186/1742-4690-9-S2-P2

Cite this article as: Rodriguez-Plata et al.: HIV-1 capture and antigen presentation by dendritic cells: enhanced viral capture does not correlate with better T-Cell activation. *Retrovirology* 2012 **9**(Suppl 2):P2.

¹AIDS Research Institute IrsiCaixa, Badalona, Spain

Full list of author information is available at the end of the article