



The prevalence of self-injury in adolescence: a systematic review and meta-analysis

Bernadett Frida Farkas¹ · Zsófia K. Takacs^{2,3} · Nóra Kollárovics¹ · Judit Balázs^{4,5}

Received: 20 November 2022 / Accepted: 18 July 2023 / Published online: 24 July 2023
© The Author(s) 2023

Abstract

In the past 10 years, there has been a growing interest in self-injurious behavior (SIB) among adolescents. The lifetime prevalence of SIB is between 16 and 22% in community sample with females more likely to engage in SIB. There are conflicting results about the global distribution of the prevalence of SIB and whether the SIB has increased in the 21st century. Our aim in the current study was to conduct a systematic search of and meta-analysis on the prevalence of SIB in adolescents over the past 5 years' worth of published papers and to examine gender, continental, and year differences. We conducted a systematic search in June 2020 of six databases (PubMed, Scopus, Web of Science, OVID Medline, PsycINFO, EBSCO) with three main search terms: “self-injurious behavior,” “prevalence,” and “adolescence.” Article inclusion criteria were (a) written in English; (b) published between January 1, 2015, and June 18, 2020; and (c) focused on a community sample. Titles and abstracts of the articles were screened first. Then, the relevant full texts were read, and those that met the inclusion criteria were collected. We used Comprehensive Meta-Analysis software was used to conduct the analyses. After the screening process 97, articles were included in the meta-analysis. The age of the samples ranged from 11.00 to 18.53 years. The overall average prevalence of nonsuicidal self-injury in the studies was 16%. There was a significant gender difference: females reported a higher prevalence than males (19.4% and 12.9%, respectively). A significantly higher prevalence was found among Asian articles than those from other continents (19.5% and 14.7%, respectively). The prevalence of SIB did not change significantly between 2013 and 2018. The current research draws attention to the high prevalence of SIB among adolescents, especially among females and those living in Asia. It is important to address this behavior, both in terms of prevention and intervention.

Keywords Adolescents · Self-injurious behavior · Prevalence · Systematic review · Meta-analysis

Abbreviations

DSH Deliberate self-harm
D-SIB Deliberate self-injurious behavior

DSM-5 Diagnostic and statistical manual of mental disorders 5th edition
NSSI Nonsuicidal self-injury
SH Self-harm
SHB Self-harm behavior
SIB Self-injurious behavior
SITB Self-injurious thoughts and behavior

✉ Bernadett Frida Farkas
farkas.bernadett@phd.semmelweis.hu;
drfarkasbernadettfrida@gmail.com

¹ Mental Health Sciences Doctoral School, Semmelweis University, Budapest, Hungary
² Faculty of Education and Psychology, Institute of Education, Eötvös Loránd University, Budapest, Hungary
³ School of Health in Social Sciences, University of Edinburgh, Edinburgh, UK
⁴ Department of Developmental and Clinical Child Psychology, Faculty of Education and Psychology, Institute of Psychology, Eötvös Loránd University, Budapest, Hungary
⁵ Oslo New University College, Oslo, Norway

Background

Self-injurious behavior (SIB) is a phenomenon whereby a person directly and deliberately damage themselves [1]. Various terms are used in the literature, such as, *nonsuicidal self-injury* (NSSI), *deliberate self-harm* (DSH), *self-cutting*, and *self-harm* [2–6]. SIB can be considered as a class of behaviors on a spectrum from NSSI—in which the person has no intent to die—to suicidal attempt (SA), which

is a form of suicidal behavior [1]. Although NSSI and SA are two distinct behaviors, it is not always easy to decide whether there was a suicidal intent behind SIB, and in this case, it is difficult to separate them [1, 3]. Moreover, the comorbidity between NSSI and suicidal behaviors is approximately 50% in traditional and 70% in clinical populations [1, 7, 8]. Previous research has indicated that the higher risk of SA is associated with the following in regard to NSSI: greater frequency, more methods, and longer duration [1].

Several meta-analyses conducted in the past 20 years have focused on the prevalence of SIB [6, 9–12]; see Table 1).

Based on the previous meta-analyses, the prevalence of SIB shows a significant increase in the 1990s, but then a stagnation from 2005 [9, 10]. Gillies et al. (2018) found that the lifetime prevalence increased over time from 1990 to 2015, and Muehlenkamp et al. (2012) found no significant difference in the prevalence of NSSI and DSH between 2005 and 2011 [6, 10]. According to previous meta-analyses,

Table 1 Previous meta-analyses

Author	Year of publication	Number of articles	Age range	Continental differences	Main results
Muehlenkamp et al. [10]	2012	52	11–24	–	Lifetime prevalence: 18% NSSI, 16.1% DSH 12-months prevalence: 19% NSSI Average lifetime prevalence did not change between 2005 and 2011
Swannell et al. [9]	2014	34	Adults: 25< Young adults: 18–24 Adolescents: 10–17	No significant difference	1990–1999: 11.7% 2000–2005: 14.7% 2006–2012: 19.3% Pooled lifetime prevalence: 19.7% Females NSSI: 19.9% Males NSSI: 14.7%
Bresin and Schoenleber [11]	2015	116	11.55–55.5 (M=20.81)	–	Females NSSI: 33.78% Males NSSI: 26.36%
Gillies et al. [6]	2018	172	12–18 (M=12.81)	Difference due to methodological factors	Lifetime prevalence: 16.9% (DSH—11.4%, NSSI—22.9%) Significant increasing over time Girls were more likely to self-harm
Lim et al. [12]	2019	66	12.59–19.78 (mean age)	Lifetime prevalence of NSSI: higher in non-Western countries (32.6%) vs. Western (19.4%) Lifetime prevalence of DSH: higher in Western countries (14.2%) vs. non-Western (12.8%) Lifetime prevalence of NSSI was higher among low- and middle-income countries vs. developed Lifetime prevalence of NSSI was highest in Australia (30.9%) lowest in Europe (18.4%) Lifetime prevalence of DSH was highest in Asia (17.4%) and lowest in North America (7.3%)	Lifetime prevalence: 22.1% NSSI, 13.7% DSH 12-months prevalence: 19.5% NSSI, 14.2% DSH

NSSI nonsuicidal self-injury, DSH deliberate self-harm, M mean age

between 1990 and 2015, the average lifetime prevalence of SIB among adolescents is between 16.9 and 19.7% [6, 9–12].

There are conflicting results about the gender differences in the prevalence of SIB [6, 9–12]. Some research has indicated that females have a lifetime prevalence of SIB that is two to three times higher than males [11], whereas other studies have found that the gender difference does not appear at all or, if it does, to a lesser extent [9, 10].

As shown in Table 1 the results are conflicting in regard to whether there is a difference in the prevalence of NSSI and DSH [6, 10, 12]. Muehlenkamp et al. (2012) did not find any significant difference between the two phenomena, however Gillies et al. (2018) and Lim et al. (2019) found a significantly higher prevalence of NSSI than DSH [6, 10, 12]. The lowest lifetime prevalence of NSSI was 18%, while the highest was 22.9% [6, 9–12]. At the same time, the lowest lifetime prevalence of DSH was 11.4%, while the highest was 16.1% [6, 9–12]. As mentioned above, unlike NSSI, DSH can be suicidal and non-suicidal as well, but it must be a non-fatal self-harm [6, 12].

There are also conflicting results about the continental differences in the prevalence of SIB. While Swannell et al. (2014) didn't find any significant continental differences in the prevalence of SIB, Gillies et al. (2018) did: Sweden had the highest, Norway had the lowest prevalence of self-harm in the meta-analysis [6, 9]. Lim et al. found that the lifetime prevalence of both NSSI and DSH were higher in non-Western countries than Western countries [12].

The above described previous meta-analyses highlight that these conflicting results can be due to the large differences among the included studies in methodological factors (e.g., sampling method, measurement, mean age of the sample), in the definitions of SIB (e.g., with or without suicidal intent), or in the place of data collection [6, 9–12].

Aims

Because of the heretofore conflicting results, our aim in this study was to follow up previous meta-analyses on the prevalence of SIB in adolescent community samples [6, 11]. We focused on data published between 2015 and 2020.

Our first hypothesis was that the prevalence of SIB did not change over time between the examined period (2015 and 2020) for both females and males. Our second hypothesis was that females reported a higher prevalence of a history of SIB than males.

Before addressing these hypotheses, we investigated the following questions: (a) What kinds of definitions of SIB are used? (b) What kinds of assessments are used to measure SIB? (c) What was the sampling method? (d) Where were the data of the included studies collected? (e) What was the mean age of the sample? (f) Was suicidal intent excluded from the definition of SIB?

Method

Literature search

We conducted a systematic literature search on June 18, 2020. We used six computerized literature databases: PubMed, Scopus, Web of Science, OVID Medline, PsycINFO, and EBSCO Discovery Service for Semmelweis University. Search terms were the following: “non-suicidal” OR nonsuicidal OR “self-injur*” OR “self-harm” OR “self harm” OR parasuic* OR “self-mutilation” AND prevalence AND adolescen* NOT “clinical trial” OR “case report” OR review. Two filters were added: (a) date between January 1, 2015, and June 18, 2020, and (b) English language. We used End-Note X9 software to remove duplicates and screen the 374 search hits.

Inclusion and exclusion criteria

To be included, studies had to report on the prevalence of SIB in adolescence in a community sample in a published article written in English. We used “adolescen*” among our search terms, and we included studies with an age range 11–18 years. However, in some articles this age range was wider (22 years being the oldest), so from these we included in the final analysis only those studies where the mean age of the sample was adolescence. For details, see Table 2.

When multiple studies reported on the same database, we included the ones with the largest sample size [13], the ones that provided data separately for males and females [14], and the ones that provided follow-up results [15, 16]. This led to the exclusion of six studies [17–22]. In addition, we contacted by email the authors of articles from which prevalence data could not be extracted. In case we did not receive sufficient statistics, we excluded the study (e.g., Carvalho et al., 2015). The methodology of this review follows the PRISMA guidelines [23].

Data extraction

Two authors (BFF, NK) coded the following information:

- (1) bibliographic information: authors, year of publication and data collection;
- (2) sample information: age range and mean age of sample, gender ratio, country, and continent the sample was recruited in, representativeness of the community sample, design;
- (3) measurement of SIB: measurement instrument, suicidal intent, terminology;
- (4) information for effect size: prevalence estimate and sample size.

Table 2 The included relevant studies

Author (year)	Country	Year of data collection	Mean age (SD)	Sample size (% female)	Sampling	Terminology	Suicidal intent dysfunction	Measurement	Prevalence
Aldrich et al. (2018) [28]	USA	2013–2015	12.86 (0.85)	121 (55.4%)	C	SITB	No	Questionnaires validated for other than SIB	Lifetime: 18.20% overall 6-months: 16.1% female, 12% male
Badoud et al. (2015) [29]	Switzerland	–	15.72 (1.74)	130 (50.8%)	C	NSSI	Yes	Questionnaires validated for other than SIB	1-year: 25.4% overall 24.24% female 26.6% male
Baetens et al. (2015) [30]	Belgium	2011	16.07 (1.12)	358 (48%)	C	NSSI	Yes	Questionnaires validated for other than SIB	1-year: 9.78%; lifetime: 13.41%
Barrocas et al. (2015) [31]	China	–	16.02 (0.61)	617 (51.4%)	C	NSSI	Yes	Not validate	Lifetime: 23.8% overall 26% male 21.7% female
Bhola et al. (2017) [32]	India	2013	17.5 (14.2)	1571 (57.8%)	RA	NSSI	Yes	Validate for SIB	1-year: 33.8%
Brausch and Woods (2019) [33]	USA	–	13.19 (1.19)	436 (52.7%)	C	NSSI	Yes	Validate for SIB	6-months: 17.2% overall 14.62% male 18.97% female
Buelens et al. (2020) [34]	Belgium	2018	15 (1.81)	2130 (54%)	C	NSSI/NSSI-D	Yes	Single-item question	Lifetime: 21.8% → 7.6% met the NSSI-D diagnosis 29.9% female, 12% male; NSSI-D diagnosis: 11.7% female, 2.9% male
Calvete et al. (2015) [35]	Spain	2010	15.32 (1.97)	1864 (51.45%)	RA	NSSI	Yes	Validate for SIB	1-year: 55.6% overall 58% female 53.3% male
Carvalho et al. (2017) [36]	Portugal	2012	16.75 (1.31)	1763 (52.9%)	C	NSSI	Yes	Validate for SIB	Lifetime: 29.5% overall 29.4% male 29.7% female
Cassels et al. (2018) [37]	UK	2005–2008	–	1238 (54.5%)	C	NSSI	Yes	Questionnaires validated for other than SIB	Lifetime: 11.89%
Castro and Kirchner (2017) [38]	Chile	–	14.98 (1.69)	965 (57%)	C	NSSI	Yes	Validate for SIB	Lifetime: 49.6% overall 41.39% male 55.45% female
Chen and Chun (2019) [39]	Taiwan	2011–2012	15.23 (1.24)	438 (100%)	C	NSSI	Yes	Validate for SIB	1-year: 36.8%
Çimen et al. (2017) [40]	Turkey	2014	15.00 (1.13)	555 (56.6%)	C	NSSI	Yes	Validate for SIB	Lifetime prevalence: 11.4%

Table 2 (continued)

Author (year)	Country	Year of data collection	Mean age (SD)	Sample size (% female)	Sampling	Terminology	Suicidal intent dis-tinction	Measurement	Prevalence
Claes et al. (2015) [41]	Belgium, Netherlands	2013	15.56 (1.32)	785 (44.5%)	C	NSSI	Yes	Validate for SIB	Lifetime prevalence: 20.1%
Copeland et al. (2019) [42]	USA	2009–2010	–	5870 (56.6%)	C	Self-cutting	No	Single-item question	1 year: 7% overall 4.67% male 11.35% female
Donath et al. (2019) [43]	Germany	2015	14.91 (0.73)	10,638 (49.8%)	RE	D-SIB	No	Single-item question	1 year: 17.8% overall 5.1% male 29.7% female
Doyle et al. (2015) [16]	Ireland	1999–2000	M = 16 years, SD = 0.715	856 (48.8%)	RE	SH	No	Validate for SIB	Lifetime: 12.1% overall 18.1% female 6.4% male
Duarte et al. (2019) [44]	Portugal	2017–2018	Study 1: 16.1 (1.8) Study 2: 15.4 (1.8)	Study 1: 620 (67.9%) Study 2: 411 (67.9%)	C	DSH	No	Validate for SIB	Lifetime: 21.1% in study 1, 26.5% in study 2
Emerson et al. (2019) [45]	Australia	2000–2001	14	9845 (47.6%)	RE	SH	No	Single-item question	1 year: 14.9%
Emery et al. (2017) [46]	Canada	–	13.38 (0.51)	639 (53%)	C	NSSI	Yes	Validate for SIB	Lifetime: 18% overall 22.7% female 13% male
Endo et al. (2017) [47]	Japan	2008–2009	15.2 (1.7)	17,347 (50.2%)	C	SH	No	Single-item question	1 year: 3.8%
Esposito et al. (2019) [48]	Italy	2016	15.60 (1.65)	640 (60.5%)	C	NSSI	Yes	Not validate	6-months: 15.3% overall 14.6% male 15.8% female
Farhat et al. (2020) [49]	USA	2000	–	2234 (43.9%)	C	SIB	Yes	Single-item question	Lifetime: 18% overall 24% female, 13.7% male
Fraser et al. (2018) [50]	New Zealand	2012–2015	15.16 (2.61)	1799 (56.5%)	C	NSSI	Yes	Validate for SIB	Lifetime: 20.6% overall 28.7% female 9.9% male
Gandhi et al. (2015) [51]	Belgium	2014	16.13 (1.47)	568 (61.8%)	C	NSSI	Yes	Validate for SIB	Lifetime: 16.5% overall 12.8% female 3.9% male
Gandhi et al. (2017) [52]	Belgium	2015–2016	15.0 (1.84)	528 (50.4%)	C	NSSI	Yes	Single-item question	Lifetime: 14.2% overall 20.8% female 7.7% male
Gandhi et al. (2018) [53]	Belgium	2012–2013	–	3880 (51%)	C	NSSI	Yes	Single-item question	Lifetime: 21% overall 26% female 17% male

Table 2 (continued)

Author (year)	Country	Year of data collection	Mean age (SD)	Sample size (% female)	Sampling	Terminology	Suicidal intent distinction	Measurement	Prevalence
Gandhi et al. (2018) [54]	Belgium	2015	16.6 (0.96)	401 (51.5%)	C	NSSI	Yes	Single-item question	Lifetime: 16.5% overall 20.77% female 11.85% male
Gandhi et al. (2019) [55]	Belgium	2015–2017	15.0 (1.85)	528 (50.4%)	C	NSSI	Yes	Single-item question	1 year: 7.55%
Garisch and Wilson (2015) [56]	New Zealand	2008–2009	16.35 (0.62)	1162 (43%)	RE	NSSI	Yes	Validate for SIB	Lifetime: 48.7% overall 49.4% female 48% male
Gaspar et al. (2019) [57]	Portugal	2014	14.8 (1.2)	3262 (54%)	RE	NSSI	No	Single-item question	1 year: 20.3% overall females 23.7% males 16.3%
Geulayov et al. (2018) [58]	UK	2015	–	5520 (51%)	RE	Non-fatal SH	No	Not validate	1 year: 5.83% overall 8.9% females 2.6% males
Gromatsky et al. (2017) [59]	USA	2013–2014	14.39 (0.63)	550 (100%)	C	NSSI	Yes	Validate for SIB	Lifetime: 7.82%
Guerreiro et al. (2015) [60]	Portugal	2009–2011	15.6 (1.7)	1713 (55.6%)	C	SH	No	Validate for SIB	Lifetime: 7.3% overall 10.5% females 3.3% males
Hamada et al. (2016) [61]	Japan	2011	13.9 (0.2)	1840 (51.4%)	C	Self-cutting	No	Single-item question	Lifetime: 8.9% overall 5.6% males 11.9% females
Han et al. (2018) [62]	China	2013	14.81	5726 (50.3%)	RA	SH	Yes	Not validate	6 months: 45.3% overall 41.6% male 49% female
Hanania et al. (2015) [63]	Jordan	–	14.53 (1.71)	952 (49.8%)	C	NSSI	Yes	Validate for SIB	Lifetime: 22.6% overall males 26.98% females 18.14%
Heerde et al. (2015) [64]	USA, Australia	2002–2003	Washington State: 14.1 Grade 7, 15.1 Grade 9; Victorian: 13.9 Grade 7, 14.9 Grade 9	3876 (51%)	RE	DSH	No	Single-item question	1 year: 1.53% in Grade 7 and 0.91% in Grade 9 for males, 4.12% and 1.34% for Grade 7 and Grade 9 for females

Table 2 (continued)

Author (year)	Country	Year of data collection	Mean age (SD)	Sample size (% female)	Sampling	Terminology	Suicidal intent dis-tinction	Measurement	Prevalence
Horváth et al. (2018) [65]	Hungary	2009–2010, 2013	Vocational school sample: 15.21 (0.77) High-school sample: 15.09 (0.75)	Vocational school sample: 140 (40%); high-school sample: 995 (59.2%)	RE, RA	D-SIB	Yes	Validate for SIB	Lifetime: 29.4% in the vocational school group, 17.2% in the high school group Vocational school sample: 25.64% males, 35.41% females High-school sample: 14.4% males, 19% females
Horváth et al. (2020) [66]	Hungary	2015–2017	15.43 (1.14)	161 (50%)	C	NSSI	Yes	Validate for SIB	Lifetime: 23.6% overall 8.64% males 33.75% females
Huang et al. (2017) [15]	Taiwan	2008–2010	16.02 (0.52)	5879 (56.7%)	C	DSH	Yes	Single-item question	Lifetime: 25.04% overall 28.96% female 19.9% male
Jantzer et al. (2015) [67]	Germany	2012	12.8 (1.95)	647 (50.7%)	C	NSSI	Yes	Single-item question	1 year: 10.97%
Jiang et al. (2016) [68]	China	2013–2014	13.17 (1.10)	813 (43.4%)	C	NSSI	Yes	Single-item question	Lifetime: 29.0% overall 27.9% male 31.3% female
Kadziela-Olech et al. (2015) [69]	Poland	2013	16.7 (1.64)	2220 (46.3%)	C	D-SIB, NSSI	Yes	Validate for SIB	D-SIB lifetime: 8.3%; NSSI lifetime: 4.8% D-SIB lifetime: 6.7% females, 9.6% males; NSSI lifetime: 6.3% males, 3.0% females
Kaess et al. (2020) [70]	10 European countries + Israel	2009–2010	14.84 (0.9)	1933 (51.47%)	RE	D-SIB	Yes	Validate for SIB	Lifetime: 24.9%; 1-year: 6.7% overall 7.04% male 6.43% female
Kang et al. (2018) [71]	China	–	15.63 (1.67)	3555 (52.0%)	RA	NSSI	Yes	Validate for SIB	6-months: 13.8% overall 16.6% female 10.4% male
Kelada et al. (2016) [72]	Australia	2014	14.49 (1.38)	117 (56.4%)	C	Self-injury	Yes	Single-item question	Lifetime: 19.7%
Kiekens et al. (2015) [73]	Netherlands, Belgium	2012	15.52 (1.34)	946 (44%)	RA	NSSI	Yes	Validate for SIB	Lifetime: 24.31% overall 24.46% male 24.26% female

Table 2 (continued)

Author (year)	Country	Year of data collection	Mean age (SD)	Sample size (% female)	Sampling	Terminology	Suicidal intent dis-tinction	Measurement	Prevalence
Kitagawa et al. (2017) [74]	Japan	2008–2009	–	18,018 (50.3%)	C	SH	Yes	Single-item question	1 year: 7.3%
Klemera et al. (2016) [75]	UK	2013–2014	15	1519 (48.8%)	RE	SH	No	Single-item question	Lifetime: 21.5% overall 31.9% females 11.4% males
Koenig et al. (2016) [76]	Germany	2010–2012	14.7	506 (52.1%)	RE	D-SIB	Yes	Validate for SIB	1-year: 8.30% overall lifetime: 30.7% male 47.45% female
Latina and Stattin (2017) [77]	Sweden	2008, 2010	13.89 (0.75)	2029 (50%)	RE	SH	Yes	Validate for SIB	6 months: 31%
Law and Shek (2016) [78]	China	–	12.53 (0.66)	2023 (52%)	RA	SH	Yes	Not validate	Lifetime: 15.3% overall 13.9% male 16.5% female
Lee (2016) [79]	South Korea	–	14.38 (1.68)	784 (48.8%)	C	SH	No	Validate for SIB	Lifetime: 12.4%
Li et al. (2019) [80]	China	2015–2016	15.36 (1.79)	22,628 (51.4%)	RE	NSSI	No	Not validate	12 months: 32.1% overall male 35.2% female 29.1%
Lin et al. (2017) [81]	Taiwan	2013	15.83 (0.38)	2170 (51.5%)	C	NSSI	Yes	Single-item question	1 year: 20.1% overall female 23.8% male 16.9%
Liu et al. (2017) [82]	Taiwan	2008–2009	15.44 (0.61)	2479 (60.3%)	C	SH	Yes	Single-item question	1 year: 10.1% overall female 11.24% male 8.32%
Liu et al. (2018) [83]	China	2015	14.97 (1.46)	11,831 (49.1%)	RE	NSSI	Yes	Validate for SIB	Lifetime: 23.7%; 1-year: overall 18.9% male 17.8% female 19.9%
Luyckx et al. (2015) [84]	Belgium	–	15.95 (1.30)	348 (100%)	C	NSSI	Yes	Validate for SIB	Lifetime: 20.7%
Lütke et al. (2017) [85]	Switzerland	2010	14.95 (0.74)	447 (48%)	C	NSSI	Yes	Validate for SIB	1 year: 5.15% male 13.08% female
Madjar et al. (2019) [86]	Israel	–	14.96 (1.33)	594 (45.6%)	C	NSSI	Yes	Validate for SIB	1-year: 19.5% male, 10.7% female
Mars et al. (2019) [87]	UK	–	16.8 (2.9)	4795 (73%)	RE	NSSI	Yes	Single-item question	Lifetime: 11.73% overall 5.7% male, 15.92% female

Table 2 (continued)

Author (year)	Country	Year of data collection	Mean age (SD)	Sample size (% female)	Sampling	Terminology	Suicidal intent dysfunction	Measurement	Prevalence
Martinez-Ferrer and Statin (2019) [88]	Sweden	–	13.94 (0.74)	987 (48.3%)	C	SH	Yes	Validate for SIB	6 months: 36%
McManus et al. (2020) [89]	UK	2000, 2007, 2014	–	2000: 103 2014: 122	RE	NSSH	Yes	Single-item question	Lifetime: 6.1% in 2000, 10.9% in 2014
Monto et al. (2018) [90]	USA	2015	–	64 671 (52.5%)	RE	NSSI	Yes	Single-item question	1-year: 23.8% female, 11.3% male
Morey et al. (2017) [91]	UK	2013	–	2000 (13–15: 54.2%; 16–18: 50.2%)	RE	SH	No	Single-item question	Lifetime: overall 15.5% females 23.1%, males 7.1%
Nguyen et al. (2020) [92]	Vietnam	2018	11	648 (47.7%)	RA	SH	No	Single-item question	Lifetime: 7.1%
Okutan (2017) [93]	Turkey	2016	17.02 (1.59)	263 (54.3%)	C	SHB	Yes	Validate for SIB	Lifetime: 44.86% overall 39.16% females, 51.67% males
Pawłowska et al. (2015) [94]	Poland	–	16.92 (1.15)	6883 (69%)	C	Self-injury	No	Not validate	Lifetime: 24.91% overall 16.24% females, 8.67% males
Pawłowska et al. (2016) [95]	Poland	–	16.91 (1.11)	5685 (30%)	C	Self-injury	No	Not validate	Lifetime: 14% overall 6.92% males, 15.74% females
Peng et al. (2019) [96]	China	2016	13.6 (1.1)	2647 (51.2%)	RE	SH	Yes	Single-item question	6-months: 1.4% females, 1.3% males
Pisinger et al. (2018) [97]	Denmark	2014	17.9 (1.5)	66,284 (62%)	RE	SH	No	Single-item question	Lifetime: 20% overall 24% females, 12% males
Plener et al. (2015) [98]	Germany	–	14.85 (0.58)	452 (46.2%)	C	NSSI	Yes	Validate for SIB	Lifetime: 20.4% overall 29.76% females, 12.97% males
Plener et al. (2016) [99]	Germany	2014	15.91	91 (57.1%)	RE	NSSI	Yes	Validate for SIB	Lifetime: 26.9% overall 1.92% males 12.82% females
Quarshie et al. (2020) [100]	Ghana	2017	16.8 (1.38)	444 (51.8%)	RE	SH	No	Single-item question	Lifetime prevalence: 23.8% males, 30% females; 1-year prevalence: 24.8% females, 19.2% males
Reigstad and Kvernmo (2017) [101]	Norway	2003–2005	–	4881 (50.1%)	C	DSH	No	Single-item question	1-year: 22.3% overall 28.8% females, 15.9% males

Table 2 (continued)

Author (year)	Country	Year of data collection	Mean age (SD)	Sample size (% female)	Sampling	Terminology	Suicidal intent dis-tinction	Measurement	Prevalence
Ren et al. (2018) [102]	Taiwan	–	15.45 (0.54)	1989 (52.0%)	RE	NSSI	Yes	Validate for SIB	1 year: 20.8% overall 24.4% females, 16.8% males
Schwartz-Mette and Lawrence (2019) [103]	USA	2016–2018	15.68 (1.49)	186 (69.9%)	C	NSSI	Yes	Single-item question	1 year: 27.4% overall 21.43% males, 30% females
Sigurdson et al. (2018) [104]	Norway	1998, 1999–2000	BL: 13.7 (0.58); FU: 14.9 (0.6)	BL: 2464 (50.8%); FU1: 2432 (50.4%)	RE	SH	No	Single-item question	Lifetime: BL—2.48% males, 7.19% females; FU—4.89% males, 11.58% females
Simioni et al. (2017) [105]	Brazil	2010–2011	–	2508 (47.2%)	RE	DSH	No	Diagnostic interview	Lifetime: 1.5%
Solis-Bravo et al. (2019) [106]	Mexico	2016	12.3 (1.3)	438 (57.2%)	C	NSSI	Yes	Validate for SIB	Lifetime: 11.5%
Somer et al. (2015) [107]	Turkey	2010–2011	16.8 (1.26)	1656 (55%)	RE	NSSI	Yes	Validate for SIB	Lifetime: 31.3% overall 33% female, 29.4% male
Stanford et al. (2017) [108]	Australia	2014–2015	14.9 (1.6)	1521 (56.4%)	C	SH	No	Single-item question	6-months: 16.8% overall 12.1% male, 20.5% female
Sutin et al. (2018) [109]	Australia	2014	14.4 (0.49)	2948 (48.3%)	RE	SH	No	Single-item question	1-year: 8.8% overall 3.68% males, 14.52% females
Tang et al. (2016) [110]	China	2013–2014	14.7 (1.9)	4405 (49.67%)	RA	NSSI	Yes	Validate for SIB	1-year: 29.2% overall 30.9% females, 27.4% males
Tang et al. (2018) [111]	China	2014–2015	15.2 (1.8)	15,623 (48.5%)	RE	NSSI	Yes	Validate for SIB	1-year: 29% overall 27.94% males, 30.50% females
Tanner et al. (2016) [112]	Australia	2010	14.20 (1.03)	2637 (58.8%)	C	NSSI	Yes	Validate for SIB	Lifetime: 7.2% males, 11.93% females
Tilton-Weaver et al. (2019) [113]	Sweden	2013–2014	13.65 (0.64)	2769 (47.3%)	C	NSSI	Yes	Validate for SIB	6-months: 5%
Tseng and Yang (2015) [114]	Taiwan	–	–	391 (54.73%)	C	NSSI	Yes	Diagnostic interview	1-year: 9.7% overall 18.7% females; 10.2% males
Victor et al. (2018) [13]	USA	2000–2014	13	2127	C	NSSI	Yes	Diagnostic interview	Lifetime: 3%;

Table 2 (continued)

Author (year)	Country	Year of data collection	Mean age (SD)	Sample size (% female)	Sampling	Terminology	Suicidal intent distinction	Measurement	Prevalence
Wan et al. (2015) [115]	China	2008	16.1 (2.8)	17,622 (51.2%)	C	NSSI	Yes	Single-item question	Lifetime: 17.0% overall 16.9% males, 17.1% females
Wan et al. (2019) [116]	China	2013–2014	15.44 (1.8)	14,820 (50.2%)	RE	NSSI	Yes	Single-item question	1-year: 26.1% overall 24.3% female, 27.9% male
Wan et al. (2020) [117]	China	2013–2014	15.59 (1.80)	9704 (52.60%)	C	NSSI	Yes	Single-item question	1-year: 38.54% overall 37.11% female, 40.13% male
Wang et al. (2016) [118]	China	–	14.63 (1.25)	5423 (52.6%)	C	NSSI	Yes	Single-item question	6-months: 18.3% overall 21.2% female 14.6% male
Zetterqvist (2016) [119]	Sweden	2011	16.56	3060 (50.5%)	RE	NSSI/NSSI-D	Yes	Validate for SIB	1-year: NSSI at least one episode: 35.1% overall 10.61% female 11.62% male NSSI-D: 6% overall 9.97% female 2.11% male
Zhang et al. (2016) [120]	China	2013–2014	15.18 (1.79)	25,378 (51.4%)	C	NSSI	Yes	Single-item question	Lifetime: 27.5% overall 28.6% male, 26.4% female
Zubrick et al. (2015) [121]	Australia	2013–2014	15.51 (1.75)	2653 (48.4%)	RE	SH	Yes	Single-item question	Lifetime: 10.9% overall 7.45% males, 17.68% females 1-year: 8% overall 4.6% males, 11.99% females

SD standard deviation, C convenience, RA randomized, RE representative, SIB self-injurious thoughts and behavior, SIB self-injurious behavior, NSSI nonsuicidal self-injury, NSSI-D nonsuicidal self-injury based on the Diagnostic and Statistical Manual of Mental Disorders 5th Edition criteria, D-SIB deliberate self-injurious behavior, SH self-harm, DSH deliberate self-harm, SHB self-harm behavior, BL baseline, FU follow-up

Interrater reliability ranged from 73 and 100%. In case a consensus could not be reached between the two coders, the other two authors were consulted (ZKT, JB).

To test our hypotheses, we preferred to include the prevalence estimates separately for males and females if a study reported on those. For longitudinal studies, prevalence at all measurement points was coded; however, they were averaged to calculate an effect size for a study before we included the data in any analyses. We made an exception when prevalence estimates were available separately for males and females at one time point but not at another. In those cases, we chose to include only the estimates at the time point when they were reported separately for males and females.

During the coding, we had to impute some scores that were not reported in the primary studies. For studies that reported only the age range, we imputed the mean age as the geometric mean of the range. For studies that did not report the year of data collection, we subtracted 2 years from the year of publication (for a similar procedure, see Protzko et al., 2020) [24].

Statistical analyses

We used the Comprehensive Meta-Analysis software to conduct the analyses [25, 26]. We applied a random effects model. When a study reported results at more than one time point, we entered all in the software, which takes the average between multiple time points before entering a study in the grand average. We made an exception when conducting meta-regression analyses regarding the year of data collection and the mean age of the sample. In these cases, we only selected the first time point from these longitudinal studies to be included. In contrast, we considered estimates for males and females when reported separately in a study as independent effect sizes in all the analyses. Outliers were inspected based on a standardized residual exceeding ± 3.29 . We inspected the results according to several moderator variables. When inspecting results according to the different continents and suicidal intent, we conducted a subgroup analysis to statistically contrast them. We only included subgroups with at least four effect sizes in this analysis (for a similar procedure see Takacs and Kassai 2019) [27].

Results

Included studies

In sum, a total of 97 articles were included in this meta-analysis; we identified 178 effect sizes (see Figs 1, 2 and Table 2).

There were six outlying effect sizes that we excluded. Altogether, we had data from 439,818 participants. The

overall average SIB prevalence in the studies was 16.0% (95% confidence interval [CI] [14.7, 17.4], $k = 172$). This was a heterogeneous effect, $Q(171) = 30,136.96$, $p < 0.001$, $I^2 = 99.43$ $\tau^2 = 0.44$.

In our assessment of publication bias, Egger's test showed significant asymmetry (intercept = -2.88 , $p = 0.046$), but the funnel plot showed a symmetric distribution based on visual inspection, which was confirmed by no imputed studies in the Duval and Tweedie's trim-and-fill procedure.

Among the 97 included articles, 74 reported prevalence data for females and males separately. There were 79 effect sizes reported for females. Two effect sizes were outliers and thus were excluded. We found an average prevalence of 19.4% for females, 95% CI [17.5, 21.4], $k = 77$. This effect was heterogeneous, $Q(76) = 8,660.74$, $p < 0.001$, $I^2 = 99.12$ $\tau^2 = 0.29$. There were 75 effect sizes reported for males. One outlying effect size was excluded. We found an average prevalence of 12.9%, 95% CI [11.3, 14.8], $k = 74$. Again, this was a heterogeneous effect, $Q(74) = 10,315.75$, $p < 0.001$, $I^2 = 99.2$ $\tau^2 = 0.43$.

Terms and definitions of SIB in the included studies

The terminology of SIB was not uniform across the included studies. All the studies defined SIB as a deliberate damage to oneself, but not all of them defined it as a nonsuicidal intent. Seventy-two articles (73.5%) made a clear distinction between suicidal and nonsuicidal intent.

There were 11 different terms for SIB in the included 97 papers. The most frequently used term was NSSI; this expression appeared in 60 articles (see Table 3).

Measurements of SIB in the included studies

Among the included studies we found diagnostic interviews, self-reported questionnaires, and single-item questions to measure SIB. Two studies measured NSSI based on DSM-5 criteria [122]. The most frequently used questionnaire was the Deliberate Self-Harm Inventory [123], which was mentioned in 13 articles. The Inventory of Statements About Self-Injury [124] was used in five studies, and the Functional Assessment of Self-Mutilation [125] also was used in five. Effect sizes based on a single item to assess SIB found an average prevalence of 11.6%, 95% CI [9.3, 14.5], $k = 31$. We found of 14.8%, 95% CI [12.8, 17.2], $k = 60$, in studies that used nonvalidated questionnaires. Questionnaires that had been validated for other constructs showed an average prevalence of 14.7%, 95% CI [9.8, 21.5], $k = 6$. Finally, questionnaires that had been validated for SIB showed the highest average percentage: 18.9%, 95% CI [16.9, 21.1], $k = 77$. For results separately for males and females, see the Supplementary Materials. Only one study used a diagnostic

interview and reported on two effect sizes. The average of these showed a similar estimate as the grand average (14.2% (95% CI [7.7, 24.8], $k=2$), more specifically, 18.7% for females and 10.2% for males).

Sampling

Of the 172 effect sizes, 99 were based on convenience sampling. These showed an average prevalence of 15.2%, 95% CI [13.4, 17.2]. Eighteen effect sizes were based on samples that applied randomization, showing a prevalence of 24.7%, 95% CI [18.9, 31.6]. For 55 sample sizes, the sample was representative of the population. Representative samples showed a pooled prevalence of 15.1%, 95% CI [13.2, 17.4]. A similar pattern was noted for females and males (see the Supplementary Materials).

Place of data collection of the included studies

From the 98 included articles, we found three collaborations in which data were collected in multiple countries; for the rest, the data were collected in single countries. When we inspected the results over all the effect sizes, we noted differences according to the continent on which the data had been collected. There were three effect sizes in two publications from South America that showed an average prevalence of 33%, 95% CI [13.7, 60.3], and we found two effect sizes in one publication from Africa that showed an average prevalence of 24.4%, 95% CI [19.1, 30.7], and data for two effect sizes published in the same article were collected in North America and Australia and Oceania as part of an international cooperation that showed a prevalence of 2.6%, 95% CI [1.3, 4.9]. These categories were excluded from the subgroup analysis as they contained less than 4 effect sizes. After we excluded these, we noted a significant difference between the prevalence estimates from the different continents (see Table 4), $Q(3)=10.97$, $p=0.012$. More specifically, prevalence estimates from Asia (19.5%, 95% CI [17.1, 22.2], $k=51$) were significantly larger than those from the other three continents (14.6%, 95% CI [13.1, 16.2], $k=114$), $Q(1)=11.20$, $p=0.001$. As shown in Table 4, the effect of continent was similar when we inspected effect sizes for female and male samples separately.

Mean age of the included samples

For assessing the effects of the mean age of the samples, we chose to focus on the first measurement point in the 17 longitudinal studies. In this analysis, seven outliers appeared that were then excluded. For an additional six effect sizes we could not extract the sample's age, and thus those were also excluded from this analysis. This resulted in 165 effect sizes. The mean age of the sample ranged from 11.00 to





Fig. 1 The selection process is summarized in the QUORUM flow-chart

18.53 years. The mean age of the sample did not have a significant effect on the effect size (coefficient = 0.067, $p=0.12$). For results separately for males and females, see the Supplementary Materials.

To make sure that longitudinal studies from which we chose to include the first estimate in this analysis did not influence the results by possibly reporting on substantially younger samples, we also ran the regression model on the cross-sectional studies only as a sensitivity analysis. This resulted in 134 effect sizes to be included. Again, the mean age of the sample did not have a significant effect on these prevalence estimates (coefficient = 0.058, $p=0.24$).

Suicidal intent

For 125 effect sizes, suicidal intent was excluded. Those showed a pooled estimate of 18.3%, 95% CI [16.7, 19.9]. This was significantly higher than what was found in studies that did not exclude suicidal intent (11.3%, 95% CI [9.3, 13.7], $k=47$), $Q(1)=20.52$, $p<0.001$. This pattern was also confirmed in only-female and only-male samples. For results

Fig. 2 The flowchart of inclusion and exclusion criteria

separately for males and females, see the Supplementary Materials.

Prevalence of SIB

We found 92 effect sizes reporting on lifetime prevalence of SIB, 72 effect sizes that estimated 1-year prevalence, and 17 that estimated 6-month prevalence. An average of 17.9%, 95% CI [16.3, 19.5], was found overall when lifetime prevalence was assessed. This estimate was 22.9 (95% CI [20.9, 25.0], $k=42$) for females and 13.7% (95% CI [11.2, 16.8], $k=39$) for males.

An overall average prevalence of 13.4%, 95% CI [11.5, 15.6] was found when assessing prevalence in the last year. This estimate was 15.9% (95% CI [12.9, 19.4], $k=32$) for females and 10.7% (95% CI [8.7, 13.2], $k=32$) for males. An overall prevalence of 16.2%, 95% CI [11.0, 23.3] was estimated when we considered only the last 6 months, 18% (95% CI [8.9, 33.2], $k=7$) for females and 13.8% (95% CI [6.2, 27.9], $k=7$) for males.

Year of data collection

In regard to assessing the effects of the year of data collection, we chose to focus on the first measurement point in the 17

Table 3 Prevalence differences in the terms of SIB

	Prevalence estimates (95% CI)		
	Overall	Only female samples	Only male samples
DSH	15.1% (11.2–20.2) (k = 8)	11.5% (6.2–20.4) (k = 4)	6.3% (3.2–11.8) (k = 4)
Deliberate self-injurious behavior (D-SIB)	16.2% (9.7–25.8) (k = 12)	20.1% (10.8–34.4) (k = 6)	12.8% (7.3–21.5) (k = 6)
Non-fatal self-harm	4.9% (1.5–15.3) (k = 2)	8.9% (7.9–10) (k = 1)	2.7% (2.1–3.3) (k = 1)
NSSI	18.4% (16.9–20) (k = 103)	20.8% (18.2–23.7) (k = 47)	17.1% (15.1–19.3) (k = 44)
NSSI based on the Diagnostic and Statistical Manual of Mental Disorders 5th Edition (DSM-5) criteria	5.3% (2.6–10.6) (k = 4)	10.6% (8.8–12.8) (k = 2)	2.5% (1.8–3.4) (k = 2)
Self-cutting	7.4% (4.8–11.2) (k = 4)	10.4% (8.2–13.2) (k = 2)	4.9% (4–6) (k = 2)
Self-injury	12.4% (8.2–18.4) (k = 5)	16.1% (15.2–17.1) (k = 2)	7.7% (6.2–9.6) (k = 2)
Self-harm (SH)	12.7% (10–16) (k = 34)	18.2% (13.5–24.1) (k = 13)	9% (5.5–14.4) (k = 13)
Self-harm behavior (SHB)	45.2% (33.4–57.6) (k = 2)	39.2% (31.5–47.4) (k = 1)	51.7% (42.8–60.5) (k = 1)
SIB	18.3% (10.2–30.6) (k = 2)	24% (21.4–26.8) (k = 1)	13.7% (11.9–15.7) (k = 1)
Self-injurious thoughts and behavior (SITB)	14.4% (9–22.2) (k = 2)	16.1% (8.9–27.4) (k = 1)	12% (5.5–24.2) (k = 1)

longitudinal studies. In this analysis, seven outliers appeared that were then excluded. Data for the primary studies were collected between 1998 and 2018. The year of data collection had a significant, positive effect on the 171 effect sizes (coefficient = 0.035, $p = 0.008$); that is, more recent studies found larger prevalence. For results presented separately for females and males, see the Supplementary Materials.

For further investigation, we restricted the year of data collection to 2013 and onward so we could assess the effect in the time constraints that corresponds to the time constraints of year of publication of the present meta-analysis (2015 and onward). Data for 119 effect sizes were collected in or after 2013. When we considered only these studies, the effect of year of data collection was not significant on the effect sizes (coefficient = -0.015 , $p = 0.72$). The same was found for the 53 effect sizes for females (coefficient = -0.005 , $p = 0.92$) and for the 51 effect sizes for males (coefficient = -0.05 , $p = 0.49$). Scatterplots are shown in the Supplementary Materials.

Risk of bias

Risk-of-bias criteria was based on the Cochrane Risk of Bias Tool [126], adapted for the studies (cohort, cross-sectional, and longitudinal).

Discussion

Because previous meta-analyses have yielded conflicting results on the prevalence of SIB in community adolescent samples [6, 11], we found it important to complete a follow-up meta-analyses with clear methodology on recently published data. In the present meta-analysis, we found that the prevalence of SIB in adolescents was 16% in studies published between 2015 and 2020. This result is comparable to the estimate of 16.9% found in a previous meta-analysis [6]. Regarding methodological differences, as can be expected, a slightly higher estimate was found when considering lifetime prevalence (17.9%) as compared with the 1-year (13.4%) or 6-month prevalence (16.2%). We also noted a significantly higher prevalence when suicidal intent was excluded (18.3%) than when it was not excluded (11.3%), and the largest prevalence was found when measurement instruments were used that had been validated for self-injurious behaviors (18.8%). In addition, methodologically more rigorous studies that focused on representative samples found an average SIB prevalence of 15.1%. This is an interesting issue, while self-harm without suicidal intent should be a subgroup of self-harm covering forms both with and without suicidal intent. Hence the first number should always be lower than the second number. A possible explanation could be that the

Table 4 Prevalence differences in continental distribution

	Prevalence estimates (95% CI)		
	Overall	Only female samples	Only male samples
Asia	19.5% (17.1–22.2) (k = 51)	22.3% (19.4–25.4) (k = 23)	19.5% (16.6–22.8) (k = 22)
Australia and Oceania	14.1% (9.5–20.5) (k = 13)	18.5% (11.8–27.8) (k = 6)	10.0% (4.3–21.5) (k = 6)
Europe	14.7% (12.9–16.8) (k = 87)	19.5% (17.0–22.4) (k = 37)	10.8% (8.5–13.79) (k = 36)
North America	13.8% (10.2–18.4) (k = 14)	14.3% (10.5–19.2) (k = 7)	11.5% (5.9–21.1) (k = 6)

studies used the same term but actually employ different criteria. In addition, differences in the prevalence of NSSI and DSH may also result from measurement differences between the two types of SIB. Previous meta-analyses have reported higher prevalence rates for multi-item instruments [6, 10], and 65.5% of NSSI measurements consisted of multiple items, compared to 60% of DSH measurements consisting of a single item. In a meta-analysis made by Swannell et al. (2014), checklist versus single-item measurement explained the 41% of variance between studies [9]. Our review shows that, among adolescents, there are no significant changes in the prevalence between ages 11.0 and 18.5 years. This result is comparable to Lim's meta-analysis [12] but does not align with Gillies and colleagues' (2018) study. We found similar prevalence estimates among studies that used convenience and representative samples. However, and surprisingly, studies that used a random sample found larger estimates. This is puzzling and needs further research.

Our first hypothesis was only partially confirmed. When we considered all data that were published between 2015 and 2018, we found that there was a significant increase between 1998 and 2018 in the prevalence of SIB. However, when we restricted our analysis to the time frame between 2013 and 2018 (to reflect the publication time window of 2015 and 2018), we found no change in prevalence, as we had expected. Previous meta-analyses have found mixed results regarding this question. Muehlenkamp and colleagues (2012) did not find any significant difference in the prevalence of SIB between 2008 and 2015, whereas Gillies and colleagues (2018) found an increase between 1990 and 2018. Our results are in line with both previous findings in that they show an increase before 2013, but no change since then. This finding can be important to both decision makers and professionals for the appropriate planning of prevention programs.

Our second hypothesis was confirmed; we found a substantial difference between the estimates for females (19.4%) and males (12.9%), with nonoverlapping confidence intervals. A similar pattern was observed when we considered only lifetime prevalence, with 22.9% for females and 13.7% for males. These results are comparable to Bresin and Schoenleber's (2015) meta-analysis, in which the prevalence was significantly higher among females. Studies that excluded suicidal intent found an average of 21% for females and 16.5% for males. Similarly, estimates based on measurement instruments that were validated for SIB showed 21.9% for females and 15.7% for males. On the other hand, studies with representative samples showed slightly lower estimates: 18.2% for females and 10.9% for males. Thus, the patterns were very similar for females and males when the effect of methodological differences in the primary studies were assessed. However, the cultural difference between countries in Asia and those on other continents was more articulated

for males (Asia = 19.5% vs. other = 10.8%) than for females (Asia = 22.3% vs. other = 18.5%). Nock and Prinstein (2005) found that NSSI often is connected to psychological distress [127], and adolescent girls usually have more psychological distress than men [128]. These results highlight that it is necessary to pay more careful attention to NSSI by female adolescents and that perhaps further attention should be given to Asian male populations.

We found some differences in the prevalence estimate as a function of methodological differences among the primary studies; however, we should note that moderators might be confounded.

The pooled estimate from Asian countries (19.5%) was significantly higher than that from other continents (14.6%). Again, this confirms earlier meta-analytic results estimating a relatively large prevalence in Asian countries (Lim et al. 2019). This difference was even more articulated for males. So, it may be that the differences in SIB between Asian and non-Asian countries are somehow connected to gender. To understand this result, further research should focus on the transcultural aspects of SIB.

We did not find a difference between NSSI (18.7%) and DSH (15.1%), unlike Gillies's results [6], but we found a substantial difference between NSSI (18.7%) and self-harm (12.7%). In contrast to females, we found a substantial difference between NSSI (17.1%) and DSH (6.3%), and between NSSI (17.1%) and self-harm (9%) among males.

Our review highlights that the highest prevalence rates were found when SIB was measured with a validated questionnaire as compared with studies that used single-item or nonvalidated questionnaires, a pattern that was also confirmed separately for male and female samples. This result is likely due to the fact that validated questionnaires are more sensitive than single-item measures [6, 10].

Our results are limited by the heterogeneity of the primary studies, that is, in regard to the sample and the measurement instruments and the conceptualization of SIBs. The findings of the present meta-analysis confirm that these differences among the primary studies have an important effect on the prevalence estimates. There is currently no consensus in the literature about the conceptualization of SIB [6, 10], which makes our work more difficult when evaluating the data. However, to provide the most precise estimate, we pooled the studies that used representative samples that reported on lifetime prevalence of SIBs excluding suicidal intent measured by a validated measurement instrument and found similar estimates. In addition, we did this to avoid a confound effect of these moderators. Moreover, although overall we found a relatively large number of studies that reported on prevalence of SIBs, it is questionable whether nonsignificant results in subgroup and meta-regression analyses are truly due to an absence of an effect or whether they are instead due to a lack of statistical power.

To our best knowledge, this is the most recent meta-analysis on the prevalence of SIB among adolescents. An overall prevalence of 16% was found, which means that one in six adolescents has a history of self-harm. Moreover, a larger estimate was found for females as compared with males: every fifth adolescent girl reported having conducted self-harm. It is interesting that estimates were largest in Asian countries with males, approaching a 20% prevalence. Further research should focus on the transcultural aspects of self-harm to understand this difference. All these results have public health importance in drawing the attention of clinicians and decision makers to adolescents who engage in SIB. Clinicians need to be aware of the high prevalence and risk factors (e.g., female gender, Asian populations) of SIB in adolescence. Prevention and intervention are very important in this age group.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00787-023-02264-y>.

Author contributions BFF made the literature search, coded the articles, made the analyses, wrote the main manuscript text and prepared the figures. ZKT supervised and reviewed the statistical analysis of the manuscript. NK made the consensus coding of the articles and the risk of bias. JB participated in the design of the study, coordinated the steps, reviewed the manuscript and the figures. All authors read and approved the final manuscript. All authors contributed equally to this work.

Funding Open access funding provided by Semmelweis University.

Data availability Not applicable.

Declarations

Conflict of interest The authors declare that they have no competing interests.

Ethical approval Not applicable.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Nock M et al (2006) Non-suicidal self-injury among adolescents: diagnostic correlates and relation to suicide attempts. *Psychiatry Res* 144:65–72
- Klonsky ED, Oltmanns TF, Turkheimer E (2003) Deliberate self-harm in a nonclinical population: prevalence and psychological correlates. *Am J Psychiatry* 160(8):1501–1508
- Klonsky ED (2007) The functions of deliberate self-injury: a review of the evidence. *Clin Psychol Rev* 27(2):226–239
- Nock M (2010) Self-injury. *Annu Rev Clin Psychol* 6(1):339–363
- Whitlock J, Eckenrode J, Silverman D (2006) Self-injurious behaviors in a college population. *Pediatrics* 117(6):1939–1948
- Gillies D et al (2018) Prevalence and characteristics of self-harm in adolescents: meta-analyses of community-based studies 1990–2015. *J Am Acad Child Adolesc Psychiatry* 57(10):733–741
- Muehlenkamp JJ, Gutierrez PM (2007) Risk for suicide attempts among adolescents who engage in non-suicidal self-injury. *Arch Suicide Res* 11(1):69–82
- Klonsky ED, Muehlenkamp JJ (2007) Self-injury: a research review for the practitioner. *J Clin Psychol* 63(11):1045–1056
- Swannell S et al (2014) Prevalence of nonsuicidal self-injury in nonclinical samples: systematic review, meta-analysis and meta-regression. *Suicide Life Threat Behav* 44: 273–303
- Muehlenkamp JJ et al (2012) International prevalence of adolescent non-suicidal self-injury and deliberate self-harm. *Child Adolesc Psychiatry Ment Health* 6(1):10
- Bresin K, Schoenleber M (2015) Gender differences in the prevalence of nonsuicidal self-injury: a meta-analysis. *Clin Psychol Rev* 38:55–64
- Lim KS et al (2019) Global lifetime and 12-month prevalence of suicidal behavior, deliberate self-harm and non-suicidal self-injury in children and adolescents between 1989 and 2018: a meta-analysis. *Int J Environ Res Public Health* 16(22): 4581
- Victor SE et al (2019) Parent and peer relationships as longitudinal predictors of adolescent non-suicidal self-injury onset. *Child Adolesc Psychiatry Ment Health* 13:1
- Gandhi A et al (2017) Age of onset of non-suicidal self-injury in Dutch-speaking adolescents and emerging adults: an event history analysis of pooled data. *Compr Psychiatry* 80:170–178
- Huang YH et al (2017) Relationship between predictors of incident deliberate self-harm and suicide attempts among adolescents. *J Adolesc Health* 60(5):612–618
- Doyle L (2015) Self-harm in young people: prevalence, associated factors and help-seeking in school-going adolescents. *Int J Ment Health Nurs* 24: 485–494
- Gandhi A et al (2016) Non-suicidal self-injury and adolescents attachment with peers and mother: the mediating role of identity synthesis and confusion. *J Child Fam Stud* 25(6):1735–1745
- Gandhi A et al (2016) Sociotropy, autonomy, and non-suicidal self-injury: the mediating role of identity confusion. *Personality Individ Differ* 99:272–277
- Gandhi A et al (2016) Reactive and regulative temperament and non-suicidal self-injury in Flemish adolescents: the intervening role of identity formation. *Personality Individ Differ* 99:254–259
- Doyle L, Sheridan A, Treacy MP (2017) Motivations for adolescent self-harm and the implications for mental health nurses. *J Psychiatr Ment Health Nurs* 24(2–3):134–142
- Huang YH et al (2017) Correlation of impulsivity with self-harm and suicidal attempt: a community study of adolescents in Taiwan. *BMJ Open* 7(12): e017949
- Scott LN et al (2015) Non-suicidal self-injury and suicidal ideation as predictors of suicide attempts in adolescent girls: a multi-wave prospective study. *Compr Psychiatry* 58:1–10
- Page MJ et al (2021) The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 372:n71
- Protzko J (2020) Kids these days! Increasing delay of gratification ability over the past 50 years in children. *Intelligence* 80:101451

25. Borenstein M, Larry H, Higgins J, Rothstein H (2015) Manual to the comprehensive meta-analysis version 3 software. Biostat, Englewood, NJ
26. Borenstein M, Larry H, Higgins J, Rothstein H (2014) Comprehensive meta-analysis version 3. Biostat, Englewood, NJ
27. Takacs ZK, Kassai R (2019) The efficacy of different interventions to foster children's executive function skills: a series of meta-analyses. *Psychol Bull* 145(7):653–697
28. Aldrich JT, Wielgus MD, Mezulis AH (2018) Low physiological arousal and high impulsivity as predictors of self-injurious thoughts and behaviors among adolescents. *J Adolesc* 62:55–60
29. Badoud D et al (2015) The French version of the reflective functioning questionnaire: validity data for adolescents and adults and its association with non-suicidal self-injury. *PLoS ONE* 10(12): e0145892
30. Baetens I et al (2015) The relationship between parental expressed emotions and non-suicidal self-injury: the mediating roles of self-criticism and depression. *J Child Fam Stud* 24(2):491–498
31. Barrocas AL et al (2015) Nonsuicidal self-injury in adolescence: longitudinal course, trajectories, and intrapersonal predictors. *J Abnorm Child Psychol* 43(2):369–380
32. Bhola P et al (2017) Predictors of non-suicidal and suicidal self-injurious behaviours, among adolescents and young adults in urban India. *Asian J Psychiatr* 29:123–128
33. Brausch AM, Woods SE (2019) Emotion regulation deficits and nonsuicidal self-injury prospectively predict suicide ideation in adolescents. *Suicide Life Threat Behav* 49(3):868–880
34. Buelens T et al (2020) Investigating the DSM-5 criteria for non-suicidal self-injury disorder in a community sample of adolescents. *J Affect Disord* 260:314–322
35. Calvete E et al (2015) Prevalence and functions of non-suicidal self-injury in Spanish adolescents. *Psicothema* 27(3):223–228
36. Carvalho BC et al (2017) Biting myself so I don't bite the dust: prevalence and predictors of deliberate self-harm and suicide ideation in Azorean youths. *Braz J Psychiatry* 39(3): 252–262
37. Cassels M et al (2018) Poor family functioning mediates the link between childhood adversity and adolescent nonsuicidal self-injury. *J Child Psychol Psychiatry* 59(8):881–887
38. Castro K, Kirchner T (2018) Coping and psychopathological profile in nonsuicidal self-injurious Chilean adolescents. *J Clin Psychol* 74(1):147–160
39. Chen WL, Chun CC (2019) Association between emotion dysregulation and distinct groups of non-suicidal self-injury in Taiwanese female adolescents. *Int J Environ Res Public Health* 16(18): 3361
40. Cimen ID, Coskun A, Etiler N (2017) Non-suicidal self-injury behaviors' features and relationship with adolescents' daily life activities and mental status. *Turk J Pediatr* 59(2):113–121
41. Claes L et al (2015) Bullying and victimization, depressive mood, and non-suicidal self-injury in adolescents: the moderating role of parental support. *J Child Fam Stud* 24(11):3363–3371
42. Copeland M et al (2019) Social ties cut both ways: self-harm and adolescent peer networks. *J Youth Adolesc* 48(8):1506–1518
43. Donath C et al (2019) Epidemiology of suicidal ideation, suicide attempts, and direct self-injurious behavior in adolescents with a migration background: a representative study. *BMC Pediatr* 19(1):45
44. Duarte E, Gouveia-Pereira M, Gomes HS (2019) Development and factorial validation of the inventory of deliberate self-harm behaviours for Portuguese adolescents. *Psychiatr Q* 90(4):761–776
45. Emerson E et al (2019) Emotional difficulties and self-harm among British adolescents with and without disabilities: cross sectional study. *Disabil Health J* 12(4):581–587
46. Emery AA, Heath NL, Rogers M (2017) Parents' role in early adolescent self-injury: an application of self-determination theory. *Sch Psychol Q* 32(2):199–211
47. Endo K et al (2017) Preference for solitude, social isolation, suicidal ideation, and self-harm in adolescents. *J Adolesc Health* 61(2):187–191
48. Esposito C, Bacchini D, Affuso G (2019) Adolescent non-suicidal self-injury and its relationships with school bullying and peer rejection. *Psychiatry Res* 274:1–6
49. Farhat LC et al (2020) Self-injurious behavior and gambling-related attitudes, perceptions and behaviors in adolescents. *J Psychiatr Res* 124:77–84
50. Fraser G et al (2018) Non-suicidal self-injury, sexuality concerns, and emotion regulation among sexually diverse adolescents: a multiple mediation analysis. *Arch Suicide Res* 22(3):432–452
51. Gandhi A et al (2015) Non-suicidal self-injury and identity distress in Flemish adolescents: exploring gender differences and mediational pathways. *Personality Individ Differ* 82:215–220
52. Gandhi A et al (2017) Directionality of effects between non-suicidal self-injury and identity formation: a prospective study in adolescents. *Personality Individ Differ* 109:124–129
53. Gandhi A et al (2018) Age of onset of non-suicidal self-injury in Dutch-speaking adolescents and emerging adults: an event history analysis of pooled data. *Compr Psychiatry* 80:170–178
54. Gandhi A et al (2018) Association between non-suicidal self-injury, parents and peers related loneliness, and attitude towards aloneness in Flemish adolescents: an empirical note. *Psychologica Belgica* 58(1):3–12
55. Gandhi A et al (2019) Maternal and peer attachment, identity formation, and non-suicidal self-injury: a longitudinal mediation study. *Child Adolesc Psychiatry Ment Health* 13: 7
56. Garisch J, Wilson M (2015) Prevalence, correlates, and prospective predictors of non-suicidal self-injury among New Zealand adolescents: cross-sectional and longitudinal survey data. *Child Adolesc Psychiatry Mental Health* 9: 28
57. Gaspar S et al (2019) Non-suicidal self-injuries and adolescents high risk behaviours: highlights from the Portuguese HBSC study. *Child Indic Res* 12(6):2137–2149
58. Geulayov G et al (2018) Incidence of suicide, hospital-presenting non-fatal self-harm, and community-occurring non-fatal self-harm in adolescents in England (the iceberg model of self-harm): a retrospective study. *Lancet Psychiatry* 5(2):167–174
59. Gromatsky MA et al (2017) The role of parental psychopathology and personality in adolescent non-suicidal self-injury. *J Psychiatr Res* 85:15–23
60. Guerreiro DF et al (2017) Self-harm in adolescents: a self-report survey in schools from Lisbon. *Portugal Arch Suicide Res* 21(1):83–99
61. Hamada S et al (2018) Association between bullying behavior, perceived school safety, and self-cutting: a Japanese population-based school survey. *Child Adolesc Mental Health* 23(3):141–147
62. Han A et al (2018) A self-harm series and its relationship with childhood adversity among adolescents in mainland China: a cross-sectional study. *BMC Psychiatry* 18(1):28
63. Hanania JW et al (2015) Non-suicidal self-injury among adolescents in Amman. *Jordan Arch Suicide Res* 19(2):260–274
64. Heerde JA et al (2015) Incidence and course of adolescent deliberate self-harm in Victoria, Australia, and Washington state. *J Adolesc Health* 57(5):537–544
65. Horváth LO et al (2018) Direct self-injurious behavior (D-SIB) and life events among vocational school and high school students. *Int J Environ Res Public Health* 15(6): 1068

66. Horvath LO et al (2020) Nonsuicidal self-injury and suicide: the role of life events in clinical and non-clinical populations of adolescents. *Front Psychiatry* 11
67. Jantzer V et al (2015) Does parental monitoring moderate the relationship between bullying and adolescent nonsuicidal self-injury and suicidal behavior? A community-based self-report study of adolescents in Germany. *BMC Public Health* 15
68. Jiang YQ et al (2016) Buffering the effects of peer victimization on adolescent non-suicidal self-injury: the role of self-compassion and family cohesion. *J Adolesc* 53:107–115
69. Kadziela-Olech H et al (2015) The prevalence of non-suicidal self-injury (NSSI) among high school students in relation to age and sex. *Psychiatr Pol* 49(4):765–778
70. Kaess M et al (2020) Life events predicting the first onset of adolescent direct self-injurious behavior—a prospective multicenter study. *J Adolesc Health* 66(2):195–201
71. Kang N et al (2018) Distress intolerance mediates the relationship between child maltreatment and nonsuicidal self-injury among chinese adolescents: a three-wave longitudinal study. *J Youth Adolesc* 47: 2220
72. Kelada L, Hasking P, Melvin G (2016) The relationship between nonsuicidal self-injury and family functioning: adolescent and parent perspectives. *J Marital Fam Ther* 42(3):536–549
73. Kiekens G et al (2015) Non-suicidal self-injury among Dutch and Belgian adolescents: personality, stress and coping. *Eur Psychiatry* 30(6):743–749
74. Kitagawa Y et al (2017) Appetite loss as a potential predictor of suicidal ideation and self-harm in adolescents: a school-based study. *Appetite* 111:7–11
75. Klemmera E et al (2017) Self-harm in adolescence: protective health assets in the family, school and community. *Int J Public Health* 62(6):631–638
76. Koenig J et al (2016) Prospective risk for suicidal thoughts and behaviour in adolescents with onset, maintenance or cessation of direct self-injurious behaviour. *Eur Child Adolesc Psychiatry* 26(3):345–354
77. Latina D, Stattin H (2017) Adolescents who self-harm: the patterns in their interpersonal and psychosocial difficulties. *J Res Adolesc* 28(4):824–838
78. Law BF, Shek DTL (2016) A 6-year longitudinal study of self-harm and suicidal behaviors among Chinese adolescents in Hong Kong. *J Pediatr Adolesc Gynecol* 29(1):S38–S48
79. Lee WK (2016) Psychological characteristics of self-harming behavior in Korean adolescents. *Asian J Psychiatr* 23:119–124
80. Li D et al (2019) Interaction of health literacy and problematic mobile phone use and their impact on non-suicidal self-injury among Chinese adolescents. *Int J Environ Res Public Health* 16(13): 2366
81. Lin M-P et al (2017) Prevalence of nonsuicidal self-injury and its risk and protective factors among adolescents in Taiwan. *Psychiatry Res* 255:119–127
82. Liu HC et al (2017) Self-harm and its association with internet addiction and internet exposure to suicidal thought in adolescents. *J Formos Med Assoc* 116(3):153–160
83. Liu X (2018) Understanding the mental distress that leads to NSSI among Chinese adolescents in Hong Kong: a three-wave longitudinal study. 2018: Dissertation abstracts international: section B: the sciences and engineering, vol 79(8-B(E))
84. Luyckx K et al (2015) Non-suicidal self-injury in female adolescents and psychiatric patients: a replication and extension of the role of identity formation. *Personality Individ Differ* 77:91–96
85. Lüdtke J et al (2017) The influence of personality traits and emotional and behavioral problems on repetitive nonsuicidal self-injury in a school sample. *Compr Psychiatry* 74:214–223
86. Madjar N et al (2019) Exploring particular facets of cognitive emotion regulation and their relationships with nonsuicidal self-injury among adolescents. *Crisis J Crisis Intervent Suicide Prevent* 40(4):280–286
87. Mars B et al (2019) Predictors of future suicide attempt among adolescents with suicidal thoughts or non-suicidal self-harm: a population-based birth cohort study. *Lancet Psychiatry* 6(4):327–337
88. Martinez-Ferrer B, Stattin H (2019) Self-harm, depressive mood, and belonging to a subculture in adolescence. *J Adolesc* 76:12–19
89. McManus S, Gunnell D (2020) Trends in mental health, non-suicidal self-harm and suicide attempts in 16–24-year old students and non-students in England, 2000–2014. *Soc Psychiatry Psychiatr Epidemiol* 55(1):125–128
90. Monto MA, McRee N, Deryck FS (2018) Nonsuicidal self-injury among a representative sample of US adolescents, 2015. *Am J Public Health* 108(8):1042–1048
91. Morey Y et al (2017) Adolescent self-harm in the community: an update on prevalence using a self-report survey of adolescents aged 13–18 in England. *J Public Health* 39(1):58–64
92. Nguyen HTL et al (2020) Relationships among cyberbullying, parental attitudes, self-harm and suicidal behavior among adolescents: results from a school-based survey in Vietnam. *BMC Public Health* 20(1)
93. Oktan V (2017) Self-harm behaviour in adolescents: body image and self-esteem. *J Psychol Couns Sch* 27(2):177–189
94. Pawłowska B et al (2015) Prevalence of self-injury, suicidal ideation, plans and attempts in adolescents aged 13 to 19 years of age. *Polish J Public Health* 125(1):55–59
95. Pawłowska B et al (2016) Prevalence of self-injury performed by adolescents aged 16–19 years. *Psychiatr Pol* 50(1):29–42
96. Peng Z et al (2019) Associations between Chinese adolescents subjected to traditional and cyber bullying and suicidal ideation, self-harm and suicide attempts. *BMC Psychiatry* 19(1):324
97. Pisinger VSC, Hawton K, Tolstrup JS (2019) School- and class-level variation in self-harm, suicide ideation and suicide attempts in Danish high schools. *Scand J Public Health* 47(2):146–156
98. Plener PL et al (2015) Immigration as risk factor for non-suicidal self-injury and suicide attempts in adolescents in Germany. *Child Adolesc Psychiatry Mental Health* 9
99. Plener PL et al (2016) The prevalence of nonsuicidal self-injury (NSSI) in a representative sample of the German population. *BMC Psychiatry* 16(1):353
100. Quarshie ENB, Waterman MG, House AO (2020) Prevalence of self-harm among lesbian, gay, bisexual, and transgender adolescents: a comparison of personal and social adversity with a heterosexual sample in Ghana. *BMC Res Notes* 13(1):271
101. Reigstad B, Kvernmo S (2017) Concurrent adversities and deliberate self-harm among indigenous Sami and majority Norwegian adolescents: the Norwegian Arctic Adolescent Health Study. *Scand J Child Adolesc Psychiatry Psychol* 5(3):92–103
102. Ren YX et al (2018) The mediating role of coping strategy in the association between family functioning and nonsuicidal self-injury among Taiwanese adolescents. *J Clin Psychol* 74(7):1246–1257
103. Schwartz-Mette RA, Lawrence HR (2019) Peer socialization of non-suicidal self-injury in adolescents' close friendships. *J Abnorm Child Psychol* 47(11):1851–1862
104. Sigurdson JF et al (2018) The longitudinal association of being bullied and gender with suicide ideations, self-harm, and suicide attempts from adolescence to young adulthood: a cohort study. *Suicide Life Threat Behav* 48(2):169–182

105. Simioni AR et al (2017) Prevalence, clinical correlates and maternal psychopathology of deliberate self-harm in children and early adolescents: results from a large community study
106. Solis-Bravo MA et al (2019) Are tattoos an indicator of severity of non-suicidal self-injury behavior in adolescents? *Psychiatry Investig* 16(7):504–512
107. Somer O et al (2015) Prevalence of non-suicidal self-injury and distinct groups of self-injurers in a community sample of adolescents. *Soc Psychiatry Psychiatr Epidemiol* 7:1163
108. Stanford S, Jones MP, Hudson JL (2017) Rethinking pathology in adolescent self-harm: towards a more complex understanding of risk factors. *J Adolesc* 54:32–41
109. Sutin AR et al (2018) Perceived body discrimination and intentional self-harm and suicidal behavior in adolescence. *Child Obes* 14(8):528–536
110. Tang J et al (2016) Stressful life events as a predictor for non-suicidal self-injury in southern Chinese adolescence: a cross-sectional study. *Medicine (Baltimore)* 95(9):e2637
111. Tang J et al (2018) Prevalence of and risk factors for non-suicidal self-injury in rural China: results from a nationwide survey in China. *J Affect Disord* 226:188–195
112. Tanner A, Hasking P, Martin G (2016) Co-occurring non-suicidal self-injury and firesetting among at-risk adolescents: experiences of negative life events, mental health problems, substance use, and suicidality. *Arch Suicide Res* 20(2):233–249
113. Tilton-Weaver L, Marshall SK, Svensson Y (2019) Depressive symptoms and non-suicidal self-injury during adolescence: latent patterns of short-term stability and change. *J Adolesc* 75:163–174
114. Tseng FY, Yang HJ (2015) Internet use and web communication networks, sources of social support, and forms of suicidal and non-suicidal self-injury among adolescents: different patterns between genders. *Suicide Life Threat Behav* 45(2):178–191
115. Wan YH et al (2015) Longitudinal effects of psychological symptoms on non-suicidal self-injury: a difference between adolescents and young adults in China. *Soc Psychiatry Psychiatr Epidemiol* 50(2):237–247
116. Wan Y et al (2019) Associations of adverse childhood experiences and social support with self-injurious behaviour and suicidality in adolescents. *Br J Psychiatry* 214(3):146–152
117. Wan YH et al (2020) Associations of coping styles with non-suicidal self-injury in adolescents: do they vary with gender and adverse childhood experiences? *Child Abuse Neglect* 104: 104470
118. Wang BY et al (2016) Developmental trajectories of nonsuicidal self-injury in adolescence and intrapersonal/interpersonal risk factors. *J Res Adolesc* 27(2):392–406
119. Zetterqvist M (2016) Nonsuicidal self-injury in adolescents: characterization of the disorder and the issue of distress and impairment. *Suicide Life Threat Behav* 47(3):321–335
120. Zhang SC et al (2016) Low health literacy and psychological symptoms potentially increase the risks of non-suicidal self-injury in Chinese middle school students. *BMC Psychiatry* 16(1):327
121. Zubrick SR et al (2015) Self-harm: prevalence estimates from the second Australian child and adolescent survey of mental health and wellbeing. *Aust N Z J Psychiatry* 50(9):911–921
122. American Psychiatric Association (2013) Diagnostic and statistical manual of mental disorders, 5th ed. Diagnostic and statistical manual of mental disorders. American Psychiatric Association, Washington DC
123. Gratz KL (2001) Measurement of deliberate self-harm: preliminary data on the deliberate self-harm inventory. *J Psychopathol Behav Assess* 23(4):253–263
124. Klonsky ED, Glenn CR (2009) Assessing the functions of non-suicidal self-injury: psychometric properties of the inventory of statements about self-injury (ISAS). *J Psychopathol Behav Assess* 31(3):215–219
125. Lloyd EE, Kelley ML, Hope T (1997) Self-mutilation in a community sample of adolescents: descriptive characteristics and provisional prevalence rates. In: Poster presented at the annual meeting of the society for behavioral medicine. New Orleans, LA
126. Higgins JPT, Sally G (2011) Cochrane handbook for systematic reviews of interventions. Higgins JPT, Sally G (eds). The Cochrane Collaboration
127. Nock MK, Prinstein MJ (2005) Contextual features and behavioral functions of self-mutilation among adolescents. *J Abnorm Psychol* 114(1):140–146
128. Hawton K, Harriss L (2008) Deliberate self-harm by under-15-year-olds: characteristics, trends and outcome. *J Child Psychol Psychiatry* 49(4):441–448