





Genome Sequence and Characteristics of Cluster C1 *Mycobacterium smegmatis* Phage EasyJones

Isabel Amaya,^a Duyen Bui,^b Ariel Egbunine,^b Ember Mushrush,^b Maggie Viland,^c  Deborah Jacobs-Sera,^c Danielle Heller,^a  Viknesh Sivanathan^a

^aDepartment of Science Education, Howard Hughes Medical Institute, Chevy Chase, Maryland, USA

^bDepartment of Biology, University of Maryland Baltimore County, Baltimore, Maryland, USA

^cDepartment of Biological Sciences, University of Pittsburgh, Pittsburgh, Pennsylvania, USA

Duyen Bui, Ariel Egbunine, and Ember Mushrush contributed equally to this work.

ABSTRACT Bacteriophage EasyJones is a myovirus infecting *Mycobacterium smegmatis* mc²155, with a genome length and gene content similar to those of phages grouped in subcluster C1. Interestingly, EasyJones contains a gene found in a subset of C1 genomes that is similar to the well-characterized immunity repressor of subcluster A1 mycobacteriophage Bxb1.

Bacteriophages are increasingly being considered as therapeutic agents for multidrug-resistant bacterial infections. Recently, several bacteriophages isolated on nonpathogenic *Mycobacterium smegmatis* cells were used to treat a patient with a disseminated *Mycobacterium abscessus* infection (1). Here, we report on EasyJones, a mycobacteriophage that was isolated, using standard methods, from soil collected from a flower bed at the University of Maryland Baltimore County (Baltimore, MD) (2). Briefly, EasyJones was extracted by washing the soil with 7H9 liquid medium, enriched in a filtered (0.2- μ m pore size) wash, and purified with multiple rounds of plating on *M. smegmatis* mc²155 cells at 37°C. Top agar overlay of EasyJones results in clear plaques with a diameter of \sim 0.5 mm after 24 h at 37°C. Negative-stain transmission electron microscopy revealed EasyJones to be a myovirus with a contracted tail and an isometric capsid measuring \sim 88 nm in both length and diameter (Fig. 1A).

Double-stranded DNA was isolated from EasyJones using the Promega Wizard DNA cleanup kit, prepared for sequencing using the NEBNext Ultra II FS kit, and sequenced using an Illumina MiSeq sequencer to yield \sim 238,000 single-end 150-bp reads, which constituted \sim 231-fold coverage of the genome. Untrimmed reads were assembled and then checked for completeness using Newbler v2.9 and Consed v29, respectively, as described previously (3), resulting in a circularly permuted genome 154,315 bp in length, with a G+C content (64.7%) like that of the host bacterium (67.4%). EasyJones was assigned to phage subcluster C1 based on nucleotide similarity to members of this subcluster, using the PhagesDB database (4) and previously described criteria (5). The genome was annotated using DNA Master v5.23.6 (<http://cobamide2.bio.pitt.edu>), Glimmer v3.02 (6), GeneMark v3.25 (7), BLAST (8), HHpred (9), ARAGON (10), and tRNAscan-SE (11), all using default parameters. The resulting annotation process revealed a total of 267 protein-coding genes, 34 tRNAs, and 1 transfer-messenger RNA. Fifty of the protein-coding genes could be assigned functions, including the lysin A, lysin B, and holin genes.

Although a gene content similarity (GCS) comparison, performed using the PhagesDB GCS tool (4), revealed that EasyJones exhibits $>$ 83% GCS to members of subcluster C1, it adds to a small but growing list of C1 phages (15/152 phages) that possess a genomic segment encoding several additional gene products (EasyJones gp48 to gp51), including a homologue (gp49) of the well-characterized immunity repressor of subcluster A1

Editor Kenneth M. Stedman, Portland State University

Copyright © 2021 Amaya et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Viknesh Sivanathan, sivanathanv@hhmi.org.

Received 6 October 2021

Accepted 29 October 2021

Published 24 November 2021




FIG 1 (A) Negative-stain transmission electron micrograph of EasyJones. (B) Phylogeny of EasyJones gp49 homologues. Protein sequences for 815 EasyJones gp49 homologues were aligned with Clustal Omega, and a SplitsTree network phylogeny was generated. Clades are shaded according to cluster. The position of EasyJones gp49 (blue circle) within a clade consisting of homologues from clusters C and F is indicated.

mycobacteriophage Bxb1 (12–14). Previous reports indicated that this genomic segment was likely acquired horizontally and that the repressor homologue does not function as a canonical immunity repressor in C1 phages (12, 14). This is consistent with the clear plaque morphology of EasyJones and the absence of any recognizable integration or partitioning systems needed to support lysogeny. Instead, this acquired repressor is thought to be maintained because it defends infected host cells from superinfection by cluster A and related phages (12, 14). A phylogenetic analysis of all 815 homologues from across clusters A, BD, C, CA, F, J, and K reveals EasyJones gp49 to be most closely related to homologues from cluster F phages, some of which also encode homologues of EasyJones gp50 and gp51 (Fig. 1B).

Data availability. The sequencing data for EasyJones is available in Sequence Read Archive (SRA) with accession no. [SRX12475160](https://www.ncbi.nlm.nih.gov/sra/SRX12475160). The genome sequence of EasyJones is available in GenBank with accession no. [MZ856343](https://www.ncbi.nlm.nih.gov/genbank/MZ856343).

ACKNOWLEDGMENTS

We thank Daniel Russell and Rebecca Garlena for sequencing and assembling the genome, Graham Hatfull for feedback on the manuscript, and the SEA-PHAGES program for support. Duyen Bui and Ember Mushrush were each supported with NIH grant T34GM136497. Ariel Egbinine was supported with NIH grant 5TL4GM118989.

REFERENCES

- Dedrick RM, Guerrero-Bustamante CA, Garlena RA, Russell DA, Ford K, Harris K, Gilmour KC, Soothill J, Jacobs-Sera D, Schooley RT, Hatfull GF, Spencer H. 2019. Engineered bacteriophages for treatment of a patient with a disseminated drug-resistant *Mycobacterium abscessus*. *Nat Med* 25: 730–733. <https://doi.org/10.1038/s41591-019-0437-z>.
- Poxleitner M, Pope W, Jacobs-Sera D, Sivanathan V, Hatfull GF. 2018. HHMI SEA-PHAGES phage discovery guide. Howard Hughes Medical Institute, Chevy Chase, MD. <https://seaphagesphagediscoveryguide.helpdocsonline.com/home>.
- Russell DA, Hatfull GF. 2018. Sequencing, assembling, and finishing complete bacteriophage genomes. *Methods Mol Biol* 1681:109–125. https://doi.org/10.1007/978-1-4939-7343-9_9.
- Russell DA, Hatfull GF. 2017. PhagesDB: the Actinobacteriophage Database. *Bioinformatics* 33:784–786. <https://doi.org/10.1093/bioinformatics/btw711>.
- Pope WH, Mavrich TN, Garlena RA, Guerrero-Bustamante CA, Jacobs-Sera D, Montgomery MT, Russell DA, Warner MH, Science Education Alliance-Phage Hunters Advancing Genomics and Evolutionary Science (SEA-PHAGES), Hatfull GF. 2017. Bacteriophages of *Gordonia* spp. display a spectrum of diversity and genetic relationships. *mBio* 8:e01069-17. <https://doi.org/10.1128/mBio.01069-17>.
- Delcher AL, Harmon D, Kasif S, White O, Salzberg SL. 1999. Improved microbial gene identification with GLIMMER. *Nucleic Acids Res* 27:4636–4641. <https://doi.org/10.1093/nar/27.23.4636>.
- Borodovsky M, McIninch J. 1993. Recognition of genes in DNA sequence with ambiguities. *Biosystems* 30:161–171. [https://doi.org/10.1016/0303-2647\(93\)90068-N](https://doi.org/10.1016/0303-2647(93)90068-N).
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J Mol Biol* 215:403–410. [https://doi.org/10.1016/S0022-2836\(05\)80360-2](https://doi.org/10.1016/S0022-2836(05)80360-2).

9. Söding J, Biegert A, Lupas AN. 2005. The HHpred interactive server for protein homology detection and structure prediction. *Nucleic Acids Res* 33:W244–W248. <https://doi.org/10.1093/nar/gki408>.
10. Laslett D, Canback B. 2004. ARAGORN, a program to detect tRNA genes and tmRNA genes in nucleotide sequences. *Nucleic Acids Res* 32:11–16. <https://doi.org/10.1093/nar/gkh152>.
11. Lowe TM, Chan PP. 2016. tRNAscan-SE on-line: integrating search and contextual analysis of transfer RNA genes. *Nucleic Acids Res* 44: W54–W57. <https://doi.org/10.1093/nar/gkw413>.
12. Pope WH, Jacobs-Sera D, Russell DA, Peebles CL, Al-Atrache Z, Alcoser TA, Alexander LM, Alfano MB, Alford ST, Amy NE, Anderson MD, Anderson AG, Ang AA, Ares M, Jr, Barber AJ, Barker LP, Barrett JM, Barshop WD, Bauerle CM, Bayles IM, Belfield KL, Best AA, Borjon A, Jr, Bowman CA, Boyer CA, Bradley KW, Bradley VA, Broadway LN, Budwal K, Busby KN, Campbell IW, Campbell AM, Carey A, Caruso SM, Chew RD, Cockburn CL, Cohen LB, Corajod JM, Cresawn SG, Davis KR, Deng L, Denver DR, Dixon BR, Ekram S, Elgin SC, Engelsens AE, English BE, Erb ML, Estrada C, Filliger LZ, Findley AM, Forbes L, Forsyth MH, Fox TM, Fritz MJ, Garcia R, George ZD, Georges AE, Gissendanner CR, Goff S, Goldstein R, Gordon KC, Green RD, Guerra SL, Guiney-Olsen KR, Guiza BG, Haghghat L, Hagopian GV, Harmon CJ, Harmson JS, Hartzog GA, Harvey SE, He S, He KJ, Healy KE, Higinbotham ER, Hildebrandt EN, Ho JH, Hogan GM, Hohenstein VG, Holz NA, Huang VJ, Hufford EL, Hynes PM, Jackson AS, Jansen EC, Jarvik J, Jasinto PG, Jordan TC, Kasza T, Katelyn MA, Kelsey JS, Kerrigan LA, Khaw D, Kim J, Knutter JZ, Ko CC, Larkin GV, Laroche JR, Latif A, Leuba KD, Leuba SI, Lewis LO, Loesser-Casey KE, Long CA, Lopez AJ, Lowery N, Lu TQ, Mac V, Masters IR, McCloud JJ, McDonough MJ, Medenbach AJ, Menon A, Miller R, Morgan BK, Ng PC, Nguyen E, Nguyen KT, Nguyen ET, Nicholson KM, Parnell LA, Peirce CE, Perz AM, Peterson LJ, Pferdehirt RE, Philip SV, Pogliano K, Pogliano J, Polley T, Puopolo EJ, Rabinowitz HS, Resiss MJ, Rhyan CN, Robinson YM, Rodriguez LL, Rose AC, Rubin JD, Ruby JA, Saha MS, Sandoz JW, Savitskaya J, Schipper DJ, Schnitzler CE, Schott AR, Segal JB, Shaffer CD, Sheldon KE, Shepard EM, Shepardson JW, Shroff MK, Simmons JM, Simms EF, Simpson BM, Sinclair KM, Sjöholm RL, Slette IJ, Spaulding BC, Straub CL, Stukey J, Sughrue T, Tang TY, Tatyana LM, Taylor SB, Taylor BJ, Temple LM, Thompson JV, Tokarz MP, Trapani SE, Troum AP, Tsay J, Tubbs AT, Walton JM, Wang DH, Wang H, Warner JR, Weisser EG, Wendler SC, Weston-Hafer KA, Whelan HM, Williamson KE, Willis AN, Wirtshafter HS, Wong TW, Wu P, Yang Yj, Yee BC, Zaidins DA, Zhang B, Zúniga MY, Hendrix RW, Hatfull GF. 2011. Expanding the diversity of mycobacteriophages: insights into genome architecture and evolution. *PLoS One* 6:e16329. <https://doi.org/10.1371/journal.pone.0016329>.
13. Jain S, Hatfull GF. 2000. Transcriptional regulation and immunity in mycobacteriophages Bxb1. *Mol Microbiol* 38:971–985. <https://doi.org/10.1046/j.1365-2958.2000.02184.x>.
14. Hatfull GF. 2018. Mycobacteriophages. *Microbiol Spectr* 6:GPP3-0026-2018. <https://doi.org/10.1128/microbiolspec.GPP3-0026-2018>.