



Corrigendum: C1q/TNF-Related Protein 9 Attenuates Atherosclerosis by Inhibiting Hyperglycemia-Induced Endothelial Cell Senescence Through the AMPKα/KLF4 Signaling Pathway

OPEN ACCESS

Edited and reviewed by:

Francesco Rossi, University of Campania Luigi Vanvitelli, Italv

*Correspondence:

Jingbo Hou jingbohou@163.com

Specialty section:

This article was submitted to Cardiovascular and Smooth Muscle Pharmacology, a section of the journal Frontiers in Pharmacology

Received: 10 November 2021 Accepted: 22 November 2021 Published: 15 December 2021

Citation:

Wang G, Han B, Zhang R, Liu Q, Wang X, Huang X, Liu D, Qiao W, Yang M, Luo X, Hou J and Yu B (2021) Corrigendum: C1q/TNF-Related Protein 9 Attenuates Atherosclerosis by Inhibiting Hyperglycemia-Induced Endothelial Cell Senescence Through the AMPK /KLF4 Signaling Pathway. Front. Pharmacol. 12:812384. doi: 10.3389/fphar.2021.812384 Gang Wang^{1,2}, Baihe Han^{1,2}, Ruoxi Zhang³, Qi Liu^{1,2}, Xuedong Wang^{1,2}, Xingtao Huang^{1,2}, Dandan Liu^{1,2}, Weishen Qiao^{1,2}, Mengyue Yang^{1,2}, Xing Luo^{1,2}, Jingbo Hou^{1,2}* and Bo Yu^{1,2}

¹The Key Laboratory of Myocardial Ischemia Organization, Chinese Ministry of Education, Harbin, China, ²Department of Cardiology Organization, The Second Affiliated Hospital of Harbin Medical University, Harbin, China, ³Department of Cardiology, Harbin Yinghua Hospital, Harbin, China

Keywords: CTRP9, atherosclerosis, senescence, hyperglycemia, AMPK, KLF4

A corrigendum on

C1q/TNF-Related Protein 9 Attenuates Atherosclerosis by Inhibiting Hyperglycemia-Induced Endothelial Cell Senescence Through the AMPKa/KLF4 Signaling Pathway

by Wang, G., Han, B., Zhang, R., Liu, Q., Wang, X., Huang, X., Liu, D., Qiao, W., Yang, M., Luo, X., Hou, J., and Yu, B. (2021). Front. Pharmacol. 12:758792. doi: 10.3389/fphar.2021.758792

In the original article, there was a mistake in the artwork for **Figure 2C** as published. We made a careless mistake about p21 protein. The correct artwork appears below.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Wang, Han, Zhang, Liu, Wang, Huang, Liu, Qiao, Yang, Luo, Hou and Yu. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.



FIGURE 2 Hyperglycemia-induced HUVEC senescence is KLF4 dependent. (A) KLF4, as measured by immunoblotting and qRT-PCR in HUVECs after transfection with siNC or siKLF4 for 48 h. (B) Cells were fixed and stained for SA- β -gal activity and the histogram represents the percentage of SA- β -gal-positive cells per microscopic field. Values represent mean ± SEM. *p < 0.05, **p < 0.01. (C) KLF4, p21, and TERT protein levels were determined by immunoblotting. (D–F) Results were normalized to controls, and histograms represent the relative intensity of KLF4, p21, and TERT. Values represent mean ± SEM (n = 3–4 per group). *p < 0.05, **p < 0.01, ***p < 0.001.