

# Broadly active zinc finger protein-guided transcriptional activation of HIV-1

Tristan A. Scott,<sup>1</sup> Denis O'Meally,<sup>1</sup> Nicole Anne Grepo,<sup>1</sup> Citradewi Soemardy,<sup>1</sup> Daniel C. Lazar,<sup>2</sup> Yue Zheng,<sup>3</sup> Marc S. Weinberg,<sup>2,4,5</sup> Vicente Planelles,<sup>3</sup> and Kevin V. Morris<sup>1</sup>

<sup>1</sup>Center for Gene Therapy, City of Hope, Beckman Research Institute and Hematological Malignancy and Stem Cell Transplantation Institute at the City of Hope, 1500 E. Duarte Rd., Duarte, CA 91010, USA; <sup>2</sup>Department of Immunology and Infectious Disease, The Scripps Research Institute, La Jolla, CA 92037, USA; <sup>3</sup>University of Utah School of Medicine, Division of Microbiology and Immunology, Department of Pathology, Salt Lake City, UT 92037, USA; <sup>4</sup>Wits-SAMRC Antiviral Gene Therapy Research Unit, Department of Molecular Medicine & Hematology, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa; <sup>5</sup>HIV Pathogenesis Research Unit, Department of Molecular Medicine and Haematology, School of Pathology, University of the Witwatersrand, Johannesburg, South Africa

**Human immunodeficiency virus type 1 (HIV-1) causes a persistent viral infection resulting in the demise of immune regulatory cells. Clearance of HIV-1 infection results in integration of proviral DNA into the genome of host cells, which provides a means for evasion and long-term persistence. A therapeutic compound that specifically targets and sustainably activates a latent HIV-1 provirus could be transformative and is the goal for the “shock-and-kill” approach to a functional cure for HIV-1. Substantial progress has been made toward the development of recombinant proteins that target specific genomic loci for gene activation, repression, or inactivation by directed mutations. However, most of these modalities are too large or too complex for efficient therapeutic application. We describe here the development and testing of a novel recombinant zinc finger protein transactivator, ZFP-362-VPR, which specifically and potently enhances proviral HIV-1 transcription both in established latency models and activity across different viral clades. Additionally, ZFP-362-VPR-activated HIV-1 reporter gene expression in a well-established primary human CD4<sup>+</sup> T cell latency model and off-target pathways were determined by transcriptome analyses. This study provides clear proof of concept for the application of a novel, therapeutically relevant, protein transactivator to purge cellular reservoirs of HIV-1.**

## INTRODUCTION

Human immunodeficiency virus type 1 (HIV-1) establishes a latent long-term infection that remains hidden from the immune system. Clearance of latent HIV-1 DNA from infected host cells was suggested to take many decades.<sup>1</sup> Activation and purging of provirus from macrophages, monocytes, and T cells, with ~1 infected cell per million cells, has been an ongoing experimental struggle, the goal being to find latency reversing agents (LRAs) that can be used along with antiretroviral therapy (ART). This “shock-and-kill” approach aims to eliminate the reservoir by inducing HIV-1 activation resulting in death of the infected cell and immune clearance of the cells presenting antigen.<sup>2</sup> Several LRAs are known to activate latent provirus in HIV-infected individuals,<sup>3</sup> but LRAs are broad effectors of cellular processes and, therefore, inherent off-target effects

are expected. The ideal LRA should also activate genetically distinct clades of HIV-1. A method that could specifically activate latent virus in quiescent CD4<sup>+</sup> T cells<sup>4</sup> and in macrophages in those HIV-infected individuals on ART, and across multiple clades of virus, could prove useful to purge cellular viral reservoirs, expose infected cells to T cell killing, and potentially lead to a functional cure.

We and others have reported the potent and broad activation of latent HIV-1 using defective CRISPR fused to transcriptional activating domains.<sup>5–7</sup> Using this system, the most effective locus in the HIV-1 long terminal repeat (LTR) for modulating transcription was shown to be the LTR-362 site,<sup>5–7</sup> as a single guide RNA (sgRNA) directed to this site (sgF2-362) with a defective CRISPR activator, which is a catalytically dead Cas9 fused to a VPR activation domain (dCas9-VPR), potently activates latent HIV.<sup>7</sup> Notably, the LTR-362 site contains a nuclear factor  $\kappa$ B (NF- $\kappa$ B) sequence doublet that is unique to HIV-1 and not found in the human genome.<sup>8</sup> This site is susceptible to non-coding RNA-directed transcriptional control<sup>9,10</sup> and represents an HIV-specific site for targeted control of proviral transcriptional activation. While dCas9-VPR can be targeted to the LTR-362 site and potently activate HIV at levels greater than available LRAs, including suberoylanilide hydroxamic acid (SAHA), tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), or prostratin,<sup>7</sup> there are limitations, such as the requirement of a guide RNA administered with the dCas9-VPR protein. An alternative approach requiring a single smaller protein delivered systemically could prove useful in the activation of latent HIV. One such approach is outlined here and encompasses generating zinc finger protein (ZFP) conjugates targeted to the promoter of HIV. ZFPs are small, versatile, and modular and can target virtually any gene of interest. The DNA-binding specificity and modularity of ZFPs, specifically Cys<sub>2</sub>His<sub>2</sub>-type ZFPs, frequently result in their

Received 30 July 2020; accepted 21 October 2020;  
<https://doi.org/10.1016/j.omtm.2020.10.018>.

**Correspondence:** Kevin V. Morris, Center for Gene Therapy, City of Hope, Beckman Research Institute and Hematological Malignancy and Stem Cell Transplantation Institute at the City of Hope, 1500 E. Duarte Rd., Duarte, CA, 91010, USA.

**E-mail:** [kmorris@coh.org](mailto:kmorris@coh.org)

use as artificial transcription factors when fused to a transcription modulation domain.<sup>11</sup> Unlike CRISPR/Cas9, ZFPs require only a single component, which is smaller compared to Cas9 (~0.6 kb versus ~4 kb), making them more compatible with delivery vectors that have size limitations, such as with the clinical applied adeno-associated viral vector (AAVs) requiring two separate vectors to deliver CRISPR systems, even without bulky activation domains and with little space for regulatory features, or novel exosome delivery platforms favoring active loading of smaller nucleic acid cargo.<sup>12</sup>

Over the last decade, much progress has been made toward the development of a bona fide therapeutic protein based on ZF technology. ZF nucleases (ZFNs) have been developed to induce insertions or deletions (indels) at specific genomic loci, such as CCR5,<sup>13</sup> to generate HIV-resistant CD34<sup>+</sup> hematopoietic stem cells (HSCs) in bone-marrow transplants, and is currently being developed as a therapeutic strategy.<sup>14</sup> Importantly, ZFPs can be bioinformatically identified and tailored to target distinct sites using well-established algorithms.<sup>15</sup>

ZFPs can be fused to other proteins, such as repressive epigenetic regulatory proteins that direct gene silencing at the target locus, to provide locus-specific functionality.<sup>16–18</sup> ZFNs have been used to target and inactivate HIV's LTR,<sup>19</sup> activate the LTR and latent virus transcription,<sup>20,21</sup> excise provirus,<sup>22</sup> and repress HIV transcription.<sup>23</sup> However, no ZFP activator has been developed to target the responsive 362 site and used to activate all clades of HIV and a latent virus target. We describe here a new ZFP that can potentially activate all clades of HIV with a characterized off-target profile. This new ZFP may prove useful in “shock-and-kill” strategies to activate latent virus in the presence of ART or facilitating HIV targeted chimeric antigen T cell killing of those cells infected with virus.

## RESULTS

### ZFP-362-VPR specifically binds and transcriptionally activates the LTR of HIV

Recently, we and others demonstrated that a defective CRISPR conjugated to activation domains was capable of activating latent HIV-1 provirus.<sup>5–7,24–26</sup> This work demonstrated that these constructs are the most potent activators found to date for HIV, exhibiting far greater activation of HIV-1 than any LRAs currently employed to activate provirus, and that one locus appeared to be the most susceptible target site for CRISPR-directed modulation of HIV-1 transcription. This locus, the LTR-362 site,<sup>5–7</sup> contains an NF- $\kappa$ B doublet that is unique to HIV-1 and not found elsewhere in the human genome<sup>8</sup> and is one of the most effective sites for small antisense RNA directed transcriptional modulation of HIV-1.<sup>9,10</sup> Because the LTR-362 site is the “Achilles heel” of HIV-1 transcription, we used ZF Tools version 3.0 to bioinformatically generate three unique ZFP amino acid sequences that theoretically bind the 362 site (Figure 1A). The new ZFPs were fused to a Tat domain used previously in a ZFP targeting *UBE3A* as a potential therapy for Angelman syndrome<sup>16</sup> and the VPR domain used previously with defective CRISPR to HIV-1.<sup>7</sup> The VPR domain has in addition to the VP64, a concatamer of VP16 for herpes simplex virus (HSV), the endogenous p65 and


Epstein-Barr virus (EBV) activator, Rta, which collectively constitute a potent transcription activation domain.<sup>27</sup> One of these three ZFPs, ZFP-362b (referred to as ZFP-362), was found to potentially activate HIV-1 expression at levels comparable to dCas-VP64+sgF2-362, in a CEM clonal T cell line has an integrated LTR-mCherry-IRES-Tat (LChIT) reporter<sup>28</sup> or pMo-HEK cells,<sup>29</sup> a HEK293 cell line that has been transduced with a lentivirus with an LTR-driven GFP (Figures 1B and 1C). Notably, this targeted activation was lost in pMo $\Delta$ 362 cells, which lack the ZFP-362 target site (Figure 1D). Next, we tested the breadth of ZFP-362 transcriptional activation. The ability of ZFP-362 to activate clades A–G of HIV-1 was assessed in co-transfected HEK293 cells with vectors containing subtype-specific LTRs driving luciferase reporter (Figure 1E). ZFP-362 demonstrated the most potent activation of subtypes A, B, D, and F and less robust activation in subtypes C, E, and G. A noteworthy observation is that the target site is conserved in subtypes A, B, D, and F, while subtypes E and G contain point mutations and deletions in the LTR-362 site (Figure S1A). Subtype C contains a triple NF- $\kappa$ B site, and the ZFP-362 binding locus aligns better between the second and third NF- $\kappa$ B motifs with a single mismatch deletion (Figure S1B). Overall, these data support the notion that ZFP-362-VPR is specifically active on the HIV-1 LTR for a wide range of HIV subtypes.

To more clearly determine the targeting of ZFP-362-VPR, a chromatin immunoprecipitation assay (ChIP)<sup>30</sup> was performed on both LTR targeted dCas9-VPR+sgF2-362 and ZFP-362-VPR treated pMo-HEK cells. Both the dCas9-VPR and ZFP-362-VPR were found to localize specifically to the LTR (Figure 1F), and ZFP-362 enrichment was lost when targeting a pMo vector lacking the 362 site (Figure S2). To verify the presence of the ZFP-362-VPR, protein expression was confirmed by western blot (Figure S3). Collectively, these data suggest that the ZFP-362-VPR fusion construct can target HIV-1 at the LTR-362 site and potentially activate viral transcription.

We have previously demonstrated that a dCas9-VPR and dCas9+sgF2-362 can activate a variety of latent HIV-1 models, whereas variable activation was observed with other LRAs.<sup>7</sup> We similarly tested the ZFP-362-VPR vector on well-established reporter cell lines for HIV latency and observed activation of clonally selected J-Lat cell lines “6.3,” “10.6,” and “15.4,” with similar potency to that of the dCas9+sgF2-362 (Figure 2). Furthermore, the ZFP-362 reliably activated the LTR compared to the most commonly used LRAs (Figure 3). These data demonstrate that ZFP-362-VPR can consistently and potentially activate HIV in different models of HIV latency and that this activation is independent of activating NF- $\kappa$ B signaling pathways.

### ZFP-VPR-362 activates HIV in a primary CD4<sup>+</sup> T cell model of latency

It was determined if the ZFP-362-VPR can activate HIV-1 in a primary HIV latency model. CD4<sup>+</sup> T cells were HIV-1 infected *in vitro* 17 days prior with a pNL4.3- $\Delta$ env-nluc replication-incompetent vector expressing nano-luciferase pseudotyped with VSVG. As active infection reduces CD4<sup>+</sup> expression, the latent population and




**Figure 1. ZFP-362-VPR-mediated transcriptional activation of HIV**

(A) A schematic is shown depicting ZFP-362-VPR, a zinc finger fused to the VPR activator domain, a Tat peptide motif, and NLS peptide sequences for nuclear targeting, which was developed to transcriptionally activate the NF- $\kappa$ B site in the HIV-1 LTR.<sup>10</sup> (B) ZFP-362-VPR activates LTR expression in LCHIT 3.2 bimodal latent HIV reporter cells at levels that are comparable to dCas-VPR. (C) ZFP-362-VPR activates LTR expression of GFP in pMO-HEK cells but (D) has no effect on pMO $\Delta$ 362 cells containing a deletion in the LTR-362 site targeted by the ZFP-362. (E) The ability of ZFP-362-VPR to activate various subtypes of HIV was determined by co-transfection of ZFP with subtype-variable LTR expressing luciferase clones. (F) ChIP analysis of ZFP-362-VPR and dCas-VPR binding to the HIV LTR. The pMO-HEK cells were transfected with the dCas-VPR+sgF2-362 or control sgRNA, or ZFP-362-VPR and ChIP assay was performed 72 hr post-transfection to determine binding to the LTR-362 site.<sup>8</sup> The experiments were performed in triplicate-treated cells, and errors bars are calculated as standard deviations. \* $p < 0.001$  from an unpaired Student's *t* test.

uninfected cells could be separated from productively infected cells on day 17. The ZFP-362-VPR vector was transfected into the CD4<sup>+</sup> T cells 72 h post-separation, and the level of luciferase activity was measured as a readout of LTR activation. There was a significant increase in luciferase levels with the ZFP-362-VPR compared to a GFP or ZFP control (Figure 4). Likewise, known activators of the LTR, PMA, or anti-CD3/CD28 stimulation, resulted in high levels of LTR activation. Although high levels of activity were observed in one donor, the levels of activity with the ZFP-362-VPR in two other donors was less pronounced (Figure S4). Nonetheless, these data demonstrate that the ZFP-362-VPR can activate the HIV-1 LTR in primary CD4<sup>+</sup> T cells.

#### Assessment of ZFP-VPR-362 off-target effects

A major concern of using LRAs is the broad off-target cellular effects that can result in cellular toxicity or oncogenic mutations. As the ZFP-362 targets the same site as the sgF2-362, we compared the effects of ZFP-362-VPR on endogenous RNA expression. The ZFP-362-VPR, dCas9+sgF2-362, or pcDNA3.1 was transfected into pMo-HEK cells, and gene expression was assessed by bulk RNA-seq analysis. Principal component analysis showed tight clustering of the sample groups, indicating experimental treatment is the major source of variation (Figure S5A). Expression plots show that ZFP-362-VPR and dCas9+sgF2-362 potentially activate the on-target GFP reporter, and CRISPR components were only detected in




**Figure 2. Activation effects of ZFP-362-VPR on different J-Lat latent infected cell models**

Three different J-Lat cell lines (J-Lat 6.3, 10.6, and 15.4) were subjected to neon transfection with dCas9-VPR+sgF2-362, ZFP-VEGF-VPR (control), or ZFP-362-VPR. The expression of GFP was determined by FACS at 72 hr post-treatment. Fold activation was the GFP MFI made relative to a mock sample set at 1. The experiments were performed in triplicate-treated cells, and errors bars are calculated as standard deviations. Statistical significance was determined using an unpaired Student's t test (\*\* $p < 0.005$ ).


dCas9-VPR+sgF2-362 cells (Figures S5B and S5C). There were 191 differentially expressed genes observed in the dCas9-VPR-sgF2-362 (Table S1) and 219 observed in the ZFP-362-VPR-treated samples (Table S2) compared to the control pcDNA3.1 vector ( $|\log_2FC| \geq 2$ , false discovery rate [FDR] = 0.05). Importantly, the ZFN-362-VPR had a comparable gene expression profile to the dCas9-VPR+sgF2-362 (Figure 5A): 358 differentially expressed genes were common to both treatments, 12 genes unique to the dCas9-VPR+sgF2-362-treated samples, and 40 genes unique to the ZFP-362-VPR-treated samples (Figure 5B; Tables S3 and S4). We looked for enriched Gene Ontology (GO) terms and KEGG pathways associated with differentially expressed genes in ZFP-362-VPR-treated samples with respect to controls, and enrichment of terms associated with neurotransmitter transport and pathways specific for cardiomyocytes was observed (Figure 5C). Similarly, enrichment for these GO terms and KEGG pathways was observed for dCas9-VPR+sgF2-362-treated samples, suggesting similar target profiles (Figure 5D). Importantly, none of the differentially expressed genes or pathways is a known oncogene or is associated with cell proliferation.

## DISCUSSION

ZFPs are versatile and modular. Several ZFPs have been developed to target and repress HIV-1, including one designed against the SP1 site in the LTR,<sup>31</sup> while others have developed ZFPs targeted to three regions in the LTR and found that they were potent stable repressors of viral expression.<sup>32</sup> However, none of these previous studies developed a ZFP to the highly conserved NF- $\kappa$ B site in the LTR. We report here a ZFP activator that specifically activates HIV-1 transcription by targeting the LTR-362 site, a site identified through CRISPR gRNA tiling across the 5' LTR and determined to be highly sensitive to promoter activation.<sup>7</sup> Moreover, these first-generation ZFPs, while functional, lacked several modifications that enhance nuclear function and trafficking. One recent and exciting example utilized a ZFP conjugated to a KRAB silencing domain and a Tat domain for targeting the brain and directed to the promoter of an antisense long non-coding RNA (lncRNA) involved in regulating genes active in Angelman syndrome.<sup>16</sup> In that study, it was conclusively shown that brain-directed targeting of a ZFP is feasible *in vivo* and contingent on



(legend on next page)



**Figure 4. ZFP-362-VPR activity in primary CD4<sup>+</sup> T cell latency model**

CD4<sup>+</sup> T cells (pseudotyped viruses pNL4.3-ΔEnv-nLuc-VSVG uninfected cells and latently infected cells) were transfected with ZFP-362-VPR and control vectors (GFP and ZFP-control), and 72 hr later, the levels of luciferase were measured. CD4<sup>+</sup> T cells were also treated with PMA (10 ng/mL) and anti-CD3/CD28 beads as positive controls. The average of triplicate-treated cells are shown with the standard deviations, and p values were determined using a one-way ANOVA and Dunnett's test. \*\*\*\*p < 0.0001; NS, not significant.

a Tat localization peptide conjugated with the ZFP.<sup>16</sup> We capitalized on these recent developments and developed ZFP-362-VPR to contain the same Tat domain used previously in the ZFP targeting in Angelman syndrome.<sup>16</sup> The use of this Tat domain may allow for the ZFP-362 to function as a soluble protein as well as to enhance nuclear targeting and retention of the ZFP-362 activator, though such a notion was not experimentally assessed in this study.

While ZFPs, transcription activator-like effector nucleases (TALENs), and several nuclease CRISPR and defective CRISPR systems containing various conjugates have been shown to target HIV-1 for applications

ranging from silencing<sup>33</sup> to excision<sup>26,34–38</sup> to activation,<sup>5,7</sup> the problem of achieving target specificity while maintaining activity across the various HIV-1 clades has remained enigmatic. Notably, ZFP-362 is capable of broadly activating all the LTRs of different clades of HIV-1 to varying levels of expression but strongly activates 4 of the 7 clades of HIV-1 with full sequence complementarity, a conservation not present in previous LTR ZFP activators,<sup>18</sup> which encourages the further characterization of ZFP-362-VPR in more complex clade-specific latency models. Additionally, there is a significant HIV-1 disease burden of “non-B” subtypes in underdeveloped regions, such as a prevalence of HIV-1 subtype C in southern Africa.<sup>39</sup> Whether next-generation gene and cellular treatments would be financially or technically feasible in these regions remains to be assessed, but as these next-generation therapeutics offer significant potential for the treatment of HIV-1, their development should be encouraged.


The ZFP activator affects a limited subset of genes that appear transcriptionally activated in an off-target manner. ZFP-362-VPR has a very similar pattern of differential gene expression compared to dCas9-VPR+sgF2-362, suggesting a comparable off-target profile for these vectors, which did not affect cell cycle or viability.<sup>7</sup> Importantly, effects on T cell proliferation and viability have been observed with traditional LRAs and represent an advantage of a more targeted approach.<sup>40</sup> Of note, we previously found that differentially expressed genes with the dCas9-VPR+sgF2-362 were significantly lower in number and magnitude compared with TNF-α,<sup>7</sup> suggesting these vectors have a better on-target profile than LRAs with broad-acting cellular pathways. Importantly, ZFP-362 expression could be regulated either through the specificity of the delivery vector or inclusion of regulatory factors ensuring tissue-specific expression and possibility minimize off-target effects a design feature not readily applicable to broad-activating small-molecule LRAs.

Nevertheless, ZFP-362-VPR affected genes enriched in pathways relevant to cardiomyocyte signaling and contraction (Figure 5). These data suggest that even though the NF-κB doublet is not present in the human genome, the dCas9-VPR+sgF2-362 and ZFP-362-VPR may interact with some NF-κB binding sites leading to off-target effects. If this was the case, not all NF-κB sites are bound and activated, as known NF-κB pathways, such as cytokine signaling or proliferation,<sup>41</sup> were not identified by enrichment analysis, and 362 site-targeting in our previous work showed no effect on cell cycle progression and viability.<sup>7</sup> However, we cannot rule out non-specific ZF binding and activation; further work is required to determine the range of motifs bound by ZFP-362-VPR, enabling us to better identify the potential physiological effects of ZFP-362-VPR-mediated dysregulation.


**Figure 3. Comparison of ZFP-362-VPR and LRAs on *in vitro* latency models**

Three different J-Lat cell lines (J-Lat 6.3, J-Lat 10.6, and J-Lat 15.4) were subjected to either electroporation with dCas9-VPR+sgF2-362 or ZFP-362-VPR or treated with various known LRAs: TNF-α, CD3/CD28 beads, prostratin, PMA, or PMA+ionomycin. DMSO and untreated (mock) cells were included as negative controls. Activation of HIV was determined 72 hr after treatment by FACS for the levels of GFP. Fold activation was the GFP percentage made relative to a control vector set at 1. The results of triplicate-treated cultures are shown with the SEM. Statistically significant differences, as determined from a one-way ANOVA and Dunnett's test are also shown (\*p < 0.05, \*\*p < 0.005, and \*\*\*\*p < 0.0001) and were analyzed compared to the mock or DMSO control for the vectors and LRAs, respectively.



**A** ZFP-362-VPR versus dCas9-VPR+sgF2-362





**B** dCas9-VPR+sgF2-362 ZFP-362-VPR



**C**



**D**



(legend on next page)

Dysregulation of NF- $\kappa$ B activation can result in cardiac remodeling and hypertrophy and contribute to heart failure,<sup>42</sup> and it remains to be determined if ZFP-362-VPR is able activate downstream NF- $\kappa$ B target sites involved in these processes. Although we did not assess ZFP-362-VPR *in vivo*, our data suggest that restricting delivery of ZFP-VPR-362 to cardiac tissue may be required, or that this approach should be used transiently to activate latent virus for a defined therapeutic window. Using delivery vectors that do not accumulate in cardiac tissue could be helpful in avoiding off-target effects. Alternatively, inclusion of tissue-specific microRNA (miRNA) target sites in the transgene has been used to prevent undesirable expression in specific tissues,<sup>43</sup> and incorporating cardiac-specific miRNA target sites into ZFP-362-VPR may restrict expression in this organ.<sup>44</sup>

ZFPs are endogenous regulators of gene transcription that can regulate transposable and viral elements,<sup>45</sup> including human ZNF10, which inhibits HIV-1 LTR function.<sup>46</sup> Interestingly, endogenous ZNF10 was found to block SP1/NF- $\kappa$ B interaction in the HIV-1 LTR and inhibit viral transcription. It is notable that the cell has evolved a ZFP regulator to this same region that ZFP-362 is directed against, suggesting that over time, perhaps HIV-1 will be silenced by the action of endogenous KRAB containing ZFPs.<sup>47</sup>

The ZFP-362 activator of HIV-1 described here may prove useful in the development of a functional cure for HIV-1 infection. For instance, the ZFP-362 activator could be used in a “shock-and-kill” strategy to activate latent virus while maintaining ART, leading to the death of the latently infected cells by viral-induced apoptotic factors or the immune system.<sup>2</sup> There is evidence that replication-competent infected CD4<sup>+</sup> T cells do not die as a result of virus-induced cytopathic effect<sup>48</sup> and may require targeted elimination. ZFP-362-VPR could be used to induce sustained expression of HIV-1 to allow for, or enhance, chimeric antigen T cell (CAR) targeting of infected cells. Notably, the lack of HIV-1 antigen has proven problematic for developing robust anti-HIV CARs.<sup>49,50</sup> However, for either of these approaches to prove viable, the issue of delivery remains to be solved. Indeed, while ZFP-, TALEN-, and CRISPR-based gene therapies are all proving exceptionally valuable in modulating HIV-1 expression and may in the next decade emerge as a bona fide means of eliminating HIV-1 infection, a means of delivery to infected target cells remains enigmatic (reviewed in Liang et al.<sup>51</sup> and Saayman et al.<sup>52</sup>). ZFPs or TALENs could be delivered using lipid nanoparticles<sup>53</sup> or cellular exosomes.<sup>54</sup> Such an approach would be amendable to CAR T cell and “shock-and-kill” based strategies; however, more work in developing these systems and determining the ability to deliver ZFP-362 remains to be carried out, as do many of the technical challenges inherent in cell delivery.

## MATERIALS AND METHODS

### Vectors

The H1sgRNA (control), dCas9-VPR, and sgF2-362 have been previously described.<sup>7</sup> The amino acid sequences of ZFP-362 and ZFP-VEGF that bind their respective target sites were identified using ZF Tools version 3.0<sup>15</sup> and fused to a VPR activation domain and cloned into pcDNA3.1 mammalian expression vector by Genscript (amino acid sequence of ZFP-362-VPR protein provided in the [Supplemental information](#)). The ZFP-362 was ordered as a gBLOCK (IDT, Coralville, IA, USA) and was cloned into a dCas9-VPR vector (Addgene #63798) to replace the dCas9 in this vector using NEBuilder HiFi DNA assembly master mix (NEB, Ipswich, MA, USA). The LTR-driven luciferases from different subtypes were obtained from the NIH AIDS reagent program (cat. nos. 4787, 4788, 4789, 4790, 4791, 4792, 4793; NIH AIDS Reagent Program, Division of AIDS, NIAID, NIH; pBlue3<sub>2</sub>LTR-luc-A from Dr. Reink Jeeninga and Dr. Ben Berkhout).<sup>55,56</sup>

### Cell lines

pMo-HEK and p $\Delta$ 362 lentiviral vectors were generated as described in Shrivastava et al.<sup>29</sup> HEK293T cells (ATCC) were transduced at a MOI of 0.1 to ensure a single integrated copy. 7 days post-transduction, GFP-positive cells were fluorescence-activated cell sorting (FACS)-sorted (FACSCalibur II, Becton Dickinson, East Rutherford, NJ, USA, at the Scripps Flow Cytometry core facility) to generate a cell line with a stably integrated HIV-1 LTR or with a  $\Delta$ 362LTR that contains mutations at the 362 site. The following J-Lat cell lines were obtained through the NIH AIDS reagent program, Division of AIDS, NIAID, NIH: J-Lat full-length clones 6.3, 10.6, and 15.4, which are collectively referred to as J-Lat cells, which were deposited by Dr. Eric Verdin.<sup>57</sup> The LChIT reporter line, a CEM T cell-based reporter system comprising of the single-copy LTR integrant driving the expression of mCherry-IRES-tat, has been described previously.<sup>7</sup>

### Cell culture

HEK293T cells (ATCC) and the derived pMo-HEK and p $\Delta$ 362 reporter lines were maintained in Dulbecco's modified Eagle's medium (Thermo Fisher Scientific, Waltham, MA, USA) supplemented with 10% fetal bovine serum (Thermo Fisher Scientific) and 50  $\mu$ g/mL penicillin/streptomycin (Thermo Fisher Scientific) at 37°C and 5% CO<sub>2</sub>. J-Lat and LChIT cell lines were both maintained in RPMI 1640 (Thermo Fisher Scientific) supplemented with 10% fetal bovine serum, 2 mmol/L l-glutamine (Thermo Fisher Scientific), and 50  $\mu$ g/mL penicillin/streptomycin (Thermo Fisher Scientific) at 37°C and 5% CO<sub>2</sub>.

### Figure 5. ZFP-362-VPR induced differential gene expression

pMo-HEK cells were transfected with the pcDNA3.1, dCas9-VPR+sgF2-362, or ZFP-362-VPR expression vectors, and total RNA was extracted 72 hr post-transfection and subjected to RNA-sequencing analysis. (A) Comparison of the log fold change of ZFP-362-VPR-treated samples with dCas9-VPR+sgF2-362 mean expression. Differentially expressed genes ( $|\log_{2}FC| > 1$ , FDR 0.05) are colored red; GFP expression is equally and potently activated in both treatments. (B) A Venn diagram of the number of differentially expressed genes unique to ZFP-362-VPR and dCas9-VPR+sgF2-362-treated samples. Bar plot of enriched KEGG pathways indicated by differentially expressed genes in the (C) ZFP-362-VPR- and (D) dCas9-VPR+sgF2-362-treated samples with respect to controls.



### CLADE-specific activation assays

HEK293 cells were transfected in triplicate using Lipofectamine 3000 (Thermo Fisher Scientific) with pcDNA-ZFP-362-VPR or pcDNA-VEGF-VPR with vectors expressing firefly luciferase off the LTRs from different subtypes of HIV-1. A vector expressing *Renilla* luciferase was included as a background control (pRL-CMV, Promega, Fitchburg, WI, USA). At 48 hr post-transfection, a dual-luciferase reporter assay was performed according to manufacturer instructions and luciferase activity detected on the Glomax Explorer system (Promega). The levels of firefly luciferase were normalized to *Renilla* luciferase and made relative to the pcDNA-VEGF-VPR control.

### Activation of the HIV-1 LTR

The protocol to assess the ability of the ZFP-362-VPR construct to activate latent HIV-1 has been described in Saayman et al.<sup>7</sup> In brief, pcDNA-ZFP-362-VPR, pcDNA-ZFP-VEGF-VPR (control), or pcDNA3.1 (mock) plasmids were transfected into the CEM LChIT cells and the J-Lat cells using the neon transfection system (Thermo Fisher Scientific) at a cell density of  $2 \times 10^7$  cells/mL with the following electroporation parameters: 3 pulses, 1,350 V, and 10 ms. The dCas9-VPR+sgF2-362 vectors, which have been previously shown to potentially activate HIV-1, were included as a comparison.<sup>7</sup> The cells were cultured without antibiotics following electroporation. To determine the activation with LRAs, J-Lat cells were treated with TNF- $\alpha$  (10 ng/mL; Gibco, cat. #PHC3015), CD3/CD28 beads (3:1 ratio beads:cells; Dynabeads human T-expander CD3/CD28 beads, cat. #11141D), prostratin (3  $\mu$ mol/L; Sigma cat. #P0077-1mg), PMA (50 ng/mL; Fisher Scientific, cat. #BP685-1) with/without ionomycin (500 ng/mL; Tocris Biosciences, cat. #1704/1). A DMSO control was included with 0.5% (v/v) DMSO dissolved in RPMI media with 10% FBS. Seventy-two hours post-treatment, the ability of these constructs to induce latent activation was assessed by FACS with 10,000 single events collected, and the data was analyzed using FlowJo vX3.05470 software.

### NF- $\kappa$ B site specificity of ZFP-362-VPR

To test the target site specificity of the pcDNA-ZFP-362-VPR to activate the LTR of HIV-1, 100,000 pMo-HEK and p $\Delta$ 362 reporter cells were seeded per well in a 24-well plate at 24 h prior to transfection. Cells were transfected in triplicate using Lipofectamine 2000 (Thermo Fisher Scientific), and 72 h post-transfection, total cellular RNA was isolated from cells using the Maxwell RSC simplyRNA cells kit (Promega) according to manufacturer's instructions. The concentrations of the isolated DNase-treated RNA samples were standardized and then reverse transcribed using Mu-MLV reverse transcriptase (Thermo Fisher Scientific) with a n-oligo-dT/random nonamer primer mix (IDT). Quantitative real-time PCR was carried out using Kapa Sybr Fast universal qPCR mix (Kapa Biosystems, Wilmington, MA, USA) on a Roche LightCycler 96 system (Roche Diagnostics, Indianapolis, IN, USA) according to provided instructions. The following PCR primer sets were used: GFP forward primer, 5'-GACAACCACTACCTGAGCAC-3'; GFP reverse primer, 5'-CAGGACCATGTGATCGCG-3'; RPLP0 forward primer, 5'-CGCAGCCAATAGACAGGAG-3'; RPLP0 reverse primer, 5'-GCCGCTGCCTTTTATAATGC-3';<sup>58</sup> and the GFP expres-

sion was analyzed using quantitative reverse transcription PCR (qRT-PCR). pcDNA-ZFP-VEGF-VPR and pcDNA3.1 were included as negative controls. The qPCR data was analyzed using LightCycler software version 1.1.1320.

### Activation of HIV-1 LTR in CD4<sup>+</sup> T cell latency model

Naive cells were isolated and activated by anti-interleukin (IL)-12 (Peprotech; cat. #500-P154G), anti-IL-4 (Peprotech; cat. #500-P24), transforming growth factor  $\beta$  (TGF- $\beta$ ) (Peprotech; cat. #100-21), and anti-CD3/CD28 beads (Invitrogen; cat. #11131D). Activated cells were infected by pseudotyped pNL4.3-deltaEnv-nLuc-VSVG viruses. At day 17, CD4<sup>+</sup> cells were isolated using a Dynabeads CD4<sup>+</sup> isolation kit (Invitrogen; cat. #11331D). At day 18, CD4<sup>+</sup> cells were treated with 10 ng/mL PMA (Sigma; cat. #P1585) or anti-CD3/CD28 beads or transfected by plasmids (GFP, ZFP-362-VPR, or ZFP-VEGF-VPR) using by neon electroporation (Invitrogen; cat. #mpk1096). After 72 hr, part of the cells were stained with fixable viability dye efluor 450 (eBioscience; cat. #65-0863-14) and anti-CD4 (Invitrogen; cat. #MHCD0405) and anti-P24 (BD; cat. #556027) antibodies and measured by flow cytometry. The cells were counted, and samples containing 2,000 live cells were lysed and measured using the Nano-Glo luciferase assay system (Promega; cat. #N1120). Ethics approval was obtained from the University of Utah to use primary tissues in the described assays (IRB 00067637).

### ChIP analysis of ZFP-362-VPR

The ZFP-362-VPR and dCas9-VPR+sgF2-362 plasmids were transfected in triplicate into 3 million pMo-HEK cells using calcium phosphate transfection. ChIP analysis was carried out on the transfected pMo-HEK cells for the myc-tagged activating complexes using anti-myc antibodies (Cell Signaling Technology, Danvers, MA, USA). The ChIP protocol was performed 72 h post-transfection using previously described protocols.<sup>10,59,60</sup> The ChIP elutes were purified using the MinElute PCR purification kit (QIAGEN, Hilden, Germany) and analyzed by qPCR. Relative enrichment at the LTR was determined using LTR specific primers that were described in Saayman et al.<sup>59</sup>

### ZFP-362-VPR and dCas9-VPR+F2gRNA effects on global RNA expression

To determine the extent to which ZFP-362-VPR and dCas9-VPR+sgF2-362 disrupted RNA expression, we compared pMo-HEK cells transfected with either construct to those transfected with a pcDNA3.1 control. The pMo-HEK were seeded at 100,000 cells per well in a 24-well plate and transfected with the ZFP-362-VPR and dCas9-VPR+sgF2-362 plasmids in quadruplicate using calcium phosphate transfection. Total cellular RNA was isolated 48 h post-transfection from cells using the Maxwell RC simplyRNA cells kit (Promega). Sequencing was performed by Genewiz on an Illumina platform with paired-end 150-bp reads. Sequence data are available via Gene Expression Omnibus (GEO: GSE150382). Raw sequencing reads were processed with the nf-core<sup>61</sup> RNASeq pipeline version 1.4. In brief, trimmed reads were mapped using spliced transcripts

alignment to a reference (STAR)<sup>62</sup> to the GRCh37 reference using the ENSEMBL (release 75) annotation. Each library was subjected to extensive quality control, including estimation of library complexity, gene body coverage, and duplication rates, among other metrics detailed in the pipeline repository. Reads were counted across genomic features using Subread featureCounts<sup>63</sup> and merged into a matrix of counts per gene for each sample. Differential expression analysis was performed with DESeq2<sup>64</sup> as implemented in the SAR-Tools package.<sup>65</sup> Only genes with at least 1 read count in 1 sample were retained for the analysis. Multiple testing was corrected using the Benjamini-Hochberg method, where p values are adjusted to control for false discovery. An adjusted p value threshold of 0.05 was used to select differentially expressed genes. Gene enrichment analyses were performed with ClusterProfiler<sup>66</sup> using the full set of genes with at least 1 count in any sample as background and differentially expressed genes as the input list. Enriched GO terms or enriched KEGG pathways were selected using an FDR threshold of 0.05 with the enrichGO function of ClusterProfiler.

### Statistical analysis

Graphing and statistical analyses were performed using GraphPad Prism version 8 (V8.1.2), and the statistical analysis used and p values are described in the figure legends.

### SUPPLEMENTAL INFORMATION

Supplemental Information can be found online at <https://doi.org/10.1016/j.omtm.2020.10.018>.

### ACKNOWLEDGMENTS

This project was supported by NIAID RO1 AI111139-01 and NIMH R01 113407-01 to K.V.M. V.P. is funded by grant AI 124843-02.

### AUTHOR CONTRIBUTIONS

Conceptualization, K.V.M., M.S.W., and T.A.S.; Methodology, T.A.S., Y.Z., and D.C.L.; Formal Analysis, T.A.S., D.C.L., N.A.G., and D.O.; Investigation, Y.Z., T.A.S., N.A.G., C.S., and D.C.L.; Writing – Original Draft, T.A.S.; Writing – Review & Editing, T.A.S. and K.V.M.; Funding Acquisition, K.V.M.; Resources, K.V.M. and V.P.; Visualization, T.A.S. and K.V.M.; Supervision, K.V.M.

### DECLARATION OF INTERESTS

K.V.M., M.S.W., T.A.S., and D.C.L. have been issued a patent on this technology, W02019079819.

### REFERENCES

- Murray, A.J., Kwon, K.J., Farber, D.L., and Siliciano, R.F. (2016). The Latent Reservoir for HIV-1: How Immunologic Memory and Clonal Expansion Contribute to HIV-1 Persistence. *J. Immunol.* 197, 407–417.
- Sadowski, I., and Hashemi, F.B. (2019). Strategies to eradicate HIV from infected patients: elimination of latent provirus reservoirs. *Cell. Mol. Life Sci.* 76, 3583–3600.
- Elliott, J.H., Wightman, F., Solomon, A., Ghneim, K., Ahlers, J., Cameron, M.J., Smith, M.Z., Spelman, T., McMahon, J., Velayudham, P., et al. (2014). Activation of HIV transcription with short-course vorinostat in HIV-infected patients on suppressive antiretroviral therapy. *PLoS Pathog.* 10, e1004473.
- Sedaghat, A.R., Siliciano, J.D., Brennan, T.P., Wilke, C.O., and Siliciano, R.F. (2007). Limits on replenishment of the resting CD4+ T cell reservoir for HIV in patients on HAART. *PLoS Pathog.* 3, e122.
- Limsirichai, P., Gaj, T., and Schaffer, D.V. (2016). CRISPR-mediated Activation of Latent HIV-1 Expression. *Mol. Ther.* 24, 499–507.
- Zhang, Y., Yin, C., Zhang, T., Li, F., Yang, W., Kaminski, R., Fagan, P.R., Putatunda, R., Young, W.B., Khalili, K., and Hu, W. (2015). CRISPR/gRNA-directed synergistic activation mediator (SAM) induces specific, persistent and robust reactivation of the HIV-1 latent reservoirs. *Sci. Rep.* 5, 16277.
- Saayman, S.M., Lazar, D.C., Scott, T.A., Hart, J.R., Takahashi, M., Burnett, J.C., Planelles, V., Morris, K.V., and Weinberg, M.S. (2016). Potent and Targeted Activation of Latent HIV-1 Using the CRISPR/dCas9 Activator Complex. *Mol. Ther.* 24, 488–498.
- Turner, A.M., Ackley, A.M., Matrone, M.A., and Morris, K.V. (2012). Characterization of an HIV-targeted transcriptional gene-silencing RNA in primary cells. *Hum. Gene Ther.* 23, 473–483.
- Suzuki, K., Shijuuku, T., Fukamachi, T., Zaunders, J., Guillemin, G., Cooper, D., and Kelleher, A. (2005). Prolonged transcriptional silencing and CpG methylation induced by siRNAs targeted to the HIV-1 promoter region. *J. RNAi Gene Silencing* 1, 66–78.
- Weinberg, M.S., Villeneuve, L.M., Ehsani, A., Amarzguioui, M., Aagaard, L., Chen, Z.X., Riggs, A.D., Rossi, J.J., and Morris, K.V. (2006). The antisense strand of small interfering RNAs directs histone methylation and transcriptional gene silencing in human cells. *RNA* 12, 256–262.
- Klug, A. (2010). The discovery of zinc fingers and their applications in gene regulation and genome manipulation. *Annu. Rev. Biochem.* 79, 213–231.
- Hung, M.E., and Leonard, J.N. (2016). A platform for actively loading cargo RNA to elucidate limiting steps in EV-mediated delivery. *J. Extracell. Vesicles* 5, 31027.
- Peterson, C.W., Wang, J., Norman, K.K., Norgaard, Z.K., Humbert, O., Tse, C.K., Yan, J.J., Trimble, R.G., Shivak, D.A., Rebar, E.J., et al. (2016). Long-term multilineage engraftment of autologous genome-edited hematopoietic stem cells in nonhuman primates. *Blood* 127, 2416–2426.
- DiGiusto, D.L., Cannon, P.M., Holmes, M.C., Li, L., Rao, A., Wang, J., Lee, G., Gregory, P.D., Kim, K.A., Hayward, S.B., et al. (2016). Preclinical development and qualification of ZFN-mediated CCR5 disruption in human hematopoietic stem/progenitor cells. *Mol. Ther. Methods Clin. Dev.* 3, 16067.
- Mandell, J.G., and Barbas, C.F., 3rd (2006). Zinc Finger Tools: custom DNA-binding domains for transcription factors and nucleases. *Nucleic Acids Res.* 34, W516–W523.
- Bailus, B.J., Pyles, B., McAlister, M.M., O'Geen, H., Lockwood, S.H., Adams, A.N., Nguyen, J.T., Yu, A., Berman, R.F., and Segal, D.J. (2016). Protein Delivery of an Artificial Transcription Factor Restores Widespread Ube3a Expression in an Angelman Syndrome Mouse Brain. *Mol. Ther.* 24, 548–555.
- Rivenbark, A.G., Stolzenburg, S., Beltran, A.S., Yuan, X., Rots, M.G., Strahl, B.D., and Blancafort, P. (2012). Epigenetic reprogramming of cancer cells via targeted DNA methylation. *Epigenetics* 7, 350–360.
- Grimmer, M.R., Stolzenburg, S., Ford, E., Lister, R., Blancafort, P., and Farnham, P.J. (2014). Analysis of an artificial zinc finger epigenetic modulator: widespread binding but limited regulation. *Nucleic Acids Res.* 42, 10856–10868.
- Reynolds, L., Ullman, C., Moore, M., Isalan, M., West, M.J., Clapham, P., Klug, A., and Choo, Y. (2003). Repression of the HIV-1 5' LTR promoter and inhibition of HIV-1 replication by using engineered zinc-finger transcription factors. *Proc. Natl. Acad. Sci. USA* 100, 1615–1620.
- Wang, P., Qu, X., Wang, X., Zhu, X., Zeng, H., Chen, H., and Zhu, H. (2014). Specific reactivation of latent HIV-1 with designer zinc-finger transcription factors targeting the HIV-1 5'-LTR promoter. *Gene Ther.* 21, 490–495.
- Wang, X., Wang, P., Fu, Z., Ji, H., Qu, X., Zeng, H., Zhu, X., Deng, J., Lu, P., Zha, S., et al. (2015). Designed transcription activator-like effector proteins efficiently induced the expression of latent HIV-1 in latently infected cells. *AIDS Res. Hum. Retroviruses* 31, 98–106.
- Qu, X., Wang, P., Ding, D., Li, L., Wang, H., Ma, L., Zhou, X., Liu, S., Lin, S., Wang, X., et al. (2013). Zinc-finger-nucleases mediate specific and efficient excision of HIV-1

- proviral DNA from infected and latently infected human T cells. *Nucleic Acids Res.* 41, 7771–7782.
23. Deng, J., Qu, X., Lu, P., Yang, X., Zhu, Y., Ji, H., Wang, Y., Jiang, Z., Li, X., Zhong, Y., et al. (2017). Specific and Stable Suppression of HIV Provirus Expression In Vitro by Chimeric Zinc Finger DNA Methyltransferase 1. *Mol. Ther. Nucleic Acids* 6, 233–242.
  24. Bialek, J.K., Dunay, G.A., Voges, M., Schäfer, C., Spohn, M., Stucka, R., Hauber, J., and Lange, U.C. (2016). Targeted HIV-1 Latency Reversal Using CRISPR/Cas9-Derived Transcriptional Activator Systems. *PLoS ONE* 11, e0158294.
  25. Ji, H., Jiang, Z., Lu, P., Ma, L., Li, C., Pan, H., Fu, Z., Qu, X., Wang, P., Deng, J., et al. (2016). Specific Reactivation of Latent HIV-1 by dCas9-SunTag-VP64-mediated Guide RNA Targeting the HIV-1 Promoter. *Mol. Ther.* 24, 508–521.
  26. Wang, Z., Pan, Q., Gendron, P., Zhu, W., Guo, F., Cen, S., Wainberg, M.A., and Liang, C. (2016). CRISPR/Cas9-Derived Mutations Both Inhibit HIV-1 Replication and Accelerate Viral Escape. *Cell Rep.* 15, 481–489.
  27. Chavez, A., Scheiman, J., Vora, S., Pruitt, B.W., Tuttle, M., P R Iyer, E., Lin, S., Kiani, S., Guzman, C.D., Wiegand, D.J., et al. (2015). Highly efficient Cas9-mediated transcriptional programming. *Nat. Methods* 12, 326–328.
  28. Weinberger, L.S., Burnett, J.C., Toettcher, J.E., Arkin, A.P., and Schaffer, D.V. (2005). Stochastic gene expression in a lentiviral positive-feedback loop: HIV-1 Tat fluctuations drive phenotypic diversity. *Cell* 122, 169–182.
  29. Shrivastava, S., Charlins, P., Ackley, A., Embree, H., Dropulic, B., Akkina, R., Weinberg, M.S., and Morris, K.V. (2018). Stable Transcriptional Repression and Parasitism of HIV-1. *Mol. Ther. Nucleic Acids* 12, 12–18.
  30. Brena, C., Chipman, A.D., Minelli, A., and Akam, M. (2006). Expression of trunk Hox genes in the centipede *Strigamia maritima*: sense and anti-sense transcripts. *Evol. Dev.* 8, 252–265.
  31. Kim, Y.S., Kim, J.M., Jung, D.L., Kang, J.E., Lee, S., Kim, J.S., Seol, W., Shin, H.C., Kwon, H.S., Van Lint, C., et al. (2005). Artificial zinc finger fusions targeting Sp1-binding sites and the trans-activator-responsive element potentially repress transcription and replication of HIV-1. *J. Biol. Chem.* 280, 21545–21552.
  32. Segal, D.J., Gonçalves, J., Eberhardy, S., Swan, C.H., Torbett, B.E., Li, X., and Barbas, C.F., 3rd (2004). Attenuation of HIV-1 replication in primary human cells with a designed zinc finger transcription factor. *J. Biol. Chem.* 279, 14509–14519.
  33. Guo, R., Wang, H., Cui, J., Wang, G., Li, W., and Hu, J.F. (2015). Inhibition of HIV-1 Viral Infection by an Engineered CRISPR Csy4 RNA Endoribonuclease. *PLoS ONE* 10, e0141335.
  34. Dampier, W., Nonnemacher, M.R., Sullivan, N.T., Jacobson, J.M., and Wigdahl, B. (2014). HIV Excision Utilizing CRISPR/Cas9 Technology: Attacking the Proviral Quasispecies in Reservoirs to Achieve a Cure. *MOJ Immunol.* 1, 00022.
  35. Ebina, H., Misawa, N., Kanemura, Y., and Koyanagi, Y. (2013). Harnessing the CRISPR/Cas9 system to disrupt latent HIV-1 provirus. *Sci. Rep.* 3, 2510.
  36. Kaminski, R., Bella, R., Yin, C., Otte, J., Ferrante, P., Gendelman, H.E., Li, H., Booze, R., Gordon, J., Hu, W., and Khalili, K. (2016). Excision of HIV-1 DNA by gene editing: a proof-of-concept in vivo study. *Gene Ther.* 23, 690–695.
  37. Wang, G., Zhao, N., Berkhout, B., and Das, A.T. (2016). CRISPR-Cas9 can inhibit HIV-1 replication but NHEJ repair facilitates virus escape. *Mol. Ther.* 24, 522–526.
  38. Yin, C., Zhang, T., Qu, X., Zhang, Y., Putatunda, R., Xiao, X., Li, F., Xiao, W., Zhao, H., Dai, S., et al. (2017). In Vivo Excision of HIV-1 Provirus by saCas9 and Multiplex Single-Guide RNAs in Animal Models. *Mol. Ther.* 25, 1168–1186.
  39. Frank, T.D., Carter, A., Jahagirdar, D., Bieh, M.H., Douwes-Schultz, D., Larson, S.L., Arora, M., Dwyer-Lindgren, L., Steuben, K.M., Abbastabar, H., et al.; GBD 2017 HIV collaborators (2019). Global, regional, and national incidence, prevalence, and mortality of HIV, 1980–2017, and forecasts to 2030, for 195 countries and territories: a systematic analysis for the Global Burden of Diseases, Injuries, and Risk Factors Study 2017. *Lancet HIV* 6, e831–e859.
  40. Clutton, G.T., and Jones, R.B. (2018). Diverse Impacts of HIV Latency-Reversing Agents on CD8+ T-Cell Function: Implications for HIV Cure. *Front. Immunol.* 9, 1452.
  41. Liu, T., Zhang, L., Joo, D., and Sun, S.-C. (2017). NF- $\kappa$ B signaling in inflammation. *Signal Transduct. Target. Ther.* 2, 17023.
  42. Kumar, R., Yong, Q.C., and Thomas, C.M. (2013). Do multiple nuclear factor kappa B activation mechanisms explain its varied effects in the heart? *Ochsner J.* 13, 157–165.
  43. Brown, B.D., Venneri, M.A., Zingale, A., Sergi Sergi, L., and Naldini, L. (2006). Endogenous microRNA regulation suppresses transgene expression in hematopoietic lineages and enables stable gene transfer. *Nat. Med.* 12, 585–591.
  44. Tian, J., An, X., and Niu, L. (2017). Role of microRNAs in cardiac development and disease. *Exp. Ther. Med.* 13, 3–8.
  45. Yang, P., Wang, Y., and Macfarlan, T.S. (2017). The Role of KRAB-ZFPs in Transposable Element Repression and Mammalian Evolution. *Trends Genet.* 33, 871–881.
  46. Nishitsuji, H., Sawada, L., Sugiyama, R., and Takaku, H. (2015). ZNF10 inhibits HIV-1 LTR activity through interaction with NF- $\kappa$ B and Sp1 binding motifs. *FEBS Lett.* 589, 2019–2025.
  47. Wang, J., Wang, J., and Tian, C.Y. (2016). Evolution of KRAB-containing zinc finger proteins and their roles in species evolution. *Yi Chuan* 38, 971–978.
  48. Shan, L., Deng, K., Shroff, N.S., Durand, C.M., Rabi, S.A., Yang, H.C., Zhang, H., Margolick, J.B., Blankson, J.N., and Siliciano, R.F. (2012). Stimulation of HIV-1-specific cytolytic T lymphocytes facilitates elimination of latent viral reservoir after virus reactivation. *Immunity* 36, 491–501.
  49. Mylvaganam, G., Yanez, A.G., Maus, M., and Walker, B.D. (2019). Toward T Cell-Mediated Control or Elimination of HIV Reservoirs: Lessons From Cancer Immunology. *Front. Immunol.* 10, 2109.
  50. Kim, G.B., Hege, K., and Riley, J.L. (2019). CAR Talk: How Cancer-Specific CAR T Cells Can Instruct How to Build CAR T Cells to Cure HIV. *Front. Immunol.* 10, 2310.
  51. Liang, C., Wainberg, M.A., Das, A.T., and Berkhout, B. (2016). CRISPR/Cas9: a double-edged sword when used to combat HIV infection. *Retrovirology* 13, 37.
  52. Saayman, S., Ali, S.A., Morris, K.V., and Weinberg, M.S. (2015). The therapeutic application of CRISPR/Cas9 technologies for HIV. *Expert Opin. Biol. Ther.* 15, 819–830.
  53. Villamizar, O., Waters, S.A., Scott, T., Saayman, S., Grepo, N., Urak, R., Davis, A., Jaffe, A., and Morris, K.V. (2019). Targeted Activation of Cystic Fibrosis Transmembrane Conductance Regulator. *Mol. Ther.* 27, 1737–1748.
  54. Kojima, R., Bojar, D., Rizzi, G., Hamri, G.C., El-Baba, M.D., Saxena, P., Ausländer, S., Tan, K.R., and Fussenegger, M. (2018). Designer exosomes produced by implanted cells intracerebrally deliver therapeutic cargo for Parkinson's disease treatment. *Nat. Commun.* 9, 1305.
  55. Jeeninga, R.E., Hoogenkamp, M., Armand-Ugon, M., de Baar, M., Verhoef, K., and Berkhout, B. (2000). Functional differences between the long terminal repeat transcriptional promoters of human immunodeficiency virus type 1 subtypes A through G. *J. Virol.* 74, 3740–3751.
  56. Klaver, B., and Berkhout, B. (1994). Comparison of 5 $\frac{1}{2}$  and 3 $\frac{1}{2}$  long terminal repeat promoter function in human immunodeficiency virus. *J. Virol.* 68, 3830–3840.
  57. Jordan, A., Bisgrove, D., and Verdin, E. (2003). HIV reproducibly establishes a latent infection after acute infection of T cells in vitro. *EMBO J.* 22, 1868–1877.
  58. Roberts, T.C., Hart, J.R., Kaikkonen, M.U., Weinberg, M.S., Vogt, P.K., and Morris, K.V. (2015). Quantification of nascent transcription by bromouridine immunocapture nuclear run-on RT-qPCR. *Nat. Protoc.* 10, 1198–1211.
  59. Saayman, S., Ackley, A., Turner, A.W., Famiglietti, M., Bosque, A., Clemson, M., Planelles, V., and Morris, K.V. (2014). An HIV-encoded antisense long noncoding RNA epigenetically regulates viral transcription. *Mol. Ther.* 22, 1164–1175.
  60. Saayman, S.M., Ackley, A., Burdach, J., Clemson, M., Gruenert, D.C., Tachikawa, K., Chivukula, P., Weinberg, M.S., and Morris, K.V. (2016). Long Non-coding RNA BGas Regulates the Cystic Fibrosis Transmembrane Conductance Regulator. *Mol. Ther.* 24, 1351–1357.
  61. Ewels, P.A., Peltzer, A., Fillinger, S., Alneberg, J., Patel, H., Wilm, A., Garcia, M.U., Tommaso, P.D., and Nahnsen, S. (2019). nf-core: Community-curated bioinformatics pipelines. *bioRxiv*. <https://doi.org/10.1101/610741>.
  62. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29, 15–21.

63. Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* 30, 923–930.
64. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* 15, 550.
65. Varet, H., Brillet-Guéguen, L., Coppée, J.-Y., and Dillies, M.-A. (2016). SARTools: A DESeq2- and EdgeR-Based R Pipeline for Comprehensive Differential Analysis of RNA-Seq Data. *PLoS ONE* 11, e0157022.
66. Yu, G., Wang, L.-G., Han, Y., and He, Q.-Y. (2012). clusterProfiler: an R package for comparing biological themes among gene clusters. *OMICS* 16, 284–287.