

POSTER PRESENTATION

Open Access

# An evaluation of outcomes, interventions and behaviour modifications following a cardiovascular MRI in HIV infected patients

N Aftab\*, C Doherty

From International Conference for Healthcare and Medical Students (ICHAMS) 2013  
Dublin, Ireland. 11-12 October 2013

## Background

The advent of antiretroviral therapy has led to increased life span which has been associated with an increase in cardiovascular risk factors and heart disease (Boccara F et al, 2013 [1]). Cardiovascular risk assessment is now advised for those with HIV. The field of behavioural cardiology is particularly applicable in HIV therapy due to the long term nature of such a condition. Therefore, it's useful to assess behaviour modifications following comprehensive cardiac evaluation, specifically CMRI.

## Methods

This is a cross-sectional study examining the medical outcomes, interventions and behaviour modifications in a cohort of 184 male (169 HIV positive compared to 21 HIV negative controls), HIV positive patients at the GUIDE Clinic. The purpose of the study was to compare these variables in 148 patients who underwent a cardiac MRI (CMRI) versus 36 patients who underwent a Framingham risk assessment. The initial study involved the identification of cardiac abnormalities by blood tests and CMRI scan, while the follow-up involved changes in lifestyle, medications as well as cardiac relevant events.

## Results

When comparing the two groups, 8.8% of the CMRI group quit smoking while only 2.8% of the Framingham group did. Of the CMRI group, 7.4% commenced new cardiac medication while commencement in the Framingham group was 8.3%. Of the CMRI group 16.2% had further cardiac investigations after their CMRI compared to 8.3% in the Framingham group.

Trinity College Dublin and The Guide Clinic, St. James Hospital, Dublin 8.  
Ireland

## Conclusions

Having a cardiac MRI seems to be more effective in producing behavioural modifications as well as more precise medical diagnoses in HIV positive patients. Ideally, Framingham Risk Assessments should be altered to include HIV specific CVD risk factors; however this model does not yet exist. Due to the increased CVD risk factors seen in HIV positive patients, cardiac assessments need to be performed more routinely with a lower threshold for further cardiac investigations.

Published: 14 January 2015

## Reference

1. Boccara F, et al: HIV and Coronary Heart Disease: Time for a Better Understanding. *Journal of the American College of Cardiology* 2013, 61(5):511-523.

doi:10.1186/1753-6561-9-S1-A30

Cite this article as: Aftab and Doherty: An evaluation of outcomes, interventions and behaviour modifications following a cardiovascular MRI in HIV infected patients. *BMC Proceedings* 2015 **9**(Suppl 1):A30.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

