


International Validation of a Novel PEACE Scale to Improve the Quality of Upper Gastrointestinal Mucosal Inspection During Endoscopy

Marcin Romańczyk, MD, PhD^{1,2,3}, Madhav Desai, MD⁴, Michal F. Kamiński, MD, PhD^{5,6,7}, Seiichiro Abe, MD, PhD⁸, Asma A. Alkandari, MD⁹, Torsten Beyna, MD, PhD¹⁰, Raf Bisschops, MD, PhD¹¹, Krzysztof Budzyń, MD^{1,2,3}, Monika Bugdol, PhD¹², Samir C. Grover, MD¹³, C. Prakash Gyawali, MD, FACP¹⁴, Haruhiro Inoue, MD, PhD¹⁵, Prasad G. Iyer, MD, FACP¹⁶, Helmut Messmann, MD, PhD¹⁷, Krish Ragunath, MD^{18,19}, Yutaka Saito, MD, PhD⁸, Sachin Srinivasan, MD²⁰, Christopher Teshima, MD, PhD²¹, Rena Yadlapati, MD, FACP²², Cesare Hassan, MD, PhD^{23,24} and Prateek Sharma, MD, FACP^{20,25}

PEACE scale's international validation

- 16 expert endoscopists (6 North America, 6 Europe and Middle East, 4 Japan and Australia)
- Teaching with 21 photographs, 4 example videos, 4 test videos
- Assessment of 39 videos



Romańczyk et al. *Clin Trans Gastroenterol.* 2024. doi: 10.14309/ctg.0000000000000786

Clinical and Translational
GASTROENTEROLOGY

¹Department of Gastroenterology, Faculty of Medicine, Academy of Silesia, Katowice, Poland; ²Department of Gastroenterology and Hepatology, School of Medicine in Katowice, Medical University of Silesia, Katowice, Poland; ³Endoterapia, H-T, Centrum Medyczne, Tychy, Poland; ⁴Interventional Gastroenterology, Department of Surgery, University of Texas Health Science Center, Houston, Texas, USA; ⁵Department of Gastroenterology, Hepatology and Clinical Oncology, Centre of Postgraduate Medical Education, Warsaw, Poland; ⁶Department of Oncological Gastroenterology, Maria Skłodowska-Curie National Research Institute of Oncology, Warsaw, Poland; ⁷Institute of Health and Society, University of Oslo, Oslo, Norway; ⁸Endoscopy Division, National Cancer Center Hospital, Tokyo, Japan; ⁹Al Jahra Hospital, Kuwait, Kuwait; ¹⁰Department of Internal Medicine, Evangelisches Krankenhaus Düsseldorf, Düsseldorf, Germany; ¹¹Department of Gastroenterology and Hepatology, Katholieke Universiteit Leuven, Leuven, Belgium; ¹²Department of Medical Informatics and Artificial Intelligence, Faculty of Biomedical Engineering, Silesian University of Technology, Gliwice, Poland; ¹³Division of Gastroenterology and Hepatology, Scarborough Health Network Research Institute, University of Toronto, Toronto, Ontario, Canada; ¹⁴Division of Gastroenterology, Washington University School of Medicine, St. Louis, Missouri, USA; ¹⁵Digestive Disease Centre, Showa University Koto Toyosu Hospital, Tokyo, Japan; ¹⁶Division of Gastroenterology and Hepatology, Mayo Clinic, Rochester, Minnesota, USA; ¹⁷Department of Gastroenterology, Faculty of Medicine, University of Augsburg, Augsburg, Germany; ¹⁸Faculty of Health Sciences, Curtin Medical School, Curtin University, Perth, Western Australia, Australia; ¹⁹Department of Gastroenterology, Royal Perth Hospital, Perth, Western Australia, Australia; ²⁰Division of Gastroenterology, Kansas City VA Medical Center, Kansas City, Missouri, USA; ²¹Center for Advanced Therapeutic Endoscopy and Endoscopic Oncology, Division of Gastroenterology and Hepatology, Department of Medicine, St Michael's Hospital, Toronto, Ontario, Canada; ²²Division of Gastroenterology, University of California San Diego, San Diego, California, USA; ²³Department of Biomedical Sciences, Humanitas University, Milan, Italy; ²⁴Endoscopy Unit, IRCCS Humanitas Research Hospital, Milan, Italy; ²⁵Division of Gastroenterology and Hepatology, University of Kansas School of Medicine, Kansas City, Kansas, USA. **Correspondence:** Marcin Romańczyk, MD, PhD. E-mail: marcin.romanczyk@htcentrum.pl.

Received October 1, 2024; accepted October 25, 2024; published online November 14, 2024

© 2024 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of The American College of Gastroenterology

- INTRODUCTION:** The performance of a high quality esophagogastroduodenoscopy (EGD) is dependent on the mucosal cleanliness. Recently, the Polprep: Effective Assessment of Cleanliness in EGD (PEACE) scale was created to assess the degree of mucosal cleanliness during EGD. The aim of this study was to validate this scoring system in a cohort of international endoscopists.
- METHODS:** In total, 39 EGD videos, with different degrees of mucosal cleanliness were retrieved from a previously conducted prospective trial. All experts rated the cleanliness of the mucosa on each video using the PEACE scale. To evaluate agreement of all scores (0–3), intraclass correlation coefficient 2.1 was used. The agreement on adequate (scores 2 and 3) and inadequate (scores 0 and 1) cleanliness was assessed using kappa values.
- RESULTS:** Videos evaluating esophagus, stomach, and duodenum cleanliness were reviewed by 16 endoscopists. The PEACE scores demonstrated good agreement (intraclass correlation coefficient 0.82, 95% CI 0.75–0.89), especially for esophagus (0.84; 95% CI 0.71–0.95) and stomach (0.81; 95% CI 0.69–0.91), while agreement was moderate for the duodenum (0.69; 95% CI 0.51–0.87). The agreement was similar between Eastern (0.86; 95% CI 0.79–0.92) and Western experts (0.80; 95% CI 0.72–0.88). Similarly, agreement regarding adequate cleanliness was comparable between Eastern (0.70; 95% CI 0.55–0.85) and Western (0.74; 95% CI 0.64–0.84) endoscopists being overall 0.75 (95% CI 0.65–0.85).
- DISCUSSION:** The PEACE scoring system is a simple and reliable scale to assess the cleanliness during EGD. The score is now validated among international experts with high concordance, justifying its use in clinical practice.

KEYWORDS: esophagogastroduodenoscopy; cleanliness; quality indicators

Clinical and Translational Gastroenterology 2025;16:e00786. <https://doi.org/10.14309/ctg.0000000000000786>

INTRODUCTION

Endoscopic procedures have become indispensable tools in the evaluation and management of gastrointestinal diseases. Technological progress has allowed for rapid improvements for both optical diagnosis and therapeutic procedures in the field of gastrointestinal endoscopy. For instance, high-definition endoscopes have replaced standard definition endoscopes in modern endoscopy (1,2). Dye-based and electronic chromoendoscopy are essential modalities for detection and evaluation of neoplastic lesions such as gastric neoplasm, gastric intestinal metaplasia, Barrett esophagus, and related neoplasia and esophageal squamous cell neoplasia (2–5). To achieve high quality diagnostic esophagogastroduodenoscopy (EGD) and to use high-definition electronic chromoendoscopy, a clean mucosa of the area of interest is required. Poor mucosal visibility during EGD could be due to various factors including the formation of bubbles, food residue, bile etc. However, this also requires efforts from the endoscopist to clear the contents and clean the mucosa to the best of their ability. To achieve mucosal cleaning, several agents have been evaluated (6–8). The preparation with simethicone/dimethicone and pronase has become the standard in EGD screening in high prevalence of gastric cancers areas (9).

In colonoscopy, where adequate preparation has been evaluated for many years, proper preparation and achieving adequate visibility is a crucial quality indicator (10). A higher clean bowel mucosal score (such as using the Boston Bowel Preparation Scale [BBPS]) has been shown to be associated with improved adenoma detection rate (11). Despite the conceived benefit of a clean mucosal inspection in the upper gastrointestinal (UGI) tract for a higher yield, there is lack of research in this area. Recently, we

proposed a novel UGI cleanliness scale—PEACE (Polprep: Effective Assessment of Cleanliness in EGD) (12,13). We showed that adequate cleanliness was associated with higher detection of pathologies in the UGI tract. In a retrospective feasibility study and subsequent prospective study, it was also shown that inadequate preparation was showed to be associated with lower detection of clinically significant UGI lesions. We also presented good interobserver and intraobserver agreement of scoring based on the still endoscopic images assessment (12). In this study we aimed to assess the interobserver agreement of the PEACE cleanliness scores among an international cohort of expert endoscopists using videos assessment.

METHODS

The study was conducted as post hoc assessment of EGD videos that were collected during the period of prospective validation study (August 2021 to October 2022) (13). The EGDs were performed using EVIS EXERA III and X1 systems (Olympus Europe, Hamburg, Germany) with GIF-EZ1500, GIF-1100, GIF-HQ190, GIF-H180J, and GIF-Q160Z endoscopes and Fujifilm EP-6000 system (Fujifilm Europe, Düsseldorf, Germany) with EG-760R endoscopes. The study was performed in accordance with the Declaration of Helsinki and conducted under Institutional Review Board agreement (No. PCN/CBN/0052/KB/15/23).

To assess the reliability of the scale, we decided to assess videos as it reflects real assessment of the cleanliness of the mucosa. We invited international endoscopy experts from different geographic regions to simulate different fashions of performing endoscopy around the world. The selected method was non-extensive training explained below followed by a single round




Figure 1. Example of PEACE scores.

assessment of videos to avoid very isolated study conditions that could not be transferred into real-world scenario.

Teaching materials and training phase

To familiarize the assessors with the PEACE scale, 21 representative and example endoscopic pictures were provided (12). In addition, 4 videos with different grades of the PEACE score were sent to the assessors. An overview of the PEACE scale along with clear instructions was provided to the assessors on how the rating should be performed:

The scoring for the PEACE scale—see Figure 1:

- 3—clean mucosa or minor amounts of transparent fluid not impeding mucosal inspection,
- 2—small amount of hazy fluid/foamy/solid content enabling inspection of most of the mucosa,
- 1—substantial amount of opaque fluid/foamy/solid content that does not allow evaluation of some parts of the mucosa,
- 0—substantial amount of fluid/foamy/content completely obscuring evaluation of the mucosa.

Instructions on how to evaluate the scale:

Authors who selected and edited the videos as well as the senior author did not participate in the assessment. To avoid significant selection bias, the author who edited the videos did not perform any EGD during the study period.




Figure 2. Study flowchart.

The overview contained information how the rating should be performed:

- Assessment of the 3 segments: esophagus, stomach, duodenum,
- Each segment scored 0–3,
- Assessment of each segment after cleansing (irrigation, suction),
- The score for each segment was the lowest score within each component of that segment (e.g. If the proximal esophagus is scored as 2, and distal esophagus scored as 3—then the score for esophagus is 2).

Finally, all assessors were asked to grade 4 test videos and the correct results were provided to them, so they could evaluate their ability to use the PEACE scale.

Video selection

The EGD videos used for the interobserver agreement study were selected from adult symptomatic patients referred for diagnostic EGD and screened for the study (13). Videos with poor technical quality of recording (such as low resolution or disturbances of images) were excluded. The videos were selected according to scoring of specific segments during the study period to provide a representation of all scores (0–3) after cleansing. Then recordings were cropped to 20–40 seconds parts representative for each segment (esophagus, stomach, and duodenum). The whole process has been summarized at Figure 2.

Sample size and statistical analysis

To achieve intraclass correlation coefficient (ICC) of 0.7 (95% CI 0.6–0.8), the minimal number of videos and assessors was 39 and 15, respectively. To evaluate interobserver agreement of all PEACE scale scores (0–3), ICC 2.1 was used. ICC values 0.5–0.75 were considered as moderate, 0.75–0.9 as good, and >0.9 as excellent (14). For the agreement of inadequate (scores 0 and 1) and adequate cleanliness (scores 2 and 3), the Conger kappa value was

calculated. Kappa values of 0.33–0.5 were considered as fair, 0.5–0.67 as moderate, 0.67–0.9 as substantial, and over 0.9 as almost perfect (15). Statistical analysis was performed in R version 4.2.3, in the RStudio environment 2023.06.0 + 421, using packages tidyverse, irr, and irrCAC.

RESULTS

Assessors and videos

A total of 39 videos were selected for assessment by 16 international experts—6 from North America (United States and Canada), 6 from Europe and Middle East, and 4 from Japan and Australia.

Scoring and reliability of PEACE scores

Of 39 videos, 13 were scored 3, 10 were scored 2, 14 were scored 1, and 2 were scored 0. Five of 39 videos (12.8%) were scored equally by all experts, 22 (56.4%) were scored within one range, 9 (23.1%) within 2, and in 3 cases (7.7%) the range was 3. The scoring has been summarized in Figure 3.

The overall reliability, based on ICC values, exceeded the assumed threshold, and was 0.82. For specific segments, the ICC was as follows: 0.84 for esophagus, 0.81 for stomach, and 0.69 for duodenum—see Table 1. The overall mean range of scores was 1.26, being highest for duodenum (1.42), followed by stomach (1.41) and esophagus (0.8).

The accordance of identification of inadequate cleanliness

The identification of adequate cleanliness (PEACE scores 2 and 3 = adequate cleanliness) was characterized with substantial agreement—kappa value of 0.75. The agreement was substantial in stomach and esophagus and moderate in duodenum (kappa value highest in stomach [0.77], then in esophagus [0.68] and lowest in duodenum [0.55]). These data have been summarized in Table 2.




Figure 3. Scores of assessed videos. The figure presents scorings of each video. Each vertical line represents video. Each dot represents one assessment. Cases not colored represents videos where concordance was 100%, green color represents videos with scores range of 1 score, orange with range 2 scores, and red with range of 3 scores.

Accordance of scores between Eastern and Western experts

The agreement among Japanese and Australian experts in comparison with European (including Middle East), United States, and Canadian experts was not significantly different—being good in both cases (ICC 0.86; 95% CI 0.79–0.92 and ICC 0.80; 95% CI 0.72–0.88, respectively). These data have been presented in Figure 4. The mean range of scores among Western experts was higher than among Eastern experts (1.15 and 0.54, respectively).

The accordance in identification of adequate and inadequate cleanliness was also not significantly different among Western experts (kappa value of Western experts 0.74; 95% CI 0.64–0.84 and kappa value of Eastern experts 0.70; 95% CI 0.55–0.85)—see Figure 5.

DISCUSSION

In this international validation study, we presented interobserver agreement of a recently created and validated PEACE scale used

for the assessment of UGI mucosal inspection and cleanliness among international endoscopy experts. We showed that despite heterogeneity of UGI tract segments and no extensive teaching, the assessment of cleanliness during EGD with PEACE scale was reliable. We also demonstrated that agreement for the scale was overall high for subgroups of Western and Eastern Experts. The good interobserver agreement may most likely be related to the impact of the scale design, which was created based on the widely used Boston Bowel Preparation Scale (16).

Our previous retrospective feasibility study of PEACE scale performance demonstrated good interobserver and intraobserver reproducibility of photographs assessment (ICC 0.8 and the Fleiss Kappa value 0.64, respectively) [9]. In this study, the interobserver agreement of UGI cleanliness assessment based on the evaluation of videos was even slightly higher. It was higher in the stomach and esophagus than in the duodenum. In previous study, on photographs, the difference was not seen. That could be

Table 1. Interobserver agreement analysis of PEACE scores (scores 0–3)

Variable	Intraclass correlation coefficient	95% CI
All scores (N = 39)	0.82	0.75–0.89
Esophagus (N = 10)	0.84	0.71–0.95
Stomach (N = 17)	0.81	0.69–0.91
Duodenum (N = 12)	0.69	0.51–0.87

The table presents interobserver agreement in the assessment of all videos and particular segments (esophagus, stomach, and duodenum). 95% CI, 95% confidence interval; PEACE, Polprep: Effective Assessment of Cleanliness in Esophagogastrroduodenoscopy.

Table 2. Interobserver agreement analysis of inadequate and adequate cleanliness according to PEACE scores

Variable	Conger's kappa value	95% CI
All scores (N = 39)	0.75	0.65–0.85
Esophagus (N = 10)	0.68	0.23–1
Stomach (N = 17)	0.77	0.62–0.93
Duodenum (N = 12)	0.55	0.32–0.79

The table presents interobserver agreement in the assessment of inadequate (PEACE scores 0 and 1) and adequate (PEACE scores 2 and 3) cleanliness of all videos and particular segments (esophagus, stomach, and duodenum). 95% CI, 95% confidence interval; PEACE, Polprep: Effective Assessment of Cleanliness in Esophagogastrroduodenoscopy.




Figure 4. PEACE scores agreement. The figure represents intraclass correlation coefficient for all raters, among Western raters (United States, Canada, and Europe with Middle East) and Eastern (Japan and Australia) raters for videos of the esophagus, stomach, and duodenum and for all videos. Each vertical line on the plot represents 95% confidence interval, and dots represent intraclass correlation coefficient values. PEACE, Polprep: Effective Assessment of Cleanliness in Esophagogastroduodenoscopy.

explained by different appearance of duodenal bulb and second part of duodenum with partially blind spot between them resulting in some differences in the assessment not observed at still photographs assessment. The strength of our study is the assessment of prospectively acquired video records by world class experts. An important aspect was the participation of experts from North America, Europe, Middle East, Japan, and Australia. The results show that despite differences in the process of

performing routine EGD, the agreement on cleanliness assessment by PEACE remains similar among different parts of the globe.

Although the topic of adequate mucosal cleanliness during EGD is of great clinical importance, no reliable tool allowing the identification of inadequately prepared patients for EGD is in common use. This issue is well accepted in colonoscopy where inadequate cleanliness is related not only with lower detection of




Figure 5. Adequate and inadequate cleanliness agreement based on the PEACE scores. The figure represents Conger kappa value for the agreement on adequate (2 and 3 PEACE scores) and inadequate cleanliness (0 and 1 PEACE scores) for all raters, among Western raters (United States, Canada, and Europe with Middle East) and Eastern (Japan and Australia) raters for videos of the esophagus, stomach, and duodenum and for all videos. Each vertical line on the plot represents 95% confidence interval, and dots represent Conger kappa values. PEACE, Polprep: Effective Assessment of Cleanliness in Esophagogastroduodenoscopy.

adenomas but also with lower rate of cecum intubation and higher discomfort felt by patients (17–19). The high reliability of the PEACE scale would allow to identify patients who require a repeat EGD due to inadequate cleanliness based on our previous study with any segment scored as 0 or 1 (12,13). This is important for a high quality EGD, as it has been well documented that there is a miss rate of neoplastic lesions (5–10%) in the UGI tract, and only a few risk factors of missed neoplasia have been identified so far (20–22). Identification of inadequately prepared individuals during all diagnostic EGDs could possibly result in better risk stratification by identification of precancerous conditions and lower miss rate of early neoplasia, especially in high-risk groups such as advanced atrophic gastritis or long Barrett esophagus. The degree of cleanliness should be assessed in the future on pre-neoplastic conditions (Barrett esophagus and advanced atrophic gastritis) and neoplastic lesions detection. In addition, the degree of demanded cleanliness needs to be assessed in the cohort of high-risk groups of patients with preneoplastic conditions during surveillance examinations. It should be also assessed along with other quality measures such as examination time. It should be mentioned that not only the identification of inadequately clean segments (scores 0 and 1) but also the assessment of full scale (scores 0–3) was associated with a high accordance.

The Toronto Upper Gastrointestinal Cleaning Score (TUGCS) was recently reported, supporting our work that scales for mucosal cleanliness of the foregut can be developed with strong validity evidence. However, the TUGCS has the notable limitations of having excluded the esophagus from assessment and having potentially introduced unnecessary complexity by having the stomach subgrouped into fundus, body, and antrum (23). It may be argued that assessing the entire stomach with one score maybe difficult. However, endoscopists are currently assessing the entire transverse colon or left colon with a single score as well. In addition, our study shows that the interobserver agreement of the stomach score was very good. Recently, the Barcelona scale was introduced also where 5 landmarks are scored from 0 to 2 (24). However, we find easy clinical implementation by similar scoring to colonoscopy scales and simple judgment of 3 investigated organs, as most important aspect of the UGI cleanliness scale.

Despite the ease and validity of the PEACE scale, our study has limitations. First, the videos were rated by international endoscopy experts and cannot be directly generalizable to the community endoscopists. This needs to be done in a future study along with assessment of degree of cleanliness on neoplastic lesions detection. Since the PEACE scale is simple to follow, does not require specific training or learning curve, and can be readily adopted without prolonging the procedure time, we do believe, that it could be adopted in clinical practice by the general gastroenterology community. Third, the PEACE scale was designed similar to the scoring of the BBPS and its utility might be lower in endoscopy units where BBPS is not routinely used. We have not measured the operator's performance that could affect the final result. In addition, we cannot exclude potential videos selection bias and bias caused by videos' cropping.

In conclusion, the interobserver agreement on UGI tract cleanliness assessment using the PEACE scale by a group of expert international endoscopists was good. This could be achieved with minimal training on the scoring system, allowing for a broad clinical implementation on this easy-to-use scale.

CONFLICTS OF INTEREST

Guarantor of the article: Marcin Romańczyk, MD, PhD.

Specific author contributions: Conception and design: M.R., M.D., P.S. Analysis and interpretation of the data: M.R., P.S., M.D., M.F.K., S.A., A.A.A., T.B., R.B., K.B., M.B., S.C.G., P.G., H.I., P.G.I., H.M., K.R., Y.S., S.S., C.T., R.Y., C.H. Drafting of the article: M.R., M.D., P.S., M.B. Critical revision of the article for important intellectual content: M.F.K., S.A., A.A.A., T.B., R.B., K.B., S.C.G., P.G., H.I., P.G.I., H.M., K.R., Y.S., S.S., C.T., R.Y., C.H.; Final approval of the article: M.R., P.S., M.D., M.F.K., S.A., A.A.A., T.B., R.B., K.B., M.B., S.C.G., P.G., H.I., P.G.I., H.M., K.R., Y.S., S.S., C.T., R.Y., C.H.

Financial support: None to report.

Potential competing interests: Samir C. Grover—equity and ownership interest in Volo Healthcare; research grants from Olympus; consulting fees and honoraria: Amgen, BioJAMP, Pfizer, and Sanofi, and AbbVie; education grants: Fresenius Kabi, BioJAMP, Celltrion, Takeda, Pfizer. Rena Yadlapati—consultant for Medtronic, Phathom Pharmaceuticals, StatLinkMD, Reckitt Benckiser Healthcare Ltd, Medscape; Research Support: Ironwood Pharmaceuticals; Advisory Board with Stock Options: RJS Mediagnostix. Christopher Teshima—consultant—Boston Scientific; speaker—Medtronic. Prasad G. Iyer—Research funding: Exact Sciences, Pentax Medical, CDx Medical, Castle Biosciences; Consultant: Exact Sciences, Pentax Medical, Medtronic, CDx Medical, Castle Biosciences. Raf Bisschops—Research grant, advisory, and consultancy fees: Pentax Medical, Fujifilm. Prateek Sharma—Consultant: Boston Scientific, Olympus Inc; Grant support: US Endoscopy, Medtronic, Fujifilm, Ironwood, Cosmo pharmaceuticals, Erbe. Michał Filip Kamiński—Consultant: Olympus, Erbe speaker: Boston Scientific, Ipsen, Recordati; Grant support: Olympus. Rest of authors—nothing to declare.

Study Highlights

WHAT IS KNOWN

- ✓ Detection of clinically significant lesions during upper gastrointestinal endoscopy depends on the degree of cleanliness.
- ✓ Until now, no scale assessing cleanliness has been properly validated.

WHAT IS NEW HERE

- ✓ Polprep: Effective Assessment of Cleanliness in Esophagogastroduodenoscopy scale allows for precise assessment of upper gastrointestinal cleanliness without extensive training.
- ✓ Its performance is unequivocal regardless the geographical distribution of endoscopists.

REFERENCES

1. Weusten BLAM, Bisschops R, Coron E, et al. Endoscopic management of Barrett's esophagus: European Society of Gastrointestinal Endoscopy (ESGE) position statement. *Endoscopy* 2017;49(2):191–8.
2. Pimentel-Nunes P, Libânio D, Marcos-Pinto R, et al. Management of epithelial precancerous conditions and lesions in the stomach (MAPS II): European Society of Gastrointestinal Endoscopy (ESGE), European Helicobacter and Microbiota Study Group (EHMSG), European Society of Pathology (ESP), and Sociedade Portuguesa de Endoscopia Digestiva (SPED) guideline update 2019. *Endoscopy* 2019;51(04):365–88.
3. Sharma P, Bergman JJGHM, Goda K, et al. Development and validation of a classification system to identify high-grade dysplasia and esophageal

- adenocarcinoma in Barrett's esophagus using narrow-band imaging. *Gastroenterology* 2016;150(3):591–8.
4. Ishihara R, Arima M, Iizuka T, et al. Endoscopic submucosal dissection/ endoscopic mucosal resection guidelines for esophageal cancer. *Dig Endosc* 2020;32(4):452–93.
 5. Esposito G, Pimentel-Nunes P, Angeletti S, et al. Endoscopic grading of gastric intestinal metaplasia (EGGIM): A multicenter validation study. *Endoscopy* 2019;51(6):515–21.
 6. Li Y, Du F, Fu D. The effect of using simethicone with or without N-acetylcysteine before gastroscopy: A meta-analysis and systemic review. *Saudi J Gastroenterol* 2019;25(4):218–28.
 7. Burke E, Harkins P, Moriarty F, et al. Does premedication with mucolytic agents improve mucosal visualization during oesophagogastroduodenoscopy: A systematic review and meta-analysis. *Surg Res Pract* 2021;2021:1570121.
 8. Wu L, Cao Y, Liao C, et al. Systematic review and meta-analysis of randomized controlled trials of Simethicone for gastrointestinal endoscopic visibility. *Scand J Gastroenterol* 2011;46(2):227–35.
 9. Yao K, Uedo N, Kamada T, et al. Guidelines for endoscopic diagnosis of early gastric cancer. *Dig Endosc* 2020;32(5):663–98.
 10. Kaminski MF, Thomas-Gibson S, Bugajski M, et al. Performance measures for lower gastrointestinal endoscopy: A European Society of Gastrointestinal Endoscopy (ESGE) quality improvement initiative. *United Eur Gastroenterol J* 2017;5(3):309–34.
 11. Sulz MC, Kröger A, Prakash M, et al. Meta-analysis of the effect of bowel preparation on adenoma detection: Early adenomas affected stronger than advanced adenomas. *PLoS One* 2016;11(6):e0154149.
 12. Romańczyk M, Ostrowski B, Kozłowska-Petriczko K, et al. Scoring system assessing mucosal visibility of upper gastrointestinal tract: The POLPREP scale. *J Gastroenterol Hepatol (Australia)* 2022;37(1): 164–8.
 13. Romańczyk M, Ostrowski B, Lesińska M, et al. The prospective validation of a scoring system to assess mucosal cleanliness during EGD. *Gastrointest Endosc* 2024;100(1):27–35.
 14. Koo TK, Li MY. A guideline of selecting and reporting intraclass correlation coefficients for reliability research. *J Chiropr Med* 2016;15(2): 155–63.
 15. Stoyan D, Pommerening A, Hummel M, et al. Multiple-rater kappas for binary data: Models and interpretation. *Biometrical J Biometrische Z* 2018;60(2):381–94.
 16. Calderwood AH, Jacobson BC. Comprehensive validation of the Boston Bowel Preparation Scale. *Gastrointest Endosc* 2009;69(5):AB120–1.
 17. Baile-Maxía S, Mangas-Sanjuan C, Medina-Prado L, et al. Diagnostic yield of early repeat colonoscopy after suboptimal bowel preparation in a fecal immunochemical test-based screening program. *Endoscopy* 2020; 52(12):1093–100.
 18. Hsu CM, Lin WP, Su MY, et al. Factors that influence cecal intubation rate during colonoscopy in deeply sedated patients. *J Gastroenterol Hepatol (Australia)* 2012;27(1):76–80.
 19. Bugajski M, Wieszczy P, Hoff G, et al. Modifiable factors associated with patient-reported pain during and after screening colonoscopy. *Gut* 2018; 67(11):1958–64.
 20. Januszewicz W, Wieszczy P, Bialek A, et al. Endoscopist biopsy rate as a quality indicator for outpatient gastroscopy: A multicenter cohort study with validation. *Gastrointest Endosc* 2019;89(6):1141–9.
 21. Januszewicz W, Witczak K, Wieszczy P, et al. Prevalence and risk factors of upper gastrointestinal cancers missed during endoscopy: A nationwide registry-based study. *Endoscopy* 2022;54(7):653–60.
 22. Kamran U, King D, Abbasi A, et al. A root cause analysis system to establish the most plausible explanation for post-endoscopy upper gastrointestinal cancer. *Endoscopy* 2023;55(2):109–18.
 23. Khan R, Gimpaya N, Vargas JJ, et al. The Toronto upper gastrointestinal cleaning score: A prospective validation study. *Endoscopy* 2023;55(2):121–8.
 24. Córdova H, Barreiro-Alonso E, Castillo-Regalado E, et al. Applicability of the Barcelona scale to assess the quality of cleanliness of mucosa at esophagogastroduodenoscopy. *Gastroenterol Hepatol* 2024;47(3):246–52.

Open Access This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.