

MEETING ABSTRACT

Open Access

# Microelements as diagnostic markers of pancreatic cancer

Marcin Lener<sup>1\*</sup>, Anna Wiechowska-Kozłowska<sup>2</sup>, Magdalena Muszyńska<sup>3</sup>, Katarzyna Jaworska-Bieniek<sup>1</sup>, Grzegorz Sukiennicki<sup>1</sup>, Katarzyna Kaczmarek<sup>1</sup>, Tomasz Gromowski<sup>1</sup>, Wojciech Marciniak<sup>3</sup>, Anna Jakubowska<sup>1</sup>, Jan Lubiński<sup>1,3</sup>

From Annual Conference on Hereditary Cancers 2014  
Szczecin, Poland. 25-26 September 2014

## Introduction

Pancreatic cancer is the eighth most commonly diagnosed cancer in the developed world and has one of the worst prognoses of any malignancy with 98% succumbing to their disease within 5 years. Little is known about the etiology of the disease despite significant new insights into the mutation signatures common this disease. Changes in microelements serum levels are reported in pancreatic cancer. In the current study we have examined the levels of Se, Cu, Fe, Zn and Mg in a moderately sized pancreatic cancer population and compared it to a healthy age-matched population.

## Material and methods

A total of 84 pancreatic cancer patients and 84 age-matched healthy controls were enrolled in the study after providing informed consent. The patients with pancreatic cancer were enrolled to the study from the Hospital of the Ministry of Internal Affairs and Administration in Szczecin, Poland. For each pancreatic cancer patient included in this study an unaffected individual registered in International Hereditary Cancer Center, Pomeranian Medical University of Szczecin, was used as a control.

Each person enrolled in the study donated ~10 ml EDTA blood for sufficient serum to be isolated and examined for the elements Se, Cu, Fe, Zn and Mg. The level of Se, Cu, Fe, Zn and Mg, in the serum was determined by mass spectrometry Inductively Coupled Plasma Mass Spectrometry (Elan DRC-e, PerkinElmer).

**Table 1 The correlation between serum Se level and the occurrence of pancreatic cancer.**

Cancer site	Quartile	Se level (µg/l)	No. of cancer / controls	OR	p-value
Pancreas	I	29.87-55.40	37/5	1	-
	II	55.90-67.97	31/11	2.62	0.1635
	III	68.06-79.20	9/33	27.13	<0.0001
	IV	79.36-137.47	7/35	37	<0.0001

**Table 2 The correlation between serum Cu level and the occurrence of pancreatic cancer.**

Cancer site	Quartile	Cu level (µg/l)	No. of cancer / controls	OR	p-value
Pancreas	I	695.06-1023.95	9/32	1	-
	II	1025.88-1175.73	14/27	1.84	0.3256
	III	1179.38-1421.96	22/19	4.11	0.0058
	IV	1448.68-2901.35	37/4	32.8	<0.0001

**Table 3 The correlation between serum Fe level and the occurrence of pancreatic cancer.**

Cancer site	Quartile	Fe level (µg/l)	No. of cancer / controls	OR	p-value
Pancreas	I	138.88-668.75	29/12	1	-
	II	672.81-921.18	19/22	2.79	0.0429
	III	931.41-1245.91	13/29	5.39	0.0004
	IV	1247.89-1631.56	22/20	2.19	0.1151

\* Correspondence: marcinlener@poczta.onet.pl

<sup>1</sup>Department of Genetics and Pathology, International Hereditary Cancer Center, Pomeranian Medical University, Szczecin, Poland  
Full list of author information is available at the end of the article

**Table 4 The correlation between serum Zn level and the occurrence of pancreatic cancer.**

Cancer site	Quartile	Zn level (µg/l)	No. of cancer / controls	OR	p- value
Pancreas	I	306.12-760.79	23/18	1	-
	II	765.29-863.96	14/27	2.46	0.0752
	III	864.59-995.08	21/21	1.27	0.6620
	IV	995.84-1890.96	25/17	0.86	0.8257

**Table 5 The correlation between serum Mg level and the occurrence of pancreatic cancer.**

Cancer site	Quartile	Mg level (µg/l)	No. of cancer / controls	OR	p- value
Pancreas	I	11964.63-18616.02	25/16	1	-
	II	18673.87-20144.41	18/23	1.99	0.1843
	III	20274.88-21673.70	17/24	2.20	0.1215
	IV	21775.80-30676.08	22/19	1.34	0.6555

## Results

### Conclusions

1. There is a very strong correlation between the level of selenium, copper in serum and the occurrence of pancreatic cancers in the Polish population.
2. The Se, Cu level in serum may be a useful diagnostic tool of pancreatic cancer.
3. Further investigations are needed to determine if Se, Cu levels can be used in:
  - a. differential diagnosis between pancreatic cancer (PC) and non-malignant pancreatic lesions
  - b. monitoring of systemic treatment efficiency in PC
  - c. identification of prognostic factors of pancreatic tumors
  - d. identification of factor of PC causative

### Authors' details

<sup>1</sup>Department of Genetics and Pathology, International Hereditary Cancer Center, Pomeranian Medical University, Szczecin, Poland. <sup>2</sup>Division of Health Care Ministry of Internal Affairs and Administration in Szczecin, Poland. <sup>3</sup>Read - Gene, S.A., Grzeczynica, Poland.

Published: 26 November 2015

doi:10.1186/1897-4287-13-S2-A2

Cite this article as: Lener et al.: Microelements as diagnostic markers of pancreatic cancer. *Hereditary Cancer in Clinical Practice* 2015 **13**(Suppl 2):A2.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

