

## REVIEW

# The 'N-factors' in pancreatic cancer: functional relevance of NF- $\kappa$ B, NFAT and Nrf2 in pancreatic cancer

A Arlt<sup>1</sup>, H Schäfer<sup>1</sup> and H Kalthoff<sup>2</sup>

Pancreatic ductal adenocarcinoma (PDAC) represents one of the deadliest malignancies, with an overall life expectancy of 6 months. Despite considerable advances in the understanding of the molecular mechanisms involved in the carcinogenesis of PDAC, the outcome of the disease was not significantly improved over the last 20 years. Although some achievements in molecular-targeted therapies have been made (that is, targeting the epidermal growth factor receptor by erlotinib), which already entered clinical settings, and despite the promising outcome of the FOLFIRINOX trial, there is an urgent need for improvement of the chemotherapy in this disease. A plethora of molecular alterations are thought to be responsible for the profound chemoresistance, including mutations in oncogenes and tumor suppressors. Besides these classical hallmarks of cancer, the constitutive or inducible activity of transcription factor pathways are characteristic changes in PDAC. Recently, three transcription factors—nuclear factor- $\kappa$ B (NF- $\kappa$ B), nuclear factor of activated T cells (NFAT) and nuclear factor-E2-related factor-2 (Nrf2)—have been shown to be crucial for tumor development and chemoresistance in pancreatic cancer. These transcription factors are key regulators of a variety of genes involved in nearly all aspects of tumorigenesis and resistance against chemotherapeutics and death receptor ligands. Furthermore, the pathways of NF- $\kappa$ B, NFAT and Nrf2 are functional, interacting on several regulatory steps, and, especially, natural compounds such as curcumin interfere with more than one pathway. Thus, targeting these pathways by established inhibitors or new drugs might have great potential to improve the outcome of PDAC patients, most likely in combination with established anticancer drugs. In this article, we summarize recent progress in the characterization of these transcription-factor pathways and their role in PDAC and therapy resistance. We also discuss future concepts for the treatment of PDAC relying on these pathways.

*Oncogenesis* (2012) 1, e35; doi:10.1038/oncis.2012.35; published online 26 November 2012

**Subject categories:** Molecular oncology

**Keywords:** chemoresistance; transcription factor; pancreatic cancer; signal transduction; apoptosis

## PANCREATIC DUCTAL ADENOCARCINOMA, CHEMORESISTANCE AND TRANSCRIPTION FACTORS

Pancreatic ductal adenocarcinoma (PDAC) represents the fourth leading cause of cancer-related death in western countries.<sup>1,2</sup> Up to 90% of PDACs develop through premalignant precursor lesions, so called the pancreatic intraepithelial neoplasias-1 to -3. The remaining cases originate in mucinous cystic neoplasia or intraductal papillary mucinous neoplasia.<sup>3,4</sup> A recent report comparing genetically engineered mouse models with human samples elegantly described an alternative pathway, which might be important especially in familial pancreatic cancer. In this model, the ductal cancer cells arise in the centroacinar–acinar region, possibly through a process of acinar–ductal metaplasia.<sup>5</sup> Several genetic alterations in pancreatic intraepithelial neoplasias and PDACs have been extensively described. The *Kras* oncogene has a central role in carcinogenesis and is mutated in nearly all PDACs. Other frequent mutations are found in the tumor suppressor genes *p16<sup>INK4</sup>*, *p53*, *SMAD4* and *BRCA2*.<sup>3,4</sup> Despite advances in the understanding of the mechanisms of carcinogenesis, therapeutic options in pancreatic cancer are associated with very limited or no improvement in life expectancy.<sup>6</sup> Because of a difficult and late

diagnosis, only 15% of the patients have a disease localized to the pancreas, allowing a potentially curative resection. The majority of patients, having a locally advanced tumor status, receive radio- and or chemotherapy. Gemcitabine or 5-fluorouracil are the most often used drugs, but such as all other therapeutical interventions they fail to significantly improve the prognosis.<sup>4</sup> Even the recommended first-line combination therapy with erlotinib and gemcitabine offers only a very limited gain in life expectancy compared with a monotherapy with gemcitabine.<sup>7</sup> Recently, clinical trials showed that FOLFIRINOX, an aggressive combination therapy of several chemotherapeutic drugs, improves the life expectancy from 6–7 months with gemcitabine therapy to 10–11 months in the FOLFIRINOX group.<sup>8</sup> Because of the higher rate of severe side effects and some problems in the group composition (a high proportion of pancreatic cancer localized in the tail of the pancreas), it still remains to be seen if FOLFIRINOX will indeed be the future gold standard in the palliative setting.<sup>9</sup>

Thus, limitations in curative and, especially, palliative treatment options of PDACs demonstrate a need for new and gemcitabine-independent strategies.<sup>10,11</sup>


<sup>1</sup>Laboratory of Molecular Gastroenterology and Hepatology, Department of Internal Medicine I, Kiel, Germany and <sup>2</sup>Division of Molecular Oncology, Institute for Experimental Cancer Research, Comprehensive Cancer Center North, Kiel, Germany. Correspondence: Dr A Arlt, Laboratory of Molecular Gastroenterology and Hepatology, Department of Internal Medicine I, UKSH-Campus Kiel, Arnold-Heller-Strasse 3, 24105 Kiel, Germany.

E-mail: aarlt@1med.uni-kiel.de

or Professor H Kalthoff, Division of Molecular Oncology, Institute for Experimental Cancer Research, Comprehensive Cancer Center North, UKSH-Campus Kiel, Arnold-Heller-Strasse 3, 24105 Kiel, Germany.

E-mail: hkalthoff@email.uni-kiel.de

Received 2 October 2012; accepted 6 October 2012



**Figure 1.** Scheme of unstimulated state. Depicted are the key factors of the three transcription factor pathways. Transcription factors (in blue) and the corresponding inhibitors (in red) as further outlined in this review. P, phosphorylation; Ub, ubiquitin.

Only deeper insight into the molecular mechanisms of carcinogenesis and the profound chemoresistance of PDACs might provide concepts for more efficient molecular-targeted therapies. Among these promising molecular targets are numerous transcription factors. Through a variety of alterations in their regulating pathways, involving constitutive activity of upstream regulators, such as *Kras*,<sup>12</sup> the epidermal growth factor receptor system<sup>13,14</sup> and/or epigenetic alterations,<sup>6,15,16</sup> a plethora of transcription factors is constitutively activated in PDACs. Furthermore, several chemotherapeutic drugs and death receptors ligands<sup>17</sup> activate antiapoptotic transcription factors, thereby counteracting their apoptotic potential.

Over the last couple of years, the transcription factors nuclear factor-κB (NF-κB), nuclear factor of activated T cells (NFATs) and nuclear factor-E2-related factor-2 (Nrf2) took center stage as promising molecular targets in the therapy of PDACs.

In this review, we will delineate the intertwined characteristics of their regulatory pathways and discuss the suitability of these factors to be used in novel therapeutic strategies.


### NF-κB PATHWAY

The transcription factor NF-κB is involved in the regulation of expression of a variety of target genes. Among these are central regulators of apoptosis, the cell cycle and metastasis.<sup>18,19</sup> NF-κB exists as a hetero- and homodimeric protein complex of members of the so-called Rel-family. RelA/p65, RelB and c-Rel harbor a transactivation domain along with the Rel homology domain. The Rel homology domain, also present in NF-κB1 (p50/p105) and NF-κB2 (p52/p100), functions as an interacting site for Rel-family members and confers DNA binding. In contrast, through the transactivation domain p65, RelB and c-Rel regulate the expression of their target genes. The most abundant NF-κB form is the heterodimer of p65 and p50. In the so-called classical NF-κB pathway, p65/p50 is anchored in the cytoplasm by the inhibitor of κB (IκB) proteins (Figure 1).<sup>6,18</sup> In response to NF-κB-activating stimuli (Figure 2), the IκB kinase complex (IKK) consisting of two catalytical kinases (IKKα and IKKβ), together with the regulatory component IKKγ/NF-κB essential modulator, is activated by phosphorylation. Once activated, IKK phosphorylates the IκBs, which are subsequently polyubiquitinated and degraded by the 26S proteasome. The NF-κB dimer then translocates to the nucleus and regulates the transcription of its target genes.<sup>6,18,19</sup>

The number of NF-κB target genes is enormous and still growing.<sup>20</sup> Among them are well-defined antiapoptotic genes like *Bcl-xL*, *cIAP*, *Bcl-2* and *c-Flip*, which are able to inhibit the apoptotic cascade at several points, conferring resistance against chemotherapeutic drugs and death receptor ligands.<sup>6,21</sup> Despite the fact that some reports indicate NF-κB activation can elicit pro-apoptotic functions,<sup>22,23</sup> published data largely support a role for NF-κB as a central antiapoptotic factor.<sup>18,24</sup> Meanwhile, there is firm evidence that the death receptor ligands TRAIL (tumor necrosis factor-related apoptosis-inducing ligand) and Fas ligand induce NF-κB,<sup>17,25–29</sup> and thereby inhibit their potential to eliminate pancreatic cancer cells. There is some controversy as to whether inducible or constitutive NF-κB activity is more important for PDAC resistance against TRAIL,<sup>30</sup> but most reports indicate that in contrast to chemotherapeutic drugs (see below), the inducible activation confers resistance against death receptor ligands. Moreover, PDAC cells gain growth advantage by auto- and paracrine death receptor ligand loops either through secretion of the death ligands by tumor cells or by infiltrating immune cells.<sup>17,31</sup> As already mentioned, some chemotherapeutic drugs like CPT-11 are able to induce NF-κB in a same manner as the tumor necrosis factor-α and TRAIL,<sup>32</sup> but the studies supporting a critical role for a constitutive NF-κB activity in the resistance of PDACs against chemotherapeutic drugs are overwhelming.<sup>6,33–37</sup> There is a multitude of mechanisms leading to this constitutive NF-κB activity. Chronic inflammation can induce NF-κB activation<sup>18,20,38</sup> as observed in the course of chronic pancreatitis and acts as a risk factor for development of PDACs.<sup>39,40</sup> In particular, proinflammatory cytokines, such as interleukin-1β,<sup>41</sup> interleukin-1α,<sup>42</sup> and interleukin-8,<sup>43,44</sup> either released by PDAC cells themselves or by immune cells, and the surrounding tissue—for example, myofibroblasts—in the course of chronic inflammation,<sup>45</sup> can lead to the constitutive activation of NF-κB in tumor cells.

As described, point mutations in *Kras* are present in up to 90% of PDACs and might be involved in constitutive NF-κB activation.<sup>46–49</sup> In addition, NF-κB itself provides feed-forward loops, which are essential for pancreatic intraepithelial neoplasia and PDAC progression.<sup>47,48</sup> Furthermore, the frequent overexpression and activation of members of the EGFR signaling pathway might contribute to NF-κB-dependent tumor progression and invasive phenotype of PDACs.<sup>50–52</sup>

So far, the NF-κB target genes involved in PDAC chemoresistance, either through interference with death receptor signaling or in association with chemotherapeutic drugs, have remained elusive. There are some reports indicating that classical inhibitors



**Figure 2.** Scheme of stimulated state. Depicted are the key factors of the three transcription factor pathways. Transcription factors (in blue), the corresponding inhibitors (in red) and activating processes (in blue) as further outlined in this review. Ca, Calcium; P, phosphorylation; Ub, ubiquitin.

of apoptosis such as cIAPs or BCL-family members are involved,<sup>36,53</sup> but also other target genes<sup>28,43,44</sup> or cross talk with other transcription factors such as STAT3<sup>54</sup> have been proposed.


These *in vitro* and *in vivo* data support the potential for NF-κB as a molecular target in PDAC therapy. As small interfering RNA-mediated approaches<sup>36</sup> are currently not feasible for broad clinical application, current clinical trials focus on upstream events related to either inducible or constitutive NF-κB activity (Figure 3). In this context, IKK inhibitors have already entered clinical settings. The clinically approved drugs thalidomide,<sup>55–57</sup> salicylates and their derivate sulfasalazine<sup>58,59</sup> have also been shown to be potent chemosensitizers in PDAC *in vitro*, *in vivo* and in clinical settings. Furthermore, more specific IKK targeting drugs, for example, PS-1145, BAY11-7082, EC-70124 or SAR113945, are in various phases of clinical trials and approval by the Food and Drug Association.<sup>60–62</sup>

Beyond these more or less NF-κB-specific strategies, proteasome inhibitors like velcade/bortezomib are promising therapeutic options in PDAC treatment.<sup>63</sup> The proteasome is involved in the NF-κB activation pathway by degradation of IκB and is therefore extensively used for pharmacological NF-κB inhibition in preclinical and clinical studies.<sup>64,65</sup> Nevertheless, the proteasome is not only part of the NF-κB pathway, but instead a central regulator of a variety of regulatory pathways involved in cancer initiation, progression and chemoresistance,<sup>66</sup> making the proteasome a promising target in PDAC therapy, but without clear conclusions on the role of NF-κB in this context.

Beyond these chemical compounds, a growing number of natural products like curcumin,<sup>6,67,68</sup> epicatechin gallate and catechin gallate<sup>44</sup> have shown the potential to block NF-κB and sensitize PDACs for apoptosis without severe side effects, and might be beneficial in combination with chemotherapeutic drugs and death ligands.<sup>69</sup> Other herbal compounds include thymoquinone,<sup>70</sup> sulforaphane,<sup>71–73</sup> dihydroartemisinin<sup>21</sup> or 3,3-diindolylmethane,<sup>53</sup> which block both constitutive and anticancer drug-induced NF-κB activity, and have been successfully tested in preclinical experiments for sensitization of PDAC cells against chemotherapy.

## THE NFAT PATHWAY

The NFAT family of transcription factors is a group of calcineurin-responsive, inducible nuclear proteins. Originally described in the context of T-lymphocyte activation, increasing evidence exists showing a crucial role of this transcription factor family in the regulation of cell growth and apoptosis.<sup>74,75</sup> Four calcium-responsive isoforms named NFATc1 (NFAT2/NFATc), NFATc2 (NFAT1/NFATp), NFATc3 (NFAT4/NFATx) and NFATc4 (NFAT3) are members of a family, which is under the control of a Ca(2+)/calcineurin signaling pathway.<sup>76</sup> Under unstimulated conditions (Figure 1), NFAT is anchored in the cytoplasm through phosphorylation of a number of serines within its highly conserved regulatory domain, which masks the nuclear localization sequence. After dephosphorylation by calcineurin, which exposes the nuclear localization sequence and masks a nuclear export sequence, NFAT enters the nucleus and regulates the transcription of target genes by dimerization with NFAT family members, but also with other transcription factors, such as activating protein-1 and NF-κB (Figure 2 for classical activation pathway and Figure 3 for interaction of the pathways). Termination of NFAT activity is mediated by multiple mechanisms, including inhibition of calcineurin and phosphorylation of NFAT by nuclear kinases. Hereby, NFAT is rephosphorylated, the nuclear export sequence unmasked and the nuclear localization sequence masked.<sup>76</sup> In addition to the still growing number of regulating kinases and phosphatases, other regulatory mechanisms including sumoylation,<sup>77</sup> ubiquitination<sup>78</sup> and *de novo* expression of NFAT members<sup>79</sup> exist. Thus, as in the case of nearly all other signaling pathways, the oversimplified linear model of a merely Ca(2+)/calcineurin-dependent signaling pathway must be revised to reflect a complex regulatory network. Recent reports indicate a crucial role of NFATc1/NFAT2 and NFATc2/NFAT1 in different steps of PDAC carcinogenesis and chemoresistance.<sup>75,80–83</sup> NFATc1/NFAT2 is activated by serum in PDAC cells and binds to a serum-responsive element within the proximal c-myc promoter, initiating p300-dependent histone acetylation, which creates a local chromatin structure permissive for the inducible recruitment of Ets-like gene (*ELK*-1). This NFATc1/NFAT2-dependent pathway



**Figure 3.** Scheme of interaction of the pathways and inhibitor strategies. Depicted are the key factors of the three transcription factor pathways. Transcription factors (in blue), the corresponding inhibitors (in red) and activating processes (in blue) as further outlined in this review. Positive/enhancing (in blue) and negative/inhibiting interactions are indicated. Furthermore, chemical and natural inhibitors (in red) as described in this review are included. Divergent effects of these inhibitors on the pathways (that is, curcumin, sulforaphane, bortezomib) are highlighted by arrows (blue for activating and red for inhibitory effects). Ca, calcium; P, phosphorylation; Ub, ubiquitin.

results in the promotion of c-myc-dependent growth of PDAC cells.<sup>82</sup> Recently, the same group elegantly delineated an NFATc2/NFAT1-driven pathway in PDACs, which, by targeting a p15(INK4b)-mediated failsafe mechanism, promotes pancreatic cancer growth.<sup>80</sup>

However, besides this well-documented role of NFAT in pancreatic cancer development and growth regulation, there are only very limited data on the suitability of NFAT as a molecular target in pancreatic cancer therapy. For other malignancies, preclinical and limited clinical data exist (Figure 3).<sup>76,84</sup> Cyclosporin A and FK506 represent two structurally unrelated, clinically established, potent inhibitors of calcineurin. These are widely used immunosuppressive agents that prevent NFAT nuclear translocation by interfering with calcineurin activation, thus resulting in the blockade of dephosphorylation of numerous other substrates in addition to NFAT.<sup>85</sup> The long-term clinical experience with these drugs in transplantation medicine demonstrates a significant increase in cancer incidence in patients subject of long-term cyclosporin A and FK506 treatment.<sup>86</sup> On the one hand, this might be due to the interference of these drugs with calcineurin- and NFAT-dependent immunosurveillance, and on the other hand, oncogenic effects of calcineurin inhibition might result from NFAT-independent mechanisms of cyclosporin A and FK506. Therefore, more selective NFAT targeting strategies are currently being developed. VIVIT, a peptide that interferes with the calcineurin–NFAT interaction, potentially blocks dephosphorylation and nuclear translocation of NFAT, and has been shown to attenuate breast cancer cell invasion.<sup>87</sup> Thus, VIVIT administration might be a specific measure to block the oncogenic effects of NFAT in PDACs. As the delivery of the VIVIT peptide to the tumor in an *in vivo* setting could be difficult other small molecule inhibitors like L-732531,<sup>88</sup> an analog of FK506, and ISATX247,<sup>89</sup> a potent and less toxic analog of cyclosporin A, are promising. However, up to now sufficient *in vivo* data of the capacity of NFAT inhibitors to reduce chemoresistance and tumorigenesis, beyond

their well-documented activities in immune suppression, are missing—especially for pancreatic cancer.

### THE NRF2 PATHWAY

Nrf2 is a transcription factor belonging to the family of cap'n'collar basic leucine zipper protein family, and represents a key regulator of the cellular defense against oxidative stress. Under homeostatic conditions (Figure 1), Nrf2 is anchored in the cytoplasm through kelch-like ECH-associated protein 1 (*Keap1*). *Keap1* is an E3 ubiquitin ligase mediating polyubiquitination of Nrf2, thereby tagging it for degradation by the 26S proteasome. In addition, *Keap1* can terminate the Nrf2 activation through retrieving the transcription factor from the nucleus. Similar to IκB in NF-κB signaling, *Keap1* represents a classical feedback regulator<sup>90,91</sup> itself, being a target gene of Nrf2.

Nrf2 is activated by numerous cellular conditions and stimuli (Figure 2). In general, through modifications of critical cysteine residues of *Keap1* and of Nrf2 the interaction of both molecules is disturbed, leading to the nuclear translocation of Nrf2.<sup>92</sup> In the nucleus, Nrf2 interacts with small v-maf musculoaponeurotic fibrosarcoma oncogene homolog (*sMAF*) regulator proteins and binds to antioxidative response elements in the promoters of a plethora of target genes.<sup>90,91</sup> Oxidative and electrophilic stress are the main inducers of Nrf2, involving, for example, ERK-, PKC-, JNK- and PI3K/Akt-dependent pathways. Accordingly, classical target genes of Nrf2 include phase-II enzymes like glutathione-S-transferase or NAD(P)H-quinone-oxidoreductase-1 (*hNQO-1*).<sup>90,91</sup> For a long time, Nrf2 was considered an attractive and quite hopeful target for chemopreventive strategies<sup>93,94</sup> to maintain cellular redox balance. Through induction of antioxidative target genes, Nrf2 could protect non-transformed cells against DNA damage and, thereby, may prevent mutagenesis. However, there is growing evidence that Nrf2 activation can lead to tumor development and enhanced chemoresistance once a malignant transformation occurred.<sup>95,96</sup> Stable overexpression of Nrf2 results in enhanced resistance of cancer cells to chemotherapeutic



agents, whereas targeting Nrf2 renders cancer cells more susceptible to these drugs.<sup>96</sup> Interestingly, the strategy of using Nrf2 inhibitors (Figure 3) to increase the efficacy of chemotherapeutic agents is not limited to certain cancer types or anticancer drugs and may be beneficial during the course of chemotherapy in general.<sup>96</sup> Besides direct regulation of expression of phase-II enzymes<sup>91</sup> and transporters for xenobiotics and drugs<sup>97</sup> accounting for direct detoxification of anticancer drugs, Nrf2 is a crucial regulator of proteasome activity by transcriptional control of several subunits of the proteasome<sup>64,98,99</sup> conferring resistance against a wide variety of apoptotic stimuli.

Through a plethora of alterations, such as gain-of-function mutations of Nrf2 itself<sup>100,101</sup> or, more frequently, through loss-of-function mutations, promoter hypermethylation, miR targeting or succination of the Nrf2 inhibitory protein Keap1,<sup>102–105</sup> Nrf2 activity is upregulated in several types of solid cancers, for example, in colonic, thyroid, endometrial, lung, ovarian, breast and pancreatic cancer.<sup>64,106–115</sup> Beyond these genetic alterations, inflammatory carcinogenesis is characterized by increased levels of metabolic and oxidative stress, leading to an exaggerated Nrf2 activity in tumors.<sup>116–118</sup> As a consequence of the increased Nrf2 activity, tumor cells acquire protection from apoptosis<sup>64,117–119</sup> and are more capable to proliferate, both conditions favoring tumorigenesis on the one hand and making tumor cells more resistant to chemo- and radiotherapy on the other hand.<sup>106,109,120</sup> Recently, a critical role of Nrf2 in oncogenesis of PDACs has been proposed.<sup>107</sup> In the K-Ras(G12D), B-Raf(V619E) and Myc(ERT2) models, and in human pancreatic cancer, the authors could show that an Nrf2-dependent antioxidant program is induced, which by lower intracellular reactive-oxygen species confers a more reduced intracellular environment and mediates oncogenesis.<sup>107</sup> Moreover, Lister *et al.*<sup>113</sup> reported that in PDAC cell lines Nrf2 is upregulated despite the absence of mutations in central mediators of the pathway, and also in the cytoplasm of tumor cells in human PDAC specimen compared with benign ductal cells. The small interfering RNA-mediated inhibition of Nrf2 activity sensitized the cell lines for chemotherapeutic drugs. Another group showed that increasing Nrf2 by overexpression or through induction on endogenous Nrf2 conferred protection of PDAC cell lines against apoptotic stimuli,<sup>108</sup> leading to the conclusion that strategies to pharmacologically manipulate the levels and/or activity of Nrf2 may have potential for PDAC therapy.<sup>108,113</sup>

Several strategies for induction of Nrf2 activity to prevent cancer development were propagated over the last couple of years<sup>91</sup> (Figure 3). Most of these strategies used food compounds like sulforaphane and triterpenoids.<sup>69,121–123</sup> As it is increasingly acknowledged that once premalignant or tumor cells acquired a persistent Nrf2 activity tumorigenesis and chemoresistance are substantially promoted, new strategies for inhibition of these pathways are under intensive investigation. Recently, retinoic acid receptor- $\alpha$  or estrogen-related receptor- $\beta$  have been described to inhibit Nrf2 activation,<sup>124,125</sup> but in PDAC cells estrogen-related receptor- $\beta$  is not detectable<sup>126</sup> and retinoic acid up to now failed to improve the outcome of PDAC patients due to profound molecular resistance mechanisms.<sup>127</sup> Thus, besides genetic modification and or small interfering RNA-mediated knock down of Nrf2,<sup>107,108,113</sup> pharmacological strategies using natural compounds have been identified to directly inhibit Nrf2. The alkaloids luteolin or trigonelline<sup>128,129</sup> would be attractive tools for sensitization of tumor cells to apoptosis, in particular, as they have already been used in clinical trials (for example, diabetes). Intriguingly, trigonelline was shown to have great potential to inhibit PDAC tumor growth *in vitro* and *in vivo*.<sup>126</sup> A recent screening of pharmacological inhibitors identified 4-(2-cyclohexylethoxy) aniline (IM3829) as highly potent in the inhibition of Nrf2.<sup>130</sup> IM3829 greatly enhances the radiosensitivity of human lung cancer cells *in vitro* and *in vivo*, and this compound may also be effective in chemosensitization of PDAC cells.

## THE THREE 'NS': INTERACTION OF THE PATHWAYS

As already mentioned NF- $\kappa$ B, NFAT and Nrf2 are influencing each other through direct interaction and/or modification of their signaling pathways (Figure 3). For instance, NF- $\kappa$ B and NFAT are members of a superfamily of transcription factors. They share similar DNA-binding domains,<sup>131</sup> and especially the regulation of COX-2 expression depends on parallel activation of both pathways.<sup>132,133</sup> In line with this, full activation of the BLYS promoter requires parallel constitutive activity of NF- $\kappa$ B and NFAT in aggressive B-cell lymphoma.<sup>134</sup> Despite the well-established role of NFAT and NF- $\kappa$ B in regulating COX-2,<sup>133,135</sup> and the importance of COX-2 for chemoresistance of PDAC,<sup>136,137</sup> up to now evidence is missing that the two pathways directly interact with each other in PDACs.

An interaction of Nrf2 and NF- $\kappa$ B is increasingly described. Most of the reports concentrated on the effects of oxidative stress induced Nrf2 activation, leading to alterations in NF- $\kappa$ B signaling. In this context, Nrf2 activation can inhibit NF- $\kappa$ B-dependent proinflammatory signaling by an unknown mechanism<sup>138</sup> and epigallocatechin-3-gallate elicits at least some of its anti-inflammatory properties by Nrf2-mediated NF- $\kappa$ B inhibition.<sup>139</sup> However, the opposite effect is evoked by Nrf2-mediated transcription of proteasome subunits that leads to an exaggerated activity of the 26S proteasome, and thereby facilitates the release of NF- $\kappa$ B from I $\kappa$ B $\alpha$ .<sup>64</sup> As dendritic cells from Nrf2 knockout mice exhibit no change in basal NF- $\kappa$ B activity,<sup>140</sup> it remains unclear if the proposed interaction of both pathways<sup>141</sup> is of functional relevance in epithelial cells of PDACs. Keap1 seems to be a central regulator of both pathways.<sup>92</sup> Through the double glycine repeat domain, Keap1 can bind to other proteins directly or indirectly, and IKK $\beta$  is destabilized by Keap1, which resulted in inhibiting NF- $\kappa$ B-derived tumor promotion.<sup>142</sup> Besides these Nrf2 effects on NF- $\kappa$ B, this central mediator of PDAC chemoresistance can influence Nrf2 activity. A recent report showed that the NF- $\kappa$ B subunit p65 directly interacts with Keap1, leading to inhibition of Nrf2 activity.<sup>143</sup>

Finally, varieties of natural compounds modulating Nrf2 also affect NF- $\kappa$ B and vice versa.<sup>69,144</sup> Especially, curcumin, which is reported to be a chemosensitizer of PDAC, affects NF- $\kappa$ B and Nrf2,<sup>145,146</sup> or sulforaphane, which is a well-known inducer of Nrf2<sup>121</sup> but has been described also as an inhibitor of NF- $\kappa$ B.<sup>71,73</sup> Taking into account the tumor- and chemoresistance-promoting role of Nrf2 just evoked over the last couple of years, it will be interesting to analyze whether the Nrf2 activation of many of these compounds is responsible for the failure of clinical trials in PDACs and whether a combination with Nrf2 inhibitors, such as trigonelline,<sup>126</sup> can improve the outcome in NF- $\kappa$ B-inhibiting strategies.

## CONCLUSIONS

Among the rising number of promising molecular targets in PDAC therapy, the transcription factor pathways of NF- $\kappa$ B, NFAT and Nrf2 are of potential interest for several reasons:

- (1) All three of these pathways have been shown to be altered in PDACs *in vitro* and *in vivo*, leading to PDAC development, migration, invasion and chemoresistance.
- (2) They interact with each other, and especially the counteraction of Nrf2 and NF- $\kappa$ B might explain some problems in targeted therapies.
- (3) A battery of natural and chemical compounds with well-known clinical safety exist, making it possible to target these pathways in upcoming clinical trials for PDAC therapy in the near future.

## CONFLICT OF INTEREST

The authors declare no conflict of interest.

## ACKNOWLEDGEMENTS

The funding by the German Research Society (DFG), German Cluster of Excellence 'Inflammation at Interfaces', Sander-Stiftung and Deutsche Krebshilfe to AA, HS and HK is greatly acknowledged.

## REFERENCES

- Jemal A, Siegel R, Ward E, Hao Y, Xu J, Murray T et al. Cancer statistics 2008. *CA Cancer J Clin* 2008; **58**: 71–96.
- Hidalgo M. Pancreatic cancer. *N Engl J Med* 2010; **362**: 1605–1617.
- Remmers N, Bailey JM, Mohr AM, Hollingsworth MA. Molecular pathology of early pancreatic cancer. *Cancer Biomark* 2010; **9**: 421–440.
- Vincent A, Herman J, Schulick R, Hruban RH, Goggins M. Pancreatic cancer. *Lancet* 2011; **378**: 607–620.
- Aichler M, Seiler C, Tost M, Siveke J, Mazur PK, Da Silva-Buttkus P et al. Origin of pancreatic ductal adenocarcinoma from atypical flat lesions: a comparative study in transgenic mice and human tissues. *J Pathol* 2012; **226**: 723–734.
- Arlt A, Muerkoster SS, Schafer H. Targeting apoptosis pathways in pancreatic cancer. *Cancer Lett* (e-pub ahead of print 13 November 2010; doi:10.1016/j.canlet.2010.10.015).
- Vulfovich M, Rocha-Lima C. Novel advances in pancreatic cancer treatment. *Expert Rev Anticancer Ther* 2008; **8**: 993–1002.
- Conroy T, Desseigne F, Ychou M, Bouche O, Guimbaud R, Becouarn Y et al. FOLFIRINOX versus gemcitabine for metastatic pancreatic cancer. *N Engl J Med* 2011; **364**: 1817–1825.
- Saif MW, Chabot J. Chemotherapy: metastatic pancreatic cancer—is FOLFIRINOX the new standard? *Nat Rev* 2011; **8**: 452–453.
- Saif MW. New developments in the treatment of pancreatic cancer. Highlights from the “44th ASCO Annual Meeting”. Chicago, IL, USA. May 30 - June 3, 2008. *JOP* 2008; **9**: 391–397.
- Shi S, Yao W, Xu J, Long J, Liu C, Yu X. Combinational therapy: new hope for pancreatic cancer? *Cancer Lett* 2012; **317**: 127–135.
- Ji B, Tsou L, Wang H, Gaiser S, Chang DZ, Daniluk J et al. Ras activity levels control the development of pancreatic diseases. *Gastroenterology* 2009; **137**: 1072–1082.
- Van den Eynde M, Baurain JF, Mazzeo F, Machiels JP. Epidermal growth factor receptor targeted therapies for solid tumours. *Acta Clin Belg* 2011; **66**: 10–17.
- Vaccaro V, Melisi D, Bria E, Cuppone F, Ciuffreda L, Pino MS et al. Emerging pathways and future targets for the molecular therapy of pancreatic cancer. *Expert Opin Ther Targets* 2011; **15**: 1183–1196.
- Fukushige S, Horii A. Road to early detection of pancreatic cancer: Attempts to utilize epigenetic biomarkers. *Cancer Lett* (e-pub ahead of print 23 March 2012; doi:10.1016/j.canlet.2012.03.022).
- Schuler S, Fritsche P, Diersch S, Arlt A, Schmid RM, Saur D et al. HDAC2 attenuates TRAIL-induced apoptosis of pancreatic cancer cells. *Mol Cancer* 2010; **9**: 80.
- Roder C, Trauzold A, Kalthoff H. Impact of death receptor signaling on the malignancy of pancreatic ductal adenocarcinoma. *Eur J Cell Biol* 2011; **90**: 450–455.
- Chaturvedi MM, Sung B, Yadav VR, Kannappan R, Aggarwal BB. NF-kappaB addiction and its role in cancer: 'one size does not fit all'. *Oncogene* 2011; **30**: 1615–1630.
- Perkins ND. The diverse and complex roles of NF-kappaB subunits in cancer. *Nat Rev Cancer* 2012; **12**: 121–132.
- Ben-Neriah Y, Karin M. Inflammation meets cancer, with NF-kappaB as the matchmaker. *Nat Immunol* 2011; **12**: 715–723.
- Wang SJ, Gao Y, Chen H, Kong R, Jiang HC, Pan SH et al. Dihydroartemisinin inactivates NF-kappaB and potentiates the anti-tumor effect of gemcitabine on pancreatic cancer both in vitro and in vivo. *Cancer Lett* 2010; **293**: 99–108.
- Jennewein C, Karl S, Baumann B, Micheau O, Debatin KM, Fulda S. Identification of a novel pro-apoptotic role of NF-kappaB in the regulation of TRAIL- and CD95-mediated apoptosis of glioblastoma cells. *Oncogene* 2011; **31**: 1468–1474.
- Martin AG. NFkappaB anti-apoptotic or pro-apoptotic, maybe both. *Cell Cycle* 2010; **9**: 3131–3132.
- Hamacher R, Schmid RM, Saur D, Schneider G. Apoptotic pathways in pancreatic ductal adenocarcinoma. *Mol Cancer* 2008; **7**: 64.
- Chen PH, Yang CR. Decoy receptor 3 expression in AsPC-1 human pancreatic adenocarcinoma cells via the phosphatidylinositol 3-kinase-, Akt-, and NF-kappa B-dependent pathway. *J Immunol* 2008; **181**: 8441–8449.
- Lemke J, Noack A, Adam D, Tchikov V, Bertsch U, Roder C et al. TRAIL signaling is mediated by DR4 in pancreatic tumor cells despite the expression of functional DR5. *J Mol Med* 2010; **88**: 729–740.
- Trauzold A, Wermann H, Arlt A, Schutze S, Schafer H, Oestern S et al. CD95 and TRAIL receptor-mediated activation of protein kinase C and NF-kappaB contributes to apoptosis resistance in ductal pancreatic adenocarcinoma cells. *Oncogene* 2001; **20**: 4258–4269.
- Trauzold A, Roder C, Sipos B, Karsten K, Arlt A, Jiang P et al. CD95 and TRAF2 promote invasiveness of pancreatic cancer cells. *Faseb J* 2005; **19**: 620–622.
- Khanbolooki S, Nawrocki ST, Arumugam T, Andtbacka R, Pino MS, Kurzrock R et al. Nuclear factor-kappaB maintains TRAIL resistance in human pancreatic cancer cells. *Mol Cancer Ther* 2006; **5**: 2251–2260.
- Brauer SJ, Buneker C, Mohr A, Zwacka RM. Constitutively activated nuclear factor-kappaB, but not induced NF-kappaB, leads to TRAIL resistance by up-regulation of X-linked inhibitor of apoptosis protein in human cancer cells. *Mol Cancer Res* 2006; **4**: 715–728.
- Trauzold A, Siegmund D, Schniewind B, Sipos B, Egberts J, Zorenkov D et al. TRAIL promotes metastasis of human pancreatic ductal adenocarcinoma. *Oncogene* 2006; **25**: 7434–7439.
- Shah SA, Potter MW, McDade TP, Ricciardi R, Perugini RA, Elliott PJ et al. 26S proteasome inhibition induces apoptosis and limits growth of human pancreatic cancer. *J Cell Biochem* 2001; **82**: 110–122.
- Arlt A, Gehrz A, Muerkoster S, Vorndamm J, Kruse ML, Folsch UR et al. Role of NF-kappaB and Akt/PI3K in the resistance of pancreatic carcinoma cell lines against gemcitabine-induced cell death. *Oncogene* 2003; **22**: 3243–3251.
- Arlt A, Vorndamm J, Breitenbroich M, Folsch UR, Kalthoff H, Schmidt WE et al. Inhibition of NF-kappaB sensitizes human pancreatic carcinoma cells to apoptosis induced by etoposide (VP16) or doxorubicin. *Oncogene* 2001; **20**: 859–868.
- Kiefel H, Bondong S, Erbe-Hoffmann N, Hazin J, Riedle S, Wolf J et al. L1CAM-integrin interaction induces constitutive NF-kappaB activation in pancreatic adenocarcinoma cells by enhancing IL-1beta expression. *Oncogene* 2010; **29**: 4766–4778.
- Kong R, Sun B, Jiang H, Pan S, Chen H, Wang S et al. Downregulation of nuclear factor-kappaB p65 subunit by small interfering RNA synergizes with gemcitabine to inhibit the growth of pancreatic cancer. *Cancer Lett* 2010; **291**: 90–98.
- Melisi D, Xia Q, Paradiso G, Ling J, Moccia T, Carbone C et al. Modulation of pancreatic cancer chemoresistance by inhibition of TAK1. *J Natl Cancer Inst* 2011; **103**: 1190–1204.
- Porta C, Riboldi E, Sica A. Mechanisms linking pathogens-associated inflammation and cancer. *Cancer Lett* 2011; **305**: 250–262.
- Uomo I, Miraglia S, Pastorello M. Inflammation and pancreatic ductal adenocarcinoma: a potential scenario for novel drug targets. *JOP* 2010; **11**: 199–202.
- Garcea G, Dennison AR, Steward WP, Berry DP. Role of inflammation in pancreatic carcinogenesis and the implications for future therapy. *Pancreatol* 2005; **5**: 514–529.
- Arlt A, Vorndamm J, Muerkoster S, Yu H, Schmidt WE, Folsch UR et al. Autocrine production of interleukin 1beta confers constitutive nuclear factor kappaB activity and chemoresistance in pancreatic carcinoma cell lines. *Cancer Res* 2002; **62**: 910–916.
- Niu J, Li Z, Peng B, Chiao PJ. Identification of an autoregulatory feedback pathway involving interleukin-1alpha in induction of constitutive NF-kappaB activation in pancreatic cancer cells. *J Biol Chem* 2004; **279**: 16452–16462.
- Zhou DH, Trauzold A, Roder C, Pan G, Zheng C, Kalthoff H. The potential molecular mechanism of overexpression of uPA, IL-8, MMP-7 and MMP-9 induced by TRAIL in pancreatic cancer cell. *Hepatobiliary Pancreat Dis Int* 2008; **7**: 201–209.
- Kurbitz C, Heise D, Redmer T, Goumas F, Arlt A, Lemke J et al. Epicatechin gallate and catechin gallate are superior to epigallocatechin gallate in growth suppression and anti-inflammatory activities in pancreatic tumor cells. *Cancer Sci* 2011; **102**: 728–734.
- Treiber M, Neuhofer P, Anetsberger E, Einwachter H, Lesina M, Rickmann M et al. Myeloid, but Not Pancreatic, RelA/p65 Is Required for Fibrosis in a Mouse Model of Chronic Pancreatitis. *Gastroenterology* 2011; **141**: e1477.
- Rachagani S, Senapati S, Chakraborty S, Ponnusamy MP, Kumar S, Smith LM et al. Activated KrasG(11)(2)D is associated with invasion and metastasis of pancreatic cancer cells through inhibition of E-cadherin. *Br J Cancer* 2011; **104**: 1038–1048.
- Maier HJ, Wagner M, Schips TG, Salem HH, Baumann B, Wirth T. Requirement of NEMO/IKKγ for effective expansion of KRAS-induced precancerous lesions in the pancreas. *Oncogene*. (e-pub ahead of print 2 July 2012; doi:10.1038/onc.2012.272).
- Ling J, Kang Y, Zhao R, Xia Q, Lee DF, Chang Z et al. KrasG12D-induced IKK2/beta/NF-kappaB activation by IL-1alpha and p62 feedforward loops is required for development of pancreatic ductal adenocarcinoma. *Cancer Cell* 2012; **21**: 105–120.
- Chiao PJ, Ling J, Kras, Pten, NF-kappaB, and inflammation: dangerous liaisons. *Cancer Discov* 2011; **1**: 103–105.
- Li Y, Vandenberg 2nd TG, Wang Z, Kong D, Ali S, Philip PA et al. miR-146a suppresses invasion of pancreatic cancer cells. *Cancer Res* 2010; **70**: 1486–1495.

- 51 Liptay S, Weber CK, Ludwig L, Wagner M, Adler G, Schmid RM. Mitogenic and antiapoptotic role of constitutive NF-kappaB/Rel activity in pancreatic cancer. *Int J Cancer* 2003; **105**: 735–746.
- 52 Zhang H, Ma G, Dong M, Zhao M, Shen X, Ma Z et al. Epidermal growth factor promotes invasiveness of pancreatic cancer cells through NF-kappaB-mediated proteinase productions. *Pancreas* 2006; **32**: 101–109.
- 53 Banerjee S, Wang Z, Kong D, Sarkar FH. 3,3'-Diindolylmethane enhances chemosensitivity of multiple chemotherapeutic agents in pancreatic cancer. *Cancer Res* 2009; **69**: 5592–5600.
- 54 Greten FR, Weber CK, Greten TF, Schneider G, Wagner M, Adler G et al. Stat3 and NF-kappaB activation prevents apoptosis in pancreatic carcinogenesis. *Gastroenterology* 2002; **123**: 2052–2063.
- 55 Infante JR, Jones SF, Bendell JC, Spigel DR, Yardley DA, Weekes CD et al. A phase I, dose-escalation study of pomalidomide (CC-4047) in combination with gemcitabine in metastatic pancreas cancer. *Eur J Cancer* 2011; **47**: 199–205.
- 56 Liu WM, Nizar S, Dalgleish AG. Gemcitabine and lenalidomide combination in a patient with metastatic pancreatic cancer: a case study. *Med Oncol* 2010; **27**: 430–433.
- 57 Shan YS, Lin PW. A phase I study of combination of intravenous gemcitabine, oxaliplatin, and 5-FU with daily oral thalidomide (GOFT) in metastatic/locally advanced pancreatic carcinoma patients. *Hepatogastroenterology* 2007; **54**: 2141–2145.
- 58 Lo M, Ling V, Low C, Wang YZ, Gout PW. Potential use of the anti-inflammatory drug, sulfasalazine, for targeted therapy of pancreatic cancer. *Curr Oncol* 2010; **17**: 9–16.
- 59 Muerkoster S, Arlt A, Witt M, Gehrz A, Haye S, March C et al. Usage of the NF-kappaB inhibitor sulfasalazine as sensitizing agent in combined chemotherapy of pancreatic cancer. *Int J Cancer* 2003; **104**: 469–476.
- 60 Lee DF, Hung MC. Advances in targeting IKK and IKK-related kinases for cancer therapy. *Clin Cancer Res* 2008; **14**: 5656–5662.
- 61 Nogueira L, Ruiz-Ontanon P, Vazquez-Barquero A, Moris F, Fernandez-Luna JL. The NFkappaB pathway: a therapeutic target in glioblastoma. *Oncotarget* 2011; **2**: 646–653.
- 62 Lin Y, Bai L, Chen W, Xu S. The NF-kappaB activation pathways, emerging molecular targets for cancer prevention and therapy. *Expert Opin Ther Targets* 2010; **14**: 45–55.
- 63 Ramaswamy B, Bekaii-Saab T, Schaaf LJ, Lesinski GB, Lucas DM, Young DC et al. A dose-finding and pharmacodynamic study of bortezomib in combination with weekly paclitaxel in patients with advanced solid tumors. *Cancer Chemother Pharmacol* 2010; **66**: 151–158.
- 64 Arlt A, Bauer I, Schafmayer C, Tepel J, Muerkoster SS, Brosch M et al. Increased proteasome subunit protein expression and proteasome activity in colon cancer relate to an enhanced activation of nuclear factor E2-related factor 2 (Nrf2). *Oncogene* 2009; **28**: 3983–3996.
- 65 Lenz HJ. Clinical update: proteasome inhibitors in solid tumors. *Cancer Treat Rev* 2003; **29**(Suppl 1): 41–48.
- 66 Hoeller D, Dikic I. Targeting the ubiquitin system in cancer therapy. *Nature* 2009; **458**: 438–444.
- 67 Kanai M, Yoshimura K, Asada M, Imaizumi A, Suzuki C, Matsumoto S et al. A phase I/II study of gemcitabine-based chemotherapy plus curcumin for patients with gemcitabine-resistant pancreatic cancer. *Cancer Chemother Pharmacol* 2011; **68**: 157–164.
- 68 Wang Z, Desmoulin S, Banerjee S, Kong D, Li Y, Deraniyagala RL et al. Synergistic effects of multiple natural products in pancreatic cancer cells. *Life Sci* 2008; **83**: 293–300.
- 69 Shanmugam MK, Nguyen AH, Kumar AP, Tan BK, Sethi G. Targeted inhibition of tumor proliferation, survival, and metastasis by pentacyclic triterpenoids: Potential role in prevention and therapy of cancer. *Cancer Lett* 2012; **320**: 158–170.
- 70 Banerjee S, Kaseb AO, Wang Z, Kong D, Mohammad M, Padhys S et al. Antitumor activity of gemcitabine and oxaliplatin is augmented by thymoquinone in pancreatic cancer. *Cancer Res* 2009; **69**: 5575–5583.
- 71 Rausch V, Liu L, Kallifatidis G, Baumann B, Mattern J, Gladkch J et al. Synergistic activity of sorafenib and sulforaphane abolishes pancreatic cancer stem cell characteristics. *Cancer Res* 2010; **70**: 5004–5013.
- 72 Lampe JW. Sulforaphane: from chemoprevention to pancreatic cancer treatment? *Gut* 2009; **58**: 900–902.
- 73 Kallifatidis G, Rausch V, Baumann B, Apel A, Beckermann BM, Groth A et al. Sulforaphane targets pancreatic tumour-initiating cells by NF-kappaB-induced antiapoptotic signalling. *Gut* 2009; **58**: 949–963.
- 74 Konig A, Fernandez-Zapico ME, Ellenrieder V. Primers on molecular pathways—the NFAT transcription pathway in pancreatic cancer. *Pancreatol* 2010; **10**: 416–422.
- 75 Baumgart S, Ellenrieder V, Fernandez-Zapico ME. Oncogenic transcription factors: cornerstones of inflammation-linked pancreatic carcinogenesis. *Gut* (e-pub ahead of print 13 October 2011; doi:10.1136/gutjnl-2011-301008).
- 76 Mancini M, Toker A. NFAT proteins: emerging roles in cancer progression. *Nat Rev Cancer* 2009; **9**: 810–820.
- 77 Nayak A, Glockner-Pagel J, Vaeth M, Schumann JE, Buttman M, Bopp T et al. Sumoylation of the transcription factor NFATc1 leads to its subnuclear relocalization and interleukin-2 repression by histone deacetylase. *J Biol Chem* 2009; **284**: 10935–10946.
- 78 Yoeli-Lerner M, Chin YR, Hansen CK, Toker A. Akt/protein kinase b and glycogen synthase kinase-3beta signaling pathway regulates cell migration through the NFAT1 transcription factor. *Mol Cancer Res* 2009; **7**: 425–432.
- 79 Jauliac S, Lopez-Rodriguez C, Shaw LM, Brown LF, Rao A, Toker A. The role of NFAT transcription factors in integrin-mediated carcinoma invasion. *Nat Cell Biol* 2002; **4**: 540–544.
- 80 Baumgart S, Glesel E, Singh G, Chen NM, Reutlinger K, Zhang J et al. Restricted heterochromatin formation links NFATc2 repressor activity with growth promotion in pancreatic cancer. *Gastroenterology* 2012; **142**: e381–e387.
- 81 Singh SK, Baumgart S, Singh G, Konig AO, Reutlinger K, Hofbauer LC et al. Disruption of a nuclear NFATc2 protein stabilization loop confers breast and pancreatic cancer growth suppression by zoledronic acid. *J Biol Chem* 2011; **286**: 28761–28771.
- 82 Koenig A, Linhart T, Schlegemann K, Reutlinger K, Wegele J, Adler G et al. NFAT-induced histone acetylation relay switch promotes c-Myc-dependent growth in pancreatic cancer cells. *Gastroenterology* 2010; **138**: e1181–e1182.
- 83 Buchholz M, Schatz A, Wagner M, Michl P, Linhart T, Adler G et al. Overexpression of c-myc in pancreatic cancer caused by ectopic activation of NFATc1 and the Ca2+/calineurin signaling pathway. *EMBO J* 2006; **25**: 3714–3724.
- 84 Medyouf H, Ghysdael J. The calcineurin/NFAT signaling pathway: a novel therapeutic target in leukemia and solid tumors. *Cell Cycle* 2008; **7**: 297–303.
- 85 Shibasaki F, Hallin U, Uchino H. Calcineurin as a multifunctional regulator. *J Biochem* 2002; **131**: 1–15.
- 86 Marcen R. Immunosuppressive drugs in kidney transplantation: impact on patient survival, and incidence of cardiovascular disease, malignancy and infection. *Drugs* 2009; **69**: 2227–2243.
- 87 Zheng J, Fang F, Zeng X, Medler TR, Fiorillo AA, Clevenger CV. Negative cross talk between NFAT1 and Stat5 signaling in breast cancer. *Mol Endocrinol* 2011; **25**: 2054–2064.
- 88 Padilla PI, Chang MJ, Pacheco-Rodriguez G, Adamik R, Moss J, Vaughan M. Interaction of FK506-binding protein 13 with brefeldin A-inhibited guanine nucleotide-exchange protein 1 (BIG1): effects of FK506. *Proc Natl Acad Sci USA* 2003; **100**: 2322–2327.
- 89 Cooper JE, Wiseman AC. Novel immunosuppressive agents in kidney transplantation. *Clin Nephrol* 2010; **73**: 333–343.
- 90 Nguyen T, Nioi P, Pickett CB. The Nrf2-antioxidant response element signaling pathway and its activation by oxidative stress. *J Biol Chem* 2009; **284**: 13291–13295.
- 91 Martin-Montalvo A, Villalba JM, Navas P, de Cabo R. NRF2, cancer and calorie restriction. *Oncogene* 2011; **30**: 505–520.
- 92 Tian H, Zhang B, Di J, Jiang G, Chen F, Li H et al. Keap1: One stone kills three birds Nrf2, IKKbeta and Bcl-2/Bcl-xL. *Cancer Lett* 2012; **325**: 26–34.
- 93 Hayes JD, McMahon M. NRF2 and KEAP1 mutations: permanent activation of an adaptive response in cancer. *Trends Biochem Sci* 2009; **34**: 176–188.
- 94 Osburn WO, Kensler TW. Nrf2 signaling: an adaptive response pathway for protection against environmental toxic insults. *Mutat Res* 2008; **659**: 31–39.
- 95 Lau A, Villeneuve NF, Sun Z, Wong PK, Zhang DD. Dual roles of Nrf2 in cancer. *Pharmacol Res* 2008; **58**: 262–270.
- 96 Wang XJ, Sun Z, Villeneuve NF, Zhang S, Zhao F, Li Y et al. Nrf2 enhances resistance of cancer cells to chemotherapeutic drugs, the dark side of Nrf2. *Carcinogenesis* 2008; **29**: 1235–1243.
- 97 Singh A, Wu H, Zhang P, Happel C, Ma J, Biswal S. Expression of ABCG2 (BCRP) is regulated by Nrf2 in cancer cells that confers side population and chemoresistance phenotype. *Mol Cancer Ther* 2010; **9**: 2365–2376.
- 98 Pickering AM, Linder RA, Zhang H, Forman HJ, Davies KJ. Nrf2-dependent induction of proteasome and Pa28alphabeta regulator are required for adaptation to oxidative stress. *J Biol Chem* 2012; **287**: 10021–10031.
- 99 Sebens S, Bauer I, Geismann C, Grage-Griebenow E, Ehlers S, Kruse ML et al. Inflammatory macrophages induce Nrf2 transcription factor-dependent proteasome activity in colonic NCM460 cells and thereby confer anti-apoptotic protection. *J Biol Chem* 2011; **286**: 40911–40921.
- 100 Hu Y, Ju Y, Lin D, Wang Z, Huang Y, Zhang S et al. Mutation of the Nrf2 gene in non-small cell lung cancer. *Mol Biol Rep* 2011; **39**: 4743–4747.
- 101 Kim YR, Oh JE, Kim MS, Kang MR, Park SW, Han JY et al. Oncogenic NRF2 mutations in squamous cell carcinomas of oesophagus and skin. *J Pathol* 2010; **220**: 446–451.
- 102 Eades G, Yang M, Yao Y, Zhang Y, Zhou Q. miR-200a regulates Nrf2 activation by targeting Keap1 mRNA in breast cancer cells. *J Biol Chem* 2011; **286**: 40725–40733.



- 103 Kinch L, Grishin NV, Brugarolas J. Succination of Keap1 and Activation of Nrf2-Dependent Antioxidant Pathways in FH-Deficient Papillary Renal Cell Carcinoma Type 2. *Cancer Cell* 2011; **20**: 418–420.
- 104 Nioi P, Nguyen T. A mutation of Keap1 found in breast cancer impairs its ability to repress Nrf2 activity. *Biochem Biophys Res Commun* 2007; **362**: 816–821.
- 105 Wang R, An J, Ji F, Jiao H, Sun H, Zhou D. Hypermethylation of the Keap1 gene in human lung cancer cell lines and lung cancer tissues. *Biochem Biophys Res Commun* 2008; **373**: 151–154.
- 106 Akhdar H, Loyer P, Rauch C, Corlu A, Guillouzo A, Morel F. Involvement of Nrf2 activation in resistance to 5-fluorouracil in human colon cancer HT-29 cells. *Eur J Cancer* 2009; **45**: 2219–2227.
- 107 DeNicola GM, Karreth FA, Humpton TJ, Gopinathan A, Wei C, Frese K et al. Oncogene-induced Nrf2 transcription promotes ROS detoxification and tumorigenesis. *Nature* 2011; **475**: 106–109.
- 108 Hong YB, Kang HJ, Spentzos D, Fountzilias E, Francoeur N, Sanisetty S, Grammatikos AP et al. Keap1 mutations and Nrf2 pathway activation in epithelial ovarian cancer. *Cancer Res* 2011; **71**: 5081–5089.
- 109 Lister A, Nedjadi T, Kitteringham NR, Campbell F, Costello E, Lloyd B et al. Nrf2 is overexpressed in pancreatic cancer: implications for cell proliferation and therapy. *Mol Cancer* 2011; **10**: 37.
- 110 Singh A, Misra V, Thimmulappa RK, Lee H, Ames S, Hoque MO et al. Dysfunctional KEAP1-NRF2 interaction in non-small-cell lung cancer. *PLoS Med* 2006; **3**: e420.
- 111 Solis LM, Behrens C, Dong W, Suraokar M, Ozburn NC, Moran CA et al. Nrf2 and Keap1 abnormalities in non-small cell lung carcinoma and association with clinicopathologic features. *Clin Cancer Res* 2010; **16**: 3743–3753.
- 112 Kim J, Cha YN, Surh YJ. A protective role of nuclear factor-erythroid 2-related factor-2 (Nrf2) in inflammatory disorders. *Mutat Res* 2010; **690**: 12–23.
- 113 Sebens S, Bauer I, Geismann C, Grage-Griebenow E, Ehlers S, Kruse ML et al. Inflammatory macrophages induce NRF2 dependent proteasome activity in colonic NCM460 cells and thereby confer anti-apoptotic protection. *J Biol Chem* 2011; **286**: 40911–40921.
- 114 Singh S, Vrishni S, Singh BK, Rahman I, Kakkar P. Nrf2-ARE stress response mechanism: a control point in oxidative stress-mediated dysfunctions and chronic inflammatory diseases. *Free Radic Res* 2010; **44**: 1267–1288.
- 115 Du ZX, Yan Y, Zhang HY, Liu BQ, Gao YY, Niu XF et al. Proteasome inhibition induces a p38 MAPK pathway-dependent antiapoptotic program via Nrf2 in thyroid cancer cells. *J Clin Endocrinol Metab* 2011; **96**: E763–E771.
- 116 Singh A, Bodas M, Wakabayashi N, Bunz F, Biswal S. Gain of Nrf2 function in non-small-cell lung cancer cells confers radioresistance. *Antioxid Redox Signal* 2010; **13**: 1627–1637.
- 117 Keum YS. Regulation of the Keap1/Nrf2 system by chemopreventive sulforaphane: implications of posttranslational modifications. *Ann NY Acad Sci* 2011; **1229**: 184–189.
- 118 Zhao CR, Gao ZH, Qu XJ. Nrf2-ARE signaling pathway and natural products for cancer chemoprevention. *Cancer Epidemiol* 2010; **34**: 523–533.
- 119 Lee JS, Surh YJ. Nrf2 as a novel molecular target for chemoprevention. *Cancer Lett* 2005; **224**: 171–184.
- 120 Wang XJ, Hayes JD, Henderson CJ, Wolf CR. Identification of retinoic acid as an inhibitor of transcription factor Nrf2 through activation of retinoic acid receptor alpha. *Proc Natl Acad Sci USA* 2007; **104**: 19589–19594.
- 121 Zhou W, Lo SC, Liu JH, Hannink M, Lubahn DB. ERbeta: a potent inhibitor of Nrf2 transcriptional activity. *Mol Cell Endocrinol* 2007; **278**: 52–62.
- 122 Arlt A, Sebens S, Krebs S, Geismann C, Grossmann M, Kruse ML et al. Inhibition of the Nrf2 transcription factor by the alkaloid trigonelline renders pancreatic cancer cells more susceptible to apoptosis through decreased proteasomal gene expression and proteasome activity. *Oncogene*. (e-pub ahead of print 29 October 2012; doi:10.1038/onc.2012.493).
- 123 Gupta S, Pramanik D, Mukherjee R, Campbell NR, Elumalai S, de Wilde RF et al. Molecular determinants of retinoic acid sensitivity in pancreatic cancer. *Clin Cancer Res* 2012; **18**: 280–289.
- 124 Tang X, Wang H, Fan L, Wu X, Xin A, Ren H et al. Luteolin inhibits Nrf2 leading to negative regulation of the Nrf2/ARE pathway and sensitization of human lung carcinoma A549 cells to therapeutic drugs. *Free Radic Biol Med* 2011; **50**: 1599–1609.
- 125 Boettler U, Sommerfeld K, Volz N, Pahlke G, Teller N, Somoza V et al. Coffee constituents as modulators of Nrf2 nuclear translocation and ARE (Epre)-dependent gene expression. *J Nutr Biochem* 2011; **22**: 426–440.
- 126 Lee S, Lim MJ, Kim MH, Yu CH, Yun YS, Ahn J et al. An effective strategy for increasing the radiosensitivity of Human lung Cancer cells by blocking Nrf2-dependent antioxidant responses. *Free Radic Biol Med* 2012; **53**: 807–816.
- 127 Serfling E, Berberich-Siebelt F, Avots A, Chuvpilo S, Klein-Hessling S, Jha MK et al. NFAT and NF-kappaB factors—the distant relatives. *Int J Biochem Cell Biol* 2004; **36**: 1166–1170.
- 128 Hai L, Kawarabayashi Y, Imai Y, Honda A, Inoue R. Counteracting effect of TRPC1-associated Ca<sup>2+</sup> influx on TNF-alpha-induced COX-2-dependent prostaglandin E2 production in human colonic myofibroblasts. *Am J Physiol Gastrointest Liver Physiol* 2011; **301**: G356–G367.
- 129 Cai T, Li X, Ding J, Luo W, Li J, Huang C. A cross-talk between NFAT and NF-kappaB pathways is crucial for nickel-induced COX-2 expression in Beas-2B cells. *Curr Cancer Drug Targets* 2011; **11**: 548–559.
- 130 Fu L, Lin-Lee YC, Pham LV, Tamayo A, Yoshimura L, Ford RJ. Constitutive NF-kappaB and NFAT activation leads to stimulation of the Bly5 survival pathway in aggressive B-cell lymphomas. *Blood* 2006; **107**: 4540–4548.
- 131 Telliez A, Furman C, Pommery N, Henichart JP. Mechanisms leading to COX-2 expression and COX-2 induced tumorigenesis: topical therapeutic strategies targeting COX-2 expression and activity. *Anticancer Agents Med Chem* 2006; **6**: 187–208.
- 132 Jendrossek V. Targeting apoptosis pathways by Celecoxib in cancer. *Cancer Lett* (e-pub ahead of print 21 February 2011; doi:10.1016/j.canlet.2011.01.012).
- 133 Lipton A, Campbell-Baird C, Witters L, Harvey H, Ali S. Phase II trial of gemcitabine, irinotecan, and celecoxib in patients with advanced pancreatic cancer. *J Clin Gastroenterol* 2010; **44**: 286–288.
- 134 Li W, Khor TO, Xu C, Shen G, Jeong WS, Yu S et al. Activation of Nrf2-antioxidant signaling attenuates NF-kappaB-inflammatory response and elicits apoptosis. *Biochem Pharmacol* 2008; **76**: 1485–1489.
- 135 Jiang J, Mo ZC, Yin K, Zhao GJ, Lv YC, Ouyang XP et al. Epigallocatechin-3-gallate prevents TNF-alpha-induced NF-kappaB activation thereby upregulating ABCA1 via the Nrf2/Keap1 pathway in macrophage foam cells. *Int J Mol Med* 2012; **29**: 946–956.
- 136 Yeang HX, Hamdam JM, Al-Huseini LM, Sethu S, Djouhri L, Walsh J et al. Loss of transcription factor nuclear factor-erythroid 2 (NF-E2) p45-related factor-2 (Nrf2) leads to dysregulation of immune functions, redox homeostasis, and intracellular signaling in dendritic cells. *J Biol Chem* 2012; **287**: 10556–10564.
- 137 Wakabayashi N, Slocum SL, Skoko JJ, Shin S, Kensler TW. When NRF2 talks, who's listening? *Antioxid Redox Signal* 2010; **13**: 1649–1663.
- 138 Kim JE, You DJ, Lee C, Ahn C, Seong JY, Hwang JI. Suppression of NF-kappaB signaling by KEAP1 regulation of IKKbeta activity through autophagic degradation and inhibition of phosphorylation. *Cell Signal* 2010; **22**: 1645–1654.
- 139 Yu M, Li H, Liu Q, Liu F, Tang L, Li C et al. Nuclear factor p65 interacts with Keap1 to repress the Nrf2-ARE pathway. *Cell Signal* 2011; **23**: 883–892.
- 140 Wang S, Penchala S, Prabhu S, Wang J, Huang Y. Molecular basis of traditional Chinese medicine in cancer chemoprevention. *Curr Drug Discov Technol* 2010; **7**: 67–75.
- 141 Goel A, Aggarwal BB. Curcumin, the golden spice from Indian saffron, is a chemosensitizer and radiosensitizer for tumors and chemoprotector and radioprotector for normal organs. *Nutr Cancer* 2010; **62**: 919–930.
- 142 Hatcher H, Planalp R, Cho J, Torti FM, Torti SV. Curcumin: from ancient medicine to current clinical trials. *Cell Mol Life Sci* 2008; **65**: 1631–1652.



*Oncogenesis* is an open-access journal published by Nature Publishing Group. This work is licensed under the Creative Commons Attribution-NonCommercial-No Derivative Works 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>