

Copy Number Variation and Differential Expression of a Protective Endogenous Retrovirus in Sheep

Barbara Viginier^{1,2}, Christine Dolmazon¹, Isabelle Lantier³, Frédéric Lantier³, Fabienne Archer¹, Caroline Leroux¹, Christophe Terzian^{1,2*}

1 UMR754 "Rétrovirus et Pathologie Comparée" INRA, Université Lyon 1, Université de Lyon, Lyon, France, **2** Ecole Pratique des Hautes Etudes, Paris, France, **3** INRA Centre de Tours, Nouzilly, France

Abstract

The Jaagsiekte sheep retrovirus exJSRV and its endogenous counterpart enJSRV co-exist in sheep. exJSRV, a betaretrovirus, is the etiological agent of ovine pulmonary adenocarcinoma, and it has been demonstrated *in vitro* that an enJSRV Gag variant bearing the R-to-W amino acid change at position 21 was able to block exJSRV budding from the cells, providing a potential protective role for the host. In this work, we developed a fast mutation detection assay based on the oligo ligation assay (OLA) that permits the quantification of the relative proportions of the R21 and W21 Gag variants present in individual genomes and in cDNA obtained from normal and exJSRV-induced lung tumors. We have shown that the W21/R21 ratio is variable within and between breeds. We also describe for the first time that putative protecting enJSRV variants were expressed in alveolar type II cells (AECII), the major target of exJSRV.

Citation: Viginier B, Dolmazon C, Lantier I, Lantier F, Archer F, et al. (2012) Copy Number Variation and Differential Expression of a Protective Endogenous Retrovirus in Sheep. PLoS ONE 7(7): e41965. doi:10.1371/journal.pone.0041965

Editor: Jean-Pierre Vartanian, Institut Pasteur, France

Received: May 28, 2012; **Accepted:** June 27, 2012; **Published:** July 24, 2012

Copyright: © 2012 Viginier et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by Institut National de la Recherche Agronomique, Ecole Pratique des Hautes Etudes, Agence National Recherche "Genomobile" and Fondation Innovation en Infectiologie. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: christophe.terzian@univ-lyon1.fr

Introduction

Endogenous retroviruses (ERVs) represent a class of retroviruses that are inherited through the germline of their hosts like cellular genes. ERVs represent a significant portion of the host genomes (i.e., 8% of the human genome) and are generally inactivated by point mutations and/or indels [1]. However, some of these endogenous retroviruses still encode functional retroviral proteins. Two main hypotheses have been proposed to understand the maintenance of the coding capacities of ERVs: (1) endogenous retroviral proteins are necessary for the replication of ERVs, and (2) endogenous retroviral proteins are beneficial to their hosts. Several important results have shed light on the benefit of ERV proteins to their hosts. One well-illustrated beneficial impact of ERVs concerns the protection of the host from infection by related exogenous retroviruses. Protection can occur at the entry level of exogenous retrovirus through receptor interference resulting from competition for receptor binding between the exogenous and endogenous envelopes [2]. Protection can also occur later during the replication cycle of the retrovirus, either before the integration of the retroviral genome into the host nucleus [3] or during the assembly of the retroviral particles [4].

The enJSRV/exJSRV retroviruses in sheep represent a powerful model for the study of interactions between an ERV, its exogenous counterpart and their host. The sheep genome contains approximately 30 copies of the endogenous retrovirus enJSRV, which is highly related to the exogenous betaretrovirus exJSRV (90%–98% identity at the amino acid level), the etiological agent of ovine pulmonary adenocarcinoma [5–8].

Among the enJSRVs, it has been shown that the enJS56A1 provirus was able to block the release of infectious JSRV particles when co-expressed in cultured cells, illustrating a novel mechanism of retroviral interference [9]. The determinant of this blocking process resides at the level of a single residue at position 21 in the Gag precursor protein. The arginine residue at position 21 (R21) of the exJSRV Gag sequence is replaced by a tryptophan (W21) in the enJS56A1 Gag sequence. This single change confers to enJS56A1 the ability to block the normal trafficking of JSRV particles to the plasma membrane. A complete set of enJSRV proviruses was characterized from a genomic library for the Texel breed, cloned and sequenced, revealing the presence of a second provirus containing non-synonymous mutations inducing the R->W change; this provirus was named enJSRV20 [10]. Interestingly, phylogenetic analysis strongly suggested that these two protecting proviruses became fixed in the sheep during its domestication [10].

Nothing is known concerning the variation of W21 enJSRV between individuals and between breeds. Therefore, we screened 42 sheep genomes and quantified the presence of W21 using an approach based on the oligo ligation assay (OLA), also known as the fast mutation detection assay [11]. Moreover, one can ask if protection against sheep exJSRV infection by W21 enJSRV, which was shown *in vitro*, is also relevant *in vivo*. Hence, previous data have shown that the enJSRVs are predominantly expressed in the placenta, and exJSRV infects and replicates in AECII [8] but it has never been shown if W21 RNA is present or not in these target cells. We then quantified the relative level of the W21 variant RNAs in AECII derived from tumoral and non-tumoral lungs.

Table 1. List of breeds analyzed. IL16 to 63 are inbred lines derived from the *Pre-Alpes du Sud* French breed.

Breed	No. of individuals
IL16	4
IL19	2
IL36	3
IL39	4
IL63	4
Texel (Tex)	2
Ile de France (IdF)	2
Bleu de Maine (BdM)	2
Rouge de l'Ouest (RdO)	2
Vendéen (Ven)	3
Solognot (Sol)	3
Pol Dorset (Pol)	3
Suffolk (Suf)	3
Thônes et Marthod (Tho)	2
Romane (Rom)	2

doi:10.1371/journal.pone.0041965.t001

Our results indicate that W21 is present at variable frequencies in the genomes of several breeds and is expressed in the lung, the target of JSRV-induced transformation. Furthermore, we observed that tumoral lung samples rarely express the W21 variant, and when expressed, that it is expressed at a low level compared to that the level in normal lung samples.

Materials and Methods

Ethics Statement

All animal experiments have been performed in compliance with our institutional and national guidelines in accordance with

the European Community Council Directive 86/609/EEC. The INRA Tours ethics committee approved the experimental protocols.

Sample Collection

Forty-one DNA samples extracted from total blood have been collected from 10 ovine breeds and five inbred lines (IL) from the experimental flock maintained at the large animal facility, INRA, Tours Nouzilly France (Table 1). IL 16 to 63 are inbred lines derived from the *Pre-Alpes* breed. Fifteen samples of total lung DNA have been prepared from 7 Manech sheep from the French Basque Region that presented with respiratory symptoms and massive tumoral lesions upon gross examination and from 8 sheep of mixed breeds from a local slaughterhouse (Corbas, France) without macroscopic pulmonary lesions. Total DNA has been extracted using the Fast DNA kit as recommended by the manufacturer (Qbiogene, France). The exJSRV statuses of the sheep have been confirmed by histopathological examination to identify tumoral lesions (Anatomy and Pathological Cytology laboratory, Hospices Civils de Lyon, Pr Françoise Thivolet-Béjui) and by PCR detection of the exJSRV genome using exJSRV-specific primers targeting the 3' end of the *env* gene and the LTR (primer sequences available upon request). From the same 15 sheep, total RNA has been extracted using the standard RNeasy protocol (Qiagen, France) from primary AECII derived from non-tumoral (n=5) or tumoral (n=7) lung tissues as previously reported (Archer et al., 2007). The derivation of primary AECII failed for 3 tissues due to the poor conditions of the tissues. For each sample, 500 ng total RNA have been reverse-transcribed into cDNAs using random hexamers and M-MLV RT (Invitrogen) according to the manufacturer's protocol and have been analyzed by PCR to determine their exJSRV status using specific primers.

DNA and cDNA Gag Amplification

The PCRs were performed using DNAs or cDNAs and the primers enJSRV₄₈₂ (5'-TCCTCGCCACTACTCTTATT-3') and enJSRV₉₅₃ (5'-GGAGGGTCGCTTTTAACC-3') to amplify specifically the Gag region of enJSRV (Fig. 1A); the corresponding




Figure 1. A. Canonical structure of the enJSRV provirus. The W/R variant at position 21 is indicated by a star. The 470 bp amplicon is represented by a gray bar. B. Ligation products obtained from the OLA detection assay of W21 and R21 variants. The sequence of the phosphate-modified downstream oligo is indicated in italics. The upstream oligo that detected the R21 to W21 mutation is labeled with HEX, whereas the two FAM-labeled oligos detected the R21 variant. Probes that were mismatched to the target by a single nucleotide at their junctions were not ligated. The ligation products were discriminated by size (60 nt for W21 coding sequence vs. 64 nt for R21 coding sequences) and color (HEX for W21 vs. FAM for R21). The W and R codons are underlined. doi:10.1371/journal.pone.0041965.g001




Figure 2. Boxplot representation of triplicate W21/R21 ratio measurements among randomly selected 10 samples. For each sample, W21/R21 measurement was repeated three times. Each dot represents a single measurement. Numbers correspond to the samples listed in Figure 1. NT1 and NT2 correspond to DNA extracted from two non-tumoral samples. doi:10.1371/journal.pone.0041965.g002

exJSRV region is not amplified using these same primers (not shown). The PCR reactions were performed in 50 µl volumes containing Taq polymerase buffer, 1.65 mM MgCl₂, 200 µM

each dNTP, 0.2 µM each primer, 100 to 500 ng of genomic DNA, and 1.25 U of Taq polymerase (Eurobio). The amplification conditions consisted of an initial 5 min denaturation at 95°C




Figure 3. Distribution of W21 variants among 41 genomes from 10 breeds and 5 inbred lines. Individuals are numbered according to Table 1. The W21/R21 ratios are indicated as percentage. doi:10.1371/journal.pone.0041965.g003

representative of Texel individuals. Thanks to the OLA approach, we are now able to estimate the R21/W21 ratio in a large sample of individuals, and do not restrict our screen to enJS65A1 but consider all enJSRV types coding the W21 Gag mutant.

We then tested the hypothesis that a significant level of W21 Gag could provide a selective advantage for the hosts. Thus, we looked for the presence of W21 *gag* expression in AECII, the primary target of exJSRV together with Clara cells in the lung.

The W21 Gag Expression Level in Sheep Lung Cells Depends on the Tumoral Status of the Sample

We compared the W21/R21 variant frequencies in DNA extracted from the 15 studied lungs (7 tumoral and 8 non-tumoral tissues) and, when available, *i.e.*, for 12 of the 15 animals, from the DNA from the corresponding *in vitro*-derived primary AECII (5 tumoral and 7 non-tumoral), established as previously reported [14]. All the tumoral alveolar type II cells harbored the JSRV provirus genome, while the non tumoral cell lines were negative [14]. We first estimated the W21/R21 ratios in the genomic DNA. The W21/R21 ratio varied from 13 to 48%, and the medians from the tumoral (T) and non-tumoral (NT) genomic samples were not significantly different (Wilcoxon test, p -value = 0.8). However, we observed a significant difference when we compared the medians between genomic DNA and cDNA extracted from tumoral and non-tumoral samples (Fig. 4): the median of W21/R21 cDNA (RNA) ratio was significantly lower than the median of W21/R21 genomic copies (DNA) in tumoral samples, whereas such a difference did not exist in non-tumoral samples (Fig. 4). This result indicates that the expression of W21 *gag* was null or lower in tumoral AECII and lung tissues than in non-tumoral samples. Whether the absence of W21 expression favors *in vivo* the replication of exJSRV could not be assessed in this study.

References

- Jern P, Coffin JM. Effects of retroviruses on host genome function. *Annu Rev Genet.* 2008;42: 709–32.
- Ikeda H, Sugimura H (1989) Fv-4 resistance gene: a truncated endogenous murine leukemia virus with ecotropic interference properties. *J Virol* 63: 5405–5412.
- Pryciak PM, Varmus HE (1992) Fv-1 restriction and its effects on murine leukemia virus integration *in vivo* and *in vitro*. *J Virol* 66: 5959–5966.
- Black LR, Aiken C (2010) TRIM5alpha disrupts the structure of assembled HIV-1 capsid complexes *in vitro*. *J Virol* 84: 6564–6569.
- Palmarini M, Sharp JM, de las Heras M, Fan H (1999) Jaagsiekte sheep retrovirus is necessary and sufficient to induce a contagious lung cancer in sheep. *J Virol* 73: 6964–6972.
- Palmarini M, Hallwirth C, York D, Murgia C, de Oliveira T, et al. (2000) Molecular cloning and functional analysis of three type D endogenous retroviruses of sheep reveal a different cell tropism from that of the highly related exogenous jaagsiekte sheep retrovirus. *J Virol* 74: 8065–8076.
- Carlson J, Lyon M, Bishop J, Vaiman A, Cribiu E, et al. (2003) Chromosomal distribution of endogenous Jaagsiekte sheep retrovirus proviral sequences in the sheep genome. *J Virol* 77: 9662–9668.
- Leroux C, Girard N, Cottin V, Greenland T, Mornex JF, et al. (2007) Jaagsiekte Sheep Retrovirus (JSRV): from virus to lung cancer in sheep. *Vet Res* 38: 211–228.
- Murcia PR, Arnaud F, Palmarini M (2007) The transdominant endogenous retrovirus enJS65A1 associates with and blocks intracellular trafficking of Jaagsiekte sheep retrovirus Gag. *J Virol* 81: 1762–1772.
- Arnaud F, Caporale M, Varela M, Biek R, Chessa B, et al. (2007) A paradigm for virus-host coevolution: sequential counter-adaptations between endogenous and exogenous retroviruses. *PLoS Pathog* 3: e170.
- Landegren U, Kaiser R, Sanders J, Hood L (1988) A ligase-mediated gene detection technique. *Science* 241: 1077–1080.
- Hallwirth C, Maeda N, York D, Fan H (2005) Variable regions 1 and 2 (VR1 and VR2) in JSRV gag are not responsible for the endogenous JSRV particle release defect. *Virus Genes* 30: 59–68.
- Armezzani A, Arnaud F, Caporale M, di Meo G, Iannuzzi L, et al. (2011) The signal peptide of a recently integrated endogenous sheep betaretrovirus envelope plays a major role in eluding gag-mediated late restriction. *J Virol* 85: 7118–7128.
- Archer F, Jacquier E, Lyon M, Chastang J, Cottin V, et al. (2007) Alveolar type II cells isolated from pulmonary adenocarcinoma: a model for JSRV expression *in vitro*. *Am J Respir Cell Mol Biol* 36: 534–540.
- Wang Y, Song Y, Liu Q, Liu C, Wang L, et al. (2010) Quantitative analysis of lentiviral transgene expression in mice over seven generations. *Transgenic Res* 19: 775–784.

Conclusions

In this study using a rapid, discriminative and quantitative molecular technique, we have shown that the W21/R21 ratio varies between individuals from different breeds. Moreover, our results indicate for the first time that W21 and R21 enJSRV are expressed in normal lung tissues and that W21 is not expressed or is expressed at a low level in tumor-derived AECIIs. Our present study did not demonstrate that the absence of W21 *gag* RNA (and therefore W21 Gag proteins) was the causal factor in the development of tumoral cells, but this study does provide new insights into the putative protective role of some endogenous retroviruses against infection by their exogenous counterparts.

Acknowledgments

We are grateful to Dr Hung Fan for the penJSRV65A and pJSR21 plasmids. We also thank Dr François Guiguen and Dr Jean-Christophe Natorp for the *post mortem* collection of tissue samples.

Author Contributions

Conceived and designed the experiments: BV CT. Performed the experiments: BV CD. Analyzed the data: BV CT. Contributed reagents/materials/analysis tools: BV IL FL FA CL. Wrote the paper: BV CL CT.