




Correction to: Composite detection rate as an upper gastrointestinal endoscopy quality measure correlating with detection of neoplasia

Marcin Romańczyk^{1,2}  · Bartosz Ostrowski¹ · Tomasz Marek¹ · Tomasz Romańczyk² · Małgorzata Błaszczyńska³ · Krzysztof Budzyń^{1,2} · Maciej Bugajski¹ · Mateusz Koziej⁴ · Maciej Kajor⁵ · Krzysztof Januszewski⁶ · Wojciech Zajęcki⁷ · Marek Waluga¹ · Marek Hartleb¹

Published online: 27 June 2022
© The Author(s) 2022

Correction to: Journal of Gastroenterology (2021) 56:651–658
<https://doi.org/10.1007/s00535-021-01790-3>

In the original publication of the article, Table 3 has been published with bold font wrongly which should be limited to the statistically significant p value (< 0.05). The correct Table 3 is given in this correction.

The original article can be found online at <https://doi.org/10.1007/s00535-021-01790-3>.

✉ Marcin Romańczyk
mromancyk@sum.edu.pl

¹ Chair and Department of Gastroenterology and Hepatology, School of Medicine in Katowice, Medical University of Silesia, Medyków 14 Street, 40-752 Katowice, Poland

² H-T. Centrum Medyczne, Tychy, Poland

³ Endoscopy Unit, District Hospitals of Chorzów Trust, Chorzów, Poland

⁴ Department of Anatomy, Jagiellonian University Medical College, Cracow, Poland

⁵ Department of Pathomorphology and Molecular Diagnostic, School of Medicine in Katowice, Medical University of Silesia, Katowice, Poland

⁶ Department of Pathomorphology, Zakład Diagnostyki Mikroskopowej, Dr Krzysztof Januszewski, Ruda Śląska, Poland

⁷ Department of Pathomorphology, District Hospitals of Chorzów Trust, Chorzów, Poland

Table 3 Logistic regression analysis of upper gastrointestinal neoplasm detection risk factors

Variables	Univariate			Multivariate			
	OR	95% CI	<i>p</i> value	OR	95% CI	<i>p</i> value	
Age	1.03	1.01–1.05	< 0.001	1.03	1.01–1.05	0.002	
Male	2.0	1.2–3.4	0.005	2.0	1.2–3.5	0.011	
HD-endoscope	0.95	0.6–1.6	0.858				
Sedation	0.8	0.4–1.3	0.326				
Out-patient	0.4	0.2–0.7	< 0.001	0.7	0.4–1.2	0.207	
CDR > 26%	4.6	2.8–8	< 0.001	5.7	1.5–22.3	0.012	
Endoscopic center*	B	Reference					
	C	1.4	0.6–3.1	0.430	0.8	0.3–2.0	0.666
	A	4.0	2.0–8.1	< 0.001			

HD high definition, *CDR > 26%* examination performed by operator with CDR higher than 26% (group 4), *OR* odds ratio, *CI* confidence interval

*In the univariate analysis center with lowest mean UGN value (center B) was the reference. In the multivariate analysis the centers were graded according to mean UGN value – center B as 1, center C as 2 and center A as 3

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not

included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.