

# Application of Radiosensitizers in Cancer Radiotherapy

This article was published in the following Dove Press journal:  
*International Journal of Nanomedicine*

Liuyun Gong<sup>1</sup>  
Yujie Zhang<sup>2,3</sup>  
Chengcheng Liu<sup>2</sup>  
Mingzhen Zhang<sup>1</sup><sup>2,3</sup>  
Suxia Han<sup>1</sup>

<sup>1</sup>Department of Oncology, The First Affiliated Hospital, Xi'an Jiaotong University, Xi'an, Shaanxi, 710061, People's Republic of China; <sup>2</sup>School of Basic Medical Sciences, Xi'an Jiaotong University, Xi'an, Shaanxi, 710061, People's Republic of China; <sup>3</sup>Key Laboratory of Environment and Genes Related to Diseases, Xi'an Jiaotong University, Ministry of Education, Xi'an, Shaanxi, 710061, People's Republic of China

**Abstract:** Radiotherapy (RT) is a cancer treatment that uses high doses of radiation to kill cancer cells and shrink tumors. Although great success has been achieved on radiotherapy, there is still an intractable challenge to enhance radiation damage to tumor tissue and reduce side effects to healthy tissue. Radiosensitizers are chemicals or pharmaceutical agents that can enhance the killing effect on tumor cells by accelerating DNA damage and producing free radicals indirectly. In most cases, radiosensitizers have less effect on normal tissues. In recent years, several strategies have been exploited to develop radiosensitizers that are highly effective and have low toxicity. In this review, we first summarized the applications of radiosensitizers including small molecules, macromolecules, and nanomaterials, especially those that have been used in clinical trials. Second, the development states of radiosensitizers and the possible mechanisms to improve radiosensitizers sensibility are reviewed. Third, the challenges and prospects for clinical translation of radiosensitizers in oncotherapy are presented.


**Keywords:** radiosensitizers, cancer radiotherapy, therapeutics, nanomedicine, mechanism

## Introduction

Cancer remains one of the greatest challenges to human health. World Health Organization (WHO) reported that about 8.8 million deaths worldwide were due to cancer in 2015, and the deaths are expected to break through 13 million in 2030 according to the report by the International Agency for Research on Cancer (IARC). To reduce the deaths from cancer, several strategies have been developed in recent years to improve cancer therapy including surgery, radiotherapy, chemotherapy, immunotherapy, targeted therapy, hormone therapy, stem cell transplant and precision medicine.<sup>1</sup> Among them, radiotherapy (RT) is considered as one important and effective modality to kill or control tumors since Marie Curie, the Nobel Prize winner, discovered radioactivity.<sup>2</sup> Typically, RT is a treatment modality to cancer cells by using high-energy photon radiation such as X-rays, gamma ( $\gamma$ )-rays, and others. RT can take effect via direct and indirect mechanisms to destroy cancer cells and tumor tissue (Figure 1).

In the direct action, radiation directly induces single-strand breaks (SSB) and double-strand breaks (DSB) in DNA, resulting in the termination of cell division and proliferation, or even cell necrosis and apoptosis. In the case of indirect action, radiation induces the generation of ROS, which can induce cellular stress in, and injure biomolecules, and ultimately alter cellular signaling pathways. Clinical studies have shown that more than half (about 70%) of patients need to receive RT,

Correspondence: Mingzhen Zhang;  
Suxia Han  
Email [mzhang21@xjtu.edu.cn](mailto:mzhang21@xjtu.edu.cn);  
[shan87@xjtu.edu.cn](mailto:shan87@xjtu.edu.cn)



**Figure 1** Schematic of the mechanism of ionizing radiation (IR) in RT. In the case of direct effect, IR directly damages the DNA, which, if unrepaired, results in cell death or permanent growth arrest. In the case of indirect effect, ROS are formed by the radiolysis of a large amount of water and oxygen, and then the ROS damage the DNA. There are many types of DNA damage, such as base change, SSB, DSB, cross-linkage with protein or with other DNA molecules.

and in some cases RT is the only kind of cancer treatment.<sup>3</sup> Therefore, there is a great need to develop approaches to improve radiosensitivity.

Innovative technologies can provide alternative strategies to improve RT efficiency. For example, image-guided radiation therapy (IGRT) is the use of imaging during radiation therapy to improve the precision and accuracy of treatment delivery. IGRT can be used to treat tumors in areas of the body that move, such as the lungs. RT machines are equipped with imaging technology to allow your doctor to image the tumor before and during treatment. By comparing these images to the reference images taken during simulation, the patient's position and/or the radiation beams may be adjusted to more precisely target the radiation dose to the tumor. To help align and target the radiation equipment, some IGRT procedures may use fiducial markers, ultrasound, MRI, X-ray images of bone structure, CT scan, 3D body surface mapping, electromagnetic transponders or colored ink tattoos on the skin.<sup>4</sup> Intensity-modulated radiation therapy (IMRT) is an advanced mode of high-precision RT that uses computer-


controlled linear accelerators to deliver precise radiation doses to a malignant tumor or specific areas within the tumor.<sup>5</sup> Although the abovementioned innovative technologies greatly improve the therapeutic effect, there are still obstacles such as cancer stem cells and tumor heterogeneity making it difficult to use RT alone to cure tumors. Radiosensitizers with the ability to increase the radiosensitivity of tumor tissue and pharmacologically decrease normal tissue toxicity are expected to be an efficient way to improve RT.<sup>6</sup>

Radiosensitizers are compounds that, when combined with radiation, achieve greater tumor inactivation than would have been expected from the additive effect of each modality. G E Adams, a pioneer in the field of RT, classified radiosensitizers into five categories: (1) suppression of intracellular thiols or other endogenous radioprotective substances; (2) formation of cytotoxic substances by radiolysis of the radiosensitizer; (3) inhibitors of repair of biomolecules; (4) thymine analogs that can incorporate into DNA; and (5) oxygen mimics that have electrophilic activity.<sup>7,8</sup> This classification was based on the mechanism of DNA damage and repair and indicated the direction for radiosensitizers at the early stage. However, with the continuous technological innovation, more and more materials and drugs with radiotherapy sensitization have been defined as radiosensitizers. In addition, some in-depth mechanisms for radiosensitization have also been discovered.<sup>9,10</sup> According to the latest research, radiosensitizers can be classified into three categories based on their structures: small molecules (Figure 2), macromolecules (Table 1), and nanomaterials (Table 2).<sup>11</sup> In the following part, the applications, the main role, and influencing factors of these three types of radiosensitizers are first summarized, especially those have currently entered clinical trials. Second, the development status and the mechanism of action of the radiosensitizer are also summarized. Third, the future development and application of the radiosensitizer was presented.

## Small Molecules

### Oxygen

Hypoxia in tumor microenvironment is one of the major limitations to radiotherapy. Tumor cells in the hypoxic microenvironment are much more resistant to radiation than in the normal oxygen microenvironment.<sup>12–14</sup> Oxygen enhancement ratio (OER) or oxygen enhancement effect in radiobiology refers to the enhancement of the



**Figure 2** Molecular structures of some representative small-molecule radiosensitizers discussed in this paper.

therapeutic or detrimental effect of ionizing radiation due to the presence of oxygen. This so-called oxygen effect is most notable when cells are exposed to an ionizing radiation dose.<sup>15,16</sup> Oxygen, a potent radiosensitizer, promotes free radical formation through its unique electronic configuration. As the most electrophilic cellular molecule, oxygen is easily reduced by electrons formed from the incident radiation. After oxygenated tumor irradiation, energy transfer results in the radiolysis of water with the initial formation of an ion radical that then forms the highly reactive hydroxyl radical after reaction with another water molecule. Oxygen leads to the formation of peroxide after reaction with the hydroxyl radical. Then, the peroxide results in permanent cellular and DNA damage.<sup>13</sup>

Accompanied with solid tumor growth, the surrounding vasculatures are not in sufficient quantities to supply oxygen to the new cells, the cancer cell mass becomes heterogeneous gradually, and necrosis occurs following ischemia. Normally, cancer cells undergo apoptosis through the p53 pathway, while those heterogeneous cells adapt to the hypoxic environment efficiently by activation of additional signaling pathways, especially the hypoxia-

inducible factor (HIF) pathway.<sup>17–19</sup> Studies showed that HIF-1 $\alpha$  was associated with vascular endothelial growth factor (VEGF) signaling pathway, glucose transport, and glycolysis pathway, which could help the tumor to build vasculature.<sup>19–21</sup> Under hypoxia, the cancer cells are more aggressive and resisted radiotherapy significantly. Thus, hypoxia often occurs in most solid tumors and leads to radioresistance both through increasing free radical scavenging and changing patterns of gene expression.<sup>22,23</sup>

More and more research has been devoted to overcoming hypoxia problems, from using high-pressure oxygen tanks and blood substitutes that carried oxygen, to using intricate, accurate approaches that proportionated differences in partial pressure of oxygen (PO<sub>2</sub>) between tumors and healthy tissue.<sup>24,25</sup> Hyperbaric oxygen is the most direct method to ameliorate hypoxia in tumor cells, while this method is inconvenient and may increase complications sometimes.<sup>26,27</sup> A new radiosensitizer, Kochi oxydol-radiation therapy for unresectable carcinomas (KORTUC), is being evaluated by a Phase I/II clinical trial (NCT02757651) for the treatment of malignant tumors

**Table 1** Some Macromolecule Radiosensitizers Discussed in This Paper

Type	Name	Mechanism of Radiosensitivity	Reference
Proteins and Peptides	HER3-ADC	Inhibiting DNA damage repair	130
	SYM004	Inhibiting DNA double strand breaks repair and inducing apoptosis	120
	Cetuximab	Increasing radiation-induced apoptosis and DNA damage	131
	Nimotuzumab	Increasing radiation-induced apoptosis and DNA damage	131
	AMG102	Inhibiting DNA damage repair and increasing radiosensitivity of glioblastoma multiforme	132
	C-reactive peptide	Used as radiotherapy targets	133
	HSP	Used as radiotherapy targets	134
	Paraoxonase-2	Used as radiotherapy targets	135
	EC1301	Assisted by HSP-70 and HMGB1	134
miRNAs	miR-621	Targeting SETDB1	141
	miR-205	Targeting zinc finger E-box binding homeobox 1 (ZEB1) and the ubiquitin-conjugating enzyme Ubc13	142
	miR-144-5p	Targeting ATF2	143
	miR-146a-5p	Activating DNA repair pathway	144
	miR-150	Acting on AKT pathway	145
	miR-99a	Targeting mTOR pathway	146
	miR-139-5p	Repressing multiple gene networks of DNA repair and ROS defense	147
	miR-320a	Inducing cancer cell apoptosis	148
siRNAs Oligonucleotides		Silencing genes related to radioresistance Regulating gene expression	151–155 156–159

that contain numerous hypoxic cancer cells and/or large quantities of antioxidative enzymes.<sup>28</sup>

## Oxygen Mimics

Oxygen mimetics, using the chemical properties of molecular oxygen as a template, have higher electron affinity and better diffusion properties to anoxic tissue than oxygen. As oxygen mimetics can theoretically substitute for oxygen in “fixing” radiation-induced damage of DNA, making it nonrepairable and hence lethal. Therefore, oxygen mimetics are considered as “true radiosensitizers”. The most representative oxygen mimetics are nitro-containing compounds and nitric oxide (NO).<sup>13</sup>

The prototype of electron-affinity radiosensitizers is nitrobenzene, and then researchers focus on nitroimidazole and its derivatives.<sup>29–31</sup> Nitroimidazoles, which undergo enzymatic and radiation-induced redox reactions. These

agents are intrinsic inactive, their effect becomes evident only in the presence of ionizing radiation to “fix” or stabilize DNA radical lesions in oxygen-deficient cells.<sup>32</sup> Misonidazole, a 2-nitroimidazole, is one of the earliest developed nitroimidazoles. In preclinical studies, misonidazole showed better radiosensitizing effect than 5-nitroimidazole or metronidazole (Flagyl®) in the majority of solid murine tumors.<sup>33–35</sup> However, the results were unsatisfactory in clinical trials, since severe neurotoxicity was caused by misonidazole.<sup>36–39</sup> Metronidazole, a 5-substituted nitroimidazole, which has less electron-affinic was proven as an inferior radiosensitizer.<sup>40,41</sup> In conclusion, because of the dose-limiting toxicity at clinically tolerable doses, misonidazole and metronidazole are not the ideal candidates in radiotherapy.<sup>42</sup>

In view of the issues discussed above, further efforts have been made to improve the pharmacokinetic

**Table 2** The List of Nanomaterials Used for Radiosensitization

Nanomaterial	Modification	Size (TEM)	Cell Line/Model	Reference
Au	GSH	<2 nm	U14 tumor models	166
Au	PEG <sub>2k</sub>	2, 5, 19 nm	PC3pip and PC3flu cells	167
Ag	PEG	18 nm	C6 cells	171
Ag	PVP	26.87±3.68 nm	U251 and C6 cells	172
Bi,Gd	PEG	11.3±1.6 nm	MCF-7 and 4T1 cells	174
Hf, Nb, Ta		100 nm		178
Gd	DTPA	3.0±1.0 nm	F98 cells	179
AGuIX		2.1±1 nm	F344 rats	181
AGuIX	DOTA	Sub-5 nm	HepG2 cells	182
Hf		7–31 nm		184
Ta	PEG	65.4±5.6 nm	Balb/c mice	188
Zn, Ga, Ge, Cr, Pr, Ta	PEG	62.8±8.6 nm	Nude mice harboring HepG2 tumors	190
Ta	PEG	119±34 nm	4T1 cells	193
Bi	PEG	3.6 nm	LO2 and 4T1 cells	197
Bi		10–70 nm	4T1 cells	200
Si,Gd,Bi	DOTA	4.5±0.9 nm	A549 cells	202
Fe,Pt	PEI,PEG	10nm	A2780 and A2780DDP cells	204
Co,Mn,Fe,Bi	PEG	11.2±1.4 nm and 14.4±2.4 nm	C6 cells	207
Zn,Fe		5–15 nm	LNCAp cells	208
Quantum dots			H460 cells	214
Ti	PAA	50–100 nm	MIAPaCa-2 cells	215
Ti	PAA	135±65 and 124±65 nm		216
R-O2-FA-CHI-SWCNTs			MDA-MB-231 and ZR-75-1 cells	220
Tf,Se		177, 192, and 312 nm	C6,A375 cells	221
GNP	PEG, RGD	20.90±0.14nm.	HeLa, Hs. 895.Sk and Hs 895.T cells	222

properties of nitroimidazoles. Second-generation nitroimidazole radiosensitizers, such as etanidazole or nimorazole, are designed to increase the hydrophilicity of the reagents and thereby reduce neurotoxicity. For example, etanidazole has better hydrophilicity than misonidazole because its side chain is modified by hydroxyl.<sup>43</sup> Although etanidazole presents lower preclinical toxicity and higher efficacy, it shows no obvious benefit for head and neck cancer patients in randomized studies.<sup>44</sup> Nimorazole, a 5-nitroimidazole, is recommended for the treatment of head and neck cancers in Denmark since its beneficial effects in several clinical trials. Moreover, it has been further explored in an EORTC international trial.<sup>45–51</sup> Notably, the DAHANCA 28 trial demonstrated that hyperfractionated, accelerated radiotherapy with concomitant cisplatin and nimorazole (HART-CN) for patients was

feasible and yielded favorable tumor control.<sup>52</sup> Other nitro compounds have also been exploited for hypoxia radiosensitization. Dinitroazetidine, RRx-001, has been evidenced as an effective radiosensitizer with low toxicity and is now being evaluated in the NCT02871843 clinic trial.<sup>53</sup>

Nitrogen oxides, in particular, NO, act as radiosensitizers through many direct and indirect mechanisms. Similar to the oxidative stress induced by oxygen, NO can “fix” or stabilize radiation-induced DNA damage through nitrosative stress pathways.<sup>54</sup> Oxidative and nitrosative stress pathways involve the generation of reactive species. For example, nitrous acid, peroxyxynitrite (ONOO<sup>-</sup>), and nitric acid produce cytotoxic effects through mechanisms including DNA cross-linking, protein nitrosylation, glutathione depletion, and inhibition of mitochondrial respiration.<sup>55–58</sup>



As an uncharged free radical, NO can diffuse across cell membranes freely and bind to soluble guanylate cyclase (sGC) to induce cyclic GMP production, thereby regulating vascular physiology.<sup>59–61</sup> Researchers have reported that 5-nitroimidazoles and sanazole can release NO.<sup>62,63</sup>

A phase I study of non-small-cell lung cancer (NSCLC) patients suggested that NO donation increased tumor perfusion and, therefore, promoted tumor growth.<sup>64</sup> However, a phase II study of prostate cancer patients claimed that low-dose NO had no direct cytotoxic effect, but could decrease hypoxia through improving blood flow in tumor tissue.<sup>65</sup> Some anticancer drugs approved by US Food and Drug Administration (FDA), such as bevacizumab, sorafenib, and etaracizumab played their roles by blocking the VEGF pathway to some extent.<sup>66</sup> VEGF is overexpressed in anoxia environment, which leads to endothelial cell proliferation and neovascularization. In angiogenesis, there is a positive and negative feedback regulation relationship between VEGF and NO, which maintains vascular homeostasis precisely.<sup>67</sup> In addition, Liebmann et al proved that pre-treatment with NO improved the survival of mice after irradiation.<sup>68</sup>

## Active Compounds from Chinese Herbs

In recent years, more and more researchers reported that active compounds from Chinese herbs such as curcumin,<sup>69–71</sup> resveratrol,<sup>72–74</sup> dihydroartemisinin<sup>75–77</sup> and paclitaxel,<sup>78–80</sup> could enhance tumor radiotherapy sensitivity (Figure 2). Curcumin is a polyphenolic active compound extracted from turmeric. Curcumin exerts anti-inflammatory effect by inhibiting the transcription factor NF- $\kappa$ B, which is involved in both tumorigenesis and radioresistance.<sup>81</sup> In a preclinical study, Chendil et al reported that when treated with RT and curcumin together, the human prostate cancer cell line, PC3 presented three-fold fewer surviving and the mechanism was supposed to have a relationship with NF- $\kappa$ B.<sup>82</sup> In addition, nanocurcumin as a radiosensitizer is being evaluated by a Phase II clinical trial (NCT02724618). Other relevant research on mutant p53 Ewing's sarcoma cells proved that radiosensitivity of curcumin was associated with other p53-response genes.<sup>83</sup>

Resveratrol is an active compound extracted from grapes, knotweed, peanuts, mulberry and other plants. Tan et al proved that resveratrol enhanced the radiosensitivity in nasopharyngeal carcinoma cells by downregulating E2F1.<sup>73</sup> Liao et al found that resveratrol enhanced

radiosensitivity in human NSCLC NCI-H838 cells by inhibiting NF- $\kappa$ B activation.<sup>84</sup> Dihydroartemisinin is a derivative of artemisinin, which can shorten the G<sub>2</sub>/M phase, while increases the G<sub>0</sub>/G<sub>1</sub> and S phase, thereby reducing the radiation resistance.<sup>85</sup> Although the relevant clinical research has not yet been carried out, researchers have demonstrated that resveratrol<sup>86–89</sup> and dihydroartemisinin<sup>90–92</sup> possessed radiosensitization on cancer cells in vitro.

Paclitaxel is widely known as a very good natural anticancer drug.<sup>93,94</sup> As a new type of antimicrotubule drug, paclitaxel can inhibit the microtubule networks formation and prevent the tumor cells proliferation to achieve radiosensitization.<sup>95</sup> Results showed that paclitaxel could obviously enhance the radiosensitivity of inoperable patients with locally advanced esophageal cancer and improve the prognosis of patients with acceptable therapeutic effect.<sup>96</sup> A three-arm randomized Phase III trial (NCT02459457)—comparison of paclitaxel-based three regimens concurrent with radiotherapy for patients with local advanced esophageal cancer and a Phase III study (NCT01591135) of comparing paclitaxel plus 5-fluorouracil vs cisplatin plus 5-fluorouracil in chemoradiotherapy for locally advanced esophageal carcinoma are underevaluated.

## Hypoxia-specific Cytotoxins

Some bioreductive agents, such as aromatic N-oxides, transition metal complexes, quinones, aliphatic N-oxides and nitro compounds, have radiosensitization effects by virtue of their preferential cytotoxicity toward hypoxic cells.<sup>11</sup> Tirapazamine (TPZ), a hypoxia-selective radiosensitizer, has shown promising results in clinical trials.<sup>97,98</sup> Under hypoxic environments, TPZ can be reduced by reductase in cells to a metabolite that produces free radical and then leads to SSB, DSB, and base damage on DNA.<sup>99</sup> A Phase I clinical trial of TPZ with cisplatin and radiotherapy in small cell lung cancer showed prolonged survival of patients.<sup>100</sup> A Phase II study of TPZ with chemoradiotherapy in locally advanced head and neck cancer reported improvements in failure-free survival and response of patients.<sup>101</sup> However, further phase III trials of TPZ with chemoradiotherapy in locally advanced head and neck cancer concluded that there was no obvious improvement in patient survival.<sup>102</sup> In addition, SN30000 (previously known as CEN-209), an analog of TPZ, with more favorable diffusion property that provides greater toxicity in hypoxic cancer cells than TPZ, is currently

under development by the Drug Development Office of Cancer Research UK.<sup>103</sup>

AQ4N, a representative to aliphatic N-oxide, can be reduced to AQ4 by cytochrome P450 isoenzymes or nitric oxide synthase 2A.<sup>104</sup> In vivo experiments showed that combined utilization of AQ4N with radiotherapy resulted in increased antitumor efficacy, as well as negligible toxicity to normal tissue compared with radiation alone.<sup>105</sup> Positive results were also evidenced in Phase I clinical trials.<sup>106</sup> A Phase I clinical trial in glioblastoma and head and neck tumor patients proved that AQ4N could be specifically activated in hypoxic regions of solid tumors.<sup>107</sup> Unfortunately, a Phase II clinical trial of AQ4N with radiotherapy and temozolomide in glioblastoma began in 2006, was in a pending status (NCT00394628).

TH-302 (evofosofamide), a similar compound that can be reduced to bromo-isophosphoramidate mustard in hypoxic conditions, has radiosensitization activity, especially in hypoxic cells.<sup>108,109</sup> In preclinical models of rhabdomyosarcoma (skeletal muscle) and NSCLC, TH-302 combined with radiotherapy treatment resulted in significant tumor growth delay.<sup>110</sup> In addition, in a study in patient-derived xenograft models of pancreatic cancer, combination treatment of TH-302 and radiotherapy was more efficient than either treatment alone.<sup>111</sup> TH-302 can specifically target the hypoxic tumor cells and induce DNA damage simultaneously in adjacent tumor tissue of the hypoxic zone, and thus holds potential radiosensitization effects in solid tumor treatment.<sup>112</sup> However, on the database of US National Institutes of Health clinical trials, only one of the 26 trials listed proposed combination treatment of TH-302 with radiotherapy (NCT02598687), and it was withdrawn because two phase III trials did not meet their primary endpoint.<sup>113</sup>

Mitomycin C, a quinone-based anticancer therapeutic, can be activated via DNA cross-linking. In preclinical study, mitomycin C showed only slight toxicity in hypoxic cells, which promotes the development of other hypoxia-sensitive quinones selection.<sup>114</sup> Among them, porfirimycin (POR) and apaziquone (EO9) are bioreductive prodrugs, represent the leading candidates.<sup>104</sup> Preclinical studies concluded that POR held higher hypoxic selectivity than mitomycin C.<sup>115</sup> Although preclinical trials proved POR had acceptable toxicity, the following Phase 3 trial demonstrated that POR had a poorer therapeutic effect than mitomycin C.<sup>116</sup> Preclinical studies indicated that EO9 had greater

antitumor property than mitomycin C, indicating EO9 can be a ideal radiosensitizer.<sup>117</sup>

## Other Chemical Radiosensitizers

Other types of chemical radiosensitizers have also seen some progress and some of them are in preclinical evaluations. For example, chemicals that influence cell signaling, suppress radioprotective substances, pseudosubstrates and targeted delivery systems are exploited. With the development of research on radioresistance mechanism, it has been found that multiple signal pathways are related to radioresistance, providing more targets for radiosensitization, such as PI3K–Akt–mTOR,<sup>118</sup> Wnt,<sup>119</sup> MAPK,<sup>120</sup> MDM2<sup>121</sup> and c-MET–PI3K–Akt.<sup>122</sup> For example, BKM120, the oral PI3K inhibitor, can inhibit the activity of PI3K/Akt by targeting the PI3K–Akt pathway, thereby increasing cell apoptosis and inhibiting DNA double-strand break repair in liver cancer cells.<sup>123</sup> BEZ235, a dual PI3K–mTOR inhibitor, can improve the radiosensitivity of colorectal cancer cells.<sup>124</sup> AMG 232, a picomolar affinity piperidinone inhibitor of MDM2, can suppress tumor growth on a mouse model.<sup>121</sup>

Suppression of radioprotective substances, such as glutathione (GSH), is another strategy of radiosensitization. Inhibition of GSH can prevent DNA damage repair and lead to increased damage in tumor cells, which improves the efficacy of radiotherapy in turn.<sup>125</sup> In addition, pseudosubstrates lead cells undergoing DNA synthesis unable to distinguish thymidine and its halogenated analogs efficiently. It is a new area of clinical research to use halogenated pyrimidine analogs, like bromodeoxyuridine (BrdUrd) and iododeoxyuridine (IdUrd), as potential clinical radiosensitizers.<sup>126</sup> One study demonstrated that electron affinities of 5-halogenated deoxyuridine led to enough ability to bind a radiation-produced secondary electron, thereby increasing the sensitivity of radiotherapy.<sup>127</sup>

In addition, research on new indications for existing drugs provides a new paradigm for the development of radiosensitizers. For instance, papaverine, an ergot alkaloid first isolated from *Papaver somniferum* in 1848, has been used for treatment of vasospasm, cerebral thrombosis, pulmonary embolism and erectile dysfunction.<sup>128</sup> Denko et al identified papaverine as an inhibitor of mitochondrial complex I and proved that papaverine could increase oxygenation and enhance radiation response.<sup>128</sup> A phase I trial (NCT03824327) study on papaverine and stereotactic body radiotherapy (SBRT) for NSCLC or lung metastases is under evaluation. In summary, small-

**Table 3** Registered Ongoing Clinical Trials (<https://Clinicaltrials.gov/>) of Small-molecule Chemical Radiosensitizers

Identifier	Drugs	Conditions	Phase	Initiation
NCT02363829	Nelfinavir	Uterine cervix cancer	I	February 2015
NCT02459457	Paclitaxel	Stage III esophageal squamous cell carcinoma	III	July 2015
NCT02598687	TH-302	Esophageal cancer	I	December 2015
NCT02724618	Curcumin	Prostate cancer	II	March 2016
NCT03066154	Docetaxel	Prostatic neoplasms	I	September 2016
NCT02757651	Hydrogen peroxide	Breast cancer	I/II	January 2017
NCT02871843	RRx-001	Glioblastoma	I	February 2017
NCT03101995	Gemcitabine	Cervical cancer	II	July 2017
NCT03824327	Papaverine Hydrochloride	Lung non-small-cell carcinoma	I	February 2019

molecule chemicals as radiosensitizers initiated in the past five years under clinical trials are summarized in Table 3.

## Macromolecules

### Proteins and Peptides

Proteins and peptides, such as antibodies and short peptides, have high affinity with antigens and receptors over-expressed on the surface of tumor cells, making them usable as radiosensitizers.<sup>129</sup> For instance, HER3-ADC, a maytansine-based antibody-drug conjugate targeting HER3, which induces cell arrest in the G<sub>2</sub>/M phase to inhibit DNA damage repair and thereby improves radiosensitivity of HER3-positive pancreatic cancer cells.<sup>130</sup> SYM004, an epidermal growth factor receptor targeting antibody, can inhibit DNA double strand breaks repair and induces apoptosis via downregulating MAPK signaling, and thereby improves radiosensitivity in tumor cells.<sup>120</sup> Cetuximab and nimotuzumab, binding the epidermal growth factor receptor (EGFR), can increase radiation-induced apoptosis and DNA damage, and thereby improve the radiosensitivity of human epidermal-like A431 cells.<sup>131</sup> The hepatocyte growth factor (HGF)/Met signaling pathway which mediates DNA double-strand break repair is upregulated in the majority of cancers. AMG102, a monoclonal antibody against HGF, can inhibit DNA damage repair and increase radiosensitivity of glioblastoma multiforme.<sup>132</sup> In addition, proteins and peptides in serum, such as C-reactive peptide,<sup>133</sup> HSP<sup>134</sup> and paraoxonase-2<sup>135</sup> contribute to radioresistance and can be used as radiotherapy targets. ECI301, a mutant derivative of macrophage inhibitory protein-1a, can be assisted by HSP-70 and HMGB1, thereby enhancing the effect of radiotherapy.<sup>134</sup> Other proteins, like DNAzyme (DZ1)<sup>136</sup>

and NKTR-214,<sup>137</sup> can also improve the effect of radiotherapy.

### miRNAs

MicroRNAs (miRNAs), which encode by endogenous genes are noncoding single-stranded RNA molecules containing about 22 nucleotides. Studies have shown that some specific miRNAs can be used to improve radiotherapy efficacy<sup>138,139</sup> and some miRNAs can be used as radiotherapy sensitization targets.<sup>140</sup> For example, miR-621 targets SETDB1 in hepatocellular carcinoma can be used as a tumor radiosensitizer directly.<sup>141</sup> miR-205 targets zinc finger E-box binding homeobox 1 (*ZEB1*) and the ubiquitin-conjugating enzyme Ubc13 to enhance the radiosensitivity of breast cancer cells.<sup>142</sup> miR-144-5p targets ATF2 to enhance radiosensitivity of NSCLC.<sup>143</sup> miR-146a-5p enhances radiosensitivity in hepatocellular carcinoma through activation of DNA repair pathway.<sup>144</sup> miR-150 modulates AKT pathway in NK/T cell lymphoma to enhance radiosensitivity.<sup>145</sup> miR-99a targets mTOR pathway to enhance the radiosensitivity of NSCLC.<sup>146</sup> miR-139-5p modulates radiotherapy resistance in breast cancer by repressing multiple gene networks of DNA repair and ROS defense.<sup>147</sup> Transcriptional activation of miR-320a induces cancer cell apoptosis under ionizing radiation conditions.<sup>148</sup> However, inhibition of miR-21-5p promotes the radiation sensitivity of NSCLC.<sup>149</sup> Inhibition of miR-630 enhances radiotherapy resistance in human glioma by directly targeting *CDC14A*.<sup>150</sup> Furthermore, a clinical study included 55 atypical meningioma patients found in seven upregulated miRNAs (miR-4286, miR-4695-5p, miR-6732-5p, miR-6855-5p, miR-7977, miR-6765-3p, miR-6787-5p) and seven downregulated miRNAs (miR-1275, miR-30c-1-3p, miR-4449, miR-4539, miR-4684-3p, miR-6129,



miR-6891-5p) in patients. Those miRNAs may induce radioresistant and radiosensitive, respectively.

## siRNAs

siRNA, known as short interfering RNA or silencing RNA, is a class of double-stranded RNA, noncoding RNA molecules, typically 20–27 base pairs in length, similar to miRNA, and operating within the RNA interference (RNAi) pathway.<sup>151</sup> HuR is a protein related to radiotherapy resistance, knockdown of HuR by siRNA resulting DNA damage and enhanced radiosensitivity.<sup>152</sup> *S100A4*, a member of the S100 family of transcription factors, modulates various activities of malignant tumor cells through different mechanisms. A short siRNA against *S100A4* enhances the radiosensitivity of human A549 cells.<sup>153</sup> NBS1 plays an important role in the radiation-induced DNA double-strand breaks repair, siRNA targets NBS1 can increase radiation sensitivity of cancer cells.<sup>154</sup> Survivin, a member of the inhibitor of apoptosis (IAP) protein family, is overexpressed in most cancers resulting in aggressive behavior of tumor and therapy resistance. Downregulation of survivin by siRNA can enhance radiosensitivity in head and neck squamous cell carcinoma.<sup>155</sup> Therefore, numerous siRNAs can be used as radiosensitizers by silencing genes related to radioresistance.

## Oligonucleotides

Similar to siRNAs, oligonucleotides also play important roles in gene expression regulation. Since they are easy to design and synthesize, antisense oligonucleotides have great potential to develop as radiosensitizers.<sup>11</sup> Telomerase expresses in many kinds of tumors (>85%), while the expression of telomerase is restricted in normal tissues. A study indicated that expression of telomerase could be inhibited by radiolabeled oligonucleotides, which targeted the RNA subunit of telomerase, thereby inducing DNA damage in telomerase-positive tumor cells.<sup>156</sup> In addition, the phosphorothioate-modified antisense oligonucleotides (PS-ASODN) against human telomerase reverse transcriptase were reported to promote radiotherapy effect in liver cancer.<sup>157</sup> Furthermore, Park et al reported that inhibition of cyclic AMP response element-directed transcription using decoy oligonucleotides enhanced tumor-specific radiosensitivity.<sup>158</sup> Yu et al demonstrated that antisense oligonucleotides targeted human telomerase RNA (hTR ASODN) could improve the radiosensitivity of nasopharyngeal carcinoma cells.<sup>159</sup>


The radiosensitization mechanism of macromolecules was summarized in Figure 3.

## Nanomaterials

### Noble Metal nanomaterials

The X-ray absorption coefficient ( $\mu$ ) represents the relationship between the X-ray absorption phenomenon ( $E$ ) and atomic number ( $Z$ ),  $\mu = \rho Z^4 / (AE^3)$ , where  $\rho$  is the density and  $A$  is the atomic mass of the element.<sup>160</sup> Therefore, the change of atomic number ( $Z$ ) causes a significant change of X-ray absorption coefficient ( $\mu$ ). Noble metal nanomaterials, such as gold (Au,  $Z=79$ ), silver (Ag,  $Z=47$ ) and platinum (Pt,  $Z=78$ ) can effectively absorb X-ray energy and interact with radiation in tumor cells, and then emit photoelectrons, auger electrons, Compton electrons and other secondary electrons. These secondary electrons not only interact with DNA directly, but also react with water to increase the production of ROS and further increase the sensitivity of tumor cells to radiation. This process is a physical sensitization mechanism.<sup>161</sup> Furthermore, functionalized noble metal nanomaterials promote the generation of ROS, transfer the cell cycle into a radiosensitive state, and inhibit p53 signaling pathway to induce cell autophagy and lysosome body function disorder, thereby increasing radiotherapy sensitivity. This process is a biochemical sensitization mechanism.<sup>162,163</sup>

Gold nanoparticles with good chemical stability, easy preparation, controllable size and shape, easy surface functionalization, high biocompatibility, and low toxicity have proven satisfactory radiosensitizing effects in various tumors.<sup>164–167</sup> Silver nanoparticles and platinum nanoparticles are also commonly used in biomedicine.<sup>168,169</sup> Research found that silver nanoparticles combined with radiotherapy could enhance the radiosensitivity of human glioma cells in vitro and extended the survival time of glioma mice.<sup>170,171</sup> Liu et al demonstrated that silver nanoparticles could induce apoptosis of cancer cells through G<sub>2</sub>/M phase arrest after radiation, and they suggested that silver nanoparticles could be used as a nanoradiosensitizer for hypoxic glioma radiotherapy.<sup>172</sup> Recently, Fathy reported that thymoquinone-capping silver nanoparticles represented a promising engineered nanoformulation for enhancing cancer radiosensitivity.<sup>173</sup> Li et al demonstrated that platinum nanoparticles could enhance radiosensitivity through increasing DNA damage, ROS stress, and cell cycle arrest.<sup>163</sup> They also proved that platinum nanoparticles could convert endogenous H<sub>2</sub>O<sub>2</sub> to O<sub>2</sub> in cancer cells,



**Figure 3** Radiosensitization mechanism of macromolecules. (A) Proteins and peptides. (a1) Direct interaction of key proteins. (a2) Loading of radioactive seeds. (a3) Radiosensitizers delivery. (a4) Conjugation with nanomaterials. (B) miRNAs can then bind with mRNAs to implement radiosensitization. (b1) Downregulation by inhibitors. (b2) Upregulation. (C) siRNAs can improve radiosensitivity by binding and degrading complementary mRNAs. (D) Oligonucleotides improve the radiosensitivity by complementary binding with DNAs.

thus significantly improving radiosensitivity without apparent toxicity to animals *in vivo*.<sup>163</sup>

## Heavy Metal Nanomaterials

Similar to noble metal nanomaterials, gadolinium (Gd,  $Z=64$ ), hafnium (Hf,  $Z=72$ ), tantalum (Ta,  $Z=73$ ), tungsten (W,  $Z=74$ ), and bismuth (Bi,  $Z=83$ ) are also metal elements with large atomic coefficients and have a great X-ray attenuation capability.<sup>174–176</sup> Based on this, numerous studies have focused on these heavy metal nanomaterials to investigate their radiotherapy sensitization. However, they usually cause damage to healthy tissues with direct contact.<sup>177</sup> Therefore, their stable forms such as oxides, sulfides, and selenides are explored as the radiosensitizers.<sup>178–180</sup>

Gadolinium-based nanoparticles are usually known as magnetic resonance imaging (MRI) contrast agents. It

should be noted that researchers discovered a family of gadolinium-based nanoparticles called AGuIX for combined MRI and radiosensitization.<sup>181</sup> Results showed that AGuIX could interact with X-rays and  $\gamma$ -rays at a certain concentration. After internalization through the enhanced permeability and retention (EPR) effect, AGuIX could be resident in the tumor for a long time before being cleared by the kidneys.<sup>182</sup> Preclinical animal experiments proved that AGuIX held obvious radiosensitization effects in several tumor models without obvious toxicity.<sup>183</sup> A Phase I clinical trial (NCT03308604) to evaluate the optimal dose of AGuIX combined with chemoradiation in patients with locally advanced cervical cancer; a Phase II clinical trial (NCT03818386) using AGuIX gadolinium-chelated polysiloxane based nanoparticles and whole brain radiotherapy in patients with multiple brain metastases; and a single-arm phase II trial (NCT04094077)

aiming to evaluate the efficacy of AGuIX during fractionated stereotactic radiotherapy of brain metastasis are being evaluated.

Hafnium, in the same family as titanium and zirconium, is chemical inertness. The oxidation state of hafnium, hafnium dioxide ( $\text{HfO}_2$ ), was usually used in radioactive protective coatings, biosensors, and X-ray contrast agents.<sup>184,185</sup> Jayaraman et al demonstrated that  $\text{HfO}_2$  nanoparticles had excellent biocompatibility.<sup>185</sup> Researchers from France discovered that  $\text{HfO}_2$  can be used as a radiosensitizer with low cytotoxicity.<sup>186</sup> A Phase I trial (NCT03589339) combining hafnium oxide nanoparticles (NBTXR3) with anti-PD-1 therapy in microsatellite instability-high solid malignant tumour and a Phase I–II clinical trial (NCT02805894) of NBTXR3 in prostate adenocarcinoma are under evaluation.

Tantalum is a nontoxic, biologically inert element with good biocompatibility.<sup>187</sup> Studies found that TaOx and  $\text{Ta}_2\text{O}_5$  could be used as CT imaging contrast agents.<sup>188–190</sup> Brown et al found  $\text{Ta}_2\text{O}_5$  nanoparticles showed a radiosensitization effect on radioresistant glioma cells.<sup>191</sup> Song et al showed hollow shell tantalum oxide (HTaOx) had a large X-ray attenuation capability and could enhance radiation therapy effects by Compton scattering and Auger effect.<sup>192</sup> In addition, TaOx can be used as functional group carrier to load drugs, thereby improving tumor hypoxic environment. For example, HTaOx loaded with catalase, which reacted with  $\text{H}_2\text{O}_2$  in the tumor microenvironment, then improved the oxygen content and overcame the radiotherapy tolerance of hypoxic tumor cells, thereby improving the radiotherapy effect.<sup>193</sup>





Tungsten and bismuth also have significant applications in medicine.<sup>194,195</sup> Hossain et al concluded that bismuth nanoparticles had stronger radiosensitizing effect than gold and platinum nanoparticles at the same physical and chemical conditions.<sup>196</sup> Yu et al found that the ultra-small semi-metallic Bi nanoparticles with LyP-1 peptide modified at 3.6 nm showed obvious radiosensitization effect.<sup>197</sup> Recently, a large number of studies shown that some nanomaterials of tungsten and bismuth had excellent photothermal absorption conversion performance and strong X-ray absorption capacity, therefore they can be used for tumor radiosensitization as well as synergistic therapy of hyperthermia and radiotherapy.<sup>198–201</sup>

In addition, research about several high Z metal elements combined together to further improve the radiosensitization effect were also explored. For example, SiBiGdNP chelated Bi and Gd in organosilane to improve

the sensitivity of radiotherapy.<sup>202</sup>  $\text{GdW}_{10}\text{O}_{36}$  contained both W and Gd to expect they had better radiotherapy sensitization effect.<sup>203</sup>

## Ferrite Nanomaterials

Ferrite-based nanomaterials can catalyze the generation of free radicals through Fenton's reaction (1) and Haber–Weiss reaction (2) to enhance the effect of radiosensitization.<sup>204</sup>



Studies proved that  $\text{Fe}_3\text{O}_4$  had a dose-enhancing effect for radiotherapy, especially superparamagnetic  $\text{Fe}_3\text{O}_4$  nanoparticles (SPIONS) possessing MRI imaging property had good application prospects in image-guided tumor radiotherapy.<sup>205</sup>

The composition of the spinel structure ferrite is usually stated as  $\text{MFe}_2\text{O}_4$ , where  $\text{M}=\text{Fe}$ , Zn, Co, Mn, Ni.<sup>206</sup> Among them,  $\text{ZnFe}_2\text{O}_4$ ,  $\text{MnFe}_2\text{O}_4$ ,  $\text{CoFe}_2\text{O}_4$  nanoparticles were widely investigated.<sup>207</sup> For example, Meidanchi et al confirmed that  $\text{ZnFe}_2\text{O}_4$  nanoparticles interacted with  $\gamma$ -rays to produce photoelectric effect resulting in a higher release level of electron in radioresistant cells.<sup>208</sup> Studies also indicated that  $\text{ZnFe}_2\text{O}_4$  nanoparticles could be used as radiosensitizers.<sup>208,209</sup> Salunkhe et al demonstrated that  $\text{MnFe}_2\text{O}_4$  and  $\text{CoFe}_2\text{O}_4$  nanoparticles could improve the therapeutic efficacy of cancer through multimodal image-guided combination therapy.<sup>210</sup>

## Semiconductor Nanomaterials

Semiconductor quantum dots have unique properties, such as quantum dimension effect, surface effect, and quantum confinement effect, making them great candidates in biomedicine applications.<sup>211</sup> Until now, numerous studies focused on using semiconductor quantum dots as photosensitizers and radiosensitizers for tumor treatment have been reported.<sup>212–214</sup> When the electronic energy levels are in the range of 1–5 eV, the semiconductor nanomaterials can absorb the photon energy and perform as photosensitizers, showing photocatalytic properties. When the electronic energy levels are at keV and MeV (X-rays and  $\gamma$ -rays), semiconductor nanomaterials can enhance absorption of high-energy photons acting as radiosensitizers and causing damage to cancer cells.<sup>212</sup> Nakayama et al synthesized a semiconductor nanomaterial PAA-TiOx to generate

hydroxyl radicals under the irradiation of X-rays, which increased DNA damage and inhibited tumor growth significantly.<sup>215</sup> Morita et al clarified the radiosensitization mechanism of PAA-TiO<sub>2</sub> nanoparticles by releasing H<sub>2</sub>O<sub>2</sub> to relieve hypoxia in tumor cells.<sup>216</sup> TiO<sub>2</sub> nanotubes have been reported to enhance the radiosensitization effect through regulating G<sub>2</sub>/M cycle arrest and reducing DNA repair of tumor cells.<sup>177</sup> The mechanism of radiosensitization of metal-based nanomaterials is shown in Figure 4.


## Nonmetallic Nanomaterials

Many nonmetallic nanomaterials also possess the function of radiosensitization.<sup>217</sup> For example, C<sub>60</sub>, fullerene, has potent anticancer activities, however, the potential toxicity to normal tissues limits its further use. Therefore, nanocrystals of C<sub>60</sub> (Nano-C<sub>60</sub>) with negligible toxicity to normal cells have been developed as a radiosensitizer.<sup>218</sup> In addition, nanodiamonds and carbon nanotubes can reduce radioresistance of tumor cells by promoting ROS generation, destroying DNA double-strands, and regulating the cell cycle.<sup>219,220</sup> Selenium (Se) nanoparticles not only work as chemotherapeutic drugs, but also improve the antitumor effect of X-rays by activating ROS to induce DNA damage in cancer cells.<sup>221</sup>

## Nanostructured Chemicals and Drug Delivery Systems

Nano-based delivery systems are efficient approaches for drug targeted transportation, which can deliver radiosensitizers, such as chemicals, oxygen carriers, siRNAs and catalases to the tumor sites and have attracted wide interest of researchers recently.<sup>222</sup> More importantly, nanobased delivery systems can precisely deliver radioactive particles like<sup>223</sup> Ac (releasing  $\alpha$ -particles), <sup>131</sup>I, and <sup>125</sup>I to tumor sites.<sup>223</sup> With the development of nanotechnology, nanobased delivery systems have great potential for radiosensitizer delivery.

However, there is still a challenge to achieve clinical translation of nanobased delivery systems, factors like physicochemical properties of the nanoformulations, radiation sources, and indications block their clinical translation.<sup>223</sup> In addition, long circulation lifetime of nanodelivery systems may increase the risk of long-term toxicity.<sup>224</sup> Another critical factor is stability in body fluid of nanodelivery systems. Because the aggregation of nanoparticles in body fluid will influence the pharmacokinetics and the cellular response and generate serious side effects such as blocking the blood vessels.<sup>222</sup> Therefore, attention should be paid to these factors when designing the nanodelivery systems. Size is also an



**Figure 4** Radiosensitization mechanism of metal-based nanomaterials. The process contains physical and biochemical sensitization mechanism.



important factor, small size and high Z nanoparticles often hold better radiosensitizing effect than larger-size ones.<sup>223</sup> In particular, the small size nanoparticles with positive charge can bind to negative charged DNA and can be eliminated by renal clearance conveniently. In addition, functional modification of nanostructures using biocompatible materials can improve their stability and targeting.<sup>225</sup>

## Conclusions and Prospects

Radiosensitizers have been developed for decades from the earliest “free radical damage and fixation” strategies to gene regulation, from chemicals to biological macromolecules and nanomaterials. Although each radiosensitizer has dialectical advantages and limitations, the mechanisms of sensitization are similar. The main mechanisms include: (I) inhibiting radiation-induced repair of DNA damage, increasing the degree of DNA damage; (II) disturbing the cell cycle and organelle function to improve cytotoxicity; and (III) inhibiting the expression of radiation resistance genes or promoting the expression of radiation sensitive genes.

Although small molecules, macromolecules, and nanomaterial radiosensitizers are being developed, and some nanoradiosensitizers have been used for clinical research (Table 4), the result still cannot meet clinical translation needs. Therefore, there is an urgent need to find new targets of radiotherapy and new mechanisms of sensitization, and after that to develop more effective radiosensitizing drugs. First of all, multitarget radiosensitizers often have more obvious efficacy than single target, researchers can focus on screening multitarget radiosensitizers or drug combinations. New approaches, in particular, nanotechnology based as radiosensitizers have shown promise. Nanomaterials with low cytotoxicity, good biocompatibility, and ease of functionalization need to be explored. In addition, other technologies, such as molecular structure analysis, molecular cloning technology, and bioinformatics analysis can accelerate the development of new radiosensitizers. Moreover, development of new drug delivery systems can

also improve radiosensitization efficacy. Finally, the application of artificial intelligence and machine learning in new drug discovery and clinical trials, may guide development of new radiosensitizers and optimization of existing radiosensitizers.

## Acknowledgment

This work was supported by Innovation Capacity Support Plan of Shaanxi Province (2018TD-002), the National Natural Science Foundation of China (No. 82000523), the Natural Science Foundation of Shaanxi province (Grant No. 2020JQ-087, 2020JQ-095), the “Young Talent Support Plan” of Xi’an Jiaotong University (YX6J001), the Fundamental Research Funds for the Central Universities (xzy012019070).

## Disclosure

The authors report no conflicts of interest in this work.

## References

- Krzyszczuk P, Acevedo A, Davidoff EJ, et al. The growing role of precision and personalized medicine for cancer treatment. *Technology (Singap World Sci)*. 2018;6(3-4):79-100. doi:10.1142/S2339547818300020
- Kulakowski A. The contribution of Marie Skłodowska-Curie to the development of modern oncology. *Anal Bioanal Chem*. 2011;400(6):1583-1586. doi:10.1007/s00216-011-4712-1
- Martin OA, Martin RF. Cancer radiotherapy: understanding the price of tumor eradication. *Front Cell Dev Biol*. 2020;8:261. doi:10.3389/fcell.2020.00261
- Franzone P, Fiorentino A, Barra S, et al. Image-guided radiation therapy (IGRT): practical recommendations of Italian Association of Radiation Oncology (AIRO). *Radiol Med*. 2016;121(12):958-965. doi:10.1007/s11547-016-0674-x
- Ge Y, Wu QJ. Knowledge-based planning for intensity-modulated radiation therapy: a review of data-driven approaches. *Med Phys*. 2019;46(6):2760-2775. doi:10.1002/mp.13526
- Farhood B, Goradel NH, Mortezaee K, et al. Melatonin as an adjuvant in radiotherapy for radioprotection and radiosensitization. *Clin Transl Oncol*. 2019;21(3):268-279. doi:10.1007/s12094-018-1934-0
- Fowler JF, Adams GE, Denekamp J. Radiosensitizers of hypoxic cells in solid tumors. *Cancer Treat Rev*. 1976;3(4):227-256. doi:10.1016/s0305-7372(76)80012-6
- Adams GE. Chemical radiosensitization of hypoxic cells. *Br Med Bull*. 1973;29(1):48-53. doi:10.1093/oxfordjournals.bmb.a070956
- Wen P, Xia J, Cao X, et al. dbCRSR: a manually curated database for regulation of cancer radiosensitivity. *Database (Oxford)*. 2018;2018. doi:10.1093/database/bay049
- Garibaldi C, Jereczek-Fossa BA, Marvaso G, et al. Recent advances in radiation oncology. *Ecancermedicalscience*. 2017;11:785. doi:10.3332/ecancer.2017.785
- Wang H, Mu X, He H, Zhang XD. Cancer radiosensitizers. *Trends Pharmacol Sci*. 2018;39(1):24-48. doi:10.1016/j.tips.2017.11.003
- Hirayama R. Mechanism of oxygen effect for photon and heavy-ion beams. *Japanese Journal of Medical Physics*. 2014;34(2):65-69.
- Oronsky BT, Knox SJ, Scicinski J. Six degrees of separation: the oxygen effect in the development of radiosensitizers. *Transl Oncol*. 2011;4(4):189-198. doi:10.1593/tlo.11166

**Table 4** Clinical Translation of Some Nanoradiosensitizers

Name	Conditions	Phase	Identifier
AGuIX	Cervical cancer	I	NCT03308604
	Brain metastases	II	NCT03818386
	Brain metastases	II	NCT04094077
NBTXR3	Microsatellite instability-high solid malignant tumor	I	NCT03589339
	Prostate adenocarcinoma	I/II	NCT02805894



14. Harrison LB, Chadha M, Hill RJ, Hu K, Shasha D. Impact of tumor hypoxia and anemia on radiation therapy outcomes. *Oncologist*. 2002;7(6):492–508. doi:10.1634/theoncologist.7-6-492
15. Zheng L, Kelly CJ, Colgan SP. Physiologic hypoxia and oxygen homeostasis in the healthy intestine. A review in the theme: cellular responses to hypoxia. *Am J Physiol Cell Physiol*. 2015;309(6):C350–C360. doi:10.1152/ajpcell.00191.2015
16. Richardson RB, Harper ME. Mitochondrial stress controls the radiosensitivity of the oxygen effect: implications for radiotherapy. *Oncotarget*. 2016;7(16):21469–21483. doi:10.18632/oncotarget.7412
17. Bel Aiba RS, Dimova EY, Görlach A, Kietzmann T. The role of hypoxia inducible factor-1 in cell metabolism—a possible target in cancer therapy. *Expert Opin Ther Targets*. 2006;10(4):583–599. doi:10.1517/14728222.10.4.583
18. Zhu H, Zhang S. Hypoxia inducible factor-1 $\alpha$ /vascular endothelial growth factor signaling activation correlates with response to radiotherapy and its inhibition reduces hypoxia-induced angiogenesis in lung cancer. *J Cell Biochem*. 2018;119(9):7707–7718. doi:10.1002/jcb.27120
19. Verdegem D, Moens S, Stapor P, Carmeliet P. Endothelial cell metabolism: parallels and divergences with cancer cell metabolism. *Cancer Metabol*. 2014;2(1):19. doi:10.1186/2049-3002-2-19
20. Boyle RG, Travers S. Hypoxia: targeting the tumour. *Anticancer Agents Med Chem*. 2006;6(4):281–286. doi:10.2174/187152006777698169
21. Maxwell PH. The HIF pathway in cancer. *Semin Cell Dev Biol*. 2005;16(4–5):523–530. doi:10.1016/j.semcdb.2005.03.001
22. Al Tameemi W, Dale TP, Al-Jumaily RMK, Forsyth NR. Hypoxia-modified cancer cell metabolism. *Front Cell Dev Biol*. 2019;7:4. doi:10.3389/fcell.2019.00004
23. Pettersen EO, Ebbesen P, Gieling RG, et al. Targeting tumour hypoxia to prevent cancer metastasis. From biology, biosensing and technology to drug development: the METOXIA consortium. *J Enzyme Inhib Med Chem*. 2015;30(5):689–721. doi:10.3109/14756366.2014.966704
24. Cabrales P, Intaglietta M. Blood substitutes: evolution from non-carrying to oxygen- and gas-carrying fluids. *ASAIO J*. 2013;59(4):337–354. doi:10.1097/MAT.0b013e318291fbaa
25. Hardavella G, Karampinis I, Frille A, Sreter K, Rousalova I. Oxygen devices and delivery systems. *Breathe (Sheffield, England)*. 2019;15(3):e108–e116. doi:10.1183/20734735.0204-2019
26. Choudhury R. Hypoxia and hyperbaric oxygen therapy: a review. *Int J Gen Med*. 2018;11:431–442. doi:10.2147/ijgm.S172460
27. Stepien K, Ostrowski RP, Matyja E. Hyperbaric oxygen as an adjunctive therapy in treatment of malignancies, including brain tumours. *Med Oncol (Northwood, London, England)*. 2016;33(9):101. doi:10.1007/s12032-016-0814-0
28. Ogawa Y, Kubota K, Ue H, et al. Phase I study of a new radiosensitizer containing hydrogen peroxide and sodium hyaluronate for topical tumor injection: a new enzyme-targeting radiosensitization treatment, Kochi oxydol-radiation therapy for unresectable carcinomas, type II (KORTUC II). *Int J Oncol*. 2009;34:609–618. doi:10.3892/ijo\_00000186
29. Chapman JD, Webb RG, Borsa J. Radiosensitization of mammalian cells by p-nitroacetophenone. *Int J Radiat Biol Relat Stud Phys Chem Med*. 1971;19(6):561–573. doi:10.1080/09553007114550741
30. Adams GE, Asquith JC, Dewey DL, Foster JL, Michael BD, Willson RL. Electron affinic sensitization. *Int J Radiat Biol Relat Stud Phys Chem Med*. 1971;19(6):575–585. doi:10.1080/09553007114550751
31. Higgins GS, O’Cathail SM, Muschel RJ, McKenna WG. Drug radiotherapy combinations: review of previous failures and reasons for future optimism. *Cancer Treat Rev*. 2015;41(2):105–113. doi:10.1016/j.ctrv.2014.12.012
32. Spisz P, Zdzrowowicz M, Makurat S, et al. Why does the type of halogen atom matter for the radiosensitizing properties of 5-halogen substituted 4-thio-2’-deoxyuridines? *Molecules*. 2019;24(15):2819. doi:10.3390/molecules24152819
33. Brown JM. Selective radiosensitization of the hypoxic cells of mouse tumors with the nitroimidazoles metronidazole and Ro 7-0582. *Radiat Res*. 1975;64(3):633–647. doi:10.2307/3574253
34. Dische S, Saunders MI, Lee ME, Adams GE, Flockhart IR. Clinical testing of the radiosensitizer Ro 07-0582: experience with multiple doses. *Br J Cancer*. 1977;35(5):567–579. doi:10.1038/bjc.1977.90
35. Brown JM. Clinical trials of radiosensitizers: what should we expect? *Int J Radiat Oncol Biol Phys*. 1984;10(3):425–429. doi:10.1016/0360-3016(84)90063-4
36. Dische S, Saunders MI, Flockhart IR, Lee ME, Anderson P. Misonidazole—a drug for trial in radiotherapy and oncology. *Int J Radiat Oncol Biol Phys*. 1979;5(6):851–860. doi:10.1016/0360-3016(79)90070-1
37. Barker HE, Paget JTE, Khan AA, Harrington KJ. The tumour micro-environment after radiotherapy: mechanisms of resistance and recurrence. *Nat Rev Cancer*. 2015;15(7):409–425. doi:10.1038/nrc3958
38. Overgaard J, Hansen HS, Andersen AP, et al. Misonidazole combined with split-course radiotherapy in the treatment of invasive carcinoma of larynx and pharynx: report from the DAHANCA 2 study. *Int J Radiat Oncol Biol Phys*. 1989;16(4):1065–1068. doi:10.1016/0360-3016(89)90917-6
39. Urtaşun RC, Chapman JD, Feldstein ML, et al. Peripheral neuropathy related to misonidazole: incidence and pathology. *Br J Cancer Suppl*. 1978;3:271–275.
40. Wardman P. Nitroimidazoles as hypoxic cell radiosensitizers and hypoxia probes: misonidazole, myths and mistakes. *Br J Radiol*. 2019;92(1093):20170915. doi:10.1259/bjr.20170915
41. Bonnet M, Hong CR, Wong WW, et al. Next-generation hypoxic cell radiosensitizers: nitroimidazole alkylsulfonamides. *J Med Chem*. 2018;61(3):1241–1254. doi:10.1021/acs.jmedchem.7b01678
42. Rosenberg A, Knox S. Radiation sensitization with redox modulators: a promising approach. *Int J Radiat Oncol Biol Phys*. 2006;64(2):343–354. doi:10.1016/j.ijrobp.2005.10.013
43. Coleman CN, Wasserman TH, Urtaşun RC, et al. Final report of the phase I trial of the hypoxic cell radiosensitizer SR 2508 (etanidazole) radiation therapy oncology group 83-03. *Int J Radiat Oncol Biol Phys*. 1990;18(2):389–393. doi:10.1016/0360-3016(90)90105-s
44. Brown JM, Wilson WR. Exploiting tumour hypoxia in cancer treatment. *Nat Rev Cancer*. 2004;4(6):437–447. doi:10.1038/nrc1367
45. Overgaard J, Hansen HS, Overgaard M, et al. A randomized double-blind phase III study of nimorazole as a hypoxic radiosensitizer of primary radiotherapy in supraglottic larynx and pharynx carcinoma. *Radiother Oncol*. 1998;46(2):135–146. doi:10.1016/s0167-8140(97)00220-x
46. Overgaard J, Eriksen JG, Nordmark M, Alsner J, Horsman MR. Plasma osteopontin, hypoxia, and response to the hypoxia sensitizer nimorazole in radiotherapy of head and neck cancer: results from the DAHANCA 5 randomised double-blind placebo-controlled trial. *Lancet Oncol*. 2005;6(10):757–764. doi:10.1016/s1470-2045(05)70292-8
47. Metwally MA, Frederiksen KD, Overgaard J. Compliance and toxicity of the hypoxic radiosensitizer nimorazole in the treatment of patients with head and neck squamous cell carcinoma (HNSCC). *Acta Oncologica (Stockholm, Sweden)*. 2014;53(5):654–661. doi:10.3109/0284186x.2013.864050

48. Bentzen J, Toustrup K, Eriksen JG, Primdahl H, Andersen LJ, Overgaard J. Locally advanced head and neck cancer treated with accelerated radiotherapy, the hypoxic modifier nimorazole and weekly cisplatin. Results from the DAHANCA 18 phase II study. *Acta Oncologica (Stockholm, Sweden)*. 2015;54(7):1001–1007. doi:10.3109/0284186x.2014.992547
49. Saksø M, Andersen E, Bentzen J, et al. A prospective, multicenter DAHANCA study of hyperfractionated, accelerated radiotherapy for head and neck squamous cell carcinoma. *Acta Oncologica (Stockholm, Sweden)*. 2019;58(10):1495–1501. doi:10.1080/0284186x.2019.1658897
50. Thomson D, Yang H, Baines H, et al. NIMRAD - a phase III trial to investigate the use of nimorazole hypoxia modification with intensity-modulated radiotherapy in head and neck cancer. *Clin Oncol (Royal College of Radiologists (Great Britain))*. 2014;26(6):344–347. doi:10.1016/j.clon.2014.03.003
51. Toustrup K, Sørensen BS, Lassen P, Wiuf C, Alsner J, Overgaard J. Gene expression classifier predicts for hypoxic modification of radiotherapy with nimorazole in squamous cell carcinomas of the head and neck. *Radiother Oncol*. 2012;102(1):122–129. doi:10.1016/j.radonc.2011.09.010
52. Saksø M, Jensen K, Andersen M, Hansen CR, Eriksen JG, Overgaard J. DAHANCA 28: a phase I/II feasibility study of hyperfractionated, accelerated radiotherapy with concomitant cisplatin and nimorazole (HART-CN) for patients with locally advanced, HPV/p16-negative squamous cell carcinoma of the oropharynx, hypopharynx, larynx and oral cavity. *Radiother Oncol*. 2020;148:65–72. doi:10.1016/j.radonc.2020.03.025
53. Oronsky B, Scicinski J, Ning S, et al. RRx-001, a novel dinitroazetidine radiosensitizer. *Invest New Drugs*. 2016;34(3):371–377. doi:10.1007/s10637-016-0326-y
54. Oronsky BT, Knox SJ, Scicinski JJ. Is Nitric oxide (NO) the last word in radiosensitization? A review. *Transl Oncol*. 2012;5(2):66–71. doi:10.1593/tlo.11307
55. Edfeldt NB, Harwood EA, Sigurdsson ST, Hopkins PB, Reid BR. Solution structure of a nitrous acid induced DNA interstrand cross-link. *Nucleic Acids Res*. 2004;32(9):2785–2794. doi:10.1093/nar/gkh606
56. Bonavida B, Khineche S, Huerta-Yepez S, Garbán H. Therapeutic potential of nitric oxide in cancer. *Drug Resist Updat*. 2006;9(3):157–173. doi:10.1016/j.drug.2006.05.003
57. Nelson EJ, Connolly J, McArthur P. Nitric oxide and s-nitrosylation: excitotoxic and cell signaling mechanism. *Biol Cell*. 2003;95(1):3–8. doi:10.1016/s0248-4900(03)00004-2
58. Meffert MK, Premack BA, Schulman H. Nitric oxide stimulates Ca(2+)-independent synaptic vesicle release. *Neuron*. 1994;12(6):1235–1244. doi:10.1016/0896-6273(94)90440-5
59. Tsai EJ, Kass DA. Cyclic GMP signaling in cardiovascular pathophysiology and therapeutics. *Pharmacol Ther*. 2009;122(3):216–238. doi:10.1016/j.pharmthera.2009.02.009
60. Lundberg JO, Gladwin MT, Weitzberg E. Strategies to increase nitric oxide signalling in cardiovascular disease. *Nat Rev Drug Discov*. 2015;14(9):623–641. doi:10.1038/nrd4623
61. Rogers N, Seeger F, Garcin E, Roberts D, Isenberg J. Regulation of soluble guanylate cyclase by matricellular thrombospondins: implications for blood flow. *Review*. 2014;5(134). doi:10.3389/fphys.2014.00134
62. Kondakova IV, Tcheredova VV, Zagrebelnaya GV, Cherdyntseva NV, Kagiya TV, Choinzonov EL. Production of nitric oxide by hypoxic radiosensitizer sanazole. *Exp Oncol*. 2004;26(4):329–333.
63. Girard M, Clairmont F, Maneckjee A, Mousseau N, Dawson B, Whitehouse L. 5-Nitroimidazoles. II: unexpected reactivity of ronidazole and dimetridazole with thiols. *Can J Chem*. 2011;71:1349–1352. doi:10.1139/v93-174
64. Ng QS, Goh V, Milner J, et al. Effect of nitric-oxide synthesis on tumour blood volume and vascular activity: a phase I study. *Lancet Oncol*. 2007;8(2):111–118. doi:10.1016/s1470-2045(07)70001-3
65. Siemens DR, Heaton JP, Adams MA, Kawakami J, Graham CH. Phase II study of nitric oxide donor for men with increasing prostate-specific antigen level after surgery or radiotherapy for prostate cancer. *Urology*. 2009;74(4):878–883. doi:10.1016/j.urolgy.2009.03.004
66. Libert N, Tourtier JP, Védrine L, Chargari C, Riou B. Inhibitors of angiogenesis: new hopes for oncologists, new challenges for anesthesiologists. *Anesthesiology*. 2010;113(3):704–712. doi:10.1097/ALN.0b013e3181ed098d
67. Pandey AK, Singhi EK, Arroyo JP, et al. Mechanisms of VEGF (Vascular Endothelial Growth Factor) inhibitor-associated hypertension and vascular disease. *Hypertension (Dallas, Tex: 1979)*. 2018;71:e1–e8. doi:10.1161/hypertensionaha.117.10271
68. Liebmann J, DeLuca AM, Coffin D, et al. In vivo radiation protection by nitric oxide modulation. *Cancer Res*. 1994;54(13):3365–3368.
69. Schwarz K, Dobiasch S, Nguyen L, Schilling D, Combs SE. Modification of radiosensitivity by curcumin in human pancreatic cancer cell lines. *Sci Rep*. 2020;10(1):3815. doi:10.1038/s41598-020-60765-1
70. Verma V. Relationship and interactions of curcumin with radiation therapy. *World J Clin Oncol*. 2016;7(3):275–283. doi:10.5306/wjco.v7.i3.275
71. Li G, Wang Z, Chong T, Yang J, Li H, Chen H. Curcumin enhances the radiosensitivity of renal cancer cells by suppressing NF-κB signaling pathway. *Biomed Pharmacother*. 2017;94:974–981. doi:10.1016/j.biopha.2017.07.148
72. da Costa Araldi IC, Bordin FPR, Cadoná FC, et al. The in vitro radiosensitizer potential of resveratrol on MCF-7 breast cancer cells. *Chem Biol Interact*. 2018;282:85–92. doi:10.1016/j.cbi.2018.01.013
73. Tan Y, Wei X, Zhang W, et al. Resveratrol enhances the radiosensitivity of nasopharyngeal carcinoma cells by downregulating E2F1. *Oncol Rep*. 2017;37(3):1833–1841. doi:10.3892/or.2017.5413
74. Mikami S, Ota I, Masui T, et al. Resveratrol-induced REG III expression enhances chemo- and radiosensitivity in head and neck cancer in xenograft mice. *Oncol Rep*. 2019;42(1):436–442. doi:10.3892/or.2019.7137
75. Kim SJ, Kim MS, Lee JW, et al. Dihydroartemisinin enhances radiosensitivity of human glioma cells in vitro. *J Cancer Res Clin Oncol*. 2006;132(2):129–135. doi:10.1007/s00432-005-0052-x
76. Li Y, Sui H, Jiang C, et al. Dihydroartemisinin increases the sensitivity of photodynamic therapy via NF-κB/HIF-1α/VEGF pathway in esophageal cancer cell in vitro and in vivo. *Cell Physiol Biochem*. 2018;48:2035–2045. doi:10.1159/000492541
77. Zhang L, Cheng LQ, Zhou Z, Lv LL, Liu GX. Effects of dihydroartemisinin on radiosensitivity of Raji cells. *Zhongguo Ying Yong Sheng Li Xue Za Zhi*. 2017;33:385–390. doi:10.12047/j.cjap.5565.2017.093
78. Momtazi-Borojeni AA, Mosafer J, Nikfar B, et al. Curcumin in advancing treatment for gynecological cancers with developed drug- and radiotherapy-associated resistance. *Rev Physiol Biochem Pharmacol*. 2019;176:107–129. doi:10.1007/112\_2018\_11
79. Chen Y, Zhu Z, Zhao W, et al. A randomized phase 3 trial comparing paclitaxel plus 5-fluorouracil versus cisplatin plus 5-fluorouracil in Chemoradiotherapy for locally advanced esophageal carcinoma-the ESO-shanghai 1 trial protocol. *Radiat Oncol (London, England)*. 2018;13(1):33. doi:10.1186/s13014-018-0979-0
80. Ren W, Sha H, Yan J, et al. Enhancement of radiotherapeutic efficacy for esophageal cancer by paclitaxel-loaded red blood cell membrane nanoparticles modified by the recombinant protein anti-EGFR-iRGD. *J Biomater Appl*. 2018;33(5):707–724. doi:10.1177/0885328218809019

81. Aggarwal BB. Nuclear factor-kappaB: the enemy within. *Cancer Cell*. 2004;6(3):203–208. doi:10.1016/j.ccr.2004.09.003
82. Chendil D, Ranga RS, Meigooni D, Sathishkumar S, Ahmed MM. Curcumin confers radiosensitizing effect in prostate cancer cell line PC-3. *Oncogene*. 2004;23(8):1599–1607. doi:10.1038/sj.onc.1207284
83. Veeraraghavan J, Natarajan M, Herman TS, Aravindan N. Curcumin-altered p53-response genes regulate radiosensitivity in p53-mutant ewing's sarcoma cells. *Anticancer Res*. 2010;30(10):4007–4015.
84. Liao HF, Kuo CD, Yang YC, et al. Resveratrol enhances radiosensitivity of human non-small cell lung cancer NCI-H838 cells accompanied by inhibition of nuclear factor-kappa B activation. *J Radiat Res*. 2005;46(4):387–393. doi:10.1269/jrr.46.387
85. Aung TN, Qu Z, Kortschak RD, Adelson DL. Understanding the effectiveness of natural compound mixtures in cancer through Their molecular mode of action. *Int J Mol Sci*. 2017;18:656. doi:10.3390/ijms18030656
86. Tak JK, Lee JH, Park JW. Resveratrol and piperine enhance radiosensitivity of tumor cells. *BMB Rep*. 2012;45(4):242–246. doi:10.5483/bmbrep.2012.45.4.242
87. Quan F, Zhao Q, Shao Y, Li H, Zhao R. Resveratrol enhances radiosensitivity of human hypopharyngeal carcinoma cell line in nude mice. *Journal of Southern Medical University*. 2014;34(11):1646–1649. Chinese.
88. Shao Y, Quan F, Li HH, Yao XB, Zhao Q, Zhao RM. The radiosensitizing effect of resveratrol on hypopharyngeal carcinoma cell line FADU and its effect on the cell cycle. *Zhongguo Zhong Xi Yi Jie He Za Zhi*. 2015;35:699–703.
89. Fang Y, DeMarco VG, Nicholl MB. Resveratrol enhances radiation sensitivity in prostate cancer by inhibiting cell proliferation and promoting cell senescence and apoptosis. *Cancer Sci*. 2012;103(6):1090–1098. doi:10.1111/j.1349-7006.2012.02272.x
90. Luo J, Chen X, Chen G, et al. Dihydroartemisinin induces radiosensitivity in cervical cancer cells by modulating cell cycle progression. *Saudi Med J*. 2013;34(3):254–260.
91. Zuo ZJ, Wang ST, Jiang LX, et al. Effect of dihydroartemisinin combined irradiation on the apoptosis of human lung cancer GLC-82 cells and its mechanism study. *Chin J Integr Med*. 2014;34(10):1220–1224.
92. Zhang ZS, Wang J, Shen YB, et al. Dihydroartemisinin increases temozolomide efficacy in glioma cells by inducing autophagy. *Oncol Lett*. 2015;10(1):379–383. doi:10.3892/ol.2015.3183
93. Barbuti A, Chen Z-S. Paclitaxel through the ages of anticancer therapy: exploring its role in chemoresistance and radiation therapy. *Cancers*. 2015;7:2360–2371. doi:10.3390/cancers7040897
94. von Eiff D, Bozorgmehr F, Chung I, et al. Paclitaxel for treatment of advanced small cell lung cancer (SCLC): a retrospective study of 185 patients. *J Thorac Dis*. 2020;12(3):782–793. doi:10.21037/jtd.2019.12.74
95. Mukhtar E, Adhami VM, Mukhtar H. Targeting microtubules by natural agents for cancer therapy. *Mol Cancer Ther*. 2014;13(2):275–284. doi:10.1158/1535-7163.Mct-13-0791
96. Xia Y, Li YH, Chen Y, et al. A phase II trial of concurrent chemoradiotherapy with weekly paclitaxel and carboplatin in advanced oesophageal carcinoma. *Int J Clin Oncol*. 2018;23(3):458–465. doi:10.1007/s10147-018-1240-4
97. Cowen RL, Williams KJ, Chinje EC, et al. Hypoxia targeted gene therapy to increase the efficacy of tirapazamine as an adjuvant to radiotherapy: reversing tumor radioresistance and effecting cure. *Cancer Res*. 2004;64(4):1396–1402. doi:10.1158/0008-5472.can-03-2698
98. Marcu L, Olver I. Tirapazamine: from bench to clinical trials. *Curr Clin Pharmacol*. 2006;1(1):71–79. doi:10.2174/157488406775268192
99. Delahoussaye YM, Hay MP, Puijnt FB, Denny WA, Brown JM. Improved potency of the hypoxic cytotoxin tirapazamine by DNA-targeting. *Biochem Pharmacol*. 2003;65(11):1807–1815. doi:10.1016/s0006-2952(03)00199-0
100. Le QT, McCoy J, Williamson S, et al. Phase I study of tirapazamine plus cisplatin/etoposide and concurrent thoracic radiotherapy in limited-stage small cell lung cancer (S0004): a Southwest Oncology Group study. *Clin Cancer Res*. 2004;10(16):5418–5424. doi:10.1158/1078-0432.Ccr-04-0436
101. Rischin D, Peters L, Fisher R, et al. Tirapazamine, cisplatin, and radiation versus fluorouracil, cisplatin, and radiation in patients with locally advanced head and neck cancer: a randomized phase II trial of the Trans-Tasman Radiation Oncology Group (TROG 98.02). *J Clin Oncol*. 2005;23:79–87. doi:10.1200/jco.2005.01.072
102. Rischin D, Peters LJ, O'Sullivan B, et al. Tirapazamine, cisplatin, and radiation versus cisplatin and radiation for advanced squamous cell carcinoma of the head and neck (TROG 02.02, HeadSTART): a phase III trial of the Trans-Tasman Radiation Oncology Group. *J Clin Oncol*. 2010;28(18):2989–2995. doi:10.1200/jco.2009.27.4449
103. Wang J, Guise C, Dachs G, et al. Identification of one-electron reductases that activate both the hypoxia prodrug SN30000 and diagnostic probe EF5. *Biochem Pharmacol*. 2014;91(4):436–446. doi:10.1016/j.bcp.2014.08.003
104. Mistry IN, Thomas M, Calder EDD, Conway SJ, Hammond EM. Clinical advances of hypoxia-activated prodrugs in combination with radiation therapy. *Int J Radiat Oncol Biol Phys*. 2017;98(5):1183–1196. doi:10.1016/j.ijrobp.2017.03.024
105. Patterson LH, McKeown SR, Ruparella K, et al. Enhancement of chemotherapy and radiotherapy of murine tumours by AQ4N, a bioreductively activated anti-tumour agent. *Br J Cancer*. 2000;82(12):1984–1990. doi:10.1054/bjoc.2000.1163
106. Steward WP, Middleton M, Benghiat A, et al. The use of pharmacokinetic and pharmacodynamic end points to determine the dose of AQ4N, a novel hypoxic cell cytotoxin, given with fractionated radiotherapy in a phase I study. *Ann Oncol*. 2007;18(6):1098–1103. doi:10.1093/annonc/mdm120
107. Albertella MR, Loadman PM, Jones PH, et al. Hypoxia-selective targeting by the bioreductive prodrug AQ4N in patients with solid tumors: results of a phase I study. *Clin Cancer Res*. 2008;14(4):1096–1104. doi:10.1158/1078-0432.Ccr-07-4020
108. Hong CR, Dickson BD, Jaiswal JK, et al. Cellular pharmacology of evofosfamide (TH-302): a critical re-evaluation of its bystander effects. *Biochem Pharmacol*. 2018;156:265–280. doi:10.1016/j.bcp.2018.08.027
109. Takakusagi Y, Kishimoto S, Naz S, et al. Radiotherapy synergizes with the hypoxia-activated prodrug evofosfamide: in vitro and in vivo studies. *Antioxid Redox Signal*. 2018;28:131–140. doi:10.1089/ars.2017.7106
110. Peeters SG, Zegers CM, Biemans R, et al. TH-302 in combination with radiotherapy enhances the therapeutic outcome and is associated with pretreatment [18F]HX4 hypoxia PET imaging. *Clin Cancer Res*. 2015;21:2984–2992. doi:10.1158/1078-0432.Ccr-15-0018
111. Lohse I, Rasowski J, Cao P, et al. Targeting hypoxic microenvironment of pancreatic xenografts with the hypoxia-activated prodrug TH-302. *Oncotarget*. 2016;7(23):33571–33580. doi:10.18632/oncotarget.9654
112. Wang Z, Dabrosin C, Yin X, et al. Broad targeting of angiogenesis for cancer prevention and therapy. *Semin Cancer Biol*. 2015;35:S224–s243. doi:10.1016/j.semcancer.2015.01.001
113. Larue RT, Van De Voorde L, Berbée M, et al. A Phase I 'window-of-opportunity' trial testing evofosfamide (TH-302), a tumour-selective hypoxia-activated cytotoxic prodrug, with preoperative chemoradiotherapy in oesophageal adenocarcinoma patients. *BMC Cancer*. 2016;16:644. doi:10.1186/s12885-016-2709-z



114. Wang Y, Gray JP, Mishin V, Heck DE, Laskin DL, Laskin JD. Distinct roles of cytochrome P450 reductase in mitomycin C redox cycling and cytotoxicity. *Mol Cancer Ther.* 2010;9(6):1852–1863. doi:10.1158/1535-7163.Mct-09-1098
115. Rockwell S, Keyes SR, Sartorelli AC. Preclinical studies of porfiromycin as an adjunct to radiotherapy. *Radiat Res.* 1988;116(1):100–113. doi:10.2307/3577481
116. Haffty BG, Wilson LD, Son YH, et al. Concurrent chemo-radiotherapy with mitomycin C compared with porfiromycin in squamous cell cancer of the head and neck: final results of a randomized clinical trial. *Int J Radiat Oncol Biol Phys.* 2005;61(1):119–128. doi:10.1016/j.ijrobp.2004.07.730
117. Burd R, Lavorgna SN, Lenaz L, et al. Radiosensitization of hypoxic tumors by the bioreductive agent apaziquone (EO9, EOquin (TM)). *Clin Cancer Res.* 2005;9008S–9008S.
118. Wang Z, Ding Y, Geng G, Zhu N. Analysis of energy efficiency retrofit schemes for heating, ventilating and air-conditioning systems in existing office buildings based on the modified bin method. *Energy Convers Manage.* 2014;77:233–242. doi:10.1016/j.enconman.2013.09.037
119. Suwala AK, Kahlert UD, Maciaczyk J. Abstract 2515: pharmacological WNT-inhibition acts synergistically with chemo- and radiotherapy by overcoming treatment-resistance in glioma stem cells. *Cancer Res.* 2016;76(14Supplement):2515. doi:10.1158/1538-7445
120. Saker J, Huang S, Park L, Pedersen M, Kragh M, Harari P. Abstract 1027: EGFR targeting antibody SYM004 causes radiosensitization in tumor cells expressing wild-type K-Ras via modulation of MAPK signaling. *Cancer Res.* 2013;73(8Supplement):1027. doi:10.1158/1538-7445
121. Prabakaran PJ, Javaid AM, Swick AD, et al. Radiosensitization of adenoid cystic carcinoma with MDM2 inhibition. *Clin Cancer Res.* 2017;23(20):6044–6053. doi:10.1158/1078-0432.Ccr-17-0969
122. Zhuang HM, Miao XL, Zhao ZQ, Zhang L. Application of nano calcium phosphate biomaterials and bone tissue engineering on in exercise-induced injury. *Adv Mater Res.* 2014;951:109–112. doi:10.4028/www.scientific.net/AMR.951.109
123. Liu WL, Gao M, Tzen KY, et al. Targeting phosphatidylinositol 3-Kinase/Akt pathway by BKM120 for radiosensitization in hepatocellular carcinoma. *Oncotarget.* 2014;5(11):3662–3672. doi:10.18632/oncotarget.1978
124. Chen Y-H, Wei M-F, Wang C-W, et al. Dual phosphoinositide 3-kinase/mammalian target of rapamycin inhibitor is an effective radiosensitizer for colorectal cancer. *Cancer Lett.* 2015;357(2):582–590. doi:10.1016/j.canlet.2014.12.015
125. Jayakumar S, Patwardhan RS, Pal D, Sharma D, Sandur SK. Dimethoxycurcumin, a metabolically stable analogue of curcumin enhances the radiosensitivity of cancer cells: possible involvement of ROS and thioredoxin reductase. *Biochem Biophys Res Commun.* 2016;478(1):446–454. doi:10.1016/j.bbrc.2016.06.144
126. Kinsella TJ, Dobson PP, Mitchell JB, Fornace AJ. Enhancement of X ray induced DNA damage by pre-treatment with halogenated pyrimidine analogs. *Int J Radiat Oncol.* 1987;13(5):733–739. doi:10.1016/0360-3016(87)90292-6
127. Wang S, Zhao P, Zhang C, Bu Y. Mechanisms responsible for high energy radiation induced damage to single-stranded DNA modified by radiosensitizing 5-halogenated deoxyuridines. *J Phys Chem B.* 2016;120:2649–2657. doi:10.1021/acs.jpcc.5b11432
128. Benej M, Hong X, Vibhute S, et al. Papaverine and its derivatives radiosensitize solid tumors by inhibiting mitochondrial metabolism. *Proc Natl Acad Sci.* 2018;115(42):10756–10761. doi:10.1073/pnas.1808945115
129. Lhuillier C, Rudqvist N-P, Elemento O, Formenti SC, Demaria S. Radiation therapy and anti-tumor immunity: exposing immunogenic mutations to the immune system. *Genome Med.* 2019;11(1):40. doi:10.1186/s13073-019-0653-7
130. Bourillon L, Bourgier C, Gaborit N, et al. An auristatin-based antibody-drug conjugate targeting HER3 enhances the radiation response in pancreatic cancer. *Int J Cancer.* 2019;145(7):1838–1851. doi:10.1002/ijc.32273
131. González JE, Barquero JF, Lee M, García O, Casaco A. Radiosensitization induced by the anti-epidermal growth factor receptor monoclonal antibodies cetuximab and nimotuzumab in A431 cells. *Cancer Biol Ther.* 2012;13(2):71–76. doi:10.4161/cbt.13.2.18439
132. Buchanan IM, Scott T, Tandle AT, et al. Radiosensitization of glioma cells by modulation of Met signalling with the hepatocyte growth factor neutralizing antibody, AMG102. *J Cell Mol Med.* 2011;15(9):1999–2006. doi:10.1111/j.1582-4934.2010.01122.x
133. Nieder C, Mannsäker B, Dalhaug A, Pawinski A, Haukland E. Palliative radiotherapy in cancer patients with increased serum c-reactive protein level. Article. *In Vivo (Brooklyn).* 2016;30(5):581–586.
134. Kanegasaki S, Matsushima K, Shiraishi K, Nakagawa K, Tsuchiya T. Macrophage inflammatory protein derivative ECI301 enhances the alarmin-associated abscopal benefits of tumor radiotherapy. *Cancer Res.* 2014;74:5070–5078. doi:10.1158/0008-5472
135. Krüger M, Amort J, Wilgenbus P, et al. The anti-apoptotic PON2 protein is Wnt-catenin-regulated and correlates with radiotherapy resistance in OSCC patients. *Oncotarget.* 2016;7(32).
136. Cao Y, Yang L, Jiang W, et al. Therapeutic evaluation of Epstein-Barr virus-encoded latent membrane protein-1 targeted DNase for treating of nasopharyngeal carcinomas. *Mol Ther.* 2014;22(2):371–377. doi:10.1038/mt.2013.257
137. Walker JM, Rolig AS, Charych DH, et al. NKTR-214 immunotherapy synergizes with radiotherapy to stimulate systemic CD8+ T cell responses capable of curing multi-focal cancer. *Immunother Cancer.* 2020;8(1). doi:10.1136/jitc-2019-000464
138. Gandellini P, Rancati T, Valdagni R, Zaffaroni N. miRNAs in tumor radiation response: bystanders or participants? *Trends Mol Med.* 2014;20(9):529–539. doi:10.1016/j.molmed.2014.07.004
139. de Jong MC, Ten Hoeve JJ, Grénman R, et al. Pretreatment microRNA expression impacting on epithelial-to-mesenchymal transition predicts intrinsic radiosensitivity in head and neck cancer cell lines and patients. *Clin Cancer Res.* 2015;21:5630–5638. doi:10.1158/1078-0432
140. Huang A, Ono Y. Estimation of wrist flexion angle from muscle thickness changes measured by a flexible ultrasonic sensor. *IEEE.* 2016;188–191.
141. Shao Y, Song X, Jiang W, et al. MicroRNA-621 acts as a tumor radiosensitizer by directly targeting SETDB1 in hepatocellular carcinoma. *Mol Ther.* 2019;27(2):355–364. doi:10.1016/j.yth.2018.11.005
142. Zhang P, Wang L, Rodriguez-Aguayo C, et al. miR-205 acts as a tumour radiosensitizer by targeting ZEB1 and Ubc13. *Nat Commun.* 2014;5:5671. doi:10.1038/ncomms6671
143. Song L, Peng L, Hua S, et al. miR-144-5p enhances the radiosensitivity of non-small-cell lung cancer cells via targeting ATF2. *Biomed Res Int.* 2018;2018:5109497. doi:10.1155/2018/5109497
144. Luo J, Si ZZ, Li T, et al. MicroRNA-146a-5p enhances radiosensitivity in hepatocellular carcinoma through replication protein A3-induced activation of the DNA repair pathway. *Am J Physiol Cell Physiol.* 2019;316(3):C299–c311. doi:10.1152/ajpcell.00189.2018
145. Wu SJ, Chen J, Wu B, Wang YJ, Guo KY. MicroRNA-150 enhances radiosensitivity by inhibiting the AKT pathway in NK/T cell lymphoma. *J Exp Clin Cancer Res.* 2018;37(1):18. doi:10.1186/s13046-017-0639-5
146. Yin H, Ma J, Chen L, et al. MiR-99a enhances the radiation sensitivity of non-small cell lung cancer by targeting mTOR. *Cell Physiol Biochem.* 2018;46(2):471–481. doi:10.1159/000488615

147. Pajic M, Froio D, Daly S, et al. miR-139-5p Modulates radiotherapy resistance in breast cancer by repressing multiple gene networks of DNA repair and ROS defense. *Cancer Res.* 2018;78(2):501–515. doi:10.1158/0008-5472.Can-16-3105
148. Hu Z, Tie Y, Lv G, Zhu J, Fu H, Zheng X. Transcriptional activation of miR-320a by ATF2, ELK1 and YY1 induces cancer cell apoptosis under ionizing radiation conditions. *Int J Oncol.* 2018;53(4):1691–1702. doi:10.3892/ijo.2018.4497
149. Song Y, Zuo Y, Qian XL, et al. Inhibition of microRNA-21-5p promotes the radiation sensitivity of non-small cell lung cancer through HSMH2. *Cell Physiol Biochem.* 2017;43(3):1258–1272. doi:10.1159/000481839
150. Zhang L, Wang C, Xue Z-X. Inhibition of miR-630 enhances the cell resistance to radiation by directly targeting CDC14A in human glioma. *Am J Transl Res.* 2017;9(3):1255–1265.
151. Gu J, Li Y, Zeng J, et al. Knockdown of HIF-1 $\alpha$  by siRNA-expressing plasmid delivered by attenuated Salmonella enhances the antitumor effects of cisplatin on prostate cancer. *Sci Rep.* 2017;7(1):7546. doi:10.1038/s41598-017-07973-4
152. Mehta M, Basalingappa K, Griffith JN, et al. HuR silencing elicits oxidative stress and DNA damage and sensitizes human triple-negative breast cancer cells to radiotherapy. *Oncotarget.* 2016;7(40):64820–64835. doi:10.18632/oncotarget.11706
153. Qi R, Qiao T, Zhuang X. Small interfering RNA targeting S100A4 sensitizes non-small-cell lung cancer cells (A549) to radiation treatment. *Onco Targets Ther.* 2016;9:3753–3762. doi:10.2147/ott.S106557
154. Ohnishi K, Scuric Z, Schiestl RH, Okamoto N, Takahashi A, Ohnishi T. siRNA targeting NBS1 or XIAP increases radiation sensitivity of human cancer cells independent of TP53 status. *Radiat Res.* 2006;166(3):454–462. doi:10.1667/rr3606.1
155. Khan Z, Khan AA, Prasad GBKS, Khan N, Tiwari RP, Bisen PS. Growth inhibition and chemo-radiosensitization of head and neck squamous cell carcinoma (HNSCC) by survivin-siRNA lentivirus. *Radiother Oncol.* 2016;118(2):359–368. doi:10.1016/j.radonc.2015.12.007
156. Jackson MR, Bavelaar BM, Waghorn PA, et al. Radiolabeled oligonucleotides targeting the RNA subunit of telomerase inhibit telomerase and induce DNA damage in telomerase-positive cancer cells. *Cancer Res.* 2019;79(18):4627–4637. doi:10.1158/0008-5472
157. Cao F, Ju X, Chen D, et al. Phosphorothioate-modified antisense oligonucleotides against human telomerase reverse transcriptase sensitize cancer cells to radiotherapy. *Mol Med Rep.* 2017;16(2):2089–2094. doi:10.3892/mmr.2017.6778
158. Park SI, Park SJ, Lee J, et al. Inhibition of cyclic AMP response element-directed transcription by decoy oligonucleotides enhances tumor-specific radiosensitivity. *Biochem Biophys Res Commun.* 2016;469(3):363–369. doi:10.1016/j.bbrc.2015.11.122
159. Yu C, Yu Y, Xu Z, et al. Antisense oligonucleotides targeting human telomerase mRNA increases the radiosensitivity of nasopharyngeal carcinoma cells. *Mol Med Rep.* 2015;11(4):2825–2830. doi:10.3892/mmr.2014.3105
160. Hernández-Rivera M, Kumar I, Cho SY, et al. High-performance hybrid bismuth-carbon nanotube based contrast agent for X-ray CT imaging. *ACS Appl Mater Interfaces.* 2017;9:5709–5716. doi:10.1021/acsami.6b12768
161. Haume K, Rosa S, Grellet S, et al. Gold nanoparticles for cancer radiotherapy: a review. *Cancer Nanotechnol.* 2016;7(1):8. doi:10.1186/s12645-016-0021-x
162. Ma N, Liu P, He N, Gu N, Wu F-G, Chen Z. Action of gold nanopikes-based nanoradiosensitizers: cellular internalization, radiotherapy, and autophagy. *ACS Appl Mater Interfaces.* 2017;9:31526–31542. doi:10.1021/acsami.7b09599
163. Li Y, Yun K-H, Lee H, Goh S-H, Suh Y-G, Choi Y. Porous platinum nanoparticles as a high-Z and oxygen generating nanozyme for enhanced radiotherapy in vivo. *Biomaterials.* 2019;197:12–19. doi:10.1016/j.biomaterials.2019.01.004
164. Yang X, Yang M, Pang B, Vara M, Xia Y. Gold nanomaterials at work in biomedicine. *Chem Rev.* 2015;115(19):10410–10488. doi:10.1021/acs.chemrev.5b00193
165. Cui L, Her S, Borst GR, Bristow RG, Jaffray DA, Allen C. Radiosensitization by gold nanoparticles: will they ever make it to the clinic? *Radiother Oncol.* 2017;124(3):344–356. doi:10.1016/j.radonc.2017.07.007
166. Zhang XD, Luo Z, Chen J, et al. Ultrasmall glutathione-protected gold nanoclusters as next generation radiotherapy sensitizers with high tumor uptake and high renal clearance. *Sci Rep.* 2015;5:8669. doi:10.1038/srep08669
167. Luo D, Wang X, Zeng S, Ramamurthy G, Burda C, Basilion JP. Prostate-specific membrane antigen targeted gold nanoparticles for prostate cancer radiotherapy: does size matter for targeted particles? *Chem Sci.* 2019;10(35):8119–8128. doi:10.1039/c9sc02290b
168. Mathur P, Jha S, Ramteke S, Jain NK. Pharmaceutical aspects of silver nanoparticles. *Artif Cells, Nanomed Biotechnol.* 2018;46(sup1):115–126. doi:10.1080/21691401.2017.1414825
169. Li S, Porcel E, Remita H, et al. Platinum nanoparticles: an exquisite tool to overcome radioresistance. *Cancer Nanotechnol.* 2017;8(1):4. doi:10.1186/s12645-017-0028-y
170. Pinel S, Thomas N, Boura C, Barberi-Heyob M. Approaches to physical stimulation of metallic nanoparticles for glioblastoma treatment. *Adv Drug Deliv Rev.* 2019;138:344–357. doi:10.1016/j.addr.2018.10.013
171. Zhao J, Liu P, Ma J, et al. Enhancement of radiosensitization by silver nanoparticles functionalized with polyethylene glycol and aptamer as1411 for glioma irradiation therapy. *Int J Nanomedicine.* 2019;14:9483–9496. doi:10.2147/IJN.S224160
172. Liu Z, Tan H, Zhang X, et al. Enhancement of radiotherapy efficacy by silver nanoparticles in hypoxic glioma cells. *Artif Cells, Nanomed Biotechnol.* 2018;46(sup3):S922–s930. doi:10.1080/21691401.2018.1518912
173. Fathy MM. Biosynthesis of silver nanoparticles using thymoquinone and evaluation of their radio-sensitizing activity. *BioNanoScience.* 2020;10(1):260–266. doi:10.1007/s12668-019-00702-3
174. Rajaei A, Wang S, Zhao L, et al. Multifunction bismuth gadolinium oxide nanoparticles as radiosensitizer in radiation therapy and imaging. *Phys Med Biol.* 2019;64(19):195007. doi:10.1088/1361-6560/ab2154
175. Fält T, Söderberg M, Wassélius J, Leander P. Material decomposition in dual-energy computed tomography separates high-Z elements from iodine, identifying potential contrast media tailored for dual contrast medium examinations. *J Comput Assist Tomogr.* 2015;39(6):975–980. doi:10.1097/rct.0000000000000298
176. Chen X, Song J, Chen X, Yang H. X-ray-activated nanosystems for theranostic applications. *Chem Soc Rev.* 2019;48(11):3073–3101. doi:10.1039/c8cs00921j
177. Liu Y, Zhang P, Li F, et al. Metal-based nano enhancers for future radiotherapy: radiosensitizing and synergistic effects on tumor cells. *Theranostics.* 2018;8(7):1824–1849. doi:10.7150/thno.22172
178. Mardare AI, Mardare CC, Kollender JP, Huber S, Hassel AW. Basic properties mapping of anodic oxides in the hafnium-niobium-tantalum ternary system. *Sci Technol Adv Mater.* 2018;19(1):554–568. doi:10.1080/14686996.2018.1498703
179. Taupin F, Flaender M, Delorme R, et al. Gadolinium nanoparticles and contrast agent as radiation sensitizers. *Phys Med Biol.* 2015;60(11):4449–4464. doi:10.1088/0031-9155/60/11/4449



180. Shahbazi M-A, Faghfour L, Ferreira MPA, et al. The versatile biomedical applications of bismuth-based nanoparticles and composites: therapeutic, diagnostic, biosensing, and regenerative properties. *Chem Soc Rev*. 2020;49(4):1253–1321. doi:10.1039/C9CS00283A
181. Le Duc G, Roux S, Paruta-Tuarez A, et al. Advantages of gadolinium based ultrasmall nanoparticles vs molecular gadolinium chelates for radiotherapy guided by MRI for glioma treatment. *Cancer Nanotechnol*. 2014;5(1):4. doi:10.1186/s12645-014-0004-8
182. Hu P, Fu Z, Liu G, et al. Gadolinium-based nanoparticles for theranostic MRI-guided radiosensitization in hepatocellular carcinoma. *Front Bioeng Biotechnol*. 2019;7:368. doi:10.3389/fbioe.2019.00368
183. Detappe A, Lux F, Tillement O. Pushing radiation therapy limitations with theranostic nanoparticles. *Nanomedicine (London, England)*. 2016;11(9):997–999. doi:10.2217/nm.16.38
184. McGinnity TL, Dominguez O, Curtis TE, Nallathambi PD, Hoffman AJ, Roeder RK. Hafnia (HfO<sub>2</sub>) nanoparticles as an X-ray contrast agent and mid-infrared biosensor. *Nanoscale*. 2016;8(28):13627–13637. doi:10.1039/C6NR03217F
185. Jayaraman V, Bhavesh G, Chinnathambi S, Ganesan S, Aruna P. Synthesis and characterization of hafnium oxide nanoparticles for bio-safety. *Mater Express*. 2014. doi:10.1166/mex.2014.1190
186. Bonvalot S, Le Pechoux C, De Baere T, et al. First-in-human study testing a new radioenhancer using nanoparticles (NBTXR3) activated by radiation therapy in patients with locally advanced soft tissue sarcomas. *Clin Cancer Res*. 2016;23:908–917. doi:10.1158/1078-0432
187. Prasad K, Bazaka O, Chua M, et al. Metallic biomaterials: current challenges and opportunities. *Materials (Basel)*. 2017;10(8):884. doi:10.3390/ma10080884
188. Jin Y, Ma X, Zhang S, et al. A tantalum oxide-based core/shell nanoparticle for triple-modality image-guided chemo-thermal synergetic therapy of esophageal carcinoma. *Cancer Lett*. 2017;397:61–71. doi:10.1016/j.canlet.2017.03.030
189. Rathnayake S, Mongan J, Torres AS, et al. In vivo comparison of tantalum, tungsten, and bismuth enteric contrast agents to complement intravenous iodine for double-contrast dual-energy CT of the bowel. *Contrast Media Mol Imaging*. 2016;11(4):254–261. doi:10.1002/cmmi.1687
190. Lu Y-C, Yang C-X, Yan X-P. Radiopaque tantalum oxide coated persistent luminescent nanoparticles as multimodal probes for in vivo near-infrared luminescence and computed tomography bioimaging. *Nanoscale*. 2015;7(42):17929–17937. doi:10.1039/C5NR05623C
191. Brown R, Tehei M, Oktaria S, et al. High-Z nanostructured ceramics in radiotherapy: first evidence of taO<sub>5</sub>-induced dose enhancement on radioresistant cancer cells in an MV photon field. *Part Part Syst Charact*. 2014;31(4):500–505. doi:10.1002/ppsc.201300276
192. Song G, Chao Y, Chen Y, et al. All-in-one theranostic nanoplat-form based on hollow TaOx for chelator-free labeling imaging, drug delivery, and synergistically enhanced radiotherapy. *Adv Funct Mater*. 2016;26(45):8243–8254. doi:10.1002/adfm.201603845
193. Song G, Chen Y, Liang C, et al. Catalase-loaded TaOx nanoshells as bio-nanoreactors combining high-Z element and enzyme delivery for enhancing radiotherapy. *Adv Mater*. 2016;28(33):7143–7148. doi:10.1002/adma.201602111
194. Bulatov A, Moskvina L, Rodinkov O, Ermakov S. Department of analytical chemistry of saint petersburg state university celebrates one hundred and fifty years anniversary: international year of the periodic table of chemical elements. *Talanta*. 2020;206:119759. doi:10.1016/j.talanta.2019.03.081
195. Chellan P, Sadler PJ. The elements of life and medicines. *Philos Trans a Math Phys Eng Sci*. 2015;373(2037). doi:10.1098/rsta.2014.0182
196. Hossain M, Su M. Nanoparticle location and material-dependent dose enhancement in X-ray radiation therapy. *J Phys Chem C*. 2012;116(43):23047–23052. doi:10.1021/jp306543q
197. Yu X, Li A, Zhao C, Yang K, Chen X, Li W. Ultrasmall semi-metal nanoparticles of bismuth for dual-modal computed tomography/photoacoustic imaging and synergistic thermoradiotherapy. *ACS Nano*. 2017;11(4):3990–4001. doi:10.1021/acsnano.7b00476
198. Qiu J, Xiao Q, Zheng X, et al. Single W18O<sub>49</sub> nanowires: a multifunctional nanoplatform for computed tomography imaging and photothermal/photodynamic/radiation synergistic cancer therapy. *Nano Res*. 2015;8(11):3580–3590. doi:10.1007/s12274-015-0858-z
199. Gong L, Yan L, Zhou R, Xie J, Wu W, Gu Z. Two-dimensional transition metal dichalcogenide nanomaterials for combination cancer therapy. *J Mater Chem B*. 2017;5(10):1873–1895. doi:10.1039/C7TB00195A
200. Cheng X, Yong Y, Dai Y, et al. Enhanced radiotherapy using bismuth sulfide nanoagents combined with photo-thermal treatment. *Theranostics*. 2017;7(17):4087–4098. doi:10.7150/thno.20548
201. Feng L, Yang D, Gai S, et al. Single bismuth tungstate nanosheets for simultaneous chemo-, photothermal, and photodynamic therapies mediated by near-infrared light. *Chem Eng J*. 2018;351:1147–1158. doi:10.1016/j.cej.2018.06.170
202. Detappe A, Thomas E, Tibbitt MW, et al. Ultrasmall silica-based bismuth gadolinium nanoparticles for dual magnetic resonance-computed tomography image guided radiation therapy. *Nano Lett*. 2017;17(3):1733–1740. doi:10.1021/acs.nanolett.6b05055
203. Yong Y, Zhou L, Zhang S, et al. Gadolinium polytungstate nanoclusters: a new theranostic with ultrasmall size and versatile properties for dual-modal MR/CT imaging and photothermal therapy/radiotherapy of cancer. *NPG Asia Mater*. 2016;8(5):e273–e273. doi:10.1038/am.2016.63
204. Ma P, Xiao H, Yu C, et al. Enhanced cisplatin chemotherapy by iron oxide nanocarrier-mediated generation of highly toxic reactive oxygen species. *Nano Lett*. 2017;17(2):928–937. doi:10.1021/acs.nanolett.6b04269
205. Vallabani NVS, Singh S. Recent advances and future prospects of iron oxide nanoparticles in biomedicine and diagnostics. *3 Biotech*. 2018;8(6):279. doi:10.1007/s13205-018-1286-z
206. Tang GD, Li ZZ, Ma L, et al. Three models of magnetic ordering in typical magnetic materials. *Phys Rep*. 2018;758:1–56. doi:10.1016/j.physrep.2018.06.009
207. Nica V, Caro C, Pérez-Muñoz JM, Leal MP, Garcia-Martin ML. Bi-magnetic core-shell CoFe(2)O(4)@MnFe(2)O(4) nanoparticles for in vivo theranostics. *Nanomaterials (Basel, Switzerland)*. 2020;10(5). doi:10.3390/nano10050907
208. Meidanchi A, Akhavan O, Khoei S, Shokri A, Hajikarimi Z, Khansari N. ZnFe<sub>2</sub>O<sub>4</sub> nanoparticles as radiosensitizers in radiotherapy of human prostate cancer cells. *Mater Sci Eng C Mater Biol Appl*. 2015;46:394–399. doi:10.1016/j.msec.2014.10.062
209. Hidayatullah M, Nurhasanah I, Budi WS. ZnFe<sub>2</sub>O<sub>4</sub> nanoparticles for potential application in radiosensitization. *J Phys Conf Ser*. 2016;694:012028. doi:10.1088/1742-6596/694/1/012028
210. Salunkhe A, Khot V, Patil SI, et al. Magneto-chemotherapy with high-magnetic-moment iron oxide nanoparticles for cancer theranostics. *ACS Appl Bio Mater*. 2020;3(4):2305–2313. doi:10.1021/acsbam.0c00077
211. Shao L, Gao Y, Yan F. Semiconductor quantum dots for biomedical applications. *Sensors (Basel)*. 2011;11(12):11736–11751. doi:10.3390/s111211736
212. Juzenas P, Chen W, Sun Y-P, et al. Quantum dots and nanoparticles for photodynamic and radiation therapies of cancer. *Adv Drug Deliv Rev*. 2008;60(15):1600–1614. doi:10.1016/j.addr.2008.08.004

213. Cline B, Delahunty I, Xie J. Nanoparticles to mediate X-ray-induced photodynamic therapy and Cherenkov radiation photodynamic therapy. *Wiley Interdiscip Rev Nanomed Nanobiotechnol.* 2019;11(2):e1541. doi:10.1002/wnan.1541
214. Yang W, Read PW, Mi J, et al. Semiconductor nanoparticles as energy mediators for photosensitizer-enhanced radiotherapy. *Int J Radiat Oncol Biol Phys.* 2008;72(3):633–635. doi:10.1016/j.ijrobp.2008.06.1916
215. Nakayama M, Sasaki R, Ogino C, et al. Titanium peroxide nanoparticles enhanced cytotoxic effects of X-ray irradiation against pancreatic cancer model through reactive oxygen species generation in vitro and in vivo. *Radiat Oncol.* 2016;11(1):91. doi:10.1186/s13014-016-0666-y
216. Morita K, Miyazaki S, Numako C, et al. Characterization of titanium dioxide nanoparticles modified with polyacrylic acid and H<sub>2</sub>O<sub>2</sub> for use as a novel radiosensitizer. *Free Radic Res.* 2016;50(12):1319–1328. doi:10.1080/10715762.2016.1241879
217. Jin J, Zhao Q. Engineering nanoparticles to reprogram radiotherapy and immunotherapy: recent advances and future challenges. *J Nanobiotechnology.* 2020;18(1):75. doi:10.1186/s12951-020-00629-y
218. Sun H, Wang X, Zhai S. The rational design and biological mechanisms of nanoradiosensitizers. *Nanomaterials (Basel, Switzerland).* 2020;10(3):504. doi:10.3390/nano10030504
219. Martel A. Recent progress in biomedical applications of nanodiamonds. *Nanosci Nanotechnol.* 2018;8:11–24. doi:10.5923/j.nn.20180801.03
220. Jia Y, Weng Z, Wang C, et al. Increased chemosensitivity and radiosensitivity of human breast cancer cell lines treated with novel functionalized single-walled carbon nanotubes. *Oncol Lett.* 2017;13(1):206–214. doi:10.3892/ol.2016.5402
221. Chan L, He L, Zhou B, et al. Cancer-targeted selenium nanoparticles sensitize cancer cells to continuous  $\gamma$  radiation to achieve synergetic chemo-radiotherapy. *Chem Asian J.* 2017;12(23):3053–3060. doi:10.1002/asia.201701227
222. Bromma K, Cicon L, Beckham W, Chithrani DB. Gold nanoparticle mediated radiation response among key cell components of the tumour microenvironment for the advancement of cancer nanotechnology. *Sci Rep.* 2020;10(1):12096. doi:10.1038/s41598-020-68994-0
223. Patra JK, Das G, Fraceto LF, et al. Nano based drug delivery systems: recent developments and future prospects. *J Nanobiotechnology.* 2018;16(1):71. doi:10.1186/s12951-018-0392-8
224. Chenthamara D, Subramaniam S, Ramakrishnan SG, et al. Therapeutic efficacy of nanoparticles and routes of administration. *Biomater Res.* 2019;23(1):20. doi:10.1186/s40824-019-0166-x
225. Blanco E, Shen H, Ferrari M. Principles of nanoparticle design for overcoming biological barriers to drug delivery. *Nat Biotechnol.* 2015;33(9):941–951. doi:10.1038/nbt.3330

## International Journal of Nanomedicine

Dovepress

### Publish your work in this journal

The International Journal of Nanomedicine is an international, peer-reviewed journal focusing on the application of nanotechnology in diagnostics, therapeutics, and drug delivery systems throughout the biomedical field. This journal is indexed on PubMed Central, MedLine, CAS, SciSearch®, Current Contents®/Clinical Medicine,

Journal Citation Reports/Science Edition, EMBase, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-nanomedicine-journal>