

Chapter 13

The Role of DUOX Isozymes in the Respiratory Tract Epithelium

Richard Harper

Abstract Increasingly, reactive oxygen species such as superoxide and hydrogen peroxide are recognized to be intentionally generated intracellularly to serve important cellular functions. A key protein family responsible for the regulated generation of reactive oxygen species in multiple cell types is the NOX/DUOX enzyme family. Two family members, DUOX1 and DUOX2, appear to be highly expressed in tissues of endodermal origin including the thyroid, respiratory tract, and gastrointestinal tract. In this chapter, we will focus our review on DUOX proteins in the respiratory tract. We will discuss a brief history of the discovery of the DUOX isoforms, the estimated hydrogen peroxide-generating capacity of DUOX in respiratory tract epithelium, putative functions of the DUOX enzymes, and some regulatory factors responsible for DUOX gene expression and oxidase activity.

Keywords DUOX1; DUOX2; respiratory tract; oxidase; peroxidase; host defense

Introduction

Recent evidence suggests that reactive oxygen species (ROS) such as superoxide ($O_2^{\cdot-}$) and hydrogen peroxide (H_2O_2) are not mere byproducts of cellular respiration, but are intentionally generated to serve important cellular functions (Terada, 2006; Finkel, 2003). A key protein family responsible for the regulated generation of ROS in multiple cell types is the NOX/DUOX enzyme family (Lambeth, 2002). The prototypical member of this family is gp91*phox* (NOX2), which is responsible for the respiratory burst in neutrophils (Gabig and Babior, 1979; Dusi et al., 1995; Hohn and Lehrer, 1975). Suh et al. were the first to characterize NOX1 (initially named Mox1; for mitogenic oxidase) as a non-mitochondrial, non-phagocytic enzyme capable of regulated ROS production (Suh et al., 1999). Since this discovery,

University of California at Davis, Genome and Biomedical Sciences Facility, 451 Health Sciences Drive, 6th Floor Davis, California USA 95616

there has been an exponentially growing body of literature characterizing the function of the non-phagocytic NOX/DUOX family members (NOX1, NOX3, NOX4, NOX5, DUOX1, and DUOX2) in various aspects of human biology and disease (Lambeth, 2004; Leto and Geiszt, 2006). In this chapter, we will focus our review on the expression and function of DUOX proteins in the respiratory tract.

Discovery and Localization of DUOX Family Members

Initial identification of the DUOX2 was made by Corinne Dupuy and colleagues using biochemical techniques to isolate the known membrane-bound NADPH oxidase in thyroid tissues (Dupuy et al., 1999). In this report, they described the isolation of a 1,210 amino acid protein, p138^{Tox}, which contained a gp91phox-like region and two EF-hand domains. EF-hand domains contain calcium binding sites and serve as regulatory sub units for enzymatic activity. The presence of EF-hand domains suggested that p138^{Tox} enzymatic activity may be regulated by calcium. Additionally, expression of p138^{Tox} was upregulated by forskolin and the gene localized to Chromosome 15q15. Shortly after this discovery, Xavier De Deken described the isolation of two human thyroid NADPH oxidases (De Deken et al., 2000). These genes, thyroid oxidase 1 (ThOX1) and thyroid oxidase 2 (ThOX2), were identified by low stringency hybridization screening of thyroid cDNA libraries using a 1.3-kilobase fragment of gp91phox. Similar to Dupuy's findings, both ThOX1 and ThOX2 contained a gp91phox-like region and two EF-hand domains, and both genes were upregulated by forskolin. In addition, increased sequence information identified novel N-terminal extensions for both proteins that contained a peroxidase-like domain. The 1,210 amino acid C-terminal portion of ThOX2 was identical to p138^{Tox}, which strongly suggested that these two proteins were the same protein. Because both proteins express dual oxidases, a heme peroxidase domain and an NADPH oxidase domain, ThOX1 and ThOX2/p138^{Tox} have been renamed DUOX1 and DUOX2, respectively.

Further characterization of DUOX enzyme expression in human tissues demonstrated *DUOX* mRNA transcripts in multiple non-thyroid human tissues including heart, kidney liver, pancreas, placenta, GI tract, prostate, testis, and the respiratory tract (Caillou et al., 2001; Edens et al., 2001; Geiszt et al., 2003; El Hassani et al., 2005). Further characterization of DUOX in the lung demonstrated that DUOX expression is likely limited to the respiratory tract epithelium (Geiszt et al., 2003). Although PCR evidence exists that several NOX family members are potentially expressed in the respiratory tract, expression levels of the other Nox family members are a thousand-fold less abundant compared to DUOX (Schwarzer et al., 2004). Based on in situ hybridization experiments, it appears that DUOX1 is predominantly located in proximal respiratory tract epithelial cells whereas DUOX2 is localized to submandibular or salivary glands during basal conditions (Schwarzer et al., 2004). However, protein confirmation of these findings has been limited due to the absence of DUOX isoform-specific antibodies.

Functional Role(s) of DUOX



The capacity for respiratory tract epithelium to generate H_2O_2 appears to be dependent upon the specific cell type utilized. For example, in unstimulated human primary tracheobronchial epithelial cells, the rate of H_2O_2 production was 0.5–2.5 nmol/10 min/10⁶ cells and this production increases to 2–5 nmol/10 min/10⁶ cells after stimulation with ATP (Moskwa et al., 2007). In HBE1 cells, a papilloma-transformed human tracheobronchial epithelial cell line (Yankaskas et al., 1993), H_2O_2 production was 0.03–0.17 nmol/10 min/10⁶ cells and this production increases to 0.26–0.45 nmol/10 min/10⁶ cells after stimulation with ATP (Wesley et al., 2007). In a human pulmonary mucoepidermoid carcinoma cell line, H_2O_2 production increased from 0.002–0.003 nmol/10 min/10⁶ cells to 0.012–0.014 nmol/10 min/10⁶ cells after PMA stimulation (Nakanaga et al., 2007). Although laboratory to laboratory variation may account for these differences, it appears that primary respiratory tract epithelial cells have the highest capacity for DUOX-mediated H_2O_2 generation. In parallel, we have observed that *DUOX1* or *DUOX2* mRNA expression and functional activity is higher in primary tracheobronchial epithelial cells compared to HBE1 cells (Xu and Haper, 2007).

It is well-accepted that DUOX2-mediated H_2O_2 production is critical for iodide oxidation during thyroid hormone synthesis (Igo et al., 1964; Degroot et al., 1972; Clark et al., 1975; Lissitzky, 1976; Rigutto et al., 2007; Song et al., 2007). Observations that mutations in *DUOX2* gene result in congenital hypothyroidism (Moreno et al., 2002; Vigone et al., 2005) confirm the available biochemical and expression data. However, similarly convincing evidence for the function of DUOX isoforms in the respiratory tract are lacking. Several compelling models to explain the expression and function of DUOX isoforms in the respiratory tract have been proposed. The location of DUOX isoforms in the respiratory tract epithelium and their homology to gp91phox strongly implicated the role of DUOX proteins in host defense.

Several mechanisms exist to provide innate host defense in the respiratory tract epithelium (Antunes and Cohen, 2007; Thacker, 2006; Grubor et al., 2006; Beisswenger and Bals, 2005; Ng et al., 2004; Zhang et al., 2000). The upper respiratory tract epithelium and conducting airways are structured to provide multiple layers of protection against invading organisms. This includes a tight epithelial barrier where the apical surface of the respiratory tract is in close apposition to prevent invasion. This apical surface is bathed in an airway surface liquid that is rich in antimicrobial peptides hostile to bacterial and viral products. Resting on top of this airway surface liquid is a mucous layer that provides a physical barrier against infection. Mucus that has ensnared bacteria and virus particles is cleared from the airway by ciliary action provided by the respiratory tract epithelial cells. The generation of hypothiocyanate in the airway surface liquid appears to be a critical part of the host defense for those bacteria and viral particles that are able to penetrate the initial mucociliary barrier (Goldman and Smith, 1973; Ratner and Prince, 2000).

Conner and others previously described the importance of lactoperoxidase (LPO) in maintaining sterility in the respiratory tract (Gerson et al., 2000; Conner

et al., 2002; Salathe et al., 1997; Wijkstrom-Frei et al., 2003). In this model, LPO catalyzes a reaction between H_2O_2 and thiocyanate (SCN^-) to generate the bactericidal product hypothiocyanate (OSCN^-) (Gerson et al., 2000; Conner et al., 2002; Salathe et al., 1997; Wijkstrom-Frei et al., 2003; Conner et al., 2007).



Absence of this abundant protein, nearly 1% of the total protein in airway surface liquid, results in significant colonization and infection of respiratory tract epithelium with bacteria. Although the source of H_2O_2 in the respiratory tract was not initially known, DUOX isoforms have since been verified to be the predominant enzymes responsible for H_2O_2 generation in the respiratory tract. In addition, Geiszt et al. demonstrated evidence to suggest that DUOX localization next to LPO production is consistent with this model (Geiszt et al., 2003).

Recently, Conner and Banfi independently, characterized the transport of SCN^- from the basolateral surface to the apical surface of the respiratory tract where enzymatic activation of SCN^- to OSCN^- will occur (Moskwa et al., 2007; Conner et al., 2007). This transport appears to be the rate-limiting step in the respiratory tract and is dependent upon the presence of functional CFTR chloride channels. The rate of SCN^- transport ($1.5\text{--}1.8\text{ nmol}/10\text{ min}/\text{cm}^2$) (Conner et al., 2007) parallels the rate of OSCN^- formation ($1.4\text{--}1.7\text{ nmol}/10\text{ min}/\text{cm}^2$) (Moskwa et al., 2007) in respiratory tract epithelium. This is consistent with the notion that SCN^- is immediately converted to OSCN^- once SCN^- is transported to the apical surface. However, two different model systems, rat versus human, were used in these reports, which limits our ability to directly compare the two studies. Of importance, LPO, DUOX-mediated H_2O_2 , and apical SCN^- are all required for bacterial killing of *S. aureus* or *P. aeruginosa* (Moskwa et al., 2007). The capacity of DUOX isoforms to generate H_2O_2 in respiratory tract epithelium ($5\text{ nmol}/10\text{ min}/\text{cm}^2$) is more than adequate to support this model.

In parallel to the putative antibacterial role for DUOX enzymes, our recent data suggest DUOX isoforms are important for innate antiviral host defense. The observation that DUOX2 expression is substantially increased by $\text{IFN-}\gamma$ and rhinovirus provides circumstantial evidence for this notion (Harper et al., 2005). In addition, respiratory tract epithelial cells are known to generate H_2O_2 in less than 1h after RV infection (Kaul et al., 2000; Biagioli et al., 1999). RV-induced H_2O_2 production is inhibited by diphenyleiodonium (DPI), which indicates the involvement of a flavoprotein, likely an NADPH oxidase, in this process. Therefore, DUOX appears to be the best candidate responsible for RV-induced H_2O_2 production in respiratory tract epithelium. The presence of EF-hand domains on DUOX isoforms suggests a mechanism by which RV infection can immediately induce DUOX-mediated H_2O_2 production (e.g. membrane-associated calcium flux (Geiszt et al., 2003; Forteza et al., 2005)). The specific contribution of DUOX isoforms in respiratory tract epithelial innate host defense against viral infection remains to be determined.

Alternative functions for DUOX isozymes in the respiratory tract have recently been suggested. Hydrogen ion concentration at the apical surface of respiratory tract epithelium serves important host defense functions. Not only does lower pH increase the efficiency of known respiratory tract peroxidases such as LPO, but an acidic environment is known to decrease viability of some respiratory viruses (Greenberg, 2007; Whitton et al., 2005; Hayden, 2004). Therefore, an attractive model for DUOX is the ability of this protein to simultaneously acidify the airway surface liquid to optimize LPO enzymatic activity and provide the H_2O_2 needed for LPO to convert SCN^- to $OSCN^-$. Recent data from Horst Fischer and colleagues support the role of DUOX in H^+ production and H^+ secretion. By using chemical inhibitors of DUOX (Schwarzer et al., 2004) and DUOX1-specific RNAi transcripts, they were able to demonstrate an approximately 50% reduction in intracellular acid production. These data suggest that under basal conditions there is DUOX-mediated turnover of NADPH to NADP + that is responsible for approximately 50% of H^+ production. Based on this model, it is presumed that activation of NADPH oxidase activity in the respiratory tract, and specifically activation of DUOX1, will result in a parallel increase in H^+ secretion. However, this presumption has not been formally tested. Similarly, it is not known if increased activation of DUOX2 will result in a similar increase in intracellular H^+ production or H^+ secretion.

DUOX1 appears to perform a specific role in signaling events related to mucin secretion and wound healing. Wesley et al. recently demonstrated that ATP-mediated P2 purinergic receptor activation was necessary for cell migration and wound closure in primary and virus-immortalized tracheobronchial epithelial cell cultures (Wesley et al., 2007). This functional activity correlated with increased DUOX1-dependent H_2O_2 production. Wound closure was significantly decreased in cells expressing siRNA against DUOX1, but not in cells expressing non-target siRNA transcripts. Together, these data strongly suggested that DUOX1-generated H_2O_2 is a necessary signal for adequate wound closure in human tracheobronchial epithelium. Although not explicitly tested, it is likely that release of intracellular ATP during injury induces purinergic receptor-mediated calcium flux, which directly triggers DUOX1 oxidase activity via EF-hand domain activation. Similarly, Shao et al. demonstrated that DUOX1 appears to be a critical protein necessary for phorbol 12-myristate 13-acetate- or human neutrophil elastase-mediated MUC5AC production in human tracheobronchial epithelial cell cultures (Shao and Nadel, 2005).

Regulation of DUOX Expression and Enzymatic Activity

As mentioned, the initial identification of DUOX isoforms demonstrated that *DUOX* mRNA is upregulated by forskolin in thyroid tissues (Dupuy et al., 1999; De Deken et al., 2000). Thyroid stimulating hormone also increased *DUOX1* and *DUOX2* mRNA expression in thyroid tissues after 24h of treatment (Pachucki et al., 2004). However, DUOX promoter activity does not appear to depend upon thyroid-specific regulators consistent with the expression of DUOX in multiple tis-

sue types (Christophe-Hobertus and Christophe, 2007). Data for the regulation of DUOX expression in the respiratory tract is similarly limited.

We recently demonstrated differential expression of *DUOX1* and *DUOX2* after treatment of human tracheobronchial epithelial cells with inflammatory cytokines. *DUOX1* increased in response to the Th2 cytokines interleukin-4 (IL-4) and interleukin-13 (IL-13). *DUOX2* mRNA expression increased significantly in response to the Th1 cytokine interferon- γ (IFN- γ), and increased modestly to interleukin-1a or interleukin-1b. Of potential importance, the nature of gene expression was significantly different between the two genes in response to their respective cytokines. *DUOX1*, for example, increased maximally three- to sixfold in response to IL-4 at varying doses of the cytokine. In comparison, *DUOX2* increased linearly with increasing doses of IFN- γ more than 15- to 20-fold. Although the fold-induction was substantially higher for *DUOX2* compared to *DUOX1*, *DUOX2* expression levels began to wane after 48 h. However, *DUOX1* levels continued to increase 72 h after cytokine treatment. These qualitative differences in regulation and the differential regulation of *DUOX1* and *DUOX2* by cytokines suggest both enzymes perform distinct functions in the respiratory tract epithelium. Localization of *DUOX1* and *DUOX2* protein in the respiratory tract epithelium, and identifying proteins that colocalized with the DUOX isoforms will be instrumental in determining the functions of each protein.

Current evidence suggests that constitutive DUOX-mediated oxidase activity is directly proportional to the total DUOX protein available in respiratory tract epithelial cells. Based on mRNA data, *DUOX1* is the predominant isoform expressed under basal conditions in airway cell cultures. Therefore, it is likely that *DUOX1* provides the majority of DUOX-mediated H_2O_2 under basal conditions. Cytokine treatments that increase *DUOX* mRNA levels, and presumably DUOX protein levels, induce a parallel increase in H_2O_2 production (Harper et al., 2005). This suggests that one method to regulate DUOX oxidase activity is to regulate the absolute level of protein in respiratory tract epithelium. Based on our previous data, it appears that maximal inducible expression of *DUOX1* or *DUOX2* result in similar levels of H_2O_2 generation in respiratory tract epithelium (Harper et al., 2005). Although *DUOX2* is more inducible by cytokine stimulation compared to *DUOX1*, it is possible that the protein levels of *DUOX2* or *DUOX1* are similar when maximally induced by their respective cytokines. Antibodies that differentiate between *DUOX1* and *DUOX2* will provide insights into the absolute protein levels for each isoform.

DUOX-mediated H_2O_2 generation can also be rapidly and robustly induced above constitutive levels in respiratory tract epithelium (Geiszt et al., 2003; Wesley et al., 2007; Forteza et al., 2005). Prior to the identification of *DUOX2* in thyroid tissue, it was established that thyroid cells required calcium for H_2O_2 generation (Ameziane-El-Hassani et al., 2005; Leseney et al., 1999; Nakamura et al., 1987). Similar to Nox5, DUOX family members contain two EF-hand domains suggesting that DUOX oxidase activity will increase in response to calcium (De Deken et al., 2000; Lambeth et al., 2007; Dupuy et al., 1988). Further evidence in thyroid and respiratory tract epithelium has confirmed that Ca^{++} influx is likely the predominant

factor regulating inducible DUOX-mediated NADPH oxidase activity. Treatment of respiratory tract epithelial cells with agonists for calcium mobilization, including ionomycin, thapsigargin, or ATP, has resulted in the rapid increase in H_2O_2 flux. This activity was blocked by nonspecific inhibitors of DUOX enzymes or siRNA against DUOX1.

There is some evidence that additional regulatory proteins are required for DUOX-mediated enzymatic activity. Schwarzer et al. demonstrated that known NOX2 regulatory proteins, p40phox, p47phox, and p67phox, are co-expressed with the DUOX isozymes in respiratory tract epithelium (Schwarzer et al., 2004). Shao et al. demonstrated that p47phox and p67phox proteins were identified in the membrane fraction of respiratory tract epithelial cells after PMA treatment. This localization corresponded to PMA-induced, DUOX1-mediated H_2O_2 production (Shao and Nadel, 2005). However, family members that are more closely related to NOX2 do not use these phox regulatory units for enzymatic activity (Banfi et al., 2003). There is direct evidence in thyroid tissues that cytosolic phox regulators, including Rac, do not provide regulatory functions for DUOX. And, patients with autosomal recessive chronic granulomatous disease, in which there are defects in cytosolic phox regulators, do not exhibit hypothyroidism. Together, these latter data suggest that the DUOX isozymes do not rely on these cytosolic phox regulatory units. However, differences in the predominant isoform found in thyroid tissue (DUOX2) versus respiratory tract epithelium (DUOX1) may be responsible for these contradictory data. Further studies with knock down of various phox regulatory units in respiratory tract epithelium will be able to definitively settle this issue.

The model that DUOX isoforms are simply H_2O_2 generators in the respiratory tract epithelium is consistent with current data, but presents potential teleological inconsistencies. For example, if the predominant function of the DUOX proteins is to generate H_2O_2 , why are *two* bulky DUOX proteins maintained in the airway rather than a single, and smaller, isoform of the NOX/DUOX family? This question can be partially addressed by the observation that DUOX1 and DUOX2 are differentially regulated in respiratory tract epithelium. Although H_2O_2 is produced by both DUOX proteins, it is possible that the functional signalosomes that assemble around DUOX1 are substantially different from the proteins that cluster around DUOX2. The presence of two different oxidases in the same cell type, allow efficient utilization of H_2O_2 as a single signal that produces two very distinct effects.

Similarly, it is unclear why two peroxidase-containing DUOX proteins are present in airway epithelium if several other heme peroxidases (e.g. LPO, MPO, and others) are abundantly present in airway surface liquid (Gerson et al., 2000; Salathe et al., 1995). Several investigators have suggested that the peroxidase domains within DUOX1 or DUOX2 are not functional (Lambeth et al., 2007; Dupuy et al., 1988; Banfi et al., 2003; Salathe et al., 1995; Geiszt and Leto, 2004). It is possible that the peroxidase domains represent the vestigial remains of an ancestral protein that has not been selected against. Or, the peroxidase domains may provide regulatory functions for the DUOX isozymes independent of heme peroxidase activity.

However, biochemical analysis of purified DUOX protein suggests that DUOX has functional heme peroxidase activity (Edens et al., 2001; Ha et al., 2005). DUOX peroxidase domains isolated from humans, *C. elegans*, and *Drosophila* demonstrated peroxidase activity. In addition, we have recently shown that *DUOX2* is required for functional peroxidase activity in respiratory tract epithelium (Harper et al., 2006). Therefore, based on physical proximity, it is likely that DUOX2-generated H_2O_2 is utilized by its own peroxidase rather than diffusing to LPO some distance away. Together, these data provide strong evidence that DUOX is not solely an H_2O_2 generator to assist the antibacterial functions of LPO but is performing distinct functions apart from other respiratory tract peroxidases. The demonstration of heme incorporation into the DUOX peroxidase domain has not been confirmed to date, and therefore, the functional capacity of the DUOX peroxidase domain remains controversial. Therefore not only are there questions related to the levels of oxidants generated via the DUOX pathway – i.e. oxidative balance but in addition there are questions related to the balance of oxidant generation between the DUOX isoforms and what these may signal.

Summary

There is growing interest in understanding the function and importance of DUOX proteins in the respiratory tract. Based on the current body of literature, it is clear that DUOX is important for antibacterial host defense functions in the respiratory tract as part of the DUOX/LPO/ SCN^- system. Evidence that DUOX isozymes are important for mucin secretion, proton secretion, and wound healing strongly supports the notion that these family members are critical for multiple functions in the respiratory tract. Regulation of DUOX enzymatic activity by calcium flux and regulation of protein expression by immune-modulating cytokines provide insights into additional potential functions for these family members. Protein isolation techniques that allow specific identification of each isoform are still required to fully characterize the role of DUOX1 and DUOX2 in the respiratory tract epithelium.

References

- Ameziane-El-Hassani R, Morand S, Boucher JL, et al. Dual oxidase-2 has an intrinsic Ca^{2+} - dependent H_2O_2 -generating activity. *J Biol Chem* 2005;280(34):30046–30054.
- Antunes MB, Cohen NA. Mucociliary clearance—a critical upper airway host defense mechanism and methods of assessment. *Current opinion in allergy and clinical immunology* 2007;7(1):5–10.
- Banfi B, Clark RA, Steger K, Krause KH. Two novel proteins activate superoxide generation by the NADPH oxidase NOX1. *J Biol Chem* 2003;278(6):3510–3513.
- Beisswenger C, Bals R. Antimicrobial peptides in lung inflammation. *Chemical immunology and allergy* 2005;86:55–71.

- Biagioli MC, Kaul P, Singh I, Turner RB. The role of oxidative stress in rhinovirus induced elaboration of IL-8 by respiratory epithelial cells. *Free Radic Biol Med* 1999;26(3-4):454-462.
- Caillou B, Dupuy C, Lacroix L, et al. Expression of reduced nicotinamide adenine dinucleotide phosphate oxidase (ThoX, LNOX, Duox) genes and proteins in human thyroid tissues. *J Clin Endocrinol Metab* 2001;86(7):3351-3358.
- Christophe-Hobertus C, Christophe D. Human Thyroid Oxidases genes promoter activity in thyrocytes does not appear to be functionally dependent on Thyroid Transcription Factor-1 or Pax8. *Mol Cell Endocrinol* 2007;264(1-2):157-163.
- Clark RA, Klebanoff SJ, Einstein AB, Fefer A. Peroxidase-H₂O₂-halide system: Cytotoxic effect on mammalian tumor cells. *Blood* 1975;45(2):161-170.
- Conner GE, Salathe M, Forteza R. Lactoperoxidase and hydrogen peroxide metabolism in the airway. *Am J Respir Crit Care Med* 2002;166(12 Pt 2):S57-S61.
- Conner GE, Wijkstrom-Frei C, Randell SH, Fernandez VE, Salathe M. The lactoperoxidase system links anion transport to host defense in cystic fibrosis. *FEBS Lett* 2007;581(2):271-278.
- De Deken X, Wang D, Many MC, et al. Cloning of two human thyroid cDNAs encoding new members of the NADPH oxidase family. *J Biol Chem* 2000;275(30):23227-23233.
- Degroot LJ, Niepomniszcze H, Nagasaka A, Hati R. Mechanism of thyroid hormone formation. *Annals of clinical research* 1972;4(3):113-120.
- Dupuy C, Deme D, Kaniewski J, Pommier J, Virion A. Ca²⁺ + regulation of thyroid NADPH-dependent H₂O₂ generation. *FEBS Lett* 1988;233(1):74-78.
- Dupuy C, Ohayon R, Valent A, Noel-Hudson MS, Deme D, Virion A. Purification of a novel flavoprotein involved in the thyroid NADPH oxidase. Cloning of the porcine and human cdnas. *J Biol Chem* 1999;274(52):37265-37269.
- Dusi S, Donini M, Rossi F. Mechanisms of NADPH oxidase activation in human neutrophils: p67phox is required for the translocation of rac 1 but not of rac 2 from cytosol to the membranes. *Biochem J* 1995;308 (Pt 3):991-994.
- Edens WA, Sharling L, Cheng G, et al. Tyrosine cross-linking of extracellular matrix is catalyzed by Duox, a multidomain oxidase/peroxidase with homology to the phagocyte oxidase subunit gp91phox. *J Cell Biol* 2001;154(4):879-891.
- El Hassani RA, Benfares N, Caillou B, et al. Dual oxidase2 is expressed all along the digestive tract. *Am J Physiol Gastrointest Liver Physiol* 2005;288(5):G933-G942.
- Finkel T. Oxidant signals and oxidative stress. *Curr Opin Cell Biol* 2003;15(2):247-254.
- Forteza R, Salathe M, Miot F, Forteza R, Conner GE. Regulated hydrogen peroxide production by Duox in human airway epithelial cells. *Am J Respir Cell Mol Biol* 2005;32(5):462-469.
- Gabig TG, Babior BM. The O₂(-)-forming oxidase responsible for the respiratory burst in human neutrophils. Properties of the solubilized enzyme. *J Biol Chem* 1979;254(18):9070-9074.
- Geiszt M, Leto TL. The Nox family of NAD(P)H oxidases: Host defense and beyond. *J Biol Chem* 2004.
- Geiszt M, Witta J, Baffi J, Lekstrom K, Leto TL. Dual oxidases represent novel hydrogen peroxide sources supporting mucosal surface host defense. *Faseb J* 2003;17(11):1502-1504.
- Gerson C, Sabater J, Scuri M, et al. The lactoperoxidase system functions in bacterial clearance of airways. *Am J Respir Cell Mol Biol* 2000;22(6):665-671.
- Goldman AS, Smith CW. Host resistance factors in human milk. *J Pediatr* 1973;82(6):1082-1090.
- Greenberg SB. Rhinovirus and coronavirus infections. *Seminars in respiratory and critical care medicine* 2007;28(2):182-192.
- Grubor B, Meyerholz DK, Ackermann MR. Collectins and cationic antimicrobial peptides of the respiratory epithelia. *Veterinary pathology* 2006;43(5):595-612.
- Ha EM, Oh CT, Bae YS, Lee WJ. A direct role for dual oxidase in *Drosophila* gut immunity. *Science* 2005;310(5749):847-850.
- Harper RW, Xu C, Eiserich JP, et al. Differential regulation of dual NADPH oxidases/peroxidases, Duox1 and Duox2, by Th1 and Th2 cytokines in respiratory tract epithelium. *FEBS Lett* 2005;579(21):4911-4917.

- Harper RW, Xu C, McManus M, Heidersbach A, Eiserich JP. Duox2 exhibits potent heme peroxidase activity in human respiratory tract epithelium. *FEBS Lett* 2006;580(22):5150–5154.
- Hayden FG. Rhinovirus and the lower respiratory tract. *Reviews in medical virology* 2004;14(1):17–31.
- Hohn DC, Lehrer RI. NADPH oxidase deficiency in X-linked chronic granulomatous disease. *J Clin Invest* 1975;55(4):707–713.
- Igo RP, Mahoney CP, Mackler B. Studies of the Biosynthesis of Thyroxine. I. Purification and Properties of a Particulate Iodide Peroxidase from Thyroid Tissue. *J Biol Chem* 1964;239:1893–1897.
- Kaul P, Biagioli MC, Singh I, Turner RB. Rhinovirus-induced oxidative stress and interleukin-8 elaboration involves p47-phox but is independent of attachment to intercellular adhesion molecule-1 and viral replication. *J Infect Dis* 2000;181(6):1885–1890.
- Lambeth JD. NOX enzymes and the biology of reactive oxygen. *Nat Rev Immunol* 2004;4(3):181–189.
- Lambeth JD. Nox/Duox family of nicotinamide adenine dinucleotide (phosphate) oxidases. *Curr Opin Hematol* 2002;9(1):11–17.
- Lambeth JD, Kawahara T, Diebold B. Regulation of Nox and Duox enzymatic activity and expression. *Free Radic Biol Med* 2007;43(3):319–331.
- Leseney AM, Deme D, Legue O, et al. Biochemical characterization of a Ca²⁺ + /NAD(P)H-dependent H₂O₂ generator in human thyroid tissue. *Biochimie* 1999;81(4):373–380.
- Leto TL, Geiszt M. Role of Nox family NADPH oxidases in host defense. *Antioxid Redox Signal* 2006;8(9–10):1549–1561.
- Lissitzky S. Biosynthesis of thyroid hormones. *Pharmacol Ther [B]* 1976;2(1):219–246.
- Moreno JC, Bikker H, Kempers MJ, et al. Inactivating mutations in the gene for thyroid oxidase 2 (THOX2) and congenital hypothyroidism. *N Engl J Med* 2002;347(2):95–102.
- Moskwa P, Lorentzen D, Excoffon KJ, et al. A novel host defense system of airways is defective in cystic fibrosis. *Am J Respir Crit Care Med* 2007;175(2):174–183.
- Nakamura Y, Ogihara S, Ohtaki S. Activation by ATP of calcium-dependent NADPH-oxidase generating hydrogen peroxide in thyroid plasma membranes. *J Biochem (Tokyo)* 1987;102(5):1121–1132.
- Nakanaga T, Nadel JA, Ueki IF, Koff JL, Shao MX. Regulation of interleukin-8 via an airway epithelial signaling cascade. *Am J Physiol Lung Cell Mol Physiol* 2007.
- Ng AW, Bidani A, Heming TA. Innate host defense of the lung: effects of lung-lining fluid pH. *Lung* 2004;182(5):297–317.
- Pachucki J, Wang D, Christophe D, Miot F. Structural and functional characterization of the two human ThOX/Duox genes and their 5'-flanking regions. *Mol Cell Endocrinol* 2004;214(1–2):53–62.
- Ratner AJ, Prince A. Lactoperoxidase. New recognition of an “old” enzyme in airway defenses. *Am J Respir Cell Mol Biol* 2000;22(6):642–644.
- Rigutto S, Hoste C, Dumont JE, Corvilain B, Miot F, De Deken X. Duox1 is the main source of hydrogen peroxide in the rat thyroid cell line PCC13. *Exp Cell Res* 2007.
- Salathe M, Guldimann P, Conner GE, Wanner A. Hydrogen peroxide-scavenging properties of sheep airway mucus. *Am J Respir Crit Care Med* 1995;151(5):1543–1550.
- Salathe M, Holderby M, Forteza R, Abraham WM, Wanner A, Conner GE. Isolation and characterization of a peroxidase from the airway. *Am J Respir Cell Mol Biol* 1997;17(1):97–105.
- Schwarzer C, Machen TE, Illek B, Fischer H. NADPH oxidase-dependent acid production in airway epithelial cells. *J Biol Chem* 2004.
- Shao MX, Nadel JA. Dual oxidase 1-dependent MUC5AC mucin expression in cultured human airway epithelial cells. *Proc Natl Acad Sci U S A* 2005;102(3):767–772.
- Song Y, Driessens N, Costa M, et al. Roles of hydrogen peroxide in thyroid physiology and disease. *J Clin Endocrinol Metab* 2007;92(10):3764–3773.
- Suh YA, Arnold RS, Lassegue B, et al. Cell transformation by the superoxide-generating oxidase Mox1. *Nature* 1999;401(6748):79–82.

- Terada LS. Specificity in reactive oxidant signaling: think globally, act locally. *J Cell Biol* 2006;174(5):615–623.
- Thacker EL. Lung inflammatory responses. *Veterinary research* 2006;37(3):469–486.
- Vigone MC, Fugazzola L, Zamproni I, et al. Persistent mild hypothyroidism associated with novel sequence variants of the DUOX2 gene in two siblings. *Human mutation* 2005;26(4):395.
- Wesley UV, Bove PF, Hristova M, McCarthy S, van der Vliet A. Airway epithelial cell migration and wound repair by ATP-mediated activation of dual oxidase 1. *J Biol Chem* 2007;282(5):3213–3220.
- Whitton JL, Cornell CT, Feuer R. Host and virus determinants of picornavirus pathogenesis and tropism. *Nat Rev Microbiol* 2005;3(10):765–776.
- Wijkstrom-Frei C, El-Chemaly S, Ali-Rachedi R, et al. Lactoperoxidase and human airway host defense. *Am J Respir Cell Mol Biol* 2003;29(2):206–212.
- Yankaskas JR, Haizlip JE, Conrad M, et al. Papilloma virus immortalized tracheal epithelial cells retain a well-differentiated phenotype. *Am J Physiol* 1993;264(5 Pt 1):C1219–C1230.
- Zhang P, Summer WR, Bagby GJ, Nelson S. Innate immunity and pulmonary host defense. *Immunol Rev* 2000;173:39–51.