

MITOGENOME ANNOUNCEMENT

 OPEN ACCESS 

## The complete mitochondrial genome of the New Zealand parasitic roundworm *Teladorsagia circumcincta* (Trichostrongyloidea: Haemonchidae) field strain NZ\_Teci\_NP

Nikola Palevich<sup>a</sup> , Paul H. Maclean<sup>a</sup>, Makedonka Mitreva<sup>b</sup>, Richard Scott<sup>a</sup> and David Leathwick<sup>a</sup>

<sup>a</sup>AgResearch Ltd., Grasslands Research Centre, Palmerston North, New Zealand; <sup>b</sup>McDonnell Genome Institute and Department of Medicine, Washington University School of Medicine, Saint Louis, Missouri, United States of America

### ABSTRACT

The complete mitochondrial genome of the New Zealand parasitic nematode *Teladorsagia circumcincta* field strain NZ\_Teci\_NP was sequenced and annotated. The 14,083 bp-long mitogenome contains 12 protein-coding genes (atp8 gene missing), two ribosomal RNAs (rRNAs), and 22 transfer RNAs (tRNAs). Phylogenetic analysis showed that *T. circumcincta* NZ\_Teci\_NP forms a monophyletic cluster with the remaining Haemonchidae species and further reinforces the high levels of diversity and gene flow observed among Trichostrongylidae.

### ARTICLE HISTORY

Received 26 May 2019  
Accepted 31 July 2019

### KEYWORDS


*Teladorsagia circumcincta*; roundworm; gastrointestinal; trichostrongyloid; nematode; phylogeny

The large and highly variable mitochondrial (mt) genomes of helminths (worms), including parasitic nematodes (roundworms), are ideal sources of molecular markers suitable for studying population genetic structures and evolution. *Teladorsagia circumcincta* NZ\_Teci\_NP was selected for genome sequencing as a representative of an anthelmintic-susceptible NZ field strain of *T. circumcincta*. The specimen was collected from the Palmerston North area ( $40^{\circ}21.3' S$ ,  $175^{\circ}36.7' E$ ) and is stored (accession number: NPX120886) and available upon request from AgResearch Ltd., Grasslands Research Centre. High molecular weight genomic DNA was isolated from multiple *T. circumcincta* adult males using a modified phenol:chloroform protocol (Palevich et al. 2017; Palevich, Kelly, et al. 2019). The Illumina MiSeq (Macrogen, Korea) platform was used to amplify the entire mitochondrial genome sequence (GenBank accession number: MN013406).

The mitogenome of NZ\_Teci\_NP (14,083 bp) is standard in size and comparable to the *T. circumcincta* (GQ888720) strain (Jex et al. 2010; Choi et al. 2017; Palevich, Maclean, et al. 2019). For example, all genes are transcribed in the same direction, there is a lack of the Atp8 gene, it contains 12 protein-coding genes (PCGs), two rRNAs, and 22 tRNAs. All 12 PCGs use standard ATN/TAN start/stop codons, respectively. The studied

genome has a high T content (46.1%) and a low C content (7.2%), resulting in a very strong A+T bias (77.3%). Gene order, sizes, and all common organization features are relatively conserved among the 43 nematode mitogenomes (usually 13.6–14.3 kb) (Jex et al. 2009; Palevich et al. 2018; Palevich, Maclean, Baten et al. 2019; Palevich, Maclean, et al. 2019).

The phylogenetic position of *T. circumcincta* was estimated using maximum-likelihood, implemented in RAxML version 8.2.11 (1000 bootstrap replications) (Stamatakis 2014), and the Bayesian inference (BI), implemented in MrBayes version 3.2.6 (default settings, four MCMC chains, 6,34–106 generations) (Huelsenbeck and Ronquist 2001) approaches. Mitogenome sequences of all 36 available nematode species were retrieved from GenBank. Analyses were performed both on the entire nucleotide sequences of the complete mitogenomes and using only the concatenated mitochondrial PCGs and rRNA genes, producing identical dendrogram topologies (Figure 1). *Teladorsagia circumcincta* formed a monophyletic cluster with the remaining Trichostrongylidae species, which then formed a sister clade with the Strongylidae family. Overall, the dendrogram topology is highly congruent with recent results (Palevich, Maclean, et al. 2019). In the pursuit of improving the phylogenetic resolution within the phylum Nematoda, future efforts should focus on the availability of



**Figure 1.** Phylogenetic analysis of the complete mt genomes for all 43 nematode species or isolates available in GenBank. The evolutionary relationships between the *T. circumcincta* field strain NZ\_Teci\_NP (highlighted by a black star) and the two major suborders of the Strongylida are represented by coloured boxes (Trichostrongylina (blue) and Strongylina (green)). Species representing the Rhabditida, Ascaridida, Aphelenchida, Tylenchida, Spirurida, and Trichocephalida have been included as outgroups. Phylogenetic analysis was conducted using maximum-likelihood and Bayesian inference (MrBayes). The numbers above the midpoint of each tree branch represent the statistical support for each node (based on posterior probability score). The phylogram provided is presented to scale (scale bar = 0.07 estimated number of substitutions per site) and GenBank accession numbers are provided (in parentheses) for all reference sequences. An identical topology was found with maximum-likelihood; all nodes were supported by >99% bootstrap re-sampling ( $n = 1000$ ).

more complete mitogenomes across all nematode species, and especially for different strains/isolates.

## Disclosure statement

No potential conflict of interest was reported by the authors.

## Funding

Research was supported and funded by the Agricultural and Marketing Research and Development Trust (AGMARDT) Postdoctoral Fellowships Programme [grant number P17001], New Zealand.

## ORCID

Nikola Palevich  <http://orcid.org/0000-0002-2101-2452>

## References

- Choi YJ, Bisset SA, Doyle SR, Hallsworth-Pepin K, Martin J, Grant WN, Mitreva M. 2017. Genomic introgression mapping of field-derived multiple-anthelmintic resistance in *Teladorsagia circumcincta*. PLoS Genet. 13:e1006857.
- Huelsenbeck JP, Ronquist F. 2001. MRBAYES: Bayesian inference of phylogenetic trees. Bioinformatics. 17:754–755.
- Jex AR, Hall RS, Littlewood DTJ, Gasser RB. 2010. An integrated pipeline for next-generation sequencing and annotation of mitochondrial genomes. Nucleic Acids Res. 38:522–533.
- Palevich N, Britton C, Kamenetzky L, Mitreva M, de Moraes Mourão M, Bennuru S, Quack T, Scholte LLS, Tyagi R, Slatko BE. 2018. Tackling hypotheticals in helminth genomes. Trends Parasitol. 34: 179–183.
- Palevich N, Maclean P, Baten A, Scott A, Leathwick D. 2019. The complete mitochondrial genome of the New Zealand parasitic roundworm *Haemonchus contortus* (Trichostrongyloidea: Haemonchidae) field strain NZ\_Hco\_NP. Mitochondrial DNA B. 2208.
- Palevich N, Kelly WJ, Ganesh S, Rakonjac J, Attwood GT. 2019. *Butyrivibrio hungatei* MB2003 Competes Effectively for Soluble Sugars Released by *Butyrivibrio proteoclasticus* B316<sup>T</sup> during Growth on Xylan or Pectin. Appl Environ Microbiol. 85:e02056–02018.
- Palevich N, Maclean PH, Baten A, Scott RW, Leathwick DM. 2019. The Genome Sequence of the Anthelmintic-Susceptible New Zealand *Haemonchus contortus*. Genome Biol Evol. 11:1965–1970. doi:10.1093/gbe/evz141
- Palevich N, Kelly WJ, Leahy SC, Altermann E, Rakonjac J, Attwood GT. 2017. The complete genome sequence of the rumen bacterium *Butyrivibrio hungatei* MB2003. Stand Genomic Sci. 12:72.
- Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. Bioinformatics. 30:1312–1313.