nature COMMUNICATIONS

https://doi.org/10.1038/s41467-019-10193-1 OPEN

Author Correction: Efficient electron transfer across hydrogen bond interfaces by proton-coupled and -uncoupled pathways

Tao Cheng[1], Dong Xue Shen[1], Miao Meng[1], Suman Mallick[1], Lijiu Cao[1], Nathan J. Patmore[2], Hong Li Zhang[1], Shan Feng Zou[1], Huo Wen Chen[1], Yi Qin[1], Yi Yang Wu[1] & Chun Y. Liu[1]

Correction to: *Nature Communications* https://doi.org/10.1038/s41467-019-09392-7, published online 4 April 2019.

The original version of this Article contained an error in the eighteenth sentence of the third paragraph of the 'Determination of H_{ab} and k_{ET} data for the Mo_2 dimers' section of the Results, which incorrectly read 'For $\mathbf{1a}^+$ with $2H_{ab} \gg k_BT$, it is inappropriate to calculate the rate constant using H_{ab} for the PCET reaction in the nonadiabatic regime.' The correct version states '$2H_{ab} >> k_BT$' in place of '$2H_{ab} \gg k_BT$'.

Also, the third sentence of the Discussion originally incorrectly read 'The linear relationship of $\ln(k_{ET})$ vs. r_{ab} gives $\beta = 1.25$ (Supplementary Fig. 9).' The correct version states 'R_{ab}' instead of 'r_{ab}'.

This has been corrected in both the PDF and HTML versions of the Article.

Published online: 02 May 2019

[1] Department of Chemistry, Jinan University, 601 Huang-Pu Avenue West, 510632 Guangzhou, China. [2] Department of Chemical Sciences, University of Huddersfield, Queensgate, Huddersfield HD1 3DH, UK. Correspondence and requests for materials should be addressed to C.Y.L. (email: tcyliu@jnu.edu.cn)