

Clinical Commentary/Current Concept Review**Forefoot Injuries in Athletes: Integration of the Movement System**Lindsay A Carroll¹, Stephen Paulseth², RobRoy L Martin^{1 a}¹ Physical Therapy, Duquesne University, ² Paulseth & Associates Physical Therapy, Inc

Keywords: taping, foot, joint mobilization, sports injuries, movement system

<https://doi.org/10.26603/001c.30021>

International Journal of Sports Physical TherapyVol. 17, Issue 1, 2022

Despite the prevalence of forefoot related problems in athletes, there are few comprehensive summaries on examination and intervention strategies for those with forefoot related symptoms. While many factors may contribute to pathology and injury, the presence of abnormal foot alignment can negatively affect lower extremity biomechanics and be associated with injuries. Physical therapists may use the characteristics associated abnormal pronation or abnormal supination to describe the movement system disorder and serve as a guide for evaluating and managing athletes with forefoot pathologies. Athletes with an abnormal pronation movement system diagnosis typically demonstrate foot hypermobility, have decreased strength of the tibialis posterior muscle, and present with a medially rotated lower extremity position. Athletes with abnormal supination movement system diagnosis typically demonstrate foot hypomobility, decreased strength of the fibularis muscles, and a laterally rotated lower extremity position. Interventions of manual therapy, taping, strengthening exercises, and neuromuscular reeducation can be directed at the identified impairments and abnormal movements. The purpose of this clinical commentary is to integrate a movement system approach in pathoanatomical, evaluation, and intervention considerations for athletes with common forefoot pathologies, including stress fractures, metatarsalgia, neuroma, turf toe, and sesamoiditis. By applying a prioritized, objective problem list and movement system diagnosis, emphasis is shifted from a pathoanatomical diagnosis-based treatment plan to a more impairment and movement focused treatment.

Level of Evidence

5

INTRODUCTION

It is well known that athletes are at risk for foot and ankle injuries. These include injuries in the forefoot, defined as the region of the foot distal to the tarsometatarsal joints. The forefoot, unlike the mid- and hindfoot, is unconstrained with movement occurring freely in all three planes. Because the forefoot is the most distal weight bearing segment of the lower extremity, it can undergo a substantial amount of stress and strain, and can be affected by footwear, terrain, and biomechanical factors in the entire lower kinetic chain. Despite the prevalence of forefoot re-

lated problems in athletes, there are few comprehensive summaries on examination and intervention strategies for those with forefoot related symptoms. The purpose of the clinical commentary is to integrate a movement system approach in pathoanatomical, evaluation, and intervention considerations for athletes with common forefoot pathologies, including stress fractures, metatarsalgia, neuroma, and sesamoiditis.

ANATOMY OF THE FOREFOOT

The forefoot is composed of five rays that are functionally

a Corresponding Author:

RobRoy L. Martin, PT, PhD, CSCS
Duquesne University
Department of Physical Therapy
111A RSHS
Pittsburgh, PA
15282
martinr280@duq.edu
Phone: 412-396-1811
Fax: 412-396-4399

divided into a medial component, including the first metatarsal and great toe (hallux), and the lateral component, consisting of metatarsals and toes two to five. The distal aspect of the medial longitudinal arch is formed by the first metatarsal. The first tarsometatarsal and Lisfranc articulations join the midfoot to the forefoot, with these joints being supported by a dense interconnection of dorsal and plantar tarsometatarsal, intermetatarsal, and Lisfranc ligaments. Distally, the hallux is joined with the first metatarsal by the first metatarsophalangeal (MTP) joint and is supported by the joint capsule and plantar, medial collateral, and lateral collateral ligaments. The five rays of the forefoot are supported by a tensed interconnection of ligaments, joint capsules, and fascia that create a transverse arch.¹ This interconnected weave of tissue includes the plantar plate, which is a fibrocartilaginous structure that runs from each metatarsal head to the respective proximal phalanx. The plantar plate also serves as an attachment for the plantar fascia and supports the transverse arch.¹

While it is recognized that abnormal function of the medial longitudinal arch can affect lower extremity biomechanics and contribute to pathology, abnormalities of the transverse arch and forefoot may also affect lower extremity biomechanics and contribute to pathology. Robberecht et al.¹ found collapse of the transverse arch to be associated with forefoot pathology. During the propulsive phase of gait, representing the last 30% of the stance phase, only the forefoot is in contact with the ground. Consequently, abnormal forefoot biomechanics may negatively affect the entire lower extremity during propulsion. Likewise, any abnormal biomechanics in the lower extremity can affect propulsion and contribute to forefoot pathology. A comprehensive examination and intervention plan for forefoot pathologies therefore needs to consider the entire lower extremity and how biomechanical abnormalities may affect movement and thus contribute to symptoms.

BIOMECHANICAL CONSIDERATIONS

While many factors may contribute to pathology and injury, the presence of abnormal foot alignment can negatively affect lower extremity biomechanics and be associated with injuries.²⁻⁸ Abnormal pronation is typically defined by excessive calcaneal eversion, plantarflexion and adduction of the talus, collapse of the medial longitudinal arch, and abduction of the forefoot on the hindfoot. Abnormal pronation has been associated with increased foot mobility, collapse of the transverse arch, and compensatory knee and hip medial rotation.^{9,10} Abnormal supination is typically defined by excessive calcaneal inversion, dorsiflexion and abduction of the talus, high medial longitudinal arch, and adduction of the forefoot on the hindfoot. This foot type is usually more rigid and may be associated with compensatory knee and hip lateral rotation.^{9,10} Altered movement patterns caused by abnormal pronation and supination may be identified during static standing, gait, and functional movement testing. The single leg squat and step-down tests are functional movement tests that can be used to assess neuromuscular control and identify potential impairments of the trunk, pelvis, hip, knee, and ankle with evidence of reliability and validity to support its use.^{11,12} Because the

step-down test may place a greater emphasis on ankle motion, it may be a better measure than the single leg squat test in those with foot and ankle pathologies.¹³ Compensatory lower extremity movements can be identified and characterized as being associated with abnormal pronation or supination during gait and functional movement assessment. Physical therapists may use the general characteristics associated with abnormal pronation or abnormal supination to describe the movement system disorder and serve as a guide for evaluating and managing athletes with common forefoot pathologies such as stress fractures, metatarsalgia, neuroma, and sesamoiditis.

FOREFOOT PATHOLOGIES

STRESS FRACTURE

Stress fractures are microscopic bone injuries resulting from repeated bouts of physiological overload without adequate time for tissue remodeling and adaptation.^{14,15} Athletes who have a sudden increase in weight bearing activities are at risk for a stress fracture, with runners and military recruits seemingly being at higher risk.¹⁶⁻²⁰ The shafts of the metatarsal bones are common locations for stress fractures, with the occurrence at the second and third metatarsals being more common than at the fourth and fifth.^{15-17,19} Athletes with a movement system diagnosis of abnormal supination may be at risk for metatarsal stress fractures because of the reduced ability to attenuate weight bearing stressors associated with a more rigid foot. An abnormal pronation movement system diagnosis can also increase the risk of sustaining a stress fracture because of atypical loading pattern associated with a more mobile foot.^{15,21,22} Increased risk for stress fracture has been associated with poor pre-participation condition, older age, female sex, Caucasian race, decreased bone density, hormonal and menstrual abnormalities, low calorie and low fat diet, inadequate sleep pattern, and collagen disease.²² Athletes with a stress fracture may complain of an insidious onset of chronic aching pain that is activity related and associated with an increase in weight bearing activity or training intensity.^{22,23} Examination should find the involved metatarsal shaft to be tender with palpation.²³ Stress fractures of the second and third metatarsals generally heal well requiring only activity modification without a reduction in weightbearing.¹⁴ Stress fractures of the proximal intermediate zone of the fifth metatarsal are considered high risk for delayed healing or non-union and require more restrictive weight bearing, partial immobilization, and may progress to surgery if healing does not occur.¹⁴ Imaging such as radiographs or MRI may be necessary to identify and grade stress fractures. Higher grade stress fractures may require 16 or more weeks of activity modification while lower grade stress fractures may improve with just three weeks of relative rest.²⁴ Treatment for stress fractures should include modifications of factors that contributed to the injury.^{14,22,25} A comprehensive lower quarter biomechanical examination and a sport-specific movement analysis will help identify and guide treatment to address contributing factors such as leg length discrepancy, abnormal foot posture, lower extremity malalignment, muscle imbalance,

ance, flexibility insufficiency, and range of motion (ROM) deficits. Athlete education should address any training errors, improper diet, or inadequate sleep patterns that are identified. Relative rest with low/non-impact aerobic activity, stretching and strengthening exercises, and immobilization in a removable boot are generally recommended until the pain resolves.²² Training can resume with a 10% increase in intensity per week after the patient has been pain free for 10–14 days.²⁶

METATARSALGIA

Metatarsalgia is a non-specific diagnosis given to athletes with pain on the plantar aspect of one or more of the metatarsal heads that is exacerbated by physical activity, barefoot walking, and/or walking in shoes with an elevated heel. This condition typically results from repetitive overloading of the metatarsal head(s) due to anatomic or biomechanical abnormalities such as first ray hypermobility, hallux abducto valgus (HAV), ankle equinus, claw or hammer toe deformities, lesser MTP joint instability, atrophy of the plantar fat pad, and/or improper footwear. A movement system diagnosis of abnormal pronation or supination may also contribute to overload the development of metatarsalgia because of altered loading of the metatarsal head. Athletes, particularly middle-aged females, may note a gradual onset of pain related to a rapid increase in training intensity, inappropriate shoe wear, or a change in running terrain. Examination should identify local tenderness at the metatarsal head and possibly a prominent metatarsal head(s). Muscle imbalance, ROM deficits, and/or biomechanical abnormalities in the lower quarter that may contribute to overloading the metatarsal heads should be identified and corrected. Assessing for and addressing any loss of ankle dorsiflexion ROM should be a primary focus. Treatment can also include orthoses, a metatarsal pad, and shoe modifications which may promote redistribution of plantar pressures and reduce pain. Taping to redistribute the plantar fat pad with or without techniques to correct hammer or claw toe deformity, when appropriate, may be beneficial (Figure 1A-B).

NEUROMA

An interdigital neuroma (Morton's neuroma) is a mechanical entrapment neuropathy of one or more of the interdigital nerves in the forefoot. The nerve may become enlarged because of fibrotic tissue and/or endoneurial edema. This condition primarily involves the third common (64%–91%) digital branch of the medial plantar nerve between the third and fourth metatarsal heads, followed by the second (18%–31%), first (0%–2.5%), and fourth (0%–6%) interdigital nerves.^{27–29} Runners and dancers are especially susceptible to interdigital neuroma due to repetitive hyperextension and longitudinal metatarsal torsional trauma at the MTP joints and resultant tissue thickening and swelling which may compress the nerve.³⁰ Narrow shoes, over-training with repetitive MTP extension are the primary risk factors for developing an interdigital neuroma.³¹ A movement system diagnosis of abnormal pronation may also contribute to the development of a neuroma. Athletes with ab-



Figure 1. Fat pad repositioning and correction for metatarsophalangeal joint extension.

A. Fat pad taping for distal displacement of fat pad. Manually reposition the fat pad to be better positioned beneath the metatarsal head. Apply two 0.75" wide strips of leukotape from distal to proximal to reposition the fat pad under the metatarsal head.

B. Fat pad repositioning with correction for metatarsophalangeal joint (MTP) extension as seen in claw or hammer toe. Flex the involved MTP joint (second toe is involved in photo below). Apply 0.75" wide strips of leukotape from the dorsal aspect of the first phalanx to the proximal aspect of the plantar surface of the foot. Tape should be crossed at the plantar aspect of the forefoot.

normal pronation and a hypermobile foot may be a higher risk because of the narrower intermetatarsal space associated with a collapsed transverse arch. Signs and symptoms of interdigital neuroma typically begin insidiously and include neurogenic pain in the plantar aspect of the forefoot.³² The pain may be associated with tenderness, cramping, burning, tingling, and/or numbness in the toes of the involved interspace. Some athletes will report a sensation of walking on a lump.^{27,33–35} During the examination, manual compression of the transverse arch and Mulder test should reproduce symptoms in athletes with interdigital neuroma.³⁶ In athletes with chronic interdigital neuroma, weakness of the intrinsic muscles may be present. Interventions such as shoe modifications, such as custom orthotics, rocker-bottom shoes, the use of a wide toe box, and metatarsal head unloading with a metatarsal pad may also be helpful at decreasing symptoms.^{27,30,33,37,38} Metatarsal mobilization (Figure 2) and taping to correct abnormal pronation and promote the transverse arch for patients with interdigital neuroma.

SESAMOIDITIS

Hallux sesamoid syndrome, or sesamoiditis, are nonspecific descriptive terms referring to pathologies, anatomical anomalies, or adaptive changes of the sesamoid bones. These injuries are associated with inflammation of the peri-

tendinous structures of the sesamoids and possible osteochondritis.³⁹⁻⁴¹ Most sesamoid injuries are overuse injuries, but direct trauma or forced extension of the hallux can cause an acute injury. Overuse of the sesamoids and the supporting structures can occur with repetitive activities such as running, jumping, tennis, and ballet. Those with an abnormal supination movement system diagnosis may be at risk to overload the sesamoids because of the associated high arch and plantar flexed first ray. Symptoms of sesamoiditis include pain that occurs with weight bearing, direct palpation, or with passive extension of the first MTP joint. Forefoot swelling,⁴⁰ tenderness, crepitus, decreased strength of the flexor hallucis longus and brevis tendons,^{41,42} decreased extension of the first MTP joint, and impaired first ray and/or first MTP joint mobility may also be present.^{39,43,44} Decreased sesamoid mobility or abnormal position of the sesamoids may also be determined during palpation by comparing sesamoid position between the involved and uninvolved sides.^{41,42} Interventions for sesamoiditis should focus on unloading the sesamoids and forefoot or protecting the first MTP joint.⁴⁵ Orthotics can be used to decrease the load on the involved sesamoid and foot and may include a cut-out for the sesamoids, metatarsal bars, a rigid shank, and/or a first metatarsal extension.³⁹ Taping of the sesamoids can help improve forefoot position and function and may decrease shear forces on the sesamoids and plantar aspect of the forefoot.⁴² If malposition or decreased sesamoid mobility is found, corrective sesamoid mobilizations and/or taping can be implemented (Figure 3).

EVALUATION AND TREATMENT TECHNIQUES

A comprehensive examination of an athlete with forefoot pathology should include a comprehensive assessment of the foot and ankle as well as static standing, gait, and functional movement evaluations. An appropriate lower quarter screen may also be needed to identify potential contributing impairments. A standard examination can consist of range of motion and strength assessment of the lumbosacral spine, hip, knee, ankle, and foot, with select special tests being used based on the athlete's history and potential differential diagnoses. Specific attention should be directed toward ankle dorsiflexion ROM, assessing for potential limitations in gastrocnemius-soleus flexibility, and talocrural joint posterior capsule mobility. The weight bearing lunge test can be used as a functional measure of tibiopedal dorsiflexion, with ROM coming from not only the talocrural joint but also the subtalar and midtarsal joints as well.⁴⁶ An assessment of great toe extension ROM in weightbearing and non-weightbearing should include evaluating mobility of the first ray tarsal-metatarsal joint, first MTP, and sesamoids. Likewise, assessment of hindfoot and forefoot ROM should include evaluating subtalar, calcaneocuboid, and talonavicular joint mobility. Because foot alignment is commonly associated with forefoot pathology,²⁻⁸ the Foot Posture Index-6 (FPI-6) can be used to assess static weight bearing alignment in the sagittal, frontal, and transverse planes and classify foot type as being normal, abnormally pronated, or abnormally supinated (Table 1).⁴⁷ A total score of 0 to +4 on the FPI-6 indicates a normal foot posture



Figure 2. Metatarsal mobilization.

Perform plantar-dorsal and dorsal-plantar glides on adjacent metatarsals to increase space and compression on the neuroma



Figure 3. Sesamoid taping to correct for laterally displaced sesamoids.

Manually reposition the laterally displaced sesamoids. Apply 0.75" leukotape from the dorsomedial aspect of the forefoot to the plantar aspect of the forefoot to stabilize.

in adults.⁸ Gait assessment, single leg squat test, and the step-down test can be used to identify abnormalities in the movement system. The findings from this comprehensive

Table 1. Summary Table for Abnormal pronation and supination disorders

Movement Disorder	Characteristics	Interventions	Forefoot Pathologies
Abnormal Pronation	FPI-6 score > +4 Medially rotated lower extremity position Decreased strength of the tibialis posterior muscle	Anti-pronation taping Single leg squat with proximally resisted hip lateral rotation Grade V mobilization to the navicular to facilitate tibialis posterior function	Stress fracture Metatarsalgia Neuroma
Abnormal Supination	FPI-6 score < 0 Laterally rotated lower extremity position Decreased strength of the fibularis muscles	Joint mobilizations to improve foot mobility; emphasize lateral subtalar glide Outward pivot exercises	Stress fracture Metatarsalgia Sesamoiditis

FPI: Foot Posture Index

examination of the entire lower quarter will identify impairments and generate a prioritized, objective problem list that can be used to develop an intervention plan within the context of the specific forefoot pathoanatomical diagnosis. The results of the FPI-6 and movement examination can identify a movement system diagnosis of abnormal pronation or supination to assist in directing intervention strategies.

Forefoot pathologies can be difficult to diagnose and often present with common impairments and movement system disorders. Using a prioritized, objective problem list and movement system diagnosis will place less emphasis on a pathoanatomical diagnosis-based treatment plan and more emphasis on the identified impairments and abnormal movements. Athletes with an abnormal pronation movement system diagnosis typically demonstrate foot hypermobility, have decreased strength of the tibialis posterior muscle, and present with a medially rotated position of the lower extremity. Treatment for those with abnormal pronation can include a grade V mobilization to the navicular to facilitate tibialis posterior function (Figure 4A-C) and anti-pronation taping to support the medial longitudinal arch (Figure 5A-D). Neuromuscular reeducation and strengthening exercises can be directed at the intrinsic and extrinsic foot muscles that support medial longitudinal and transverse arches. These exercises can also work to correct the medially rotated lower extremity and stabilize the hip and lumbosacral spine. The single leg squat with proximally resisted hip lateral rotation (Figure 7) can be used to facilitate the tibialis posterior, hip lateral rotators, hip abductors, and lumbosacral spine stabilizers. Athletes with abnormal supination movement system diagnosis typically demonstrate foot hypomobility, decreased strength of the fibularis muscles, and a laterally rotated lower extremity position. Treatment for those with abnormal supination can include joint mobilizations to improve foot mobility, with an emphasis on improving lateral subtalar glide (Figure 8). Exercises to facilitate fibularis activity and foot pronation while engaging the trunk and hip musculature can include the outward pivot (Figure 9A-B). The characteristics, select treatment techniques, and forefoot pathologies associated with abnormal pronation and supination movement system disorders summarized in the Table. Joint mobilization, taping technique, and exercise should be appropriately se-

**Figure 4. Navicular whip joint mobilization**

- Hand position for navicular whip. Place thumbs over the plantar aspect of the navicular.
- Starting position. Apply a dorsally directed force with the thumbs and begin moving the ankle into plantarflexion.
- Ending position. As the ankle nears end-range plantarflexion, apply a grade V force to the navicular in a plantar to dorsal direction.

lected based the athlete's impairment and movement system diagnosis while considering their unique treatment goals and desired outcome.

CONCLUSION

Forefoot injuries are common in athletes because of the stress and strain that occur during competition and training. Many biomechanical factors can contribute to forefoot symptoms and therefore a thorough examination, that includes a functional movement assessment, should be per-

formed in order to identify contributing factors throughout the entire lower quarter. An evaluation should include a comprehensive history with a description of training and competition regimen, as well as a systematic examination for the entire quarter to identify impairments and generate a prioritized objective problem list. Using a movement system diagnosis of abnormal pronation or supination may also help in directing treatment to correct the associated abnormal movements.

CONFLICTS OF INTEREST

The authors have no conflicts to disclose.

DISCLOSURES

This study received no funding

Submitted: July 02, 2021 CST, Accepted: September 04, 2021
CST



Figure 5. Anti-pronation taping

A. Step 1: Plantarflex the first ray and apply a strip of athletic tape from the dorsal aspect of the distal first metatarsal, around the plantar aspect of the foot, to the dorsal-lateral forefoot. Apply a second strip of tape from the dorsal-medial aspect of the great toe, around the calcaneus, to the lateral aspect of the forefoot.

B. Step 2, medial view: Starting at the distal strip that was applied in figure 6B, wrap strips of 1" athletic tape around the sole of the foot, beginning at the dorsal-lateral foot and ending at the dorsomedial foot. Lift the foot into a supinated position as you apply each strip. Continue applying supination strips until about half of the heel is covered with tape.

C. Step 2, superolateral view: Do not overlap the supination strips on the superior aspect of the foot.

D. Step 3: With the patient in standing, apply strips of tape to the superior aspect of the foot to connect the supination strips applied in the previous step.



Figure 6. Single leg squat with proximal resistance for hip lateral rotators.

A. Starting position: The patient is in single leg stance on the affected extremity, holding a resistance band in the contralateral upper extremity. The patient should stabilize to maintain a neutral hip and pelvis position with elevated medial longitudinal arch throughout the exercise.

B. The patient begins the exercise by performing a row with the contralateral upper extremity so that the hip lateral rotators are engaged.

C. The patient maintains the row position from 7B and performs a single leg squat while not allowing trunk leaning or rotation, pelvis rotation or tilting, medial rotation or adduction of the hip, valgus at the knee, or loss of balance.



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-NC-4.0). View this license's legal deed at <https://creativecommons.org/licenses/by-nc/4.0> and legal code at <https://creativecommons.org/licenses/by-nc/4.0/legalcode> for more information.

REFERENCES

1. Robberecht J, Decroocq L, Schramm M, Gabay A, Maestro M. Transverse laxity of the forefoot. *Foot Ankle Surg*. Published online 2021. [doi:10.1016/j.fas.2021.01.006](https://doi.org/10.1016/j.fas.2021.01.006)
2. Tong JW, Kong PW. Association between foot type and lower extremity injuries: systematic literature review with meta-analysis. *J Orthop Sports Phys Ther*. 2013;43(10):700-714. [doi:10.2519/jospt.2013.4225](https://doi.org/10.2519/jospt.2013.4225)
3. Perez-Morcillo A, Gomez-Bernal A, Gil-Guillen VF, et al. Association between the Foot Posture Index and running related injuries: A case-control study. *Clin Biomech (Bristol, Avon)*. 2019;61:217-221. [doi:10.1016/j.clinbiomech.2018.12.019](https://doi.org/10.1016/j.clinbiomech.2018.12.019)
4. Menz HB, Dufour AB, Riskowski JL, Hillstrom HJ, Hannan MT. Association of planus foot posture and pronated foot function with foot pain: the Framingham foot study. *Arthritis Care Res (Hoboken)*. 2013;65(12):1991-1999. [doi:10.1002/acr.22079](https://doi.org/10.1002/acr.22079)
5. Neal BS, Griffiths IB, Dowling GJ, et al. Foot posture as a risk factor for lower limb overuse injury: a systematic review and meta-analysis. *J Foot Ankle Res*. 2014;7(1):55. [doi:10.1186/s13047-014-0055-4](https://doi.org/10.1186/s13047-014-0055-4)
6. Cain LE, Nicholson LL, Adams RD, Burns J. Foot morphology and foot/ankle injury in indoor football. *J Sci Med Sport*. 2007;10(5):311-319. [doi:10.1016/j.jsams.2006.07.012](https://doi.org/10.1016/j.jsams.2006.07.012)
7. Burns J, Keenan AM, Redmond A. Foot type and overuse injury in triathletes. *J Am Podiatr Med Assoc*. 2005;95(3):235-241. [doi:10.7547/0950235](https://doi.org/10.7547/0950235)
8. Redmond AC, Crane YZ, Menz HB. Normative values for the Foot Posture Index. *J Foot Ankle Res*. 2008;1(1):6. [doi:10.1186/1757-1146-1-6](https://doi.org/10.1186/1757-1146-1-6)
9. Cornwall MW, McPoil TG. Relationship between static foot posture and foot mobility. *J Foot Ankle Res*. 2011;4:4. [doi:10.1186/1757-1146-4-4](https://doi.org/10.1186/1757-1146-4-4)
10. Riegger-Krugh C, Keysor JJ. Skeletal malalignments of the lower quarter: correlated and compensatory motions and postures. *J Orthop Sports Phys Ther*. 1996;23(2):164-170. [doi:10.2519/jospt.1996.23.2.164](https://doi.org/10.2519/jospt.1996.23.2.164)
11. McGovern RP, Martin RL, Christoforetti JJ, Kivlan BR. Evidence-based procedures for performing the single leg squat and step-down tests in evaluation of non-arthritic hip pain: a literature review. *Int J Sports Phys Ther*. 2018;13(3):526-536.
12. Ressman J, Grooten WJA, Rasmussen Barr E. Visual assessment of movement quality in the single leg squat test: a review and meta-analysis of inter-rater and intrarater reliability. *BMJ Open Sport Exerc Med*. 2019;5(1):e000541. [doi:10.1136/bmjsem-2019-000541](https://doi.org/10.1136/bmjsem-2019-000541)
13. Carroll LA, Kivlan BR, Martin RL, Phelps AL, Garcia CR. The single leg squat test: a “top-down” or “bottom-up” functional performance test? *Int J Sports Phys Ther*. Published online 2021. [doi:10.26603/001c.21317](https://doi.org/10.26603/001c.21317)
14. Mandell JC, Khurana B, Smith SE. Stress fractures of the foot and ankle, part 1: biomechanics of bone and principles of imaging and treatment. *Skeletal Radiol*. 2017;46(8):1021-1029. [doi:10.1007/s00256-017-2640-7](https://doi.org/10.1007/s00256-017-2640-7)
15. Asano LY, Duarte A Jr, Silva AP, Brazilian Medical Association. Stress fractures in the foot and ankle of athletes. *Rev Assoc Med Bras*. 2014;60(6):512-517. [doi:10.1590/1806-9282.60.06.006](https://doi.org/10.1590/1806-9282.60.06.006)
16. Matheson GO, Clement DB, McKenzie DC, Taunton JE, Lloyd-Smith DR, MacIntyre JG. Stress fractures in athletes. A study of 320 cases. *Am J Sports Med*. 1987;15(1):46-58. [doi:10.1177/036354658701501017](https://doi.org/10.1177/036354658701501017)
17. Barrack MT, Gibbs JC, De Souza MJ, et al. Higher incidence of bone stress injuries with increasing female athlete triad-related risk factors: a prospective multisite study of exercising girls and women. *Am J Sports Med*. 2014;42(4):949-958. [doi:10.1177/0363546513520295](https://doi.org/10.1177/0363546513520295)
18. Zadpoor AA, Nikooyan AA. The relationship between lower-extremity stress fractures and the ground reaction force: a systematic review. *Clin Biomech (Bristol, Avon)*. 2011;26(1):23-28. [doi:10.1016/j.clinbiomech.2010.08.005](https://doi.org/10.1016/j.clinbiomech.2010.08.005)
19. Rizzone KH, Ackerman KE, Roos KG, Dompier TP, Kerr ZY. The Epidemiology of Stress Fractures in Collegiate Student-Athletes, 2004-2005 Through 2013-2014 Academic Years. *J Athl Train*. 2017;52(10):966-975. [doi:10.4085/1062-6050-52.8.01](https://doi.org/10.4085/1062-6050-52.8.01)
20. Bhatnagar A, Kumar M, Shivanna D, Bahubali A, Manjunath D. High Incidence of Stress Fractures in Military Cadets During Training: A Point of Concern. *J Clin Diagn Res*. 2015;9(8):RC01-3. [doi:10.7860/JCDR/2015/12535.6282](https://doi.org/10.7860/JCDR/2015/12535.6282)

21. Raikin SM, Slenker N, Ratigan B. The association of a varus hindfoot and fracture of the fifth metatarsal metaphyseal-diaphyseal junction: the Jones fracture. *Am J Sports Med.* 2008;36(7):1367-1372. doi:10.1177/0363546508314401
22. Asano LY, Duarte A Jr, Silva AP, Brazilian Medical Association. Stress fractures in the foot and ankle of athletes. *Rev Assoc Med Bras (1992).* 2014;60(6):512-517. doi:10.1590/1806-9282.60.06.006
23. Shindle MK, Endo Y, Warren RF, et al. Stress fractures about the tibia, foot, and ankle. *J Am Acad Orthop Surg.* 2012;20(3):167-176. doi:10.5435/JAAOS-20-03-167
24. Astur DC, Zanatta F, Arliani GG, Moraes ER, Pochini Ade C, Ejnisman B. Stress fractures: definition, diagnosis and treatment. *Rev Bras Ortop.* 2016;51(1):3-10. doi:10.1016/j.rboe.2015.12.008
25. Pohl MB, Mullineaux DR, Milner CE, Hamill J, Davis IS. Biomechanical predictors of retrospective tibial stress fractures in runners. *J Biomech.* 2008;41(6):1160-1165. doi:10.1016/j.jbiomech.2008.02.001
26. Raasch WG, Hergan DJ. Treatment of stress fractures: the fundamentals. *Clin Sports Med.* 2006;25(1):29-36, vii. doi:10.1016/j.csm.2005.08.013
27. Hodes A, Umans H. Metatarsalgia. *Radiol Clin North Am.* 2018;56(6):877-892. doi:10.1016/j.rcl.2018.06.004
28. Matthews BG, Hurn SE, Harding MP, Henry RA, Ware RS. The effectiveness of non-surgical interventions for common plantar digital compressive neuropathy (Morton's neuroma): a systematic review and meta-analysis. *J Foot Ankle Res.* 2019;12:12. doi:10.1186/s13047-019-0320-7
29. de Oliveira HAV, Natour J, Vassalli M, Rosenfeld A, Jennings F, Jones A. Effectiveness of customized insoles in patients with Morton's neuroma: a randomized, controlled, double-blind clinical trial. *Clin Rehabil.* 2019;33(12):1898-1907.
30. Ferkel E, Davis WH, Ellington JK. Entrapment Neuropathies of the Foot and Ankle. *Clin Sports Med.* 2015;34(4):791-801. doi:10.1016/j.csm.2015.06.002
31. Bencardino J, Rosenberg ZS, Beltran J, Liu X, Marty-Delfaut E. Morton's neuroma: is it always symptomatic? *AJR Am J Roentgenol.* 2000;175(3):649-653.
32. Gougoulas N, Lampridis V, Sakellariou A. Morton's interdigital neuroma: instructional review. *EFORT open reviews.* 2019;4(1):14-24. doi:10.1302/2058-5241.4.180025
33. Pomeroy G, Wilton J, Anthony S. Entrapment neuropathy about the foot and ankle: an update. *J Am Acad Orthop Surg.* 2015;23(1):58-66. doi:10.5435/jaas-23-01-58
34. Sault JD, Morris MV, Jayaseelan DJ, Emerson-Kavchak AJ. Manual therapy in the management of a patient with a symptomatic Morton's Neuroma: A case report. *Man Ther.* 2016;21:307-310.
35. Spina R, Cameron M, Alexander R. The effect of functional fascial taping on Morton's neuroma. *Australas Chiropr Osteopathy.* 2002;10(1):45-50.
36. Charen DA, Markowitz JS, Cheung ZB, Matijakovich DJ, Chan JJ, Vulcano E. Overview of Metatarsalgia. *Orthopedics.* 2019;42(1):e138-e143. doi:10.3928/01477447-20181206-06
37. Davis F. Therapeutic Massage Provides Pain Relief to a Client with Morton's Neuroma: A Case Report. *Int J Ther Massage Bodywork.* 2012;5(2):12-19.
38. Deniz S, Purtuloglu T, Tekindur S, et al. Ultrasound-guided pulsed radio frequency treatment in Morton's neuroma. *J Am Podiatr Med Assoc.* 2015;105(4):302-306. doi:10.7547/13-128.1
39. Omev ML, Micheli LJ. Foot and ankle problems in the young athlete. *Med Sci Sports Exerc.* 1999;31(7 Suppl):S470-86. doi:10.1097/00005768-199907001-00008
40. Hockenbury RT. Forefoot problems in athletes. *Med Sci Sports Exerc.* 1999;31(7 Suppl):S448-58. doi:10.1097/00005768-199907001-00006
41. Sanders TG, Rathur SK. Imaging of painful conditions of the hallux sesamoid complex and plantar capsular structures of the first metatarsophalangeal joint. *Radiol Clin North Am.* 2008;46(6):1079-1092, vii. doi:10.1016/j.rcl.2008.09.01
42. Anwar R, Anjum SN, Nicholl JE. Sesamoids of the foot. *Curr Orthop.* 2005;19(1):40-48. doi:10.1016/j.cuor.2005.01.001
43. Grebing BR, Coughlin MJ. The effect of ankle position on the exam for first ray mobility. *Foot Ankle Int.* 2004;25(7):467-475. doi:10.1177/10711007040250705
44. Menz HB, Zammit GV, Munteanu SE, Scott G. Plantarflexion strength of the toes: age and gender differences and evaluation of a clinical screening test. *Foot Ankle Int.* 2006;27(12):1103-1108. doi:10.1177/107110070602701217

45. York PJ, Wydra FB, Hunt KJ. Injuries to the great toe. *Curr Rev Musculoskelet Med*. 2017;10(1):104-112. [doi:10.1007/s12178-017-9390-y](https://doi.org/10.1007/s12178-017-9390-y)

46. Smith MD, Lee D, Russell T, Matthews M, MacDonald D, Vicenzino B. How Much Does the Talocrural Joint Contribute to Ankle Dorsiflexion Range of Motion During the Weight-Bearing Lunge Test? A Cross-sectional Radiographic Validity Study. *J Orthop Sports Phys Ther*. 2019;49(12):934-941. [doi:10.2519/jospt.2019.8697](https://doi.org/10.2519/jospt.2019.8697)

47. Redmond AC, Crosbie J, Ouvrier RA. Development and validation of a novel rating system for scoring standing foot posture: the Foot Posture Index. *Clin Biomech (Bristol, Avon)*. 2006;21(1):89-98. [doi:10.1016/j.clinbiomech.2005.08.002](https://doi.org/10.1016/j.clinbiomech.2005.08.002)