

SUMO Proteins are not Involved in TGF- β 1-induced, Smad3/4-mediated Germline α Transcription, but PIASy Suppresses it in CH12F3-2A B Cells

Sang-Hoon Lee^{1#}, Pyeung-Hyeun Kim^{2#}, Sang-Muk Oh³, Jung-Hwan Park³, Yung-Choon Yoo¹, Junglim Lee¹ and Seok-Rae Park^{1*}

¹Department of Microbiology, College of Medicine, Konyang University, Daejeon 302-718, ²Department of Molecular Bioscience, School of Biomedical Science, Kangwon National University, Chuncheon 200-701, ³Department of Biochemistry, College of Medicine, Konyang University, Daejeon 302-718, Korea

TGF- β induces IgA class switching by B cells. We previously reported that Smad3 and Smad4, pivotal TGF- β signal-transducing transcription factors, mediate germline (GL) α transcription induced by TGF- β 1, resulting in IgA switching by mouse B cells. Post-translational sumoylation of Smad3 and Smad4 regulates TGF- β -induced transcriptional activation in certain cell types. In the present study, we investigated the effect of sumoylation on TGF- β 1-induced, Smad3/4-mediated GL α transcription and IgA switching by mouse B cell line, CH12F3-2A. Overexpression of small ubiquitin-like modifier (SUMO)-1, SUMO-2 or SUMO-3 did not affect TGF- β 1-induced, Smad3/4-mediated GL α promoter activity, expression of endogenous GL α transcripts, surface IgA expression, and IgA production. Next, we tested the effect of the E3 ligase PIASy on TGF- β 1-induced, Smad3/4-mediated GL α promoter activity. We found that PIASy overexpression suppresses the GL α promoter activity in cooperation with histone deacetylase 1. Taken together, these results suggest that SUMO itself does not affect regulation of GL α transcription and IgA switching induced by TGF- β 1/Smad3/4, while PIASy acts as a repressor.

[Immune Network 2014;14(6):321-327]

Keywords: IgA, germline α transcripts, SUMO, PIASy, HDAC1

INTRODUCTION

Under the influence of antigens and cytokines, Ig class switch recombination (CSR) on the heavy chain locus in IgM⁺ mature B cells allows the cells to produce other Ig isotypes (IgG, IgA, and IgE). IL-4, IFN- γ , and TGF- β 1 induce IgG1/IgE, IgG2a/IgG3, and IgA/IgG2b class switching by mouse B cells, respectively (1). Each transcription of germline transcripts (GLTs) in the switch regions is a prerequisite for each Ig CSR process (2). For instance, selective induction of GL α transcription by TGF- β 1 initiates IgA CSR by increasing the accessibility of activation-induced cytidine deaminase, which is an essential enzyme for Ig CSR (3), to a non-transcribed DNA strand of the switch region. TGF- β 1 induces IgA class switching through Smad3/4-mediated GL α transcription (4). We previously reported that many proteins such as Runx3, p300, Smurf, Arkadia, Tiul1, TGIF, and Stat1 are involved in the regulation of TGF- β 1-induced, Smad3/4-mediated GL α transcription by B cells (5-8).

The Smad proteins are key intracellular effectors of TGF- β signaling pathways that encompass a diverse set of cellular responses, including cell proliferation and differentiation. Post-translational modifications of Smad proteins regulate

Received on October 7, 2014. Revised on November 26, 2014. Accepted on December 2, 2014.

© This is an open access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Corresponding Author. Seok-Rae Park, Department of Microbiology, College of Medicine, Konyang University, 28, Gasuwon-dong, Wonang maedul 1-gil, Seo-gu, Daejeon, Korea. Tel: 82-42-600-6497; Fax: 82-42-600-6314; E-mail: srpark@konyang.ac.kr

#These authors contributed equally to this work.

Abbreviations: CSR, class switch recombination; GL α , germline α transcripts; SUMO, small ubiquitin-like modifier; Ubc, ubiquitin carrier protein; PIAS, protein inhibitor of activated STAT; HDAC, histone deacetylase; TSA, Trichostatin A

their function. Phosphorylation of receptor-regulated Smads (Smad2 and Smad3) induces hetero-oligomeric formation. Ubiquitination and acetylation of Smads are also required for Smads-dependent gene regulation. Another type of post-translational modification, sumoylation, has been shown to be important for TGF- β signaling pathways (9). Sumoylation is a process in which a small ubiquitin-like modifier (SUMO), a ubiquitin-related polypeptide, covalently binds to lysine residues of the target proteins and thereby regulates the function of proteins engaged in diverse processes. Three SUMO proteins, SUMO-1, SUMO-2, and SUMO-3, are expressed in higher eukaryotes. SUMO-2 and SUMO-3 share 97% similarity with each other, but only 50% similarity with the most studied SUMO-1 protein (10). Sumoylation is mediated through a process involving an E2 ubiquitin carrier protein (Ubc9) and five protein inhibitor of activated STAT (PIAS) family E3 ubiquitin ligases (PIAS1, PIAS3, PIASx α , PIASx β , and PIASy). This post-translational modification can affect target protein interaction, localization, and function (11).

SUMO-1 induces Smad4 sumoylation through TGF- β -induced p38 MAP kinase pathway activation and facilitates Smad-dependent transcriptional activation, and PIAS1/PIASx β -mediated sumoylation of Smad4 is also regulated by the p38 MAP kinase pathway (12). Smad4 is sumoylated in a process involving Ubc9 and PIASy (13-15), PIAS1 (12,16), and PIASx β (12). The sumoylation of Smad4 alters its subnuclear localization, enhances its stability, and increases Smad4-mediated transcriptional activation (13-15). Furthermore, the interaction between PIAS1 and Smad4 is positively regulated by TGF- β , thus creating a positive feedback loop (16). Association of Smad3 and Smad7 with PIASy leads to increased sumoylation of Smad3 (17). PIASy suppresses TGF- β /Smad3-mediated signaling by stimulating sumoylation and nuclear export of Smad3 (18). PIASy also can inhibit TGF- β /Smad-mediated transcriptional responses by interacting with Smads and histone deacetylase 1 (HDAC1) (19). On the other hand, PIAS3 activates Smad transcriptional activity through its interaction with Smads and p300 (20). In addition, sumoylation of type I TGF- β receptor enhances receptor function by facilitating the recruitment and phosphorylation of Smad3, consequently regulating TGF- β -induced transcription (21).

Thus, sumoylation of Smad3/4 by SUMO-1, Ubc9, and PIAS proteins regulates TGF- β /Smads-mediated transcriptional responses. Nonetheless, it is not known if the sumoylation-related molecules are involved in the regulation of GL α transcription by TGF- β 1 in B cells. We report here for the first

time the effects of SUMO proteins, Ubc9, and PIASy on TGF- β 1-induced, Smad3/4-mediated GL α transcription and IgA switching by a mouse B cell line, CH12F3-2A.

MATERIALS AND METHODS

Cell culture and reagents

The mouse B lymphoma cell line CH12F3-2A (surface Ig μ ⁺) (22) was provided by Dr. T. Honjo (Kyoto University, Kyoto, Japan). The cells were cultured at 37°C in a humidified CO₂ incubator (Forma Scientific, Marietta, OH, USA) in RPMI-1640 medium (WelGENE, Daegu, Korea) supplemented with 10% fetal bovine serum (PAA Laboratories, Etobicoke, ON, Canada). The cells were stimulated with TGF- β 1 (0.5 ng/ml, R&D Systems, Minneapolis, MN, USA) and trichostatin A (TSA) (Sigma-Aldrich, St Louis, MO, USA).

Expression and reporter plasmids

Genes encoding Smad3 (23) and Smad4 (24) subcloned into Flag-pcDNA3, were provided by Dr. M. Kawabata (The Cancer Institute, Tokyo, Japan). The expression plasmid for SUMO-1 (pcDNA3-HA-SUMO-1) (12) was obtained from Dr. T. Ohshima (Tokushima Bunri University, Kagawa, Japan). Dr. F. Liu (The State University of New Jersey, NJ, USA) provided pcDNA3-HA-SUMO-2, pcDNA3-HA-SUMO-3, pcDNA3-HA-Ubc9, and pCMV5-FLAG-PIASy (25). The expression plasmid for HDAC1 (pK7-HDAC1-GFP) (26) was obtained from AddGene (Cambridge, MA, USA). The GL α -448 to +72 luciferase promoter was subcloned into pGL3-Basic Vector (Promega, Madison, WI, USA) (27) and was named pGL3-GL α [-448/+72].

Transfection and luciferase reporter assays

Transfection was performed by electroporation with a Gene Pulser II (Bio-Rad, Hercules, CA, USA) (4). Reporter plasmids were co-transfected with expression plasmids and pCMV β gal (Stratagene, La Jolla, CA, USA), and luciferase and β -gal assays were performed as described previously (4).

SDS-PAGE and immunoblotting

CH12F3-2A cells were transfected as above with the appropriate expression vectors. After 24 h incubation, total cell lysates were subjected to SDS-PAGE under reducing conditions, and proteins were transferred to PVDF membranes (Bio-Rad). Specific immunodetection was carried out by incubation of the membranes with an anti-HA antibody (Sigma-Aldrich) fol-

lowed by peroxidase-conjugated goat anti-mouse IgG antibody (Pierce, Thermo Scientific, Rockford, IL, USA), and visualized by chemiluminescence (Supersignal detection kit, Pierce).

RT-PCR

RNA preparation, reverse transcription, and PCR were performed as described previously (4). PCR primers were synthesized by Bioneer (Daejeon, Korea): GLT α (forward 5'-CTA-CATAGGGAAGATAGCCT-3', reverse 5'-TAATCGTGAATCAGGCAG-3', product size: 206 bp) and β -actin (forward 5'-CATGTTTGAGACCTTCAACACCCC-3', reverse 5'-GCCATCTCCTGCTCGAAGTCTAG-3', product size, 320 bp). All reagents for RT-PCR were purchased from iNtRON Biotechnology (Seongnam, Korea). PCR for β -actin was performed in parallel to normalize cDNA concentrations within each set of samples. Aliquots of the PCR products were resolved by electrophoresis on 2% agarose gels.

Flow cytometry analysis

Cultured cells were washed with HBSS (WelGENE) and resuspended in DMEM (WelGENE) containing 5% FBS, 0.1% NaN₃ at a density of 1×10^6 cells/ml. FITC-conjugated goat anti-mouse IgA (eBioscience, San Diego, CA, USA) and PE-conjugated goat anti-mouse IgM (eBioscience) were added to the cell suspension and placed at 4°C for 30 min. Cells were washed with HBSS three times and resuspended in PBS containing 1% formalin. Cytofluorometric analysis was carried out using a FACSCalibur (Becton Dickinson, Mountain View, CA, USA).

Isotype-specific ELISA

Antibodies produced in B cell cultures were detected by using isotype-specific ELISA. Affinity purified anti-isotype specific antibodies were added at 1.2 μ g/ml in 0.05 M sodium bicarbonate buffer (pH 9.3) to 96-well U bottomed polyvinyl microplates (Falcon, Becton Dickinson & Co., Oxnard, CA, USA). Plates were washed with PBS containing 0.05% Tween-20 (PBST) followed by overnight incubation at 4°C, and blocked for 1 h with 0.5% BSA solution. After washing, 50 μ l of standard myeloma proteins and culture supernatants were added to each well and incubated for 1 h at 37°C. Following another washing step, horseradish-peroxidase (HRPO) conjugated anti-isotype specific antibodies (Southern Biotechnology, Birmingham, AL, USA) were added to each well and incubated for 1 h. Plates were then washed and TMB substrate (BD

Biosciences) was added. After incubation, 0.05 M sulfuric acid was added to each well and the colorimetric reaction was measured at 450 nm with an Absorbance Microplate Reader.

Statistical analysis

Statistical differences between experimental groups were determined by analysis of variances. Values with $p < 0.05$ and $p < 0.01$ by an unpaired two tailed Student's *t*-test were considered significant.

RESULTS AND DISCUSSION

SUMO overexpression does not affect GL α transcription and IgA expression

In this study, we used a well-characterized mouse CH12F3-2A B lymphoma cell line to investigate IgA isotype switching (22,28). Smad3 plays a key role in TGF- β 1-induced GL α transcription. Therefore, we first tried to address whether SUMO overexpression affects TGF- β 1-induced, Smad3-mediated GL α promoter activity in CH12F3-2A cells. As shown in Fig. 1A (left panel), Smad3 overexpression enhanced TGF- β 1-induced GL α promoter activity. However, overexpression of SUMO-1, SUMO-2, or SUMO-3 did not alter TGF- β 1-induced, Smad3-mediated GL α promoter activity. We confirmed through detection of sumoylated proteins following transfection of SUMO-1 and SUMO-3 expression vectors that overexpression of SUMO proteins results in sumoylation of certain proteins (Fig. 1A, right panel). Smad4 physically interacts with Ubc9, and SUMO-1 overexpression enhances TGF- β -induced transcriptional responses by increasing intranuclear Smad4 level (13,14). Smad4 sumoylation was shown to increase in the presence of Ubc9 and PIASy, which enhances Smad4-mediated transcriptional activation (15). In contrast, it was reported that SUMO-1 and Ubc9 can inhibit a TGF- β -responsive transcriptional activity via sumoylation of Smad4 (25). The net effect of Smad4 sumoylation can therefore be either stimulatory or inhibitory, depending on distinct transcriptional effects of the target promoter that is analyzed. We next determined whether Ubc9 and SUMO-1 overexpression regulates Smad3/4-mediated GL α promoter activity induced by TGF- β 1. As shown in Fig. 1B (left panel), Ubc9 hardly affected TGF- β 1-induced, Smad3- or Smad4-mediated GL α promoter activity. In addition, Ubc9 and SUMO-1 affected neither the TGF- β 1-induced GL α promoter activity nor Smad3/4-mediated promoter activity (Fig. 1B, right panel). Next, we investigated the effect of SUMO-1 overexpression on the ex-




Figure 1. Effects of SUMO and Ubc9 on TGF- β 1-induced, Smad3/4-mediated GL α promoter activity. (A) CH12F3-2A cells were transfected with GL α promoter reporter construct (pGL3-GL α [-448/+72], 10 μ g) and the indicated expression plasmids (5 μ g each). TGF- β 1 (0.5 ng/ml) was added, and luciferase activity was determined 16 h later. Transfection efficiency was normalized to β -gal activities. Data shown are average luciferase activities of three independent transfections with SEM (bars) (left panel). Sumoylated proteins were detected by immunoblotting analysis for HA from extracts of CH12F3-2A cells transfected with SUMO-1 and SUMO-3 (right panel). (B) CH12F3-2A cells were transfected with pGL3-GL α [-448/+72] (10 μ g) and the indicated expression plasmids (5 μ g each) including Ubc9 (5 μ g). Induction and analysis were performed as shown in (A). * p <0.05, ** p <0.01.

pression of endogenous GLT α . As shown in Fig. 2A, Smad3 overexpression enhanced TGF- β 1-induced GLT α expression, while SUMO-1 overexpression did not alter the expression. Furthermore, SUMO-1 overexpression had no effect on TGF- β 1-induced, Smad3-mediated surface IgA expression and IgA production by CH12F3-2A cells (Fig. 2B and 2C). We further tested the effects of SUMO-2 and SUMO-3 overexpression on TGF- β 1/Smad3-mediated GL α transcription and IgA expression, but found no significant effects (data not shown). In summary, these results suggest that SUMO and Ubc9 proteins do not affect on TGF- β 1-induced, Smad3/4-mediated GL α transcription and IgA secretion by B cells.

PIASy and HDAC1 cooperatively suppress TGF- β 1-induced, Smad3/4-mediated GL α promoter activity

Smad4 sumoylation by the PIASy redistributes Smad4 to subnuclear speckles, which result in enhanced Smad4-mediated transcriptional activation by TGF- β (15). In contrast, PIASy inhibits TGF- β /Smad3-mediated transcriptional response either by stimulating sumoylation of Smad3 and nuclear export of Smad3 (18), or by interacting with Smad3/4 and HDAC1 (19). Here we first tested the effect of PIASy on GL α promoter activity in B cells. Overexpression of PIASy diminished TGF- β 1-induced GL α promoter activity in a dose-dependent manner (Fig. 3A, left panel). Overexpression of the Flag-




Figure 2. SUMO-1 is not involved in TGF- β /Smad3-induced GL α transcription and IgA isotype switching. CH12F3-2A cells were transfected with Smad3 (5 μ g) and SUMO-1 (5 μ g), and the cells were stimulated with TGF- β 1 (0.5 ng/ml). (A) After 36 h of culture, RNA was isolated, and levels of GLT α and β -actin mRNA were measured by RT-PCR. (B) Following 3 days incubation, cells were stained with FITC-labeled anti-mouse IgA and PE-labeled anti-mouse IgM and analyzed by flow cytometry. (C) Culture supernatants were harvested after incubation for 4 days, and IgA levels were determined by ELISA. Data are means \pm SEM of three cultures.

PIASy protein was detected by immunoblot using an anti-Flag antibody (data not shown). In addition, PIASy suppressed TGF- β 1-induced, Smad3/4-mediated GL α promoter activity (Fig. 3A, right panel). Thus, PIASy act as a negative regulator of Smad3/4-mediated GL α transcription by TGF- β 1 in B cells. Next, we blocked HDAC with the effective inhibitor TSA to determine if the deacetylase is involved in the suppression of the GL α promoter activity. As shown in Fig. 3B (upper panel), TSA dose-dependently increased TGF- β 1-induced GL α promoter activity and restored PIASy-mediated suppression of GL α activity. Conversely, PIASy inhibited TSA-mediated fortification of GL α promoter activity. Nevertheless, TSA could still enhance GL α promoter activity even when PIASy was overexpressed, albeit to a lesser degree (Fig. 3B, upper panel). This result suggests that although both PIASy and HDAC can suppress GL α transcription, they can act independently from each other. Namely, PIASy and HDAC may have their own inhibitory mechanisms that do not rely on each other rather than interrelated common mechanism. In addition,

overexpression of HDAC1 further decreased PIASy-mediated suppression of GL α promoter activity (Fig. 3B, lower left panel). Overexpression of the HDAC1-GFP protein was measured by flow cytometric detection of GFP (data not shown). Furthermore, TSA recovered suppression of GL α promoter activity by PIASy in the presence of TGF- β 1 (Fig. 3B, lower right panel). Thus, PIASy can cooperate with HDAC1 to suppress TGF- β 1-induced, Smad3/4-mediated GL α transcription in mouse B cells. These suppressions by PIASy and HDAC1 may be due to 1) repression of GL α transcriptional activity by PIASy-mediated SUMO modification of certain proteins including Smad3/4, since SUMO modification often represses transcriptional activity of a number of transcription factors, such as Sp3, c-Jun, c-Myb, AP2, Elk-1, and p300 (29), and 2) recruitment of transcriptional repressor HDAC1 into the Smad3/4-PIASy complex, because PIASy can suppress Smad-mediated transcriptional responses by interacting with Smads and HDAC (19). However, these possibilities remain to be tested. In addition, we are currently exploring the effects of




Figure 3. Effects of PIASy and HDAC1 on TGF- β 1-induced, Smad3/4-mediated GL α promoter activity. (A) CH12F3-2A cells were transfected with pGL3-GL α [-448/+72] (10 μ g) and the indicated doses of PIASy alone (left panel) or PIASy (30 μ g) and Smad3/4 (5 μ g each) (right panel). Induction and analysis were performed as shown in Fig. 1A. (B) CH12F3-2A cells were transfected with pGL3-GL α [-448/+72] (10 μ g) and empty vector (pcDNA3, 30 μ g) or PIASy (30 μ g). TGF- β 1 (0.5 ng/ml) and the indicated doses of TSA were added (upper panel). PIASy (30 μ g) and HDAC1 (15 μ g) were transfected along with the GL α promoter reporter (10 μ g) into CH12F3-2A cells, and cells were treated with TGF- β 1 (lower left panel). The GL α promoter reporter (10 μ g), Smad3/4 (5 μ g each), and PIASy (30 μ g) were transfected into CH12F3-2A cells, and TGF- β 1 (0.5 ng/ml) and TSA (5 ng/ml) were added (lower right panel). Luciferase activity was determined 16 h later. * p <0.05, ** p <0.01.

PIASy and HDAC1 on the endogenous expression of GL α transcripts and IgA expression induced by TGF- β 1/Smad3/4 in B cells.

Overall, our novel findings provide new basic evidence that SUMO and Ubc9 are unlikely to affect TGF- β /Smad3/4-mediated GL α transcription by B cells, while PIASy and HDAC1

act as repressor molecules in the transcription.

ACKNOWLEDGEMENTS

This research was supported by Basic Science Research Program through the National Research Foundation of Korea

(NRF) funded by the Ministry of Education, Science and Technology (MEST) (NRF-2010-0025648).

CONFLICTS OF INTEREST

The authors have no financial conflict of interest.

REFERENCES

1. Stavnezer, J. 1996. Immunoglobulin class switching. *Curr. Opin. Immunol.* 8: 199-205.
2. Stavnezer, J. 2000. Molecular processes that regulate class switching. *Curr. Top. Microbiol. Immunol.* 245: 127-168.
3. Park, S. R. 2012. Activation-induced cytidine deaminase in B cell immunity and cancers. *Immune Netw.* 12: 230-239.
4. Park, S. R., J. H. Lee, and P. H. Kim. 2001. Smad3 and Smad4 mediate transforming growth factor-beta1-induced IgA expression in murine B lymphocytes. *Eur. J. Immunol.* 31: 1706-1715.
5. Park, S. R., E. K. Lee, B. C. Kim, and P. H. Kim. 2003. p300 cooperates with Smad3/4 and Runx3 in TGFbeta1-induced IgA isotype expression. *Eur. J. Immunol.* 33: 3386-3392.
6. Choi, S. H., G. Y. Seo, E. H. Nam, S. H. Jeon, H. A. Kim, J. B. Park, and P. H. Kim. 2007. Opposing effects of Arkadia and Smurf on TGFbeta1-induced IgA isotype expression. *Mol. Cells* 24: 283-287.
7. Park, K. H., E. H. Nam, G. Y. Seo, S. R. Seo, and P. H. Kim. 2009. Tiul1 and TGIF are involved in downregulation of TGFbeta1-induced IgA isotype expression. *Immune Netw.* 9: 248-254.
8. Park, S. R., M. H. Jung, S. H. Jeon, M. H. Park, K. H. Park, M. R. Lee, and P. H. Kim. 2010. IFN-gamma down-regulates TGF-beta1-induced IgA expression through Stat1 and p300 signaling. *Mol. Cells* 29: 57-62.
9. Miyazono, K., Y. Kamiya, and K. Miyazawa. 2008. SUMO amplifies TGF-beta signalling. *Nat. Cell Biol.* 10: 635-637.
10. Melchior, F. 2000. SUMO--nonclassical ubiquitin. *Annu. Rev. Cell Dev. Biol.* 16: 591-626.
11. Welchman, R. L., C. Gordon, and R. J. Mayer. 2005. Ubiquitin and ubiquitin-like proteins as multifunctional signals. *Nat. Rev. Mol. Cell Biol.* 6: 599-609.
12. Ohshima, T. and K. Shimotohno. 2003. Transforming growth factor-beta-mediated signaling via the p38 MAP kinase pathway activates Smad-dependent transcription through SUMO-1 modification of Smad4. *J. Biol. Chem.* 278: 50833-50842.
13. Lin, X., M. Liang, Y. Y. Liang, F. C. Brunnicardi, and X. H. Feng. 2003. SUMO-1/Ubc9 promotes nuclear accumulation and metabolic stability of tumor suppressor Smad4. *J. Biol. Chem.* 278: 31043-31048.
14. Lin, X., M. Liang, Y. Y. Liang, F. C. Brunnicardi, F. Melchior, and X. H. Feng. 2003. Activation of transforming growth factor-beta signaling by SUMO-1 modification of tumor suppressor Smad4/DPC4. *J. Biol. Chem.* 278: 18714-18719.
15. Lee, P. S., C. Chang, D. Liu, and R. Derynck. 2003. Sumoylation of Smad4, the common Smad mediator of transforming growth factor-beta family signaling. *J. Biol. Chem.* 278: 27853-27863.
16. Liang, M., F. Melchior, X. H. Feng, and X. Lin. 2004. Regulation of Smad4 sumoylation and transforming growth factor-beta signaling by protein inhibitor of activated STAT1. *J. Biol. Chem.* 279: 22857-22865.
17. Imoto, S., K. Sugiyama, R. Muromoto, N. Sato, T. Yamamoto, and T. Matsuda. 2003. Regulation of transforming growth factor-beta signaling by protein inhibitor of activated STAT, PIASy through Smad3. *J. Biol. Chem.* 278: 34253-34258.
18. Imoto, S., N. Ohbayashi, O. Ikeda, S. Kamitani, R. Muromoto, Y. Sekine, and T. Matsuda. 2008. Sumoylation of Smad3 stimulates its nuclear export during PIASy-mediated suppression of TGF-beta signaling. *Biochem. Biophys. Res. Commun.* 370: 359-365.
19. Long, J., I. Matsuura, D. He, G. Wang, K. Shuai, and F. Liu. 2003. Repression of Smad transcriptional activity by PIASy, an inhibitor of activated STAT. *Proc. Natl. Acad. Sci. U. S. A.* 100: 9791-9796.
20. Long, J., G. Wang, I. Matsuura, D. He, and F. Liu. 2004. Activation of Smad transcriptional activity by protein inhibitor of activated STAT3 (PIAS3). *Proc. Natl. Acad. Sci. U. S. A.* 101: 99-104.
21. Kang, J. S., E. F. Saunier, R. J. Akhurst, and R. Derynck. 2008. The type I TGF-beta receptor is covalently modified and regulated by sumoylation. *Nat. Cell Biol.* 10: 654-664.
22. Nakamura, M., S. Kondo, M. Sugai, M. Nazarea, S. Imamura, and T. Honjo. 1996. High frequency class switching of an IgM⁺ B lymphoma clone CH12F3 to IgA⁺ cells. *Int. Immunol.* 8: 193-201.
23. Zhang, Y., X. Feng, R. We, and R. Derynck. 1996. Receptor-associated Mad homologues synergize as effectors of the TGF-beta response. *Nature* 383: 168-172.
24. Hahn, S. A., M. Schutte, A. T. Hoque, C. A. Moskaluk, L. T. da Costa, E. Rozenblum, C. L. Weinstein, A. Fischer, C. J. Yeo, R. H. Hruban, and S. E. Kern. 1996. DPC4, a candidate tumor suppressor gene at human chromosome 18q21.1. *Science* 271: 350-353.
25. Long, J., G. Wang, D. He, and F. Liu. 2004. Repression of Smad4 transcriptional activity by SUMO modification. *Biochem. J.* 379: 23-29.
26. Tou, L., Q. Liu, and R. A. Shivdasani. 2004. Regulation of mammalian epithelial differentiation and intestine development by class I histone deacetylases. *Mol. Cell Biol.* 24: 3132-3139.
27. Park, M. H., S. R. Park, M. R. Park, Y. H. Kim, and P. H. Kim. 2011. Retinoic acid induces expression of Ig germ line α transcript, an IgA isotype switching indicative, through retinoic acid receptor. *Genes Genom.* 33: 83-88.
28. Jang, Y. S., S. H. Choi, S. R. Park, H. A. Kim, J. B. Park, and P. H. Kim. 2004. Characterization of mouse B lymphoma cells (CH12F3-2A) for the study of IgA isotype switching. *Immune Netw.* 4: 216-223.
29. Verger, A., J. Perdomo, and M. Crossley. 2003. Modification with SUMO. A role in transcriptional regulation. *EMBO Rep.* 4: 137-142.