

POSTER PRESENTATION

Open Access

# Effects of a reduced efficacy of the KCC2 co-transporter in temporal lobe epilepsy: single neuron and network study

Anatoly Buchin<sup>1,2\*</sup>, Gilles Huberfeld<sup>3,4</sup>, Richard Miles<sup>5</sup>, Anton Chizhov<sup>6</sup>, Boris Gutkin<sup>1,7</sup>

From 24th Annual Computational Neuroscience Meeting: CNS\*2015  
Prague, Czech Republic. 18-23 July 2015

Epilepsy is one of the most common neurological disorders. Seizures in about 40% of patients with temporal lobe epilepsies are pharmaco-resistant [1]. In surgically removed hippocampal tissue from these patients, the KCC2 cotransporter is absent or non-functional in about 20 % of subicular pyramidal cells [2]. KCC2 normally assures the maintenance of low intra-neuronal chloride levels [3] and also regulates potassium levels [4]. Chloride concentration changes in the remaining pyramidal cells due to intensive GABAergic input during seizures could reverse the effects of GABA currents from inhibitory to excitatory [5,6]. Such changes may shift a pyramidal cell into a periodic bursting regime associated with ictal discharges. Using a detailed biophysical model of a single cell incorporating these mechanisms of ionic homeostasis and a neural network model, we show that decreasing the activity of KCC2 pump leads to repetitive seizure-like firing in the pathologic network due to increased extracellular potassium and intracellular chloride (Fig. 1). This model provides insights into how a dysregulation of pyramidal cell chloride homeostasis due to reduced levels of the KCC2 cotransporter may lead to seizures in the epileptic human subiculum.

## Acknowledgements

This work has been supported by the following grants: ANR-10-LABX-0087 IEC, ANR-10-IDEX-0001-02 PSL, ERC-322721, FRM FDT20140930942. Especially we would like to thank Giri Krishnan for useful discussions.

## Authors' details

<sup>1</sup>École normale supérieure, Laboratoire des Neurosciences Cognitives, Group for Neural Theory, Paris, France. <sup>2</sup>Peter the Great St.-Petersburg Polytechnic University, St.-Petersburg, Russia. <sup>3</sup>Neurophysiology Department, Pitie-Salpêtrière Hospital, UPMC, Paris, France. <sup>4</sup>Epilepsie de l'Enfant et Plasticité Cérébrale, INSERM U1129, Paris, France. <sup>5</sup>Institut du Cerveau et de la Moelle Epinière, Cortex et Epilepsie Group, Paris, France. <sup>6</sup>Ioffe Physical Technical Institute, Computational Physics Laboratory, St.-Petersburg, Russia. <sup>7</sup>Higher School of Economics, Moscow, Russia.

Published: 18 December 2015

## References

1. Beghi E, Berg A, Carpio A, Forsgren L, Hesdorffer DC, Hauser WA, Tomson T: Comment on epileptic seizures and epilepsy: definitions proposed by the International League Against Epilepsy (ILAE) and the International Bureau for Epilepsy (IBE). *Epilepsia* 2005, 46(10):1698-1699.
2. Huberfeld G, Wittner L, Clemenceau S, Baulac M, Kaila K, Miles R, Rivera C: Perturbed chloride homeostasis and GABAergic signaling in human temporal lobe epilepsy. *J Neuroscience* 2007, 27(37):9866-9873.
3. Blaeser P, Airaksinen MS, Rivera C, Kaila K: Cation-chloride cotransporters and neuronal function. *Neuron* 2009, 61(6):820-838.
4. Payne JA: Functional characterization of the neuronal-specific K-Cl cotransporter: implications for [K+] oregulation. *American Journal of Physiology-Cell Physiology* 1997, 273(5):C1516-C1525.
5. Khalilov I, Dzhala V, Ben-Ari Y, Khazipov R: Dual role of GABA in the neonatal rat hippocampus. *Developmental Neuroscience* 1999, 21(3-5):310-319.
6. Jedlicka P, Deller T, Gutkin BS, Backus KH: Activity-dependent intracellular chloride accumulation and diffusion controls GABA-A receptor-mediated synaptic transmission. *Hippocampus* 2011, 21(8):885-898.


doi:10.1186/1471-2202-16-S1-P5

Cite this article as: Buchin et al.: Effects of a reduced efficacy of the KCC2 co-transporter in temporal lobe epilepsy: single neuron and network study. *BMC Neuroscience* 2015 16(Suppl 1):P5.

\* Correspondence: anat.buchin@gmail.com

<sup>1</sup>École normale supérieure, Laboratoire des Neurosciences Cognitives, Group for Neural Theory, Paris, France

Full list of author information is available at the end of the article



**Figure 1** A raster plot of pyramidal cell population firing; B raster plot of interneuron firing; C spatial distribution of extracellular potassium; D intracellular chloride distribution.