

**1055. Antibody Response to Intradermal and High Dose Influenza Vaccine in 2012-13 Among Adults Who Did and Did Not Respond to Standard Dose Vaccine in 2011-12**

Maria Sundaram, MSPH<sup>1</sup>; Huong Mclean, MPH, PhD<sup>1</sup>; Jennifer Meece, PhD<sup>2</sup>; Richard K. Zimmerman, MD MPH<sup>3</sup>; Mary Patricia Nowalk, PhD<sup>3</sup>; Chyongchiou J Lin, PhD<sup>4</sup>; Thomas Friedrich, PhD<sup>5</sup>; Sarah Spencer, PhD<sup>5</sup>; Edward Belongia, MD<sup>1</sup>; <sup>1</sup>Center for Clinical Epidemiology and Population Health, Marshfield Clinic Research Foundation, Marshfield, WI; <sup>2</sup>Integrated Research and Diagnostic Laboratory, Marshfield Clinic Research Foundation, Marshfield, WI; <sup>3</sup>Family Medicine, University of Pittsburgh, Pittsburgh, PA; <sup>4</sup>Pathobiological Sciences, University of Wisconsin School of Veterinary Medicine, Madison, WI; <sup>5</sup>Influenza Division, Centers for Disease Control and Prevention, Atlanta, GA

**Session:** 124. Vaccines: Influenza  
**Friday, October 10, 2014: 12:30 PM**

**Background.** We compared antibody responses to influenza vaccine in adults  $\geq 50$  years old in two consecutive seasons. In 2011-12, all participants received trivalent inactivated influenza vaccine (IIV3). In 2012-13, those age 50-64 received intradermal (ID) and those age  $\geq 65$  received high-dose vaccine (HD). In 2012-13, the WHO-recommended vaccine strains were modified for H3N2 and B.

**Methods.** Participants were recruited in 2011 and vaccinated in the fall of 2011 and 2012. In both years, pre- and post (21-28 days)-vaccination sera were tested for antibody response to vaccine [measured by hemagglutination inhibition (HI) titer]. We compared geometric mean titers (GMT) and titer fold rise from pre- to post-vaccination in 2012, among 2011-12 responders ( $\geq 4$ -fold rise in HI titer from pre-to post-vaccination) and 2011-12 non-responders ( $<4$ -fold rise in HI titer) to the corresponding vaccine strain.

**Results.** 183 adults received IIV3 in 2011 and either HD (N = 82) or ID (N = 101) in 2012. For the 2011-12 vaccine, 56 (31%) responded to the H1N1 strain, 68 (37%) responded to H3N2, and 28 (15%) responded to B. Among HD recipients, post-vaccination GMT for H3N2 was higher among 2011-12 responders vs non-responders ( $p = 0.04$ ). Fold rise did not differ significantly for any vaccine strains between responders vs non-responders.

Strain	2012-13 Response	HD recipients		ID recipients	
		2011-12		2011-12	2011-12
		Non- responders	Responders		
H1N1	Post-vaccination	93 (57, 160)	128 (80, 320)	78 (40, 226)	105 (40, 160)
	Median fold rise (IQR)	4 (2, 8)	4 (1, 8)	2 (1, 4)	2 (2, 4)
H3N2	Post-vaccination	154 (80, 320)*	242 (160, 452)	120 (57, 320)	118 (40, 226)
	Median fold rise (IQR)	4 (2, 8)	4 (2, 11)	2 (1, 4)	2 (1, 8)
B	Post-vaccination	185 (80, 320)	180 (80, 320)	137 (80, 320)	163 (80, 320)
	Median fold rise (IQR)	3 (1, 8)	2 (2, 11)	2 (1, 6)	3 (1, 16)

\*  $p < 0.05$  for comparison of 2011-12 responders and non-responders.

**Conclusion.** Prior response to 2011-12 vaccine antigen was not associated with response to the new H3N2 or B antigens in 2012-13, nor was it associated with response to the H1N1 antigen used in both the 2011-12 and 2012-13 vaccines.

**Disclosures.** M. Sundaram, MedImmune, LLC: Project scientist on funded grant, Research grant; H. Mclean, MedImmune, LLC: Grant Investigator, Research grant R. K. Zimmerman, Sanofi: Grant Investigator, Research grant; Pfizer: Grant Investigator, Research grant M. P. Nowalk, Sanofi: Grant Investigator, Research support; Pfizer: Grant Investigator, Research support; Merck: Grant Investigator, Research support; MedImmune: Consultant, Consulting fee C. J. Lin, MedImmune, LLC: Consultant, Consulting fee Sanofi Pasteur: Grant Investigator, Research grant; Merck: Grant Investigator, Research grant E. Belongia, MedImmune, LLC: Grant Investigator, Research grant