

ORIGINAL ARTICLE

Open Access



Knocking out Analysis of the *CpxP* gene using Crispr/Cas9 in *Escherichia coli* MG1655

Xiaoliang He¹, Yuwen Ren, Wanli Meng, Xinran Yu and Xiaohui Zhou*

Abstract

Based on the analysis of *cpxP* genes among *Escherichia coli* strains, *cpxP* gene-targeting short guide RNA (sgRNA) was designed and inserted into the pGL3-MGP-RNA. The donor sequences (MG-HR) for homologous repair were designed and cloned by PCR. MG-HR and pGL3-MGP-RNA were transformed into *E. coli* MG1655 (pCas9). The *cpxP* gene expression cassette was amplified by PCR and subcloned into pBBR1MCS-2. Then the pBBR-*cpxP* was independently transformed into *E. coli* MG1655. The results of motility experiment suggest that *cpxP* gene had a significant effect on the movement ability of *E. coli* strain. The CpxP protein had a significant inhibition of bacterial activity. The latest 81 CpxP proteins sequences were selected and analyzed by multi-sequence alignment and molecular cluster. The CpxP proteins were roughly divided into three categories. Our results suggest that the CpxP protein was involved in bacterial motility, infection and pathogenicity.

Keywords: Knocked out, *cpxP*, Crispr/Cas9, Homologous repair, *Escherichia coli*

Key points

The *E. coli* MG1655- Δ *cpxP* were successfully obtained.

The *cpxP* gene had a significant effect on the movement ability of *E. coli* strain.

The CpxP proteins had a significant inhibition of bacterial activity.

Introduction

The Cpx system is one of the most common two-component regulatory systems in Gram-negative bacteria. It consists of the membrane-anchored sensor kinase CpxA, the cytosolic response regulator CpxR, and the peripheral spatial helper protein CpxP (Dong et al. 1993; Ruiz and Silhavy 2005). The CpxP proteins can inhibit activation of CpxA and are indispensable for the quality control system of P pili that are protein filaments expressed by uropathogenic *Escherichia coli* (Fällman et al. 2005; Nevesinjac and Raivio 2005). The structure of CpxP was interdigitated with two monomers like “left hands” forming a cap-shaped dimer. The structure revealed an

antiparallel dimer of intertwined α -helices with a highly basic concave surface (Thede et al. 2011). The CpxP proteins inhibit the kinase CpxA through direct interaction between its concave polar surface and the negatively charged sensor domain on CpxA (Zhou et al. 2011).

The CRISPR-Cas system was used recently as efficient genome engineering technology in several prokaryotes and eukaryotes, including (but not limited to) *Escherichia coli* (Jiang et al. 2013), *Saccharomyces cerevisiae* (DiCarlo et al. 2013), yeast (DiCarlo et al. 2015), *Streptomyces spp.* (Cobb et al. 2014), higher plants (Shan et al. 2013), *Bombyx mori* (Wang et al. 2013), *Drosophila* (Yu et al. 2013), insects (Gantz and Bier 2015), *Anopheles stephensi* (Gantz et al. 2015), *Anopheles gambiae* (Hammond et al. 2016), mouse (Grunwald et al. 2019) and human cell lines (Cong et al. 2013; Mali et al. 2013; Zhang et al. 2013). The CRISPR/Cas9 system was also used to remove plasmid harbouring *mcr-1* from *Escherichia coli* (Dong et al. 2019).

This study constructed an expression vector pGL3-MGP-RNA including gene-targeting short guide RNA and cloned a donor sequences (MG-HR) for homologous repair. The cell-envelope stress modulator *cpxP* gene of MG1655 strain was successfully knocked out

*Correspondence: zhouxh2003@aliyun.com

School of Biological Science and Engineering, Hebei University of Science and Technology, No. 26 Yuxiang Street, Shijiazhuang 050018, Hebei, China

by CRISPR-Cas9 system-based gene editing strategy. Likewise, the system can efficiently edit a large plasmid, target genes on the bacterial chromosome, or be adapted to introduce functional gene cargos alongside the gRNA cassette.

Materials and methods

Plasmid construction

The pCas9 and pGL3-U6-SgRNA-PGK-Puromycin Plasmid (Youbio, Hunan, China) was extracted from *E. coli* DH5 α by Plasmid extraction kit (GenStar, Beijing, China). The pCas9 Plasmid was transfected into *E. coli* MG1655 (Tiangen, Beijing, China) to acquire *E. coli* MG1655 (pCas9). The transformed *E. coli* MG1655 was screened using Kanamycin (50 mg/L) that was added to the LB medium. Genomic DNA was extracted from *E. coli* MG1655 by Bacterial genome extraction kit (GenStar, Beijing, China).

The MG-HR-S and MG-HR-X were amplified by PCR using the genomic DNA as a template. The primers were designed with prime primer 5.0. The primers were synthesized by the Beijing Invitrogen Biotechnology Company. The primers TCTGGTGTGTCTGGC GAAGT and TGCTAATTCGTGGAGCTTATGCCA GCG TTGAGGCCATG were used to amplify MG-HR-S. The primers TAAGCTCCACGAATTAGCATCAGC AGATGCGAGATCTTAT and CTATGGC AAGGAA AACAGGGT were used to amplify MG-HR-X. MG-HR-S was connected to MG-HR-X by PCR to acquire a MG-HR.

The MGP-sgRNA and pGL3 were amplified by PCR using the pGL3-U6-SgRNA-PGK-Puromycin Plasmid as a template.

The primers CGGGATCCTTGACAGCTAGCT CAGTCCTAGGTATAATACTAG TTCAGGCGATAA CTGGCATCCGTTTTAGAGCTAGAAAT and GGG GTACCGG AACCACGCCAGAGCAG were used to amplify MGP-sgRNA.


The primers GGGGTACCGCTCACTGACTCGCTG CGCT and CGGGATCCGC TTAATGCGCCGCTAC AGG were used to amplify pGL3.

The *cpxP* gene expression cassette comprising said promoter, a *cpxP* gene and said terminator was amplified by PCR using the genomic DNA as a template.

cpxP-F: CCAAGCTTACGCGGTCTAATTCA CTGCC 3'

cpxP-R: CGCGGATCCAGACAGGGATGGTGTCTA TGGC 3'

The PCR was carried out by using the Gene Amp PCR system 9700 (Applied Biosystems). PCR products were confirmed on 1.0% agarose gels and recycled by the agarose gel extraction kit (Macherey–Nagel, Germany).



Transformation of recombinant plasmid

Recycled PCR products of the MGP-sgRNA and pGL3 were digested by KpnI and BamHI (Thermo Fisher, USA), and then connected by T4 DNA Ligase (Thermo Fisher, USA) to acquire a pGL3-MGP-sgRNA.

400 μ g pGL3-U6-sgRNA-PGK-puromycin, 400 μ g pGL3-MGP-RNA, 400 μ g pGL3-MGP-RNA and 1.6 μ g MG-HR were separately transformed into *E. coli* MG1655 (pCas9). The transformed *E. coli* MG1655 (pCas9) were screened using Kanamycin (50 mg/L) and Ampicillin (100 mg/L) that were added to the LB medium.

Recycled PCR products of the *cpxP* gene expression cassette were digested by BamHI and HindIII (Thermo Fisher, USA), purified by the agarose gel extraction kit. The purified PCR products were directly subcloned




Fig. 2 Antibacterial effect. **a** Antibacterial effect to *E. coli* K88. (1) 200 μ l ddH₂O. (2) 200 μ l CpxP proteins. (3) 200 μ l Kan (2.5 mg/ml). **b** Antibacterial effect to *E. coli* MG1655. (1) 200 μ l ddH₂O. (2) 200 μ l CpxP proteins. (3) 200 μ l Kan (2.5 mg/ml)

Table 1 Inhibitory circle of CpxP after dispersed to two kinds of pathogen vibrio

	Inhibitory circle (mm)	
	<i>E. coli</i> K88	<i>E. coli</i> MG1655
ddH ₂ O	0	0
CpxP proteins	27 \pm 1.15	29 \pm 1.25
Kan	37 \pm 1.23	34 \pm 1.19

Values are the mean \pm SD (n = 4)

into pBBR1MCS-2 (Youbio, Hunan, China) (No. pBBR-*cpxP*). Then the pBBR-*cpxP* was independently transformed into *E. coli* MG1655 to acquire the overexpression transformants and *E. coli* MG1655- Δ *cpxP* to acquire a revertant. The transformed *E. coli* MG1655 were screened using Kanamycin (50 mg/L).

Motility experiment

Escherichia coli MG1655, overexpression transformants, the knocked out mutants and revertants were inoculated on LB medium for overnight. Then the cultures (2 μ l) were added to the LB semi-solid medium for overnight.

Bacteriostatic experiment

200 μ l ddH₂O, 200 μ l Kan (2.5 mg/ml) and 200 μ l CpxP proteins (8 mg/ml) were independently added to the LB solid medium by 6 mm Oxford cup containing 100 μ l *E. coli* MG1655 and *E. coli* K88 (10⁵ cfu/

ml) for 16 h at 4 °C. Then they were kept at 37 °C for bacteriostatic experiment.

Molecular clustering

To evaluate and analyze CpxP sequence resource preliminary, we had found 81 CpxP sequences from public database (<http://www.ncbi.nlm.nih.gov/>) and done multi-sequence alignment and molecular cluster by Clustal X and Treeview (He et al. 2015).

Results

Cloning of *cpxP* gene fragment


The *cpxP* gene fragments were cloned by PCR using MG-HR-S-F and MG-HR-S-R, MG-HR-X-F and MG-HR-X-R as primers. The results showed that the lengths of *cpxP* gene fragment were 424 bp and 377 bp respectively. The lengths of MG-HR-S and MG-HR-X were 782 bp (Additional file 1: Figure S1).

Identification of pGL3-MGP-sgRNA plasmid

To confirm whether the MGP-sgRNA and pGL3 were connected by T4 DNA Ligase, the recombinant plasmids (pGL3-MGP-sgRNA) were digested with KpnI and BamHI. According to the electrophoresis, the recombinant plasmids (pGL3-MGP-sgRNA) were successfully obtained (Additional file 1: Figure S2).

Identification of the knockout of *cpxP* gene

The knockout of *cpxP* gene was identified by PCR using MG-HR-S-F and MG-HR-X-R as primers, the genomic DNA of *E. coli* MG1655 (pCas9) as a template. According



to the electrophoresis, the lengths of PCR products were 888 bp without knockout and 782 bp after knockout of *cpXP* gene (Additional file 1: Figure S3A). The PCR products were sent to the Beijing Invitrogen Biotechnology Company for sequencing. The sequences were done multi-sequence alignment by DNASTAR (Additional file 1: Figure S3B). These results suggest that the *E. coli* MG1655- Δ *cpXP* were successfully obtained.

Identification of pBBR-*cpXP* plasmid

PCR products of the *cpXP* gene expression cassette were digested by BamHI and HindIII, and cloned into pBBR1MCS-2. According to the electrophoresis, the recombinant plasmids (pBBR1MCS-2), the overexpression transformants and revertant were successfully obtained (Additional file 1: Figure S4).

Motility experiment

Many studies have confirmed that the motility of bacteria plays an important role in the pathogenicity of bacteria during the early interaction with the host. To investigate the effect of the *cpXP* gene on the motility of *E. coli*, the mobility of the strain was determined. The results showed that the diffusion diameters of *E. coli* MG1655- Δ *cpXP* is significantly greater than *E. coli* MG1655 (Student's t-test, $P < 0.05$) (Fig. 1). The diffusion diameters of the overexpression transformants are significantly smaller than *E. coli* MG1655 (Student's t-test, $P < 0.05$). These results suggest that *cpXP* gene had a significant effect on the movement ability of *E. coli* strain.

Bacteriostatic experiment

To investigate the affect of the CpxP proteins on the antibacterial, the diameters of inhibition zone were determined. The results (Fig. 2) showed that the inhibition effect of CpxP proteins was significantly greater than ddH₂O (Student's t-test, $P < 0.01$), and there was significant difference compared with Kan (Student's t-test, $P < 0.05$) (Table 1). These results suggest that CpxP proteins had a significant inhibition of bacterial activity.

Molecular clustering

Multi-sequence alignment and molecular cluster was carried out for 81 CpxP proteins sequences. The results indicated that CpxP were roughly divided into three categories (Fig. 3). The protein sequence of the *E. coli* CpxP (BAJ45639) exhibited 100%, 100% and 89.2% homology with the *Shigella flexneri* (AUU33472), *Shigella sonnei* (AMG18626) and *Salmonella enterica* (AMG25549) homologous proteins, respectively. All the bacterial

names corresponding to the gi numbers and sequences were showed (Additional file 1: Table S1).

Discussion

The cell-envelope stress modulator *cpXP* (periplasmic protein) gene has been investigated for many years, but there are few studies on its function. The CpxP proteins can inhibit activation of CpxA and are indispensable for the quality control system of P pili.

In this report, the results showed that the *cpXP*-overexpression *E. coli* MG1655, the knocked out mutants *E. coli* MG1655- Δ *cpXP* and the revertants *E. coli* MG1655- Δ *cpXP* (pBBR-*cpXP*) were obtained. Deprivation of the *cpXP* gene resulted in significant enhancement in the mobility of *E. coli* strains. The overexpression of the *cpXP* gene also resulted in significant attenuation in the mobility of *E. coli* strains. The mobility of *E. coli* revertants strains was lower than *E. coli* MG1655. The mobility of bacteria had important pathological significance, moreover, and mainly played its role during the early stage of the infection (Mao and He 1998). In an experimental urinary tract infection of the mouse, colonization of the urinary bladder by isogenic strains of *Salmonella enterica* serovar Typhimurium was found to depend on the motility of the bacteria (Siitonen and Nurminen 1992). Our results suggest that the deprivation of the *cpXP* gene resulted in significant enhancement in the mobility and infection of *E. coli* strains.

Multi-sequence alignment and molecular cluster indicated that the CpxP proteins had a high homology at protein level with *Shigella flexneri*, *Shigella sonnei* and *Salmonella enteric* which are the main pathogenic bacteria in China. Our results suggest that the overexpressions of the *cpXP* gene may significantly reduce the pathogenicity of these bacteria.

Supplementary information

Supplementary information accompanies this paper at <https://doi.org/10.1186/s13568-020-01099-z>.

Additional file 1. The electrophoregram and sequences of *cpXP*.

Acknowledgements

The research was supported by Natural Science Foundation of China (Nos. 31100044, 31270077), "100 Talent Program" of Hebei Province (No. E2012100005), Scientific Research Foundation for the Returned Overseas Chinese Scholars (1792-47) and Hebei Provincial Natural Science Fund (No. C2014208093, 17223608D, Key Project No. C2020208020).

Authors' contributions

XH and XZ conceived and designed research. YR, WM and XY conducted experiments. XZ contributed the reagents. XH and YR analyzed data. XH wrote the manuscript. All authors read and approved the final manuscript.

Funding

Not applicable.

Availability of data and materials

Not applicable.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

All authors declare that he has no conflict of interest.

Received: 19 May 2020 Accepted: 27 August 2020

Published online: 26 September 2020

References

- Cobb RE, Wang Y, Zhao H (2014) High-efficiency multiplex genome editing of *Streptomyces* species using an engineered CRISPR/Cas system. *ACS Synth Biol* 4(6):723–728
- Cong L, Ran FA, Cox D, Lin S, Barretto R, Habib N, Hsu PD, Wu X, Jiang W, Marraffini LA, Zhang F (2013) Multiplex genome engineering using CRISPR/Cas systems. *Science* 339:819–823
- DiCarlo JE, Norville JE, Mali P, Rios X, Aach J, Church GM (2013) Genome engineering in *Saccharomyces cerevisiae* using CRISPR-Cas systems. *Nucleic Acids Res* 41:4336–4343
- DiCarlo JE, Chavez A, Dietz SL, Esvelt KM, Church GM (2015) Safeguarding CRISPR-Cas9 gene drives in yeast. *Nat Biotechnol* 33:1250–1255
- Dong J, Iuchi S, Kwan HS, Lu Z, Lin ECC (1993) The deduced amino-acid sequence of the cloned CpxR gene suggests the protein is the cognate regulator for the membrane sensor, CpxA, in a two-component signal transduction system of *Escherichia coli*. *Gene* 136(1):227–230
- Dong H, Xiang H, Mu D, Wang D, Wang T (2019) Exploiting a conjugative CRISPR/Cas9 system to eliminate plasmid harbouring the mcr-1 gene from *Escherichia coli*. *Int J Antimicrob Agents* 53(1):1–8
- Fällman E, Schedin S, Jass J, Uhlin BE, Axner O (2005) The unfolding of the pili quaternary structure by stretching is reversible, not plastic. *EMBO Rep* 6(1):52–56
- Gantz VM, Bier E (2015) The mutagenic chain reaction: a method for converting heterozygous to homozygous mutations. *Science* 348:442–444
- Gantz VM, Jasinskiene N, Tatarenkova O, Fazekas A, James AA (2015) Highly efficient Cas9-mediated gene drive for population modification of the malaria vector mosquito *Anopheles stephensi*. *Proc Natl Acad Sci USA* 112:E6736–E6743
- Grunwald HA, Gantz VM, Poplawski G, Xu XS, Bier E, Cooper KL (2019) Super-Mendelian inheritance mediated by CRISPR-Cas9 in the female mouse germline. *Nature* 566:105–109
- Hammond A, Galizi R, Kyrou K, Simoni A, Siniscalchi C, Katsanos D (2016) A CRISPR-Cas9 gene drive system targeting female reproduction in the malaria mosquito vector *Anopheles gambiae*. *Nat Biotechnol* 34:78–83
- He XL, Zhou XH, Yang Z, Xu L, Yu YX, Jia LL, Li GQ (2015) Cloning, expression and purification of d-tagatose 3-epimerase gene from *Escherichia coli* JM109. *Protein Express Purif* 114:77–81
- Jiang W, Bikard D, Cox D, Zhang F, Marraffini LA (2013) RNA-guided editing of bacterial genomes using CRISPR-Cas systems. *Nat Biotechnol* 31:233–239
- Mali P, Yang L, Esvelt KM, Aach J, Guell M, DiCarlo JE, Norville JE, Church GM (2013) RNA-guided human genome engineering via Cas9. *Science* 339:823–826
- Mao GZ, He LY (1998) Relationship of wild type strain motility and interaction with host plants in *Ralstonia solanacearum*. *Bacterial wilt disease*. Springer, Berlin, pp 184–191
- Nevesinjac AZ, Raivio TL (2005) The cpx envelope stress response affects expression of the type iv bundle-forming pili of enteropathogenic *Escherichia coli*. *J Bacteriol* 187(2):672–686
- Ruiz N, Silhavy TJ (2005) Sensing external stress: watchdogs of the *Escherichia coli* cell envelope. *Curr Opin Microbiol* 8:122–126
- Shan Q, Wang Y, Li J, Zhang Y, Chen K, Liang Z, Zhang K, Liu J, Xi JJ, Qiu JL, Gao C (2013) Targeted genome modification of crop plants using a CRISPR-Cas system. *Nat Biotechnol* 31:686–688
- Sitonen A, Nurminen M (1992) Bacterial motility is a colonization factor in experimental urinary tract infection. *Infect Immun* 60(9):3918–3920
- Theede GL, Arthur DC, Edwards RA, Buelow DR, Wong JL, Raivio TL (2011) Structure of the periplasmic stress response protein CpxP. *J Bacteriol* 193(9):2149–2157
- Wang Y, Li Z, Xu J, Zeng B, Ling L, You L, Chen Y, Huang Y, Tan A (2013) The CRISPR/Cas System mediates efficient genome engineering in *Bombyx mori*. *Cell Res* 23:1414–1416
- Yu Z, Ren M, Wang Z, Zhang B, Rong YS, Jiao R, Gao G (2013) Highly efficient genome modifications mediated by CRISPR/Cas9 in *Drosophila*. *Genetics* 195:289–291
- Zhang Q, Rho M, Tang H, Doak TG, Ye Y (2013) CRISPR-Cas systems target a diverse collection of invasive mobile genetic elements in human microbiomes. *Genome Biol* 14:R40
- Zhou X, Keller R, Volkmer R (2011) Structural basis for two-component system inhibition and pilus sensing by the auxiliary CpxP protein. *J Biol Chem* 286(11):9805–9814

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](https://www.springeropen.com)
