



Analysis of circRNA-miRNA-mRNA Regulatory Network in Peripheral Blood of Radiation Workers

Dose-Response:
An International Journal
April-June 2022:1–10
© The Author(s) 2022
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/115593258221088745
journals.sagepub.com/home/dos


Jin Gao^{1,2,†} , Tinxi Lan^{1,2,†}, Xumin Zong^{1,2,†}, Gensheng Shi^{1,2}, Shuqing He^{1,3}, Na Chen^{1,2}, Fengmei Cui^{1,2}, and Yu Tu^{1,2,*}

Abstract

The health of radiation workers has always been our focus. Epidemiological investigation shows that long-term exposure to low-dose ionizing radiation can affect human health, especially cancer and cardiovascular disease, and there are many studies on it. However, up to now, there have been few reports on the research of blood and biological samples from radiation workers. In this study, radiation workers and healthy control groups were strictly screened, and the transcriptome of mRNA and circRNA was sequenced by extracting their peripheral venous blood. At the same time, appropriate data sets were selected in the GEO database for bioinformatics analysis, and circRNA-miRNA-mRNA network was constructed. We identified 9 different circular ribonucleic acids, 3 tiny ribonucleic acids, and 2 central genes (NOD 2 and IRF 7). These differentially expressed genes and non-coding RNA are closely related to ionizing radiation damage, and play an important role as biological markers. In conclusion, this study may provide new insights into the role of the circRNA-miRNA-mRNA regulatory network in the health of radiation workers, and provides a new strategy for the future study of radiation biology.

Keywords

radiologist, RNA deep sequencing, bioinformatics, biomarkers of ionizing radiation, regulatory network

Introduction

The peripheral blood examination of radiation workers is an indispensable examination in the current occupational health examination. By analyzing a series of indicators such as blood routine, blood biochemical changes, chromosome aberration, and micronucleus rate (quoted), it can be judged whether the workers suffer from radiation diseases and whether they are suitable for the current radiation posts.^{1,2} However, in most cases, long-term exposure to low-dose ionizing radiation may be within normal range of peripheral blood examination, which may lead to neglect of potential disease risk.³ The risks of disease caused by long-term low-dose ionizing radiation usually have a long incubation period. If the changes of peripheral blood transcriptomics are used to judge the exposure level of workers, the possibility of continuing to engage in radioactive work and the possibility of suffering from certain diseases, it will be better to monitor the health status of radiation work.^{3,4} In recent years, with the development of transcriptome

technology, it is possible to detect non-coding RNA and mRNA in blood samples.⁵ Therefore, in order to better protect the health of radiation workers, it is necessary to systematically analyze the changes of transcriptome of blood samples of radiation workers, hoping to find possible potential markers and better determine their health status.

¹ State Key Laboratory of Radiation Medicine and Protection, School of Radiation Medicine and Protection, Soochow University, Suzhou, China

² Collaborative Innovation Center of Radiation Medicine of Jiangsu Higher Education Institutions, Suzhou, China

³ School of Public Health, Medical College of Soochow University, Suzhou, China

Received 2 February 2022; accepted 17 February 2022

[†]Jin Gao, Tinxi Lan, Xumin Zong contributed equally to this work.

Corresponding Author:

Yu Tu, Department of Radiation Medicine, School of Radiation Medicine and Protection, Soochow University, Suzhou 215123, China.
Email: tuyu@suda.edu.cn



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE

and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

CircRNA is a specific endogenous non-coding RNA, which has evolved and preserved in eukaryotic species and is widely expressed in high abundance human cells. The closed-loop structure makes the cyclic ribonucleic acids more stable and protects them from degradation by ribonuclease.⁶ With the increasingly extensive application of high-throughput sequencing and microarray technology, *noncoding-RNA* have been proved to be related to ionizing radiation is increasingly.⁷⁻⁹ It was found that circRNA could be used as microRNA sponge, protein bait and transporter to regulate gene expression.¹⁰ Existing evidence shows that both cyclic ribonucleic acid and micro ribonucleic acid are involved in the pathogenesis and progress of ionizing radiation damage, and play an important role in them.¹¹⁻¹³

At present, there are few studies on the characteristics of regulatory network of circRNA-miRNA-mRNA in blood samples of radiation workers. Therefore, this study was designed to investigate the potential circRNA-miRNA-mRNA regulatory network in the peripheral blood of radiation workers. Genome-wide cyclic ribonucleic acid and gene microarray analysis was performed to screen out differentially expressed cyclic ribonucleic acid and genes. Through the ionizing radiation mined in GEO database and the change of micro ribonucleic acid in peripheral blood, the differentially expressed micro ribonucleic acid can be obtained. Prediction and construction of regulatory network of circRNA-miRNA-mRNA. Gene function enrichment analysis of differential genes was conducted to identify their potential function. Our results provide new information on the role of the regulatory network of circRNA-miRNA-mRNA in the health examination of radiation workers and provide a theoretical basis for the interaction mechanism of circRNA, miRNA, and mRNA as biological dose markers and possible disease effects for radiation workers.

Methods

Study Objects

According to the questionnaire scheme in *Specification of survey technique for radiation epidemiology*, select the appropriate control group and radiation group in the designated hospitals where the industrial radiation workers have occupational health check.¹⁴ With the informed consent form signed, 5 mL of peripheral blood was extracted from the enrolled personnel for subsequent transcriptomic sequencing and RT-qPCR.

Preparation and Sequence of RNA Library

According to the instructions, ribosomal RNA (rRNA) was removed from samples by using Nebner Extranadplenion Kit (New England Bioabs, Inc., Massachusetts, USA). The sequencing library was constructed by using Nebnext-miRNA library preparation kit (New England BioABS, Inc., Massachusetts, USA). The library was controlled and quantified by the bioanalyzer 2100 system (Agilent Technologies, USA), and 150 bp double-ended sequencing was carried out by Illumina HiSeq instrument.

Differentially Expressed CircRNAs and Differentially Expressed Genes Data Analysis

Sequencing was carried out by an Illumina HiSeq 4000 sequencer, and two-terminal readings were obtained. Q30 was used for quality control, and the cut adapt software (v1.9.3) was used to disconnect and disconnect low quality readings to obtain high quality readings. (1). circRNA: High-quality reads were compared to the reference genome/transcriptome using STAR software (v2.5.1b), and circRNA was detected and identified using DCC software (v0.4.4). The identified circRNA was also annotated with the circBase database (<http://circbase.org/>) and Circ2Traits.¹⁵ EdgeR software (v3.16.5) was used for data standardization and differential expression. The circRNA passing through fold change >2 and $P < .05$ was selected as the differentially expressed circRNAs (DECs). (2). mRNA: hisat2 software (v2.0.4) was used to compare high-quality reads to the human reference genome (UCSC HG19). Then, under the guidance of gtf gene annotation file, the FPKM value of mRNA at gene level was obtained as the expression spectrum of mRNA using cuffdiff software (v2.2.1, part of the cufflinks software suite). Fold change >2 and $P < .05$ were calculated between the 2/group samples to screen for differentially expressed genes (DEGs).

Gene Function Analysis

In order to further understand the function and main functional pathways of mRNA, all the differential functional pathways of mRNA are annotated and analyzed. The mRNA obtained from the above screening was put into David database (<https://david.cifcrf.gov/>) and the species was selected as "*Homo sapiens*." The threshold $P < .05$ was set for GO enrichment and KEGG pathway annotation analysis, and the enrichment analysis results were visualized.

DEMs in Peripheral Blood After Ionizing Radiation

Microarray data sets of miRNA expression profiles in peripheral blood following ionizing radiation were obtained from the GEO database (<https://www.ncbi.nlm.nih.gov/geo/>), GSE55954. The miRNA expression profiles of human peripheral blood mononuclear cells in the experimental design at 20 h after exposure to 60 Gy of Cs-137 γ -ray were selected. The experiment was carried out independently for 4 times, and the system error was eliminated. The data were analyzed to extract differentially expressed miRNA using the GEO2R program provided by the GEO database. The miRNAs with fold change >2 and $P < .05$ were selected as differentially expressed miRNAs (DEMs).

Predicting the Target miRNA of DECs and Predicting the Target Genes of DEMs

Using circMir2.0 and mirTargets2.0 software (Nanjing medical university). The software included Starbase (<https://starbase.sysu.edu.cn>), MIRREDB (<http://mirdb.org/cgi-bin>),

and Target (<http://www.targetscan.org>) databases to predict circRNA-miRNA and miRNA-mRNA interactions and selected predicted results with an interaction score greater than 80 and a $P < .05$ for final validation.

Construction of circRNA-miRNA-mRNA Network in Peripheral Blood of Radiation Workers

We extracted the circRNA/miRNA/mRNA competition constraint. The circRNA-miRNA and miRNA-mRNA with potential binding were selected from the predicted differential genes, and the selected DECs, DEMIRs, and DEGs were mapped to construct a circRNA-miRNA-mRNA interaction network, which was visualized using Cytoscape3.8.2 software.¹⁶

RNA Extraction and RT-qPCR

Use HI SCRIPT II first strand cDNA synthesis kit (+GDNA Wiper) kit (Vazyme) to reverse transcribe circRNA and mRNA, and the HI SCRIPT II 1st strand cDNA Synthesis Kit (+GDNA Wiper) kit (Vazyme) were used in reverse transcribed. Use the miRNA first strand cDNA synthesis kit (Bystem-loop) p (Vazyme) to synthesize miRNA reverse transcription kit cDNA. RT-qPCR was used to evaluate the expression of circRNA, miRNA and mRNA, and verified by Chamq universal sybr qpcr master mix kit with PCR instrument (ViiA 7, Life Technologies). GAPDH was used as the internal control for circRNAs and mRNAs, and U6 was used as the internal control for miRNAs. Primers for circRNA and miRNA were designed by CircInteract (<https://circinteractome.nia.nih.gov/>) and miRNA Design V1 (Vazyme), respectively, and primers for mRNA were found in PrimerBank. The cycle threshold ($2^{-\Delta\Delta Ct}$) was compared for calculation. The primers applied are displayed in Table S1.

Statistical Analysis

All statistical analyses were conducted with SPSS 20.0. Independent t test was used to compare continuous variables with normal distribution, Mann-Whitney U test was used to compare continuous variables with skewed distribution, and chi-square test was used to test the ratio. All statistical tests are bidirectional.

Result

Basic Information of People Involved in Gene Sequencing

Transcriptome sequencing: ten normal staff who came to the hospital for physical examination and 12 exposed people were selected. The average total dose of these 12 people at work was 63.3 mSv, and the average working life was 9 years. All the 22 men were male. Except for salt intake, age, smoking

history, eating habits and other influencing factors had no statistical difference ($P < .05$). There were 10 people in the control group and 10 people in the radiotherapy group, male and female were equally divided. The radiation group received an average external radiation dose of 21.4 mSv, and the average working life was 5 years. There was no statistical difference in baseline. The above radiation workers are all involved in X-ray operation. The details are shown in Table 1.

RNA and Library Quality Control

The RNA concentration of each sample was measured using a NanoDrop ND-1000 instrument (Thermo Fisher Scientific, Waltham, MA, USA). The OD₂₆₀/OD₂₈₀ value was used as the RNA purity index. RNA purity was acceptable when the OD₂₆₀/OD₂₈₀ value ranged from 1.8 to 2.1; Degenerated agarose gel electrophoresis was used to measure RNA integrity and gDNA contamination, as shown in Figure 1. The quality of the library was measured by Agilent 2100 Bioanalyzer (Table 2).

Differential Transcriptome Results


According to the criteria of foldchange>2 and $P < .05$ between samples, the differential results were screened. Compared with the control group, there were 1149 differentially expressed circRNAs in the radiation group in this study, of which 639 were up-regulated and 510 were down-regulated. A total of 873 differentially expressed genes were identified, of which 68 were up-regulated and 805 were down-regulated. Compared with the control group, in the GSE55954 dataset selected from GEO database, there were 229 differentially expressed miRNA in peripheral blood after irradiation, of which 117 were up-regulated and 112 were down-regulated (Figure 2).

Functional Enrichment Analysis

To understand the biological action and potential function of 873 differential genes, GO analysis was performed, including biological process (BP), cellular component (CC) and molecular function (MF), and KEGG signaling pathway enrichment analysis. GO analysis showed that in terms of BP, these overlapping genes were mainly enriched in the positive regulation of cellular metabolic process, apoptotic process involving in patterning of blood vessels. positive regulation of ERN protein response, cell-substrate junction assembly, positive regulation of cyto-me signaling pathway. For CC, the overlapping genes were mainly enriched in pore complex, MOZ/MORF histone acetyltransferase complex, actomyosin, cyclin-dependent protein kinase holoenzyme complex, and stress fiber. In the case of MF, these overlapping genes were mainly enriched in N-acyl-s phospholipase D activity, phospholipase D activity, G-quadruplex DNA binding, 3-chloroallyl aldehyde dehydrogenase activity, and vinculin binding.

Table 1. Baseline Survey Results.

Type	Data for Transcriptome Sequencing			Data for RT-qPCR		
	Control	Radiation	P	Control	Radiation	P
Age	48.60 ± 4.79	50.50 ± 5.67	.411	29.95 ± 1.78	30.18 ± 2.45	.819
Smoke	50.00%	91.70%	.056	40.00%	50.00%	.653
Alcohol	50.00%	58.60%	.696	40.00%	50.00%	.653
Tea	70.00%	91.70%	.293	20.00%	20.00%	.999
Rice	10.00%	0	.455	40.00%	30.00%	.639
Vegetable	40.00%	66.70%	.436	40.00%	30.00%	.639
Oil	70.00%	58.30%	.064	60.00%	50.00%	.653
Sugar	20.00%	41.70%	.424	50.00%	40.00%	.653
Chili	20.00%	41.70%	.485	60.00%	50.00%	.653
Salted	50.00%	75.00%	.44	60.00%	80.00%	.329
Fruit	30.00%	66.70%	.128	50.00%	50.00%	.999
Salt	60.00%	33.30%	.033	70.00%	60.00%	.639
Meat	40.00%	66.70%	.431	60.00%	70.00%	.639

**Figure 1.** Agarose gel electrophoresis of total RNA.**Table 2.** RNA Library Quality Control.

Sample ID	Sample Name	OD _{260/280} Ratio	Conc. (ng/gl)	Volume (gl)	Quantity (gg)
1	A1	2.04	149.26	9	1.34
2	A2	1.86	139.39	9	1.25
3	A3	1.80	182.05	9	1.64
4	A4	1.83	180.54	9	1.62
5	A5	1.86	122.99	9	1.11
6	A6	1.84	224.62	9	2.02
7	A7	1.83	167.14	9	1.50
8	A8	1.88	278.76	9	2.51
9	A9	1.89	184.34	9	1.66
10	A10	2.07	204.26	9	1.84
11	B2	1.86	120.01	9	1.08
12	B3	1.87	210.60	9	1.90
13	B4	1.90	204.18	9	1.84
14	B5	1.89	152.58	9	1.37
15	B7	1.84	106.18	9	.96
16	B9	1.84	250.71	9	2.26
17	B10	1.85	116.05	9	1.04
18	B11	1.87	142.57	9	1.28
19	B12	1.87	113.77	9	1.02
20	B13	1.86	113.03	9	1.02
21	B14	1.86	121.76	9	1.10
22	B15	1.81	144.55	9	1.30




Figure 2. Summary of variance expression data. (A): circRNA volcano map of differential expression between the control and radiology staff blood samples. (B): Volcano map of differentially expressed mRNA between blood samples of radiation workers in the control group and irradiated group. (C): Volcano diagram showing differential expression of miRNA in peripheral blood after CS-137 γ -ray compared with that in the control group. (D): Summary of up and down regulated genes.

Enrichment analysis of KEGG signaling pathway revealed that DEGs was mainly enriched in their types of O-glycan biosynthesis, Pertussis, Leishmaniasis, *Staphylococcus aureus* infection, and Propanoate metabolism (Figure 3).

Construction of circRNA-miRNA-mRNA Regulatory Network

We obtained the first 20 hub genes using the cytoHubba plug-in of Cytoscape3.8.2 software (Figure 4B), and screened the functional module with the MCODE plug-in to select the first module with the highest score (Figure 4C), and selected a total of 9 genes screened by 2 methods to construct a CIRC RNA-miRNA-mRNA regulatory network (Figure 4D). Finally, we obtained a circRNA-miRNA-mRNA regulatory network

consisting of 9 DECs, 3 overlapping miRNAs and 2 target mRNA (Figure 4E). For further details see Table S2.

According to the results of RT-qPCR, when compared with the control group, only hsa_circ_0093865 and hsa_circ_0005940 showed no statistical differences in expression levels, and other results were consistent with the results of transcriptomics (Figure 5).

Discussion

Occupational health examination of radiation workers is very important for finding out whether there are defects in their health status in time. For classical examination items such as chromosome micronucleus and chromosome aberration, it can be observed whether the dose is still suitable for the current




Figure 3. Summary of gene function enrichment. (A): Biological process analysis. (B): Cellular component analysis. (C): Molecular function analysis. (D): KEGG pathway analysis.

post.¹⁷ However, it is not clear whether the occupational health of radiation workers will be affected. Epidemiological results show that long-term low-dose ionizing radiation may affect the health of radiation workers, such as cancer, cardiovascular, and cerebrovascular diseases, etc.^{18,19} Therefore, it is very important to find key molecules and markers. At present, the general peripheral blood examination focuses more on judging the health status of the subjects through the changes of blood cells and blood biochemistry, while the potential genetic changes and changes of non-coding RNA are rarely involved. More and more researchers are involved in the research of non-coding RNA, because these non-coding RNA are involved in the regulation of many diseases, including the research of radioactive diseases and radioactive biosimeters.

Researchers found that circRNA was associated with many biomarkers of ionizing radiation injury, such as mouse brain and primary neuron injury, mouse bone marrow injury, and radiation-related esophageal injury.^{20,21,11} In addition, it is also found that circRNA is related to a variety of radiation

diseases, such as hsa_circ_0001649 participating in the following regulation of migration and proliferation of cholangiocarcinoma cells after radiotherapy, and hsa_circRNA_100367 regulating radiosensitivity of esophageal cancer through Wnt pathway.^{22,23} The miRNA is an evolutionary conservative small molecule with a length of 18–25 nucleotides. In recent years, there are many studies related to miRNA ionizing radiation, such as miR-29 participating in ionizing radiation-induced fibrosis through the overexpression of inhibitory type I collagen, miR-92b participating in the regulation of susceptibility of liver cancer to ionizing radiation therapy, miR-375-3 p as a marker of acute radiation syndrome, and miR-34 a as a biomarker of ionizing radiation.²⁴⁻²⁷ These studies show that the research of non-coding ribonucleic acid is very important for ionizing radiation damage and health concerns of radiation workers.

Through literature search, we explored the latest research progress and discovery of *NOD2*, *IRF7*, miR-671-5p, and miR-654-5p in circRNA-miRNA-miRNA network that was finally discovered. Nucleotide-binding oligomerisation




Figure 4. Regulation process of constructing circRNA/miRNA/mRNA network. (A): Analysis of differential gene protein interaction network. (B): top20 hub genes of cytoHubba. (C): Screening first module of MCODE. (D): Total circRNA/miRNA/mRNA network regulation. (E): Regulation of circRNA/miRNA/mRNA network acquired by up-and down-regulation of binding genes. Validation of correlation network regulation in blood samples of radiation workers.




Figure 5. RT-qPCR results. (A): the expression level of circRNAs. (B): expression level of miRNAs. (C): mRNA expression level. * means $P < .05$, and ns means $P > .05$.

domain protein 2 (*NOD2*) participates in studies of ionizing radiation by influencing the regeneration of intestinal epithelial cells damaged by radiation. At the same time, *NOD2* alleviates radiation-induced damage through the agonist of ATR-mediated DNA damage response pathway, Morabutanol, and also inhibits oxidative stress and apoptosis by blocking the participation of wow/ROCK-mediated TAK1/*NOD2* in NF- κ B pathway.²⁸⁻³⁰ Meanwhile, *NOD2* also plays an important role in Parkinson's disease and intestinal disease.^{31,32} Interferon regulatory factor 7 (IRF7) is highly expressed in the apoptosis of melanoma cell after ultraviolet irradiation, and it is used as a reactive molecular indicator of radiation-induced injury of human umbilical vein endothelial cells.^{33,34} Of course, IRF7 has been proved to play a role in virus-induced cellular gene transcription activation, and its encoded protein plays an important role in the innate immune response against deoxyribonucleic acid and ribonucleic acid virus.³⁵ The miR-671-5p can be used as a biomarker to detect the therapeutic effect of radiotherapy on colorectal cancer,³⁶ furthermore, the invention can also be used as a new biomarker for early breast cancer detection and a target for breast cancer treatment; It also inhibits the progression of glioblastoma in circ_0001946/miR-671-5p/CDR1 network regulation.^{37,38} miR-654-5p can target and regulate EPSTI1 to slow the progression of breast cancer; in addition, it can inhibit the MYC, WNT, and AKT pathways from affecting the development of ovarian cancer.^{39,40} In our study, all circRNAs and miR-3616-3p were both reported for the first time, and we hope to find out their functions through further experiments at the later stage.

In conclusion, we have constructed a circRNA-miRNA-mRNA network by comparing the expression of circRNAs and mRNAs in peripheral blood of radiation workers and healthy control, and the ionizing radiation-related miRNAs obtained from the GEO database. Subsequently, I constructed a circRNA-miRNA-miRNA network consisting of 9 circRNAs (And the expression levels of the 7 circRNAs were consistent with transcriptomics), 4 miRNAs and 2 hub genes using a variety of different bioinformatics methods. This research can provide some help for the development of biological dosimeters for future dose estimation of radiation workers and the possible mechanism of radiation-related diseases. Of course, this needs further experimental verification in the later period.

Acknowledgments

We thank CNNC 401 Hospital for the great help. Thanks to my dear Duobao for her support.

Authors' Note

Conceived and designed the experiment: Yu Tu. Performed the experiments: Jin Gao, Tinxi Lan, Xumin Zong, Gensheng Shi, Shuqing He, Na chen, Fengmei Cui. Analysed the data: Jin Gao Tinxi Lan and Xumin Zong. Wrote the paper: Jin Gao.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by the Natural Science Research Projects of Colleges and Universities in Jiangsu Province (grant numbers 20KJB310007), the Priority Academic Program Development of Jiangsu Higher Education Institutions (PAPD), the Collaborative Innovation Center of Radiological Medicine of Jiangsu Higher Education Institutions, China and the Nuclear Energy Development Project, China (No. 2016-1295).

ORCID iD

Jin Gao  <https://orcid.org/0000-0003-3519-5832>

Supplemental Material

Supplemental material for this article is available online.

References

1. ICRP. The 2007 Recommendations of the International Commission on Radiological Protection. ICRP publication 103. *Ann ICRP*, 2007;37(2-4):1-332. doi:10.1016/j.icrp.2007.10.003.
2. Su YW, Wang JY, Zhang Y, et al. Effect of low dose ionizing radiation on peripheral blood cells of radiation workers in nuclear power industry. *Zhonghua Lao Dong Wei Sheng Zhi Ye Bing Za Zhi*. 2020;38(8):633-635. doi:10.3760/cma.j.cn121094-20190314-00100.
3. Leuraud K, Richardson DB, Cardis E, et al. Risk of cancer associated with low-dose radiation exposure: Comparison of results between the INWORKS nuclear workers study and the A-bomb survivors study. *Radiat Environ Biophys*. 2021;60(1):23-39. doi:10.1007/s00411-020-00890-7.
4. Kumar D, Kumari S, Salian SR, et al. Genetic instability in lymphocytes is associated with blood plasma antioxidant levels in health care workers occupationally exposed to ionizing radiation. *Int J Toxicol*. 2016;35(3):327-335. doi:10.1177/1091581815625593.
5. Tomita K, Fukumoto M, Itoh K, et al. MiR-7-5p is a key factor that controls radioresistance via intracellular Fe(2+) content in clinically relevant radioresistant cells. *Biochem Biophys Res Commun*. 2019;518(4):712-718. doi:10.1016/j.bbrc.2019.08.117.
6. Chen L, Zhou H, Guan Z. CircRNA_000543 knockdown sensitizes nasopharyngeal carcinoma to irradiation by targeting miR-9/platelet-derived growth factor receptor B axis. *Biochem Biophys Res Commun*. 2019;512(4):786-792. doi:10.1016/j.bbrc.2019.03.126.
7. Yadav M, Bhayana S, Liu J, et al. Two-miRNA-based fingerstick assay for estimation of absorbed ionizing radiation dose. *Sci Transl Med*. 2020;12(552):eaaw5831. doi:10.1126/scitranslmed.aaw5831.

8. Borghini A, Vecoli C, Mercuri A, et al. Low-dose exposure to ionizing radiation deregulates the brain-specific MicroRNA-134 in interventional cardiologists. *Circulation*. 2017;136(25):2516-2518. doi:10.1161/CIRCULATIONAHA.117.031251.
9. Deng L, Liang H, Xu M, et al. STING-dependent cytosolic dna sensing promotes radiation-induced type I interferon-dependent antitumor immunity in immunogenic tumors. *Immunity*. 2014;41(5):843-852. doi:10.1016/j.immuni.2014.10.019.
10. Yu L, Liu Y. circRNA_0016624 could sponge miR-98 to regulate BMP2 expression in postmenopausal osteoporosis. *Biochem Biophys Res Commun*. 2019;516(2):546-550. doi:10.1016/j.bbrc.2019.06.087.
11. Luo J, Zhang C, Zhan Q, et al. Profiling circRNA and miRNA of radiation-induced esophageal injury in a rat model. *Sci Rep*. 2018;8(1):14605. doi:10.1038/s41598-018-33038-1.
12. May JM, Bylicky M, Chopra S, et al. Long and short non-coding RNA and radiation response: A review. *Transl Res*. 2021;233:162-179. doi:10.1016/j.trsl.2021.02.005.
13. He N, Sun Y, Yang M, et al. Analysis of circular RNA expression profile in HEK 293T cells exposed to ionizing radiation. *Dose Response*. 2019;17(2):155932581983779. doi:10.1177/1559325819837795.
14. Shen YP, Qu SG, Chen DW, et al. *T/CNS 7—2018 Specification of survey technique for radiation epidemiology*. Beijing: China atomic energy authority; 2018.
15. Ghosal S, Das S, Sen R, et al. Circ2Traits: A comprehensive database for circular RNA potentially associated with disease and traits. *Front Genet*. 2013;4:283. doi:10.3389/fgene.2013.00283.
16. Shannon P, Markiel A, Ozier O, et al. Cytoscape: A software environment for integrated models of biomolecular interaction networks. *Genome Res*. 2003;13(11):2498-2504. doi:10.1101/gr.1239303.
17. Cool DA, Kase KR, Boice JD. NCRP report no.180-management of exposure to ionizing radiation: NCRP radiation protection guidance for the United States. *J Radiol Prot*. 2019;39(3):966-977. doi:10.1088/1361-6498/ab1826.
18. Boice JD, Cohen SS, Mumma MT, et al. Mortality from leukemia, cancer and heart disease among U.S. nuclear power plant workers, 1957-2011. *Int J Radiat Biol*. 2022;2022:1-22. doi:10.1080/09553002.2021.1967507.
19. Kreuzer M, Auvinen A, Cardis E, et al. Low-dose ionising radiation and cardiovascular diseases—Strategies for molecular epidemiological studies in Europe. *Mutat Res Rev Mutat Res*. 2015;764:90-100. doi:10.1016/j.mrrev.2015.03.002.
20. Mfossa ACM, Puthenparampil HT, Inalegwu A, et al. Exposure to ionizing radiation triggers prolonged changes in circular RNA abundance in the embryonic mouse brain and primary neurons. *Cells*. 2019;8(8):778. doi:10.3390/cells8080778.
21. Zhang J, Jiang J, Huang R, et al. Circular RNA expression profiles are significantly altered in mice bone marrow stromal cells after total body irradiation. *Leuk Res*. 2018;70:67-73. doi:10.1016/j.leukres.2018.05.010.
22. Xu Y, Yao Y, Zhong X, et al. Downregulated circular RNA hsa_circ_0001649 regulates proliferation, migration and invasion in cholangiocarcinoma cells. *Biochem Biophys Res Commun*. 2018;496(2):455-461. doi:10.1016/j.bbrc.2018.01.077.
23. Liu J, Xue N, Guo Y, et al. CircRNA_100367 regulated the radiation sensitivity of esophageal squamous cell carcinomas through miR-217/Wnt3 pathway. *Aging (Albany NY)*. 2019;11(24):12412-12427. doi:10.18632/aging.102580.
24. Yano H, Hamanaka R, Nakamura-Ota M, et al. Regulation of type I collagen expression by microRNA-29 following ionizing radiation. *Radiat Environ Biophys*. 2018;57(1):41-54. doi:10.1007/s00411-017-0723-4.
25. Wang J, Zhao H, Yu J, et al. MiR-92b targets p57kip2 to modulate the resistance of hepatocellular carcinoma (HCC) to ionizing radiation (IR)-based radiotherapy. *Biomed Pharmacother*. 2019;110:646-655. doi:10.1016/j.biopha.2018.11.080.
26. Chiba M, Monzen S, Iwaya C, et al. Serum miR-375-3p increase in mice exposed to a high dose of ionizing radiation. *Sci Rep*. 2018;8(1):1302. doi:10.1038/s41598-018-19763-7.
27. Halimi M, Shahabi A, Moslemi D, et al. Human serum miR-34a as an indicator of exposure to ionizing radiation. *Radiat Environ Biophys*. 2016;55(4):423-429. doi:10.1007/s00411-016-0661-6.
28. Lee C, Choi C, Kang HS, et al. NOD2 supports crypt survival and epithelial regeneration after radiation-induced injury. *Int J Mol Sci*. 2019;20(17):4297. doi:10.3390/ijms20174297.
29. Liu L, Qu H, Qin H, et al. NOD2 agonist murabutide alleviates radiation-induced injury through DNA damage response pathway mediated by ATR. *J Cell Physiol*. 2019;234(11):21294-21306. doi:10.1002/jcp.28734.
30. Liang J, Zeng X, Halifu Y, et al. Blocking RhoA/ROCK inhibits the pathogenesis of pemphigus vulgaris by suppressing oxidative stress and apoptosis through TAK1/NOD2-mediated NF-kappaB pathway. *Mol Cell Biochem*. 2017;436(1-2):151-158. doi:10.1007/s11010-017-3086-x.
31. Xu M, Zhang S, Zhao H, et al. Association between nucleotide-binding oligomerization domain protein 2 (NOD2) gene polymorphisms and Parkinson's disease (PD) susceptibility. *Artif Cells Nanomed Biotechnol*. 2019;47(1):2134-2138. doi:10.1080/21691401.2019.1617730.
32. Schäffler H, Schneider N, Hsieh C-J, et al. NOD2 mutations are associated with the development of intestinal failure in the absence of Crohn's disease. *Clin Nutr*. 2013;32(6):1029-1035. doi:10.1016/j.clnu.2013.02.014.
33. Zhang K, Anumanthan G, Scheaffer S, et al. HMGB1/RAGE mediates UVB-induced secretory inflammatory response and resistance to apoptosis in human melanocytes. *J Invest Dermatol*. 2019;139(1):202-212. doi:10.1016/j.jid.2018.05.035.
34. Furusawa Y, Zhao Q-L, Hattori Y, et al. Comprehensive and computational analysis of genes in human umbilical vein endothelial cells responsive to X-irradiation. *Genom Data*. 2016;8:126-130. doi:10.1016/j.gdata.2016.05.007.
35. Yao C, Qin J, Wu J-C, et al. Transcription factor networks involved in cell death in the dorsal root ganglia following

- peripheral nerve injury. *Neural Regen Res.* 2018;13(9):1622-1627. doi:[10.4103/1673-5374.237183](https://doi.org/10.4103/1673-5374.237183).
36. Tan X, Li Z, Ren S, et al. Dynamically decreased miR-671-5p expression is associated with oncogenic transformation and radiochemoresistance in breast cancer. *Breast Cancer Res.* 2019; 21(1):89. doi:[10.1186/s13058-019-1173-5](https://doi.org/10.1186/s13058-019-1173-5).
37. Zhu Y, Peng Q, Lin Y, et al. Identification of biomarker microRNAs for predicting the response of colorectal cancer to neoadjuvant chemoradiotherapy based on microRNA regulatory network. *Oncotarget.* 2017;8(2):2233-2248. doi:[10.18632/oncotarget.13659](https://doi.org/10.18632/oncotarget.13659).
38. Li X, Diao H. Circular RNA circ_0001946 acts as a competing endogenous RNA to inhibit glioblastoma progression by modulating miR-671-5p and CDR1. *J Cell Physiol.* 2019; 234(8):13807-13819. doi:[10.1002/jcp.28061](https://doi.org/10.1002/jcp.28061).
39. Tan YY, Xu XY, Wang JF, et al. MiR-654-5p attenuates breast cancer progression by targeting EPSTI1. *Am J Cancer Res.* 2016;6(2):522-532.
40. Majem B, Parrilla A, Jiménez C, et al. MicroRNA-654-5p suppresses ovarian cancer development impacting on MYC, WNT and AKT pathways. *Oncogene.* 2019;38(32):6035-6050. doi:[10.1038/s41388-019-0860-0](https://doi.org/10.1038/s41388-019-0860-0).