Management of malignant pleural effusion with different pleural catheters

Sir,

We were greatly impressed with recent article by Dixit *et al.*^[1] We congratulate the authors for their successful study.

We have been using in recent years the small-bore wire-guided chest drains (Pleurocan, 10F, B. Braun Melsungen, Germany) at increasing rates in our clinic for the treatment of malignant pleural effusions (MPEs). We prefer pleurocan in many MPE cases since it involves a less invasive procedure, maintains patient comfort by causing less pain, and enables making chemical pleurodesis in a similar way.^[2,3] In our practice, after we place a catheter, we connect its end to a urine collection bag or a vacuum drainage bottle. In this way, we ensure both active drainage of the fluid and shortening of the time required for the occurrence of spontaneous pleurodesis. We have been carrying out a study in our clinic in the meantime with fifty patients who were administered pleurocan and in that study, various complications occurred, including loculated effusion in 3 (6%), empyema in 2 (4%), drain blockage in 2 (4%), drain malposition in 1 (2%), and chronic pain in 1 (2%) subject. Despite all these complications, we achieved 94% success rate and 26% spontaneous pleurodesis and we could enable this patient group with relatively short expected survey to spend this time comfortably in a home setting. Moreover, pleurocan is more cost-effective with an average price of €55 in a developing country like Turkey. An indwelling pleural catheter costs around €1750 in our country and vacuum bottles are used in greater numbers, especially in patients with large amounts of drainage, a factor further increasing the cost. Finally, we believe that the use of small-bore wire-guided chest drains in MPE cases is a reliable, well-tolerated, cheap, and effective method.

We congratulate the authors once more for their contribution to this subject. If the authors could report their views and recommendations about different catheter interventions used in MPE in general, they will increase the worth of study and will further encourage surgeons like us who deal with MPE.

Financial support and sponsorship Nil.

Conflicts of interest

There are no conflicts of interest.

Sami Karapolat, Alaaddin Buran, Celal Tekinbas

Department of Thoracic Surgery, Karadeniz Technical University Medical School, Trabzon, Turkey E-mail: samikarapolat@yahoo.com

REFERENCES

- Dixit R, Agarwal KC, Gokhroo A, Patil CB, Meena M, Shah NS, et al. Diagnosis and management options in malignant pleural effusions. Lung India 2017;34:160-6.
- Horsley A, Jones L, White J, Henry M. Efficacy and complications of small-bore, wire-guided chest drains. Chest 2006;130:1857-63.
- 3. Cafarotti S, Dall'Armi V, Cusumano G, Margaritora S, Meacci E, Lococo F, et al. Small-bore wire-guided chest drains: Safety, tolerability, and effectiveness in pneumothorax, malignant effusions, and pleural empyema. J Thorac Cardiovasc Surg 2011;141:683-7.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

Access this article online	
Quick Response Code:	Website: www.lungindia.com
	DOI: 10.4103/lungindia.lungindia_173_17

How to cite this article: Karapolat S, Buran A, Tekinbas C. Management of malignant pleural effusion with different pleural catheters. Lung India 2018;35:92.

© 2017 Indian Chest Society | Published by Wolters Kluwer - Medknow