

REVIEW

Silent sinus syndrome: systematic review and proposal of definition, diagnosis and management

Sindrome del seno silente: revisione sistematica e proposta di definizione, diagnosi e trattamento

Cecilia Rosso¹, Alberto Maria Saibene¹, Giovanni Felisati², Carlotta Pipolo²

¹ Department of Otorhinolaryngology, Santi Paolo e Carlo Hospital, Università degli Studi di Milano, Milan, Italy; ² Department of Otorhinolaryngology, Santi Paolo e Carlo Hospital, Department of Health Sciences, Università degli Studi di Milano, Milan, Italy

SUMMARY

Silent sinus syndrome (SSS) is a rare disease consisting of a collapse of maxillary sinus walls with concomitant orbital floor descent. Due to its rareness, the literature highlights some confusion on its definition, diagnosis and management. A PRISMA-compliant systematic review was performed on SSS with focus on definition, diagnosis and therapeutic management. Twenty-eight studies were selected, with 276 patients evaluated. The analysis revealed that the leading definition of SSS includes evidence of both enophthalmos and maxillary atelectasia. Although the definition of SSS accepts only spontaneous sinus collapse, the presence of sinonasal diseases and history of facial trauma are starting to be included in the criteria. Most studies (n = 21) considered CT scans satisfactory for diagnosis of SSS, while 7 also performed MR. The majority of SSS were successfully treated with isolated functional endoscopic sinus surgery (n = 17), sparing orbital reconstruction as a rescue procedure in case of non-satisfactory long-term resolution of signs. Although the literature is starting to coordinate on diagnosis of SSS, our review revealed the necessity of consensus on its definition and management.

KEY WORDS: silent sinus syndrome, maxillary atelectasia, enophthalmos, imploding antrum syndrome, silent sinus syndrome management

RIASSUNTO

La sindrome del seno silente (SSS) è una rara patologia caratterizzata da collasso delle pareti del seno mascellare con discesa del pavimento orbitario. A causa della rarità di questa sindrome, la letteratura mostra confusione circa la sua definizione, diagnosi e trattamento. È stata condotta una revisione sistematica fedele alle procedure PRISMA sulla SSS con focus su definizione, criteri diagnostici e trattamento (28 studi, 276 pazienti). L'analisi ha mostrato come la definizione predominante della SSS includa l'evidenza sia di enoftalmo che di atelettasia mascellare. Sebbene la definizione originaria accetti solo la presenza di collasso del seno spontaneo, ultimamente la presenza di disturbi nasosinusal e una storia pregressa di trauma facciale vengono inclusi nei criteri diagnostici. La maggior parte degli studi (28) ha considerato le immagini TC sufficienti per la diagnosi radiologica, mentre 7 hanno richiesto anche la RMN.

Il trattamento prevalente è la FESS (Functional Endoscopic Sinus Surgery) (17 studi), lasciando l'intervento di ricostruzione orbitaria come procedura di salvataggio in caso di mancata risoluzione clinica. Sebbene la letteratura inizi a coordinarsi riguardo la diagnosi di SSS, è necessario un consenso riguardo alla definizione e alla gestione terapeutica.

PAROLE CHIAVE: sindrome del seno silente, atelettasia mascellare, sindrome del seno imploso, enoftalmo, gestione sindrome del seno silente

Introduction

Silent sinus syndrome (SSS) or imploding antrum syndrome is a very rare condition, usually consisting of asymptomatic spontaneous collapse of the sinus walls and floor of the orbit ¹. Due to its rarity, the literature has often

Received: April 2, 2021
Accepted: November 14, 2021
Published online: June 30, 2022

Correspondence

Cecilia Rosso
Department of Otorhinolaryngology, San Paolo Hospital, University of Milan, via A. di Rudini 8, 20154 Milan, Italy
E-mail: rosso.cecilia1@gmail.com

How to cite this article: Rosso C, Saibene AM, Felisati G, et al. Silent sinus syndrome: systematic review and proposal of definition, diagnosis and management. *Acta Otorhinolaryngol Ital* 2022;42:305-316. <https://doi.org/10.14639/0392-100X-N1598>

© Società Italiana di Otorinolaringoiatria e Chirurgia Cervico-Facciale



OPEN ACCESS

This is an open access article distributed in accordance with the CC-BY-NC-ND (Creative Commons Attribution-Non-Commercial-NoDerivatives 4.0 International) license. The article can be used by giving appropriate credit and mentioning the license, but only for non-commercial purposes and only in the original version. For further information: <https://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>

highlighted the confusion around its definition, diagnosis and proper management.

The aetiology of SSS remains controversial. Patients typically deny preexisting sinus disease or orbitofacial trauma. Most authors postulate that collapse of the inferior orbital wall is induced by negative pressure generated by resorption of gas after a natural ostium occlusion that may occur during the first or second decade of life^{2,3}.

Clinical appearance usually consists in asymptomatic enophthalmos and altered facial appearance. Nevertheless, diplopia, sinusitis, rhinorrhoea, post-nasal drip, facial pressure, or pain may also be present^{1,4}.

SSS and Chronic Maxillary sinus Atelectasia (CMA) are terms which have been used interchangeably; the latter can be defined as a persistent and progressive decrease in maxillary sinus volume secondary to inward bowing of the antral walls⁵.

Some authors define CMA as SSS, while others sustain that the two are separate clinical entities^{6,7}. For example, according to Ende et al, it is possible to consider three stages of CMA: the last, group III, is defined by clinical facial deformity with the onset of ocular disturbances, and may be also defined as SSS⁸. Brandt et al. also argued that SSS should be considered a subtype of CMA⁹.

The original definition of SSS dates to 1994 and included only spontaneous enophthalmos not associated with prior trauma or surgery². Nevertheless, many authors have started accepting wider inclusion criteria involving more aetiologies (idiopathic, post traumatic, and iatrogenic) and sustain that they should be included in the definition of SSS¹⁰.

Treatment is also a subject under continuous re-evaluation. At first, surgical treatment consisted in Caldwell-Luc sinus surgery with inferior meatotomy and concurrent transconjunctival repair of the orbital floor. In 1993, Blackwell et al. described endoscopic maxillary antrostomy (FESS: functional endoscopic sinus surgery) in conjunction with a transconjunctival orbital floor repair (OR) in three patients and reported resolution of maxillary disease on follow-up and no recurrence of enophthalmos¹.

Nowadays, the leading treatment of SSS, in order to restore the natural ventilation of the maxillary sinus, appears to be FESS¹¹. Instead, the main controversy in literature remains the necessity and timing of the orbital floor reconstruction via a subconjunctival approach. Some authors prefer a one-stage approach, with endoscopic antrostomy and reconstruction of the orbital floor performed at the same time^{10,12}. Others believe that a two-stage approach is more convenient, with a delay between antrostomy and the orbital reconstruction of 6 months¹³. Still others consider FESS alone to be the best choice of therapy with less invasiveness and satisfactory results, with no need of OR⁶.

To shed light on these controversies, we performed a systematic review of the literature on SSS, with particular focus on definition, diagnosis and surgical approach.

Materials and methods

A PRISMA-compliant systematic literature review¹⁴ was carried out in December 2020 on the Web of Science, PubMed and Scopus databases, using a search strategy for “(Silent Sinus) AND (Maxillary)” and “(Silent Sinus Syndrome) AND (Maxillary)”.

We included studies focused on SSS with the following criteria.

Inclusion criteria

- Age 1-100 years.
- Silent sinus syndrome (SSS).
- Chronic maxillary atelectasia (CMA).

Exclusion criteria

- Studies whose main purpose was unrelated to SSS characteristics and management.
- No human patients involved.
- Language other than English, Italian, French, German and Spanish.
- Article accepted but not published.
- Article type: case reports with less than 3 patients, reviews, comments, letters to the editor, book chapters.

Population, Intervention, Comparison, Outcomes, and Study (PICOS) criteria

PICOS criteria¹⁵ for the present review were as follows:

- Patients with CMA or SSS diagnosis.
- Intervention: evaluation of definition, diagnostic criteria and treatment.
- Comparison: comparison of different definitions, diagnosis and therapeutical options (FESS + OR, FESS alone, OR alone, antibiotic therapy, wait & see).
- Outcome: proposal of shared definition, diagnostic criteria and treatment.
- Study design: Retrospective case studies and case series (more than 3 patients) were enrolled in the review.

Data extraction and quality assessment

Two of the authors (CR and CP) independently screened the retrieved studies based on title and abstract; when uncertainty existed in the abstract evaluation, we retrieved and assessed the full text. After completion of all searches, duplicates were removed. Evaluation through full-text screening was then carried out. Critical appraisal led to the selection of 28 studies (Fig. 1). Both




Figure 1. PRISMA flow-chart.

retrospective and prospective studies were included, while case reports and small case series were excluded because of their intrinsically lower level of evidence (the minimum number of patients was arbitrarily set at 3). Published reviews on SSS were similarly excluded, but their reference list was reviewed to identify possible additional studies. A manual search in the reference lists of these articles was performed to identify potentially relevant papers missed during the database search. Differing opinions were resolved by consensus between the two authors. Data extracted and analysed included study design, sample size, mean patient age, diagnostic criteria, associated factors, instrumental diagnosis, surgical approach and timeline, outcomes, complications and follow-up time.

Results

The systematic review retrieved 28 original articles^{3,4,6,7,10-13,16-19,21,23-37}. Twenty-one were retrospective case studies (RCS), while 7 were case reports reporting at least 3 cases. A total of 276 patients were evaluated with a mean of 9.8 cases per study (range 3-57). Mean age was 40.4 years, although 3 papers did not report the age of participants (Tab. I). There was a wide variety of diagnostic criteria among studies (Tab. II). The most frequently required findings for diagnosis were evidence of enophthalmos (En)

(25 studies), maxillary atelectasia (MA) (23 studies) and hypoglobus (15 studies), while post-traumatic, facial asymmetry and diplopia evidence were less required (respectively, in 3, 3 and 1 studies) as seen in Table II. Moreover, 7 studies did not directly define diagnostic criteria, which were extrapolated from the manuscript.

Nine studies included patients with sinonasal symptoms; 6 included post-traumatic cases, and 1 had post-surgical SSS (Tab. I). Twenty studies diagnosed SSS with CT alone, while 7 performed both CT and MR imaging. One study did not define what imaging was performed (Tab. I).

Surgical approaches consisted in combined FESS and OR in 72 patients, mostly in a one-step surgery. Four studies performed both types of timing-choices (one step or two steps 6 months apart), deciding according to the individual case. FESS alone was performed in 116 patients; other less common therapies were a wait & see approach (57 patients), OR alone (8 patients) and antibiotic therapy (1 patient) (Tab. III). A study by Khon et al. did not define their therapeutical approach in the 22-patient case series²⁹.

All patients had clinical resolution (CR) or clinical improvement (CI). Only Lin and Brown reported 2 patients who had FESS surgery alone that did not resolve enophthalmos, who then underwent subsequent OR^{13,16}. Complications are reported in Table I.

Discussion

SSS is a relatively rare disease that most ENT specialists and ophthalmologists know as an entity, but the sporadic evidence leads to ambiguity on many of its aspects, from definition, to diagnostic criteria to treatment.

Definition

Our systematic review revealed that among the 28 studies evaluated, the most required diagnostic criteria were evidence of enophthalmos (25 studies) and maxillary atelectasia (MA) (23 studies), with 21 requiring both signs. Hypoglobus observation follows as essential criteria in 15 studies, also with evidence of orbital floor remodeling at CT scans (11 studies) and the absence of sinonasal diseases (10 studies) (Tab. II).

Therefore, even if there is much misunderstanding around the definition of SSS, it is possible to assume that most authors agree in having both enophthalmos and MA as necessary criteria. Only 2 studies required only MA in the absence of enophthalmos, considering CMA and SSS as a same entity^{4,17}.

Post-traumatic silent sinuses were originally excluded from the definition of SSS^{18,19}, and some recent papers continue

Table 1. Baseline data of studies included.

Study (year)	Study design	Sample size	Mean age (range) in years	Diagnostic criteria	Associated factors	Diagnosis	Surgical approach	Surgical timeline	Outcome	Complications	Follow-up (months)
Behbehani ²⁵ (2006)	RCS	5	36.7 (32-42)	En Hypoglobus MA	Post-traumatic (1) Chronic sinusitis (4)	CT scans	FESS and OR (5)	One step (5)	100% CI or CR	Residual 1 mm En (2) Transient infraorbital hypoesthesia (1)	24
Bossolesi ¹⁸ (2008)	Case reports	4	42 (38-45)	En Opacified maxillary sinus Absence of major sinus pathology Absence of previous trauma, surgery or congenital facial deformity	Chronic sinusitis (2)	End CT MR	FESS and OR (4)	One step (4)	100% CR	None	12-24
Brown ¹⁶ (2017)	RCS and review	6	43 (35-52)	En MA Hypoglobus	Chronic sinusitis (1) Post-traumatic (6)	CT	FESS (1) OR (1) FESS and OR (4)	One step (1) Two steps (3)	100% CI	Residual 2 mm En (1) minor diplopia (1)	Not defined
Chariba ¹¹ (2014)	RCS	13	34 (13-61)	En MA	Chronic sinusitis (5)	CT	FESS (13)	N/A	100% CI	Orbital breach (2)	30
Chavez-Montoya ²⁶ (2017)	RCS	3	44 (37-48)	MA Opacified maxillary sinus	Nasal polyposis (1)	CT	FESS (1) FESS and OR (1) Wait & see (1)	One step (1)	100% CI	-	24
Claròs ¹² (2015)	Case reports	3	36.3 (25-45)	Not defined En Hypoglobus Low pressure in maxillary sinus	-	CT	FESS + OR (3)	One step (3)	100% CR	None	Not defined
Claròs ¹⁰ (2019)	RCS	13	38 (25-53)	Not defined En Hypoglobus Orbital floor resorption MA Absence of trauma and congenital facial deformity	-	CT	FESS + OR (13)	One step (13)	100% CI	None	Not defined
Cobb ²¹ (2012)	Case reports	3	44 (30-60)	En MA Maxillary opacity	Post-traumatic (2) Post-surgical (1)	CT MR	FESS and OR (3)	One step (2) Two steps (1)	100% CI	None	12 (1) 36 (1) 48 (1)
De Dordodot ⁷ (2017)	RCS	4	44 (12-60)	Facial asymmetry En and/or hypoglobus MA Absence of sinonasal symptoms	-	CT	FESS (3) FESS and OR (1)	Two steps (1)	100% CI	Slight enophthalmos in some patients (No. not defined)	Not defined
Eyigor ⁶ (2016)	RCS	16	42.37 (20-66)	MA En Absence of trauma or sinusitis	-	CT MR	FESS (16)	N/A	Ongoing follow-up	Ongoing follow-up	Ongoing
Farneti ³ (2017)	RCS and review	6	10 (7-14)	Absence of sinusitis Remodeling of orbital floor at CT/MR scans Opacified maxillary sinus Absence of trauma or congenital deformities	-	CT	FESS (6)	N/A	100% CR	Residual headache (1)	18-135
Freiser ¹³ (2020)	RCS	57	12.5 (3.7-18)	Not defined En Hypoglobus MA	-	CT	FESS (19) Wait & see (38)	N/A	100% CI or CR	Not defined	Not defined
Gaudino ²³ (2013)	RCS	6	44 (22-67)	En Diplopia Opacified sinus MA	-	CT MR	FESS (1) FESS and OR (2) Wait & see (3)	One step (1) Two steps (1)	2 FESS: no significant CI 2 FESS and OR: CI	Not defined	Not defined

continues ►

Table 1. Baseline data of studies included (*follows*).

Study (year)	Study design	Sample size	Mean age (range) in years	Diagnostic criteria	Associated factors	Diagnosis	Surgical approach	Surgical timeline	Outcome	Complications	Follow-up (months)
Illner ²⁷ (2002)	RCS	5	47 (39-65)	Not defined En Hypoglobus Maxillary sinus completely developed Maxillary sinus opacified Infundibulum occluded	-	CT/MR	FESS (3) Antibiotic therapy (1) Wait & see (1)	N/A	Not defined	Not defined	Not defined
Kashima ²⁸ (2016)	RCS	11	39.5 (23-62)	Not defined MA En Hypoglobus	Not defined	Not defined	FESS and OR (11)	One step (11)	100% CI	Residual enophthalmos (1) 1 mm over correction (1) Lower eye-lid retraction (1) Nasolacrimal duct obstruction (1)	9
Kohn ²⁹ (2013)	RCS	22	41.2 (22-70)	En Hypoglobus MA Orbital changes at CT scans	Chronic rhinosinusitis (9) Post-traumatic (10)	CT/MR	Not defined	Not defined	Not defined	Not defined	Not defined
Korn ¹⁷ (2009)	Case reports	5	Not defined	MA Orbital floor resorption	Chronic rhinosinusitis (2)	CT	FESS and OR (5)	One step (5)	100% CI	Transient infraorbital Hypoesthesia (1) Residual diplopia (1)	Not defined
Lee ³⁰ (2018)	Case reports	3	44.6 (37-55)	En Hypoglobus MA Opacified maxillary sinus	Post-traumatic (1)	CT MR	FESS (1) OR (1) Wait & see (1)	N/A	100% CI	Residual diplopia (1) Residual enophthalmos (1)	Not defined
Lin ³¹ (2015)	RCS	9	Not defined	MA Orbital floor resorption En Absence of sinusitis	Post-traumatic (1)	CT	FESS (7) FESS and OR (2)	Two steps (2)	2 FESS alone did not resolved enophthalmos and so underwent OR	None	21.4
Martinez-Capoccioni ³² (2016)	RCS	20	44.2 (28-67)	En and/or Hypoglobus Endoscopic findings of MA Altered facial appearance Maxillary contraction and orbital enlargement at CT scans	Nasal obstruction (9)	CT	FESS (15) Wait & see (5)	N/A	100% CI	None	6-18
Rose ³³ (2003)	RCS	14	41.3 (25-78)	Not defined En Hypoglobus Remodeling of orbital floor Absence of nasal diseases	-	CT	OR (6) Wait & see (8)	N/A	100% CI	None	5-33
Sesenna ³⁴ (2010)	Case reports	3	39 (28-46)	En Hypoglobus MA Absence of sinusitis, trauma and congenital deformities Orbital floor remodeling at CT scans	-	CT	FESS and OR (3)	One step (2)	100% CR	None	10-16
Sivasubramaniam ³⁵ (2011)	RCS	18	Not defined (19-54)	En and/or Hypoglobus MA Altered facial appearance Absence of sinusitis	-	CT	FESS (18)	N/A	78% CR 17% CI	Residual enophthalmos (1)	15-120
Thomas ²² (2003)	RCS	4	32 (27-35)	En Absence of sinusitis MA Opacified maxillary sinus Inferior bowing of orbital floor	-	CT	FESS and OR (2) FESS (2)	Two steps (2)	100% CR	Residual enophthalmos after FESS alone (2) which required second step surgery with OR	Not defined

continues ►

Table I. Baseline data of studies included (*follows*).

Study (year)	Study design	Sample size	Mean age (range) in years	Diagnostic criteria	Associated factors	Diagnosis	Surgical approach	Surgical timeline	Outcome	Complications	Follow-up (months)
Vander Meer ⁴ (2001)	RCS	4	42 (38-47)	En MA Absence of sinusitis, trauma or congenital deformities Remodeling of orbital floor at CT scans	-	CT	FESS and OR (4)	One step (4)	100% CR	none	2-36
Virgin ³⁶ (2008)	RCS	5	42 (22-65)	En MA	-	CT	FESS (4) FESS and OR (1)	Two steps (1)	100% CR	None	24
Wan ³⁷ (2000)	Case reports	3	41 (38-44)	Not defined En Hypoglobus MA Remodeling orbital floor	Chronic sinusitis (1)	CT	FESS (3)	N/A	100% CR	none	N/A
Wise ¹⁹ (2007)	RCS Focused only on CT findings	11	46.5 (11-70)	En Opacified maxillary sinus Orbital floor remodelling Absence of chronic sinusitis, trauma or congenital deformities	-	CT	FESS (3) FESS and OR (8)	Not defined	Not defined	none	Not defined

RCS: retrospective case study; En: enophthalmos; MA: maxillary atelectasia; FESS: functional endoscopic sinus surgery; OR: orbital reconstruction; CI: clinical improvement; CR: clinical resolution; End: nasal endoscopy.

considering post-traumatic cases not definable as SSS due to their lack of spontaneous development^{10,20}.

Nevertheless, a retrospective analysis of 6 cases and literature review showed how traumatic SSS management follows the same principles as for spontaneous SSS²¹. In fact, our analysis revealed that recent papers started to include enophthalmos with MA secondary to traumatic events in the group of SSS, explaining that clinical and radiological presentations are comparable, as well as surgical treatment^{13,21,22}.

Moreover, the association of SSS with sinonasal symptoms is debated. Ten studies considered the evidence of sinonasal symptoms as an exclusion criterion because MA secondary to chronic sinusitis with sinus dysventilation and sinonasal symptoms is, by those authors, considered as CMA and not as SSS⁹. At the same time, the majority of studies do not clarify this point, and include among SSS subjects who have associated factors such as sinusitis, chronic nasal congestion and other sinonasal symptoms of obstruction, even though these are not considered diagnostic criteria (Tab. II). Finally, 7 studies did not specify diagnostic criteria, which were extrapolated from the clinical description of cases. This risks producing inaccurate studies given that the literature lacks in consensus in SSS diagnosis and the reader may find its definition confusing.

From our analysis, it appears that the leading definition of SSS should include the evidence of enophthalmos and maxillary atelectasia. We believe that presence of sinonasal diseases and history of facial trauma may be reasonable factors to be included in anamnestic data because of the comparable clinical presentation and treatment.

DEFINITION PROPOSAL

Our proposal for definition is:

- criteria to diagnose SSS are contemporary presence of enophthalmos and maxillary atelectasia;
- minor associated factors may be the presence of sinonasal diseases and history of facial trauma.

Diagnosis

Most studies only required CT for imaging investigation, since the relevance was to investigate maxillary bony walls atelectasia with possible resorption of the orbital floor. Nevertheless, 7 groups also considered it useful to add MR scans (Tab. I) to evaluate:

- dislocation of extra and intra-conical fat, extrinsic ocular muscles and optic nerve;
- differentiation between mixed signal central secretions and high signal peripheral;
- thick edematous mucosal lining within the maxillary sinus^{6,18,23}.

DIAGNOSTIC PROPOSAL

Only CT scan is required. MR imaging can be associated in selected cases, and specifically to evaluate a marked hypoglobus. We propose to add evaluation of extraocular muscle movement and potential diplopia at first ENT clinical evaluation, in order to select symptomatic and therefore the most critical cases^{10,20}.

Treatment

The review also showed disagreement on management strategies. Seventeen authors proposed combined FESS and OR surgery as the leading therapy to obtain resolution

of anatomical impairment and diplopia, if present. Among these, the majority (n = 9) performed both the procedures simultaneously, while 4 preferred to first carry out FESS surgery and to observe a possible progressive improvement of the enophthalmos to decide whether or not to perform additional OR. Eventually, 3 studies performed FESS and either simultaneous or delayed OR according to the criticality of the case (Tab. III).

The reasons which guide each surgical team towards one choice or another are multiple. The main factor is undoubtedly the severity of enophthalmos and hypoglobus, although no study defined a quantitative cut-off which could help in the surgical decision. OR approach is supported by the fact that, although there have been reports of resolution of the progression of enophthalmos by antrostomy alone, it is unclear whether other aesthetic deformities caused by SSS as hypoglobus or superior sulcus deformity also respond. Behbehani et al. believe that delaying orbital implant placement in cases with significant enophthalmos and hypoglobus is unjustified since complications like diplopia or infection are rare with this procedure. Furthermore, simultaneous implant placement also obviates the need for additional anaesthesia and hospitalisation²⁵. Cobb et al. maintain that FESS alone may stop the descent of the orbital wall, but there would be no reason to expect that the position of the orbital floor, and thus the globe, would be reversed²¹.

On the contrary, Thomas et al. support delayed repair of the orbital floor as in some patients enophthalmos improves with antrostomy alone²². Moreover, OR has not been shown to provide any significant restoration in the orbital muscle functions, and because of that diplopia is not corrected⁶.

A total of 116 patients underwent FESS surgery alone, being considered as a necessary and sufficient procedure to resolve MA and enophthalmos (Tab. III). If we consider the entire case series of the review, FESS alone appears as the leading therapeutic choice, with only 7 reporting residual enophthalmos (6%), and 4 requiring subsequent OR (3.4%). Moreover, Numa et al. undertook a review of 84 cases and concluded that for patients with SSS diagnosis, uncinectomy alone may be sufficient²⁴.

FESS + OR follows with 72 cases treated with this management. Wait & see (57 patients) may be a valid alternative in asymptomatic cases or young population¹³. OR alone (8 patients) and antibiotic therapy (1 patient) remain marginal therapeutic options (Tab. III).

This leads us to the conclusion that most of SSS may be successfully treated with isolated FESS surgery, sparing OR as a rescue procedure in case of non-satisfactory resolution of enophthalmos or diplopia.

TREATMENT PROPOSAL

FESS is the leading treatment for SSS and orbital reconstruction should be performed only in selected and symptomatic cases or, if needed, in a second approach when no resolution is seen. The proposed timepoint for a potential postponed OR, according to the literature, is 6 months.

Our review clearly shows the need to develop consensus regarding the definition of SSS and most of all its management. Even if the literature has started to find marginal consensus in recent years, it appears necessary to define shared diagnostic criteria, as well as a shared approach to the best treatment choices with the lowest rate of invasiveness and morbidity. The literature would also

Table II. Summary of diagnostic criteria for SSS.

Not directly defined	Study (year)	En	Hypoglobus	MA	Opacified maxillary Sinus	No previous trauma, surgery or congenital deformities	No sinonasal symptoms	Post-traumatic	Facial asymmetry	Diplopia	Orbital floor remodeling at CT
	Behbehani ²⁵ (2006)	X	X	X							
	Bossolesi ¹⁶ (2008)	X			X	X	X				
	Brown ¹⁶ (2017)	X	X	X				X			
	Chariba ¹¹ (2014)	X		X							
	Chavez-Montoya ²⁶ (2017)			X	X						
x	Claròs ¹² (2015)	X	X	X							
x	Claròs ¹⁰ (2019)	X	X	X		X					
	Cobb ²¹ (2012)	X		X				X			
	De Dordodot ⁷ (2017)	X	X	X			X		X		
	Eyigor ⁶ (2016)	X		X		X					
	Farneti ³ (2017)				X	X	X				X
x	Freiser ¹³ (2020)	X	X	X							
	Gaudino ²³ (2013)	X		X	X					X	
x	Illner ²⁷ (2002)	X	X		X						
x	Kashima ²⁸ (2016)	X	X	X							
	Kohn ²⁹ (2013)	X	X	X							X
	Korn ¹⁷ (2009)			X							X
	Lee ³⁰ (2018)	X	X	X	X						
	Lin ³¹ (2015)	X		X			X	X			X
	Martinez-Capoccioni ³² (2016)	X	X	X					X		X
x	Rose ³³ (2003)	X	X				X				X
	Sesenna ³⁴ (2010)	X	X	X		X	X				X
	Sivasubramaniam ³⁵ (2011)	X	X	X			X		X		
	Thomas ²² (2003)	X		X	X		X				X
	Vander Meer ⁴ (2001)	X		X		X	X				X
	Virgin ³⁶ (2008)										
	Wan ³⁷ (2000)	X		X							
x	Wise ¹⁹ (2007)	X	X	X							X
7	TOTAL	24	15	23	7	6	9	3	3	1	10

En: enophthalmos; MA: maxillary atelectasia.

benefit from prospective studies on the best outcome in SSS management.

Conclusions

Literature about SSS is controversial and confusing. Our systematic review illustrates that the leading definition of SSS includes the evidence of enophthalmos and maxillary

atelectasia by CT. Hypoglobus, presence of sinonasal diseases and history of facial trauma may or may not be associated, although the clinical relevance and management seems to be comparable. FESS alone appears to be the first choice of treatment, since at post-operative follow-up, orbital floor retraction tends to spontaneously reverse with clinically satisfactory results.

Table III. Summary of therapeutic strategies for SSS.

Study (year)	FESS + OR	FESS	OR	Wait & see	Antibiotic therapy
Behbehani ²³ (2006)	X (5)				
Bossolesi ¹⁶ (2008)	X (4)				
Brown ¹⁴ (2017)	X (4)	X (1)	X (1)		
Chariba ¹¹ (2014)		X (13)			
Chavez-Montoya ²⁴ (2016)	X (1)	X (1)		X (1)	
Claròs ¹² (2015)	X (3)				
Claròs ¹⁰ (2019)	X (13)				
Cobb ¹⁹ (2012)	X (3)				
De Dordodot ⁷ (2017)	X (1)	X (3)			
Eyigor ⁶ (2016)		X (16)			
Farneti ³ (2017)		X (6)			
Freiser ¹³ (2020)		X (19)		X (38)	
Gaudino ²¹ (2012)	X (2)	X (1)		X (3)	
Illner ²⁵ (2002)		X (3)		X (1)	X (1)
Kashima ²⁶ (2016)	X (11)				
Kohn ²⁷ (2013)	-	-	-	-	-
Korn ¹⁷ (2009)	X (5)				
Lee ²⁸ (2019)		X (1)	X (1)	X (1)	
Lin ²⁹ (2015)	X (2)	X (7)			
Martinez-Capoccioni ³⁰ (2016)		X (15)		X (5)	
Rose ³¹ (2003)			X (6)	X (8)	
Sesenna ³² (2010)	X (3)				
Sivasubramaniam ³³ (2011)		X (18)			
Thomas ²⁰ (2002)	X (2)	X (2)			
Vander Meer ⁴ (2001)	X (4)				
Virgin ³⁴ (2008)	X (1)	X (4)			
Wan ³⁵ (2000)		X (3)			
Wise ¹⁷ (2007)	X (8)	X (3)			
TOTAL	17	17	3	7	1
TOTAL PER CASES	72	116	8	57	1

Acknowledgements

We gratefully thank the Librarians from the Università degli Studi di Milano, Biblioteca del Polo Centrale. Without their help, we would not have been able to locate and analyse a significant number of the studies included in this systematic review.

Conflict of interest statement

The authors declare no conflict of interest.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Author contributions

CR made a substantial contribution to the conception and design of the article, to the acquisition, analysis and interpretation of data. AMS and GF critically revised the article and gave the final approval of the version to be published. GP made a substantial contribution to the conception and design of the article and gave the final approval of the version to be published.

References

- Blackwell KE, Calcaterra TC, Goldberg RA. Atelectasis of the maxillary sinus with enophthalmos and midface depression. *Ann Otol Rhinol Laryngol* 1993;102:429-432. <https://doi.org/10.1177/000348949310200604>

- 2 Soparkar CNS, Patrinely JR, Cuaycong MJ, et al. The silent sinus syndrome: a cause of spontaneous enophthalmos. *Ophthalmology* 1994;101:772-778. [https://doi.org/10.1016/s0161-6420\(94\)31267-x](https://doi.org/10.1016/s0161-6420(94)31267-x)
- 3 Farneti P, Sciarretta V, Macri G, et al. Silent sinus syndrome and maxillary sinus atelectasis in children. *Int J Pediatr Otorhinolaryngol* 2017;98:150-157. <https://doi.org/10.1016/j.ijporl.2017.05.005>
- 4 Vander Meer JB, Harris G, Toohill RJ, et al. The silent sinus syndrome: a case series and literature review. *Laryngoscope* 2001;111:975-978. <https://doi.org/10.1097/00005537-200106000-00008>
- 5 Hens G, Hermans R, Jorissen M. Chronic maxillary atelectasis. *B ENT* 2005;1:25-29.
- 6 Eyigör H, Çekici B, Çoban DT, et al. Is there a correlation between the clinical findings and the radiological findings in chronic maxillary sinus atelectasis? *J Cranio-Maxillofacial Surg* 2016;44:820-826. <https://doi.org/10.1016/j.jcms.2016.04.004>
- 7 De Dorlodot C, Collet S, Rombaux P, et al. Chronic maxillary atelectasis and silent sinus syndrome: two faces of the same clinical entity. *Eur Arch Otorhinolaryngol* 2017;274:3367-3373. <https://doi.org/10.1007/s00405-017-4622-8>
- 8 Ende K, Mah L, Kass ES. Progression of late-stage chronic maxillary atelectasis. *Ann Otol Rhinol Laryngol* 2002;111:759-762. <https://doi.org/10.1177/000348940211100819>
- 9 Brandt MG, Wright ED. The silent sinus syndrome is a form of chronic maxillary atelectasis: a systematic review of all reported cases. *Am J Rhinol* 2008;22:68-73. <https://doi.org/10.2500/ajr.2008.22.3118>
- 10 Clarós P, Sobolewska AZ, Cardesa A, et al. Silent sinus syndrome: combined sinus surgery and orbital reconstruction-report of 15 cases. *Acta Otolaryngol* 2019;139:64-69. <https://doi.org/10.1080/00016489.2018.1542161>
- 11 Chariba I, Lazard DS, Sain-Oulhen C, et al. Correlation between the rate of asymmetry volume of maxillary sinuses and clinical symptomatology in the silent sinus syndrome: a retrospective study about 13 cases. *Rev Laryngol Otol Rhinol* 2014;135:197-200.
- 12 Clarós P, Ahmed H, Minka Ngom EG, et al. Le silent sinus syndrome: reconstruction orbitaire par implant Medpor [The silent sinus syndrome: A reconstruction of the orbital floor with Medpor implant]. *Rev Laryngol Otol Rhinol* 2015;136:37-40. PMID: 26749604.
- 13 Freiser ME, McCoy J, Shaffer AD, et al. Silent sinus syndrome in children. *Int J Pediatr Otorhinolaryngol*. 2020;134:110034. <https://doi.org/10.1016/j.ijporl.2020.110034>
- 14 Moher D, Liberati A, Tetzlaff J, et al. PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *J Clin Epidemiol* 2009;62:1006-1012. <https://doi.org/10.1016/j.jclinepi.2009.06.005>
- 15 Amir-Behghadami M, Janati A. Population, Intervention, Comparison, Outcomes and Study (PICOS) design as a framework to formulate eligibility criteria in systematic reviews. *Emerg Med J* 2020;37:387. <https://doi.org/10.1136/emermed-2020-209567>
- 16 Brown SJ, Hardy TG, McNab AA. 'Silent Sinus Syndrome' following orbital trauma: a case series and review of the literature. *Ophthal Plast Reconstr Surg* 2017;33:209-212. <https://doi.org/10.1097/IOP.0000000000000713>
- 17 Korn BS, Weisman RA, Kikkawa DO, et al. Multidisciplinary approach to management of silent sinus syndrome. *Laryngoscope* 2009;119 (Suppl 1);S180. <https://doi.org/10.1002/lary.20470>
- 18 Bossolesi P, Autelitano L, Brusati R, et al. The silent sinus syndrome: diagnosis and surgical treatment. *Rhinology* 2008;46:308-316.
- 19 Wise SK, Wojno TH, DelGaudio J. M. Silent sinus syndrome: lack of orbital findings in early presentation. *Am J Rhinol* 2007;21:489-494. <https://doi.org/10.2500/ajr.2007.21.3055>
- 20 Brown SJ, McNab AA. Ethmoid silent sinus syndrome following orbital trauma. *Ophthal Plast Reconstr Surg* 2017;33:e60-e61. <https://doi.org/10.1097/IOP.0000000000000746>
- 21 Cobb AR, Murthy R, Cousin G, et al. Silent sinus syndrome. *Br J Oral Maxillofac Surg* 2012;50:e81-e85. <https://doi.org/10.1016/j.bjoms.2011.10.001>
- 22 Thomas RD, Graham SM, Carter KD, et al. Management of the orbital floor in silent sinus syndrome. *Am J Rhinol* 2003;17:97-100.
- 23 Gaudino S, Di Lella GM, Piluddu E, et al. CT and MRI diagnosis of silent sinus syndrome. *Radiol Med* 2013;118:265-275. <https://doi.org/10.1007/s11547-012-0822-x>
- 24 Numa WA, Desai U, Gold DR, et al. Silent sinus syndrome: a case presentation and comprehensive review of all 84 reported cases. *Ann Otol Rhinol Laryngol* 2005;114:688-694. <https://doi.org/10.1177/000348940511400906>
- 25 Behbehani R, Vacarezza N, Bilyk JR, et al. Simultaneous endoscopic antrostomy and orbital reconstruction in silent sinus syndrome. *Orbit* 2006;25:97-101. <https://doi.org/10.1080/01676830600671516>
- 26 Chavez-Montoya R, Araujo-Ramírez O, Castillo-López IY, et al. Impinging antrum syndrome: three cases with different management approaches. *Cir y Cir (English Ed.)* 2017;85:529-534. <https://doi.org/10.1016/j.circir.2016.08.005>
- 27 Illner A, Davidson HC, Harnsberger HR, et al. The silent sinus syndrome: clinical and radiographic findings. *Am J Roentgenol* 2002;178:503-506. <https://doi.org/10.2214/ajr.178.2.1780503>
- 28 Kashima T, Goldberg RA, Kohn JC, et al. One-stage bone strip reconstruction technique with balloon sinus dilatation surgery for chronic maxillary atelectasis. *Clin Ophthalmol* 2016;10:2363-2368. <https://doi.org/10.2147/OPHTH.S103809>
- 29 Kohn JC, Rootman DB, Xu D, et al. Infratemporal fossa fat enlargement in chronic maxillary atelectasis. *Br J Ophthalmol* 2013;97:1005-1009. <https://doi.org/10.1136/bjophthalmol-2013-303299>
- 30 Lee DS, Murr AH, Kersten RC, et al. Silent sinus syndrome without opacification of ipsilateral maxillary sinus. *Laryngoscope* 2018;128:2004-2007. <https://doi.org/10.1002/lary.27108>
- 31 Lin GC, Sedaghat AR, Bleier BS, et al. Volumetric analysis of chronic maxillary atelectasis. *Am J Rhinol Allergy* 2015;29:166-169. <https://doi.org/10.2500/ajra.2013.27.4173>
- 32 Martínez-Capoccioni G, Varela-Martínez E, Martín-Martín C. Silent sinus syndrome an acquired condition and the essential role of otorhinolaryngologist consultation: a retrospective study. *Eur Arch Otorhinolaryngol* 2016;273:3183-3188. <https://doi.org/10.1007/s00405-016-3965-x>
- 33 Rose GE, Lund VJ. Clinical features and treatment of late enophthalmos after orbital decompression: a condition suggesting cause for idiopathic 'impinging antrum' (silent sinus) syndrome. *Ophthalmology* 2003;110:819-826. [https://doi.org/10.1016/S0161-6420\(02\)01994-2](https://doi.org/10.1016/S0161-6420(02)01994-2)
- 34 Sesenna E, Oretti G, Anghinoni ML, et al. Simultaneous management of the enophthalmos and sinus pathology in silent sinus syndrome: a report of three cases. *J Cranio-Maxillofac Surg* 2010;38:469-472. <https://doi.org/10.1016/j.jcms.2009.12.003>
- 35 Sivasubramaniam R, Sacks R, Thornton M. Silent sinus syndrome: dynamic changes in the position of the orbital floor after restoration of normal sinus pressure. *J Laryngol Otol* 2011;125:1239-1243. <https://doi.org/10.1017/S0022215111001952>
- 36 Virgin F, Ling FTK, Kountakis SE. Radiology and endoscopic findings of silent maxillary sinus atelectasis and enophthalmos. *Am J Otolaryngol* 2008;29:167-170. <https://doi.org/10.1016/j.amjoto.2007.05.005>
- 37 Wan MK, Francis IC, Carter PR, et al. Spectrum of presentation of Silent Sinus Syndrome. *J Neuroophthalmol* 2000;20:207-212. <https://doi.org/10.1097/00041327-200020030-00010>