



Data Article

Dataset of endometrial blood flow from pregnant and non-pregnant mares on day 7 and 8 post-ovulation

Pilar Nieto-Olmedo^a, Gemma Gaitskell-Phillips^b,
 Francisco E Martín-Cano^b, Jose Manuel Ortiz-Rodríguez^b,
 Fernando J Peña^b, Cristina Ortega-Ferrusola^{b,*}

^aCEFIVA-Centro de Fertilización In vitro de Asturias, Spain

^bLaboratory of Equine Reproduction and Equine Spermatology, Veterinary Teaching Hospital, University of Extremadura, Cáceres, Spain

ARTICLE INFO

Article history:

Received 26 February 2020

Revised 8 April 2020

Accepted 17 April 2020

Available online 25 April 2020

Keywords:

Uterine blood flow

Endometrial angiogenesis

Pregnant mare

Doppler ultrasonography

Embryo transfer

Barren mare

Pregnancy diagnosis mare

ABSTRACT

This article provides the dataset for the use of power Doppler ultrasound to assess the equine uterus from the recent research article titled "Power Doppler can detect the presence of 7-8 days conceptuses prior to flushing in an equine embryo transfer program"(1). The vascularization of the endometrium was objectively assessed in mares by quantification of pixels in bitmap format (BMP) using computer assisted analysis of images. Fifty-two mares were examined on days 7 (26 mares) and 8 (26 mares) post-ovulation prior to performing flushing procedures for embryo recovery. Receiver operating characteristic (ROC) curves and Youden's J statistics were used to evaluate the value of the suggested variable in terms of its diagnostic value for identification of early pregnancy and to establish cut-off values allowing differentiation between pregnant and non-pregnant mares on days 7 and 8 post-ovulation.

© 2020 Published by Elsevier Inc.

This is an open access article under the CC BY-NC-ND license. (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

* Corresponding author: Cristina Ortega Ferrusola.

E-mail addresses: reproduccionequina@gmail.com, cristinaof@unex.es (C. Ortega-Ferrusola).

Specifications table

Subject	Veterinary Medicine
Specific subject area	Diagnostic imaging using Doppler ultrasound for early detection of pregnancy in mares
Type of data	Table Figure
How data were acquired	Ultrasonographic examination was performed using MyLabFive Vet equipment (Esaote S.p.A, Genova, Italy) and a linear 5–7.5 MHz probe. Rectal ultrasound examination was performed and the ultrasound probe was positioned transversely in the middle point of each uterine horn. Two still images were obtained per horn from the ultrasound examination. Signals for blood flow in all endometrial, myometrial and perimetrial vessels were visualized using Power flow modes and blood flow was measured in pixels. Still Doppler images were assessed using computer analysis with ImageJ v1.48 software (National Institute of Health, USA). ImageJ is a Java-based image processing program which is capable of calculating pixel value statistics for user-defined selections.
Data format	Raw Analyzed
Parameters for data collection	Fifty-two mares (Pure Bred Spanish horses), none of which showed signs of uterine pathology were used, with an age range of between 2 and 18 years old. Endometrial blood flow was evaluated, and embryo recovery was subsequently performed on days 7 and 8 post-ovulation. Mares were all sedated with romifidine (0.02 mg/kg) (Sedivet®, Barcelona, Spain), before ultrasonographic assessments were commenced.
Description of data collection	Rectal ultrasound examination was performed, and the ultrasound probe was positioned transversely in the middle point of each uterine horn. Pixel quantification was used to accurately and objectively assess the degree of uterine vascular perfusion. Blind analysis was performed on a total of 208 images (four images per mare) using computer analysis with ImageJ v1.48 software (National Institute of Health, USA). Detection of blood vessels was carried out on the images with application of a color threshold which was restricted to the uterine horn under examination.
Data source location	Institution: Faculty of Veterinary Medicine-University of Extremadura City/Town/Region: Cáceres, Extremadura Country: Spain
Data accessibility	With the article
Related research article	Pilar Nieto-Olmedo ^a ; Francisco E Martín-Cano ^b ; Gemma Gaitskell-Phillips ^a ; Jose Manuel Ortiz-Rodríguez ^a ; Fernando J Peña ^a ; Cristina Ortega-Ferrusola ^a Power Doppler can detect the presence of 7–8 day conceptuses prior to flushing in an equine embryo transfer program. Theriogenology 145, 1-9. https://doi.org/10.1016/j.theriogenology.2020.01.015

Value of the data

- These data provide useful information concerning clinical applicability of power doppler ultrasound for discrimination between pregnant and non-pregnant mares on days 7 and 8 post-ovulation before performing embryo recovery.
- These data could be beneficial in clinical practice. The technique could be used in routine clinical practice by veterinarians in order to maximize embryo recovery rates from donor mares. Cut-off values established using this data can be used to predict pregnancy diagnosis prior to embryo collection.
- In addition, these data could be useful for further research into the study of early embryonic death and the changes which occur in uterine tissues immediately prior to it.
- The data may also prove useful with regard to development of strategies for implementation of timely preventative treatments for mares which are known to have suffered from early embryonic death in the past and have been unable to maintain pregnancies.

Uterine blood flow




Fig. 1. This graph represents the differences in endometrial blood flow between pregnant and non-pregnant mares on days 7 and 8 post-ovulation. The results are expressed in pixels as a mean \pm SD. There were significant differences detected between pregnant and non-pregnant mares ($p < 0.001$). Vascularization between left and right horns showed no differences in either group.




Fig. 2. Receiver operating characteristic (ROC) curves for the blood flow area parameter (pixels) in mares on days 7 and 8 post-ovulation. AUC: Area under the Curve. This curve was then used to identify pregnant mares on days 7 and 8 post-ovulation. When analyzed using Youden's test, data showed that the uterine blood flow area in pregnant mares was greater than 1134 pixels on day 7 with a sensitivity of 69.1% and a specificity of 94.4% and an AUC: 0.836. After analysis, the cut off value for pregnant mares on day 8 post-ovulation was 912 pixels, with a sensitivity of 86.8% and a specificity of 100% (AUC: 0.96). Consequently, evaluation of endometrial blood flow in pixels presented a greater predictive value on day 8 post-ovulation.

- These data can be used as a basis for the development of further experiments into the changes which occur to uterine blood flow after entry of the equine embryo into the uterus which could lead to further development of the technique as a diagnostic complement for early pregnancy diagnosis with future potential to avoid unnecessary flushing procedures in non-pregnant mares.
- Both the raw and analyzed data in pixels are presented in this article. These data could be exploited in order to train veterinarians who wish to learn how to perform the technique and

Table 1

Descriptive statistics of endometrial blood flow in non-pregnant and pregnant mares expressed in pixels. Mean, standard deviation (SD), maximum, minimum and range.

	Non-pregnant mares					Pregnant mares				
	Mean	SD	Minimum	Maximum	Range	Mean	SD	Minimum	Maximum	Range
Day 7	727,67	260,32	250	1338	1088	1389,57	589,07	313	2819	2506
Day 8	508,36	220,35	119	904	785	1796,60	836,74	144	3998	3854

Table 2

Raw data file of uterine blood flow in non-pregnant mares on day 7. Nine barren mares were evaluated and 36 images (four images per mare) were assessed in the study. Pixel quantification was used to assess the degree of uterine vascular perfusion. The table shows the number of mares, the age of the mare, the uterine horn which was assessed and the respective value for blood flow in pixels. The mean, standard deviation (SD) and co-efficient of variation (CV) (%) between the two measurements from each uterine horn are also shown in the table.

Uterine Blood flow non-pregnant mare (Day 7) N° of mare	Age	Uterine horn	Blood flow pixel	Mean	SD	CV (%)
3	4	LEFT	400	455,00	77,78	17,09
		LEFT	510			
		RIGHT	250			
21	4	RIGHT	527	362,00	121,62	33,60
		LEFT	448			
		LEFT	276			
23	6	RIGHT	970	817,00	216,37	26,48
		RIGHT	664			
		LEFT	466			
26	5	LEFT	828	647,00	255,97	39,56
		RIGHT	948			
		RIGHT	1028			
29	2	LEFT	307	598,00	411,54	68,82
		LEFT	889			
		RIGHT	627			
45	10	RIGHT	719	602,00	84,85	14,10
		LEFT	662			
		LEFT	542			
47	11	RIGHT	544	588,50	62,93	10,69
		RIGHT	633			
		LEFT	786			
49	11	LEFT	616	701,00	120,21	17,15
		RIGHT	1115			
		RIGHT	692			
51	9	LEFT	993	1050,00	80,61	7,68
		LEFT	1107			
		RIGHT	1218			
51	9	RIGHT	1338	1278,00	84,85	6,64
		LEFT	814			
		LEFT	914			
51	9	RIGHT	865	864,00	70,71	8,18
		RIGHT	793			
		LEFT	640			
51	9	LEFT	587	613,50	37,48	6,11
		RIGHT	786			
		RIGHT	694			

used as a reference to ensure they are performing the procedure correctly. The raw data highlights the extent of variation that can be expected from measurements of uterine blood flow via Doppler images.

Table 3

Raw data file of uterine blood flow in pregnant mares on day 7.

Seventeen pregnant mares (positive flushing) were assessed and 68 images (four images per mare) were evaluated in the experiment. Pixel quantification was used to assess the degree of uterine vascular perfusion. The table shows the number of mares, the age of each mare, the uterine horn assessed and the values for blood flow in pixels. The mean, standard deviation (SD) and co-efficients of variation (CV) (%) between the two measurements from each uterine horn are also shown in the table.

Uterine Blood flow pregnant mare (Day 7)			Blood flow pixel	Mean	SD	CV (%)
Nº of mare	Age	Uterine horn				
1	18	LEFT	1814	1656,50	222,74	13,45
		LEFT	1499			
		RIGHT	2618			
4	6	RIGHT	1669	626,50	224,15	35,78
		RIGHT	468			
		LEFT	785			
		LEFT	949			
5	11	RIGHT	752	1447,50	9,19	0,64
		RIGHT	1454			
		LEFT	1441			
		LEFT	1204			
6	11	RIGHT	1759	1481,50	392,44	26,49
		RIGHT	1853			
		LEFT	1668			
		LEFT	1258			
9	9	RIGHT	1956	1607,00	493,56	28,49
		RIGHT	2012			
		LEFT	2102			
		LEFT	1578			
11	7	RIGHT	1241	2057,00	63,64	3,09
		RIGHT	1578			
		LEFT	2102			
		LEFT	1241			
15	8	RIGHT	1182	1631,50	635,69	38,96
		RIGHT	2081			
		LEFT	1242			
		LEFT	1710			
16	7	RIGHT	1941	1936,50	6,36	0,33
		RIGHT	1932			
		LEFT	1982			
		LEFT	2534			
20	7	RIGHT	1912	1532,50	536,69	35,02
		RIGHT	1153			
		LEFT	1307			
		LEFT	1829			
22	6	RIGHT	2624	2103,00	736,81	35,04
		RIGHT	1582			
		LEFT	1900			
		LEFT	1459			
24	5	RIGHT	1799	1679,50	311,83	18,57
		RIGHT	794			
		LEFT	1391			
		LEFT	499			
25	5	RIGHT	1258	1291,50	47,38	3,67
		RIGHT	1325			
		LEFT	776			
		LEFT	942			
33	3	RIGHT	1478	859,00	117,38	13,66
		RIGHT	1433			
		LEFT	858			
		LEFT	1916			
34	2	RIGHT	746	1455,50	31,82	2,19
		RIGHT	1984			
		LEFT	903			
		LEFT	653			
			1412	1289,00	173,95	13,49

(continued on next page)

Table 3 (continued)

Uterine Blood flow pregnant mare (Day 7) N° of mare	Age	Uterine horn	Blood flow pixel	Mean	SD	CV (%)
35	2	LEFT	1166	1482,00	113,14	7,63
		RIGHT	1402			
		RIGHT	1562	403,50	127,99	31,72
		LEFT	494			
		LEFT	313			
		RIGHT	655			
36	2	RIGHT	1000	827,50	243,95	29,48
		LEFT	727			
		LEFT	496	611,50	163,34	26,71
		RIGHT	820			
		RIGHT	470			
		LEFT	2045			
41	6	LEFT	1178	1611,50	613,06	38,04
		RIGHT	1326			
		RIGHT	799	1062,50	372,65	35,07
		LEFT	1178			

1. Data Description

This dataset provides a complete set of measurements from images showing blood flow to the equine uterus in the mare at 7 and 8 days post-ovulation which were obtained using power Doppler ultrasound. Quantification of pixels was used to objectively assess each image whilst in bitmap format (BMP) aided by computer assisted image analysis.

The dataset is shown in the following tables and figures:

[Figs. 1 and 2.](#)
[Tables 1-5](#)

2. Experimental Design, Materials, and Methods

The aim of this experiment was to assess whether evaluation of endometrial blood flow in mares could be rated as a good tool for identification of early pregnancy. In total, fifty-two mares of a range of different ages (2-18 years old) were used to obtain this data. Ultrasonographic examination with Power Doppler was used to assess uterine blood flow prior to uterine flushing procedures for embryo recovery on both days 7 and 8 after ovulation. Two transverse images were captured per uterine horn and analysis of images was subsequently performed at a later date using Image J 1.48 software.

Prior to ultrasonographic examination all mares were sedated with romifidine (0.02 mg/kg) (Sedivet® Boehringer Ingelheim, Barcelona, España). MyLabFive Vet equipment (Esaote S.p.A, Genova, Italy) with a linear 5-7.5 MHz probe was used to complete the assessment. The specific settings used were: gain: 70%, PRF: 1,4 KHz at a depth of 9 cm. Examination was performed per rectum and the ultrasound probe was placed transversely at a mid-point of both uterine horns in order to capture two still images per horn. Signals indicating blood flow in all the endometrial, myometrial and perimetrial vessels were visualized using Power flow modes due to their capacity for greater sensitivity for detection of weak signals from small vessels. Degree of uterine vascular perfusion was objectively assessed using quantification of pixels in bitmap (BMP) format. Blind analysis was performed on a total of 208 images (four images per mare). Subsequent computer analysis of Doppler images was executed using ImageJ v1.48 software (National Institute of Health, USA). User-defined selections and intensity-thresholded objects can be analyzed with ImageJ, which is a Java-based image processing program. Analysis of endometrial vascular perfusion was performed using spot meter techniques, measuring blood flow area. A color

Table 4

Raw data file showing uterine blood flow in non-pregnant mares on day 8. Nine barren mares were evaluated and 36 images (four images per mare) were assessed in the study. Pixel quantification was used to assess the degree of uterine vascular perfusion. The table shows the number of mares, the age of each mare, the uterine horn assessed and the values for blood flow in pixels. The mean, standard deviation (SD) and co-efficients of variation (CV) (%) between the two measurements from each uterine horn are also shown in the table.

Uterine Blood flow non-pregnant mare (Day 8)						
Nº of mare	Age	Uterine horn	Blood flow pixel	Mean	SD	CV (%)
2	6	LEFT	140	152,50	17,68	11,59
		LEFT	165			
17	7	RIGHT	510	325,00	261,63	80,50
		RIGHT	140			
		LEFT	292	328,00	50,91	15,52
		LEFT	364			
38	3	RIGHT	284	337,00	74,95	22,24
		RIGHT	390			
		LEFT	595	536,50	82,73	15,42
		LEFT	478			
40	3	RIGHT	369	337,00	45,25	13,43
		RIGHT	305			
		LEFT	563	341,00	313,96	92,07
		LEFT	119			
43	3	RIGHT	317	323,00	8,49	2,63
		RIGHT	329			
		LEFT	490	440,50	70,00	15,89
		LEFT	391			
46	10	RIGHT	427	398,00	41,01	10,30
		RIGHT	369			
		LEFT	686	713,00	38,18	5,36
		LEFT	740			
48	11	RIGHT	614	647,00	46,67	7,21
		RIGHT	680			
		LEFT	889	783,50	149,20	19,04
		LEFT	678			
50	11	RIGHT	904	728,50	248,19	34,07
		RIGHT	553			
		LEFT	729	699,50	41,72	5,96
		LEFT	670			
52	9	RIGHT	473	673,50	283,55	42,10
		RIGHT	874			
		LEFT	544	670,00	178,19	26,60
		LEFT	796			
		RIGHT	660	717,00	80,61	11,24
		RIGHT	774			

Table 5

Raw data file showing uterine blood flow in pregnant mares on day 8. Seventeen pregnant mares (positive flushings) were assessed and 68 images (four images per mare) were evaluated in the study. Pixel quantification was used to assess the degree of uterine vascular perfusion. The table shows the number of mares, the age of each mare, the uterine horn assessed and the values for blood flow in pixels. The mean, standard deviation (SD) and co-efficients of variation (CV) (%) between the two measurements from each uterine horn are also shown in the table.

Uterine Blood flow non-pregnant mare (Day 8)						
Nº of mare	Age	Uterine horn	Blood flow pixel	Mean	SD	CV (%)
2	6	LEFT	140	152,50	17,68	11,59
		LEFT	165			
17	7	RIGHT	510	325,00	261,63	80,50
		RIGHT	140			
		LEFT	292	328,00	50,91	15,52
		LEFT	364			

(continued on next page)

Table 5 (continued)

Uterine Blood flow non-pregnant mare (Day 8)						
N° of mare	Age	Uterine horn	Blood flow pixel	Mean	SD	CV (%)
38	3	RIGHT	284	536,50	74,95	22,24
		RIGHT	390			
		LEFT	595			
		LEFT	478			
40	3	RIGHT	369	341,00	313,96	92,07
		RIGHT	305			
		LEFT	563			
		LEFT	119			
43	3	RIGHT	317	440,50	8,49	2,63
		RIGHT	329			
		LEFT	490			
		LEFT	391			
46	10	RIGHT	427	398,00	41,01	10,30
		RIGHT	369			
		LEFT	686			
		LEFT	740			
48	11	RIGHT	614	713,00	38,18	5,36
		RIGHT	680			
		LEFT	889			
		LEFT	678			
50	11	RIGHT	904	783,50	149,20	19,04
		RIGHT	553			
		LEFT	729			
		LEFT	670			
52	9	RIGHT	473	673,50	283,55	42,10
		RIGHT	874			
		LEFT	544			
		LEFT	796			
		RIGHT	660	717,00	80,61	11,24
		RIGHT	774			

threshold restricted to the uterine horn in question was applied first for blood vessel detection. Following this, the total area for the region of interest (blood vessels) was calculated.

Uncited references

[1]

Author Contribution Section

PNO: Conceptualization, conceived the study and performed experiments, methodology and data curation; FEMC: performed the analysis of data with Image J (Software), validation and reviewed the draft. GGP: data curation and English review (native speaker) writing - review & editing, Software. JMO: Methodology, data curation. FJP: Conceptualization, funding acquisition, English review and statistical analysis, COF: Conceptualization, conceived the study, supervision, formal analysis and roles/writing - original draft.

Competing Interests

The authors declare that they have no known competing financial interests or personal relationships which have, or could be perceived to have, influenced the work reported in this article.

Acknowledgments

The authors received financial support for this study from the Ministerio de Economía y Competitividad-FEDER, Madrid, Spain, grants, AGL2017-83149-R and Junta de Extremadura-FEDER (GR 18008 and IB16030). JMOR holds a PhD grant from the Junta de Extremadura- FEDER (PD 18005), Mérida, Spain. GGPh holds a PhD grant from the Ministry of Science, Madrid, Spain (PRE2018-083354).

References

- [1] P Nieto-Olmedo, FE Martín-Cano, G Gaitskell-Phillips, JM Ortiz-Rodríguez, FJ Pena, C Ortega-Ferrusola, Power Doppler can detect the presence of 7-8 day conceptuses prior to flushing in an equine embryo transfer program, *Theriogenology* 145 (2020) 1–9.