



ORAL PRESENTATION

Open Access

# Telomere length, proviral load and neurologic impairment in HTLV-1-and HTLV-2-infected humans

Benjamin Usadi<sup>1,2</sup>, Roberta Bruhn<sup>1</sup>, Jue Lin<sup>3</sup>, Tzong-Hae Lee<sup>1</sup>, Elizabeth Blackburn<sup>3</sup>, Edward L Murphy<sup>3,1\*</sup>

From 16th International Conference on Human Retroviruses: HTLV and Related Viruses  
Montreal, Canada. 26-30 June 2013

## Background

Telomeres shorten with aging and short or damaged telomeres have been implicated in degenerative conditions. We hypothesized that telomere length might be altered in chronic HTLV-1 and -2 infection and could be a marker of HTLV-associated disease and viral dynamics.

## Methods

45 HTLV-1, 45 HTLV-2, and 45 seronegative subjects were selected from the larger HTLV Outcomes Study (HOST) cohort, and stratum-matched on age, sex and race/ethnicity. The telomere-to-single copy gene (T/S) ratio and HTLV-1 and -2 proviral load were measured using real-time PCR on the same PBMC samples. Unpaired t-tests, linear regression and logistic regression were used to test associations.

## Results

Ln T/S ratio was inversely associated with age among seronegatives ( $p=.006$ ) but HTLV-1 and -2 subjects did not show an inverse age association. There was no difference in mean T/S ratio between HTLV-1 (1.02), HTLV-2 (1.03) and matched seronegative (0.99) subjects. In HTLV-1 subjects, there was a borderline inverse association ( $p=0.07$ ) between T/S ratio and log<sub>10</sub> proviral load which did not persist after multivariate adjustment ( $p=0.17$ ). Among HTLV-2 subjects only, Ln T/S ratio was significantly associated ( $p=0.026$ ) with increased odds of vibration-sensation impairment.

## Conclusions

We found no evidence for an overall difference in telomere length between HTLV cases and controls but

there was a weak association between HTLV-1 proviral load and telomere length. The association between telomere length and impaired vibration sense in the HTLV-2-positive group is intriguing, and suggests avenues for future investigation of previously described neuropathy in that group.

## Authors' details

<sup>1</sup>Blood Systems Research Institute, San Francisco, CA, USA. <sup>2</sup>University of California, Berkeley, Berkeley, CA, USA. <sup>3</sup>University of California San Francisco, San Francisco, CA, USA.

Published: 7 January 2014

doi:10.1186/1742-4690-11-S1-O15

**Cite this article as:** Usadi et al.: Telomere length, proviral load and neurologic impairment in HTLV-1-and HTLV-2-infected humans. *Retrovirology* 2014 **11**(Suppl 1):O15.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



<sup>3</sup>University of California San Francisco, San Francisco, CA, USA  
Full list of author information is available at the end of the article