



# State-of-the-art review of secondary pulmonary infections in patients with COVID-19 pneumonia

Woon H. Chong<sup>1</sup> · Biplab K. Saha<sup>2</sup> · Ananthakrishnan Ramani<sup>3</sup> · Amit Chopra<sup>1</sup>

Received: 9 October 2020 / Accepted: 3 March 2021 / Published online: 11 March 2021  
© Springer-Verlag GmbH Germany, part of Springer Nature 2021

## Abstract

**Background** The incidence of secondary pulmonary infections is not well described in hospitalized COVID-19 patients. Understanding the incidence of secondary pulmonary infections and the associated bacterial and fungal microorganisms identified can improve patient outcomes.

**Objective** This narrative review aims to determine the incidence of secondary bacterial and fungal pulmonary infections in hospitalized COVID-19 patients, and describe the bacterial and fungal microorganisms identified.

**Method** We perform a literature search and select articles with confirmed diagnoses of secondary bacterial and fungal pulmonary infections that occur 48 h after admission, using respiratory tract cultures in hospitalized adult COVID-19 patients. We exclude articles involving co-infections defined as infections diagnosed at the time of admission by non-SARS-CoV-2 viruses, bacteria, and fungal microorganisms.

**Results** The incidence of secondary pulmonary infections is low at 16% (4.8–42.8%) for bacterial infections and lower for fungal infections at 6.3% (0.9–33.3%) in hospitalized COVID-19 patients. Secondary pulmonary infections are predominantly seen in critically ill hospitalized COVID-19 patients. The most common bacterial microorganisms identified in the respiratory tract cultures are *Pseudomonas aeruginosa*, *Klebsiella* species, *Staphylococcus aureus*, *Escherichia coli*, and *Stenotrophomonas maltophilia*. *Aspergillus fumigatus* is the most common microorganism identified to cause secondary fungal pulmonary infections. Other rare opportunistic infection reported such as PJP is mostly confined to small case series and case reports. The overall time to diagnose secondary bacterial and fungal pulmonary infections is 10 days (2–21 days) from initial hospitalization and 9 days (4–18 days) after ICU admission. The use of antibiotics is high at 60–100% involving the studies included in our review.

**Conclusion** The widespread use of empirical antibiotics during the current pandemic may contribute to the development of multidrug-resistant microorganisms, and antimicrobial stewardship programs are required for minimizing and de-escalating antibiotics. Due to the variation in definition across most studies, a large, well-designed study is required to determine the incidence, risk factors, and outcomes of secondary pulmonary infections in hospitalized COVID-19 patients.

**Keywords** Severe acute respiratory syndrome coronavirus 2 · SARS-CoV-2 · Coronavirus disease 2019 · COVID-19 · Secondary infection · Superimposed infection · Superinfection · Bacterial infection · Fungal infection

## Introduction

Since coronavirus disease 2019 (COVID-19) was first recognized in December 2019, it has resulted in the ongoing worldwide pandemic. COVID-19 is caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), an enveloped RNA beta-coronavirus. SARS-CoV-2 shares a similar genetic identity with severe acute respiratory syndrome coronavirus (SARS-CoV) and belongs to the sarbecovirus subgenus of the Coronaviridae family [1]. COVID-19 primarily presents as a respiratory tract infection with

✉ Woon H. Chong  
chongw@amc.edu

<sup>1</sup> Department of Pulmonary and Critical Care Medicine, Albany Medical Center, 43 New Scotland Avenue, Albany, NY, USA

<sup>2</sup> Department of Pulmonary and Critical Care, Ozarks Medical Center, West Plains, MO, USA

<sup>3</sup> Department of Infectious Disease, Albany Medical Center, Albany, NY, USA

symptoms varying from mild flu-like illness to acute respiratory distress syndrome (ARDS) [2, 3]. Viral-related respiratory infections belonging to the same family of coronaviruses such as SARS-CoV and Middle East respiratory syndrome coronavirus (MERS-CoV) have been reported to be associated with secondary bacterial and fungal infections [4–7]. However, secondary pulmonary infections in COVID-19 patients are not well described and raised an important knowledge gap. Furthermore, other infectious and non-infectious complications have been described in hospitalized COVID-19 patients strongly associated with underlying COVID-19 infection such as pneumothorax, myocarditis, and even device-related secondary infections (e.g., central venous catheter, foley catheter). [8–10]. The aim of this review is to explore the incidence of secondary bacterial and fungal pulmonary infections in hospitalized patients with COVID-19 infection. We also discuss the bacterial and fungal microorganisms identified, the time to diagnose secondary pulmonary infections, and the frequency of antibiotic use in hospitalized COVID-19 patients with suspected or confirmed secondary pulmonary infections. There is a lack of data in terms of well-defined risk factors or predictors, and associated outcomes of secondary pulmonary infections in hospitalized patients with COVID-19 infection and, therefore, will not be a major focus of this review.

## Method

A literature search was performed through MEDLINE, Pubmed, and Google Scholar using keywords of “coronavirus disease 2019 (COVID-19),” “severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2),” “secondary infection,” “superimposed infection,” “superinfection,” “bacterial infection,” “fungal infection,” “bacterial pneumonia,” “fungal pneumonia,” “bacteremia,” “fungemia,” “hospital-acquired pneumonia (HAP),” and “ventilator-associated pneumonia (VAP)” from January 1st, 2020 to December 31st, 2020. Our selection criteria comprised of articles with confirmed diagnoses of secondary bacterial and fungal pulmonary infections (defined as new microorganisms identified 48 h after admission) using respiratory tracts with corresponding blood cultures for similar microorganisms thought to be respiratory in origin in hospitalized adult COVID-19 patients. Respiratory tract cultures were defined as cultures obtained from sputum, endotracheal aspirates, and bronchoalveolar lavage (BAL). We also included articles in which the diagnoses of secondary pulmonary infections were suspected based on the description of cultures obtained that were respiratory in nature or microorganisms that are recognized to be respiratory in origin. Articles published in the English language were selected, and any cited references were reviewed to identify relevant literature in


the English language that comprised of observational studies, case reports, and series that met our selection criteria that described secondary pulmonary infections in hospitalized COVID-19 patients. We excluded articles involving COVID-19 infections in children and pregnant women; non-hospitalized COVID-19 patients; patients with pulmonary co-infections (defined as infections diagnosed at the time of admission) by non-SARS-CoV-2 viruses, bacteria, and fungal microorganisms; secondary pulmonary infections from microorganisms that were known to be colonizers such as candida; studies that included both secondary infections and co-infections from a non-pulmonary source in hospitalized COVID-19 patients; and the diagnosis of secondary pulmonary infections made during the post-mortem examination of deceased COVID-19 patients. We screened 114 studies and included 49 studies that described secondary pulmonary infections in hospitalized adult COVID-19 patients that met our criteria (Fig. 1). Of the 12 studies published in the non-English language (defined as non-English language articles and had no English translation versions/options) that described secondary pulmonary infections in hospitalized COVID-19 patients, 6 were published in Mandarin, 4 were published in Spanish, and the remaining 2 were published in French. The diagnosis of COVID-19 was made by reverse transcriptase-polymerase chain reaction (RT-PCR) in all cases from respiratory tract specimens that include nasal and pharyngeal swabs, sputum, endotracheal aspirates, and bronchoalveolar lavage (BAL).

## Results

### Incidence of secondary pulmonary infections

Among the 49 studies identified (Table 1), 28 (57%) studies were observational studies done on hospitalized COVID-19 patients; the remainder, 21 (43%), were small case series and case reports. Of the 28 observational studies, 78.6% were retrospective and 21.4% were prospective in nature. The majority of observational studies originated from China in 25% (7/28) of cases followed by 17.9% (5/28) in Spain, 14.3% (4/28) in France, 7.1% in Netherlands and USA, respectively, and the remainder in Belgium, Denmark, England, Germany, Italy, Mexico, Pakistan, and Switzerland. A total of 5,047 hospitalized patients with COVID-19-related pneumonia were identified in the 28 observational studies included in our review (Table 1). The incidence of secondary bacterial pulmonary infections in hospitalized COVID-19 patients reported was 16% (580/3,633) and ranged between 4.8–42.8% in 14 observational studies, whereas the incidence of secondary fungal infections in hospitalized COVID-19 patients was 6.3% (171/2,703) and ranged between 0.9 and 33.3% according to 18 observational studies

**Fig. 1** Flowchart for Studies Selected in Review of Hospitalized COVID-19 Patients With Secondary Pulmonary Infections



(Table 1). The majority of hospitalized COVID-19 patients who developed secondary bacterial and fungal infections were critically ill where they required ICU admission and invasive mechanical ventilation (IMV). Studies by Chang et al. , Rouze et al. , and Torrego et al. were the only observational studies that examined the incidence of secondary bacterial pulmonary infection based solely on BAL findings in COVID-19 patients requiring IMV with moderate to severe ARDS [11–13]. However, for examining the incidence of secondary fungal pulmonary infections, respiratory cultures obtained from BAL were only used in observational studies by Bartoletti et al. , Rutsaert et al. , and Van Biesen et al. [14–16].

### Microbiology of secondary pulmonary infections

Out of the 28 observational studies, 14.3% (4/28) of studies had no descriptions of the specific bacterial or fungal microorganisms identified (Table 2). The most common bacterial microorganisms identified in the respiratory tract cultures among the nine observational studies (Table 2) that reported the type and frequency of secondary bacterial infection were 21.1% (75/355) *Pseudomonas aeruginosa*, 17.2% (61/355) *Klebsiella* species, 13.5% (48/355) *Staphylococcus aureus*, 10.4% (37/355) *Escherichia coli*, and 3.1% (11/355) *Stenotrophomonas maltophilia*. However, the fungal microorganisms identified in 18 observational studies included in our review were predominantly *Aspergillus* species in which *Aspergillus fumigatus* was most frequently isolated in all

studies. Other less common *Aspergillus* species identified were *Aspergillus flavus*, *Aspergillus calidoustus*, *Aspergillus citrinoterreus*, *Aspergillus niger*, *Aspergillus terreus*, and *Aspergillus versicolor*. One observational study by Fekkar et al. reported an uncommon finding of *Mucor* species and *Fusarium proliferatum* in respiratory tract cultures of critically ill COVID-19 patients [17]. Other rare opportunistic fungal infections such as *Pneumocystis jirovecii* (PJP) had been observed in four case reports/series included in our review [18–21].

### The time to diagnosis of secondary pulmonary infections and use of antibiotics

The average time taken to diagnose secondary bacterial and fungal pulmonary infections from hospital and ICU admission among the 18 observational studies described was 10 days (ranged 2–21 days) and 9 days (ranged 4–18 days), respectively (Table 2). The reported use of empirical antibiotics was 60–100% during the current pandemic between 11 observational studies (Table 2). Furthermore, although specific data on antibiotic resistance patterns lacked in the majority of observational studies included in our review, limited observational studies had reported of detection of multidrug-resistant (MDR) microorganisms such as extended-spectrum beta-lactamase (ESBL) *Klebsiella pneumoniae*, ESBL *Escherichia coli*, MDR *Pseudomonas aeruginosa*, carbapenem-resistant *Klebsiella pneumoniae*, and Methicillin-resistant *Staphylococcus aureus* (MRSA)

**Table 1** Characteristics of Secondary Pulmonary Infections Studies in Hospitalized COVID-19 Patients With Incidence of Secondary Bacterial and Fungal Infections

Study	Region	Study Type	Total COVID-19 Patients (N)	Age	ICU (%)	IMV (%)	Hospital Mortality* (%)	Secondary bacterial infections (%)	Secondary fungal infections (%)
Observational Studies									
Alanio et al. [43]	France	Retrospective, ICU	27	63	100	100	44.4	NR	33.3
Barrasa et al. [25]	Spain	Retrospective, ICU	48	63 (mean)	100	93.7	35.4	12.5	NR
Bartoletti et al. [14]	Italy	Prospective, ICU	108	63* (median)	100	100	44.0	NR	27.7
Chang et al. [12]	USA	Retrospective, ICU	412	62 (median)	100	100	NR	15.5	NR
Du et al. [56]	China	Prospective, Hospital	179	58 (median)	NR	NR	11.7	5.6	NR
Dupont et al. [69]	France	Prospective, ICU	106	69* (median)	100	100	35.3	NR	17.9
Fekkar et al. [17]	France	Retrospective, ICU	145	55 (median)	100	100	57.1	NR	4.8
Feng et al. [32]	China	Retrospective, Hospital/ICU	410	53 (median)	14.8	8.0	8.0	9.0	NR
Fu et al. [23]	China	Retrospective, Hospital/ICU	101	69* (mean)	35.6	100	NR	5.0	0.9
Gangneux et al. [70]	France	Prospective, ICU	45	70* (mean)	100	100	28.6	NR	15.6
Garcia-Vidal et al. [22]	Spain	Retrospective, Hospital/ICU	989	62 (median)	11.9	NR	9.8	4.3	0.7
Helleberg et al. [71]	Denmark	Retrospective, ICU	25	58	100	100	100	NR	8.0
Huang et al. [30]	China	Retrospective, Hospital/ICU	41	49 (median)	46.0	5.0	15.0	10.0	NR
Karmen-Tuohy et al. [58]	USA	Retrospective, Hospital	63	60 (median)	20.6	15.8	25.3	6.3	NR
Lamoth et al. [72]	Switzerland	Retrospective, ICU	118	62* (mean)	100	100	33.3	NR	3.8
Machado et al. [35]	Spain	Prospective, ICU	239	63* (mean)	100	100	100	NR	2.5
Nasir et al. [26]	Pakistan	Retrospective, Hospital/ICU	147	71 (median)	15.6	40	60.0	4.8	6.1
Roman-Montes et al. [73]	Mexico	Retrospective, ICU	144	48* (mean)	100	NR	57.1	NR	9.7
Rouze et al. [13]	Germany	Retrospective, ICU	568	64 (mean)	100	100	NR	36.1	NR
Rutsaert et al. [15]	Belgium	Retrospective, ICU	34	66	100	100	58.8	NR	20.5
Segrelles-Calvo et al. [74]	Spain	Retrospective, ICU	215	60 (median)	100*	100*	71.4	NR	3.3

**Table 1** (continued)

Study	Region	Study Type	Total COVID-19 Patients (N)	Age	ICU (%)	IMV (%)	Hospital Mortality* (%)	Secondary bacterial infections (%)	Secondary fungal infections (%)
Torrego et al. [11]	Spain	Retrospective, ICU	93	NR	100	NR	NR	19.3	NR
Van Arkel et al. [75]	Netherlands	Retrospective, ICU	31	64	100	83.9	67.7	NR	19.4
Van Biesen et al. [16]	Netherlands	Retrospective, ICU	42	68 (mean)	100	100	22.2	NR	21.4
Wang et al. [57]	China	Retrospective, Hospital/ICU	339	71 (median)	NR	23.6	19.1	42.8	NR
White et al. [34]	England	Prospective, ICU	135	57 (median)	100	72.0	52.0	NR	18.5
Yang et al. [24]	China	Retrospective, ICU	52	60 (mean)	100	71.0	61.5	7.7	3.8
Zhou et al. [31]	China	Retrospective, Hospital/ICU	191	56 (median)	26.0	17.0	28.2	15.0	NR
Case Series/Reports				Age* (mean)	ICU* (%)	IMV* (%)	Death* (%)	Secondary bacterial infections (%)	Secondary fungal infections (%)
Abdalla et al. [76]	Qatar	Case Series, ICU	2	66	100	100	100	NR	100
Blanco et al. [19]	Spain	Case Series, Hospital	5	38	40.0	20.0	0.0	NR	20.0
Falces-Romero et al. [77]	Spain	Case Series, Hospital	10	70	70.0	70.0	70.0	NR	100
Koehler et al. [78]	Germany	Case Series, ICU	19	63	100	100	57.8	NR	26.3
Lahmer et al. [79]	Germany	Case Series, ICU	2	75	100	100	100	NR	100
Lescure et al. [80]	France	Case Series, Hospital	5	47	60.0	40.0	40.0	20.0	20.0
Sharifipour et al. [81]	Iran	Case Series, ICU	19	67	100	100	94.7	100	NR
Antinori et al. [82]	Italy	Case Report, ICU	1	56	100	100	100	NR	100
Blaize et al. [83]	France	Case Report, ICU	1	74	100	100	100	NR	NR
Duployez et al. [84]	France	Case Report, ICU	1	30 s	100	100	100	NR	NR
Fernandez et al. [85]	Argentina	Case Report, ICU	1	85	100	100	100	100	100
Ghelfenstein et al. [86]	France	Case Report, ICU	1	56	100	100	100	NR	100
Kelly et al. [21]	England	Case Report, ICU	1	50 s	100	100	100	NR	100
Mang et al. [20]	Germany	Case Report, ICU	1	52	100	100	0.0	NR	100
Meijer et al. [87]	Netherlands	Case Report, ICU	1	74	100	100	100	NR	100

**Table 1** (continued)

Case Series/Reports				Age* (mean)	ICU* (%)	IMV* (%)	Death* (%)	Secondary bacterial infections (%)	Secondary fungal infections (%)
Menon et al. [18]	USA	Case Report, ICU	1	83	100	100	0.0	NR	NR
Mohamed et al. [47]	Ireland	Case Report, ICU	1	66	100	100	100	100	100
Nasri et al. [88]	Iran	Case Report, ICU	1	42	100	100	100	NR	100
Prattes et al. [89]	Austria	Case Report, ICU	1	70	100	100	100	NR	100
Schein et al. [90]	France	Case Report, Hospital	1	87	0	0	100	NR	100
Sharma et al. [91]	Australia	Case Report, ICU	1	66	100	100	0	NR	100

\*Among those with secondary infections

ICU = intensive care unit, IMV = invasive mechanical ventilation, NR = not reported

from the respiratory tract and blood cultures in critically ill COVID-19 patients [22–26].

## Discussion

In hospitalized COVID-19 patients, the incidence of secondary pulmonary infections was low at 16% (4.8–42.8%) for bacterial infections and lower for fungal infections with an incidence of 6.3% (0.9–33.3%). However, the frequency of empirical antibiotic therapy was high at 60–100% among several observational studies included. The most common bacterial microorganisms identified in the respiratory tract cultures were 21.1% *Pseudomonas aeruginosa*, 17.2% *Klebsiella* species, 13.5% *Staphylococcus aureus*, 10.4% *Escherichia coli*, and 3.1% *Stenotrophomonas maltophilia*. *Aspergillus fumigatus* was the most common fungal microorganism identified to cause secondary pulmonary infections. Other rare opportunistic infection such as PJP was mostly confined to small case series and case reports. The overall time to diagnose secondary bacterial and fungal pulmonary infections was 10 days (2–21 days) and 9 days (4–18 days), respectively, from the time of hospital and ICU admission.

In contrast, the incidence of secondary bacterial pulmonary infections during the 2009 Influenza A pandemic is up to 7% in critically ill patients [27]. However, for secondary fungal pulmonary infections, the incidence is as high as 14% in critically ill patients with seasonal influenza [28], 29. A retrospective study by Rouze et al. reported that secondary bacterial pulmonary infections were 1.6 times more likely to occur in critically ill COVID-19 patients compared to influenza patients. Four observational studies did not report any specific type of

microorganism identified [3, 30–32]. In these studies, secondary pulmonary infections were minor secondary outcomes identified while assessing the many characteristics, risk factors, and outcomes of hospitalized COVID-19 patients. Furthermore, although 18 observational studies described secondary fungal pulmonary infections predominantly *Aspergillus fumigatus*, there was an absence of a standardized definition with the heterogeneity of diagnostic criteria used to differentiate between true infection versus colonization [33–35]. The microorganisms identified to cause secondary bacterial pulmonary infections in hospitalized COVID-19 patients are similar to microorganisms isolated during seasonal/pandemic influenza and even during the 2003 SARS outbreak [13, 36, 37]. The identification of gram-negative microorganisms in hospitalized COVID-19 patients is consistent with the type of pathogens commonly associated with hospital-acquired pneumonia involving *Pseudomonas aeruginosa*, *Klebsiella* species, *Stenotrophomonas maltophilia*, and *Acinetobacter baumannii* that does not necessarily suggest a specific preference for gram-negative infections in COVID-19 [38–41]. The time taken for the diagnosis of secondary pulmonary infections is highly variable between 2 and 21 days from hospital admission and 4–18 days from ICU admission according to the 18 observational studies included in our review (Table 2). This differs in contrast to secondary bacterial infections that are diagnosed earlier in patients with influenza infection, which are 3–6 days from the initial presentation [36, 42]. For secondary invasive pulmonary aspergillosis in influenza patients, the median time to diagnosis is between 5 and 10 days after ICU admission [15, 29]. Although all observational studies included described respiratory tract cultures obtained more than 48 h after

**Table 2** Studies in Hospitalized COVID-19 Patients With Types of Microorganisms Reported, Rate of Antibiotic Use, and Time to Diagnosis of Secondary Pulmonary Infections

Study	Top Five Bacterial Microorganisms From Respiratory Tract Culture	Fungal Microorganisms From Respiratory Tract Culture	Empirical Antibiotic Therapy (%)	Median Day to Secondary Pulmonary Infection Diagnosis From Admission
<b>Observational Studies</b>				
Alantio et al. [43]	NR	100% <i>Aspergillus fumigatus</i>	NR	NR
Barrasa et al. [25]	50% <i>Pseudomonas aeruginosa</i> , 16% <i>Enterococcus faecium</i> , 16% <i>Haemophilus influenzae</i> , 16% Methicillin-Resistant <i>Staphylococcus aureus</i> (MRSA)	NR	87.5	NR
Bartoletti et al. [14]	NR	78.9% <i>Aspergillus fumigatus</i> , 15.8% <i>Aspergillus niger</i> , 5.3% <i>Aspergillus flavus</i>	NR	4 (ICU)
Chang et al. [12]	36% <i>Klebsiella</i> spp., 23% MSSA, 11% <i>Escherichia coli</i> , 11% <i>Serratia</i> spp., 11% <i>Stenotrophomonas</i> spp.	NR	NR	NR
Du et al. [56]	<i>Acinetobacter baumannii</i> , <i>Escherichia coli</i> , <i>Klebsiella pneumoniae</i> , <i>Staphylococcus</i> species**	NR	NR	NR
Dupont et al. [69]	NR	87.5% <i>Aspergillus fumigatus</i> , 6.3% <i>Aspergillus flavus</i> , 6.3% <i>Aspergillus calidoustus</i>	NR	11 (ICU)
Fekkar et al. [17]	NR	66.7% <i>Aspergillus fumigatus</i> , 22.2% <i>Mucor</i> spp., 11.1% <i>Fusarium proliferatum</i>	NR	18 (ICU)
Feng et al. [32]	NR	NR	67.1	NR
Fu et al. [23]	29% <i>Pseudomonas aeruginosa</i> , 29% <i>Burkholderia cepacia</i> , 14% <i>Escherichia coli</i> , 14% Extended-spectrum beta-lactamase (ESBL) <i>Klebsiella pneumoniae</i> , 14% <i>Stenotrophomonas maltophilia</i>	100% <i>Aspergillus fumigatus</i>	NR	12
Gangneux et al. [70]	NR	100% <i>Aspergillus fumigatus</i>	NR	NR
Garcia-Vidal et al. [22]	33% multidrug-resistant (MDR) <i>Pseudomonas aeruginosa</i> , 25% ESBL <i>Escherichia coli</i> , 21% ESBL <i>Klebsiella pneumoniae</i> , 21% <i>Staphylococcus aureus</i>	100% <i>Aspergillus fumigatus</i>	61.9	11
Helleberg et al. [71]	NR	100% <i>Aspergillus fumigatus</i>	NR	3 (ICU)
Huang et al. [30]	NR	NR	100	>2
Karmen-Tuohy et al. [58]	29% <i>Stenotrophomonas maltophilia</i> , 29% <i>Pseudomonas aeruginosa</i> , 14% <i>Escherichia coli</i> , 14% <i>Klebsiella pneumoniae</i> , 14% <i>Staphylococcus aureus</i>	NR	NR	>6
Lamoth et al. [72]	NR	100% <i>Aspergillus fumigatus</i>	100	6 (ICU)
Machado et al. [35]	NR	66.6% <i>Aspergillus fumigatus</i> , 16.7% <i>Aspergillus citrinotereus</i> , 16.7% <i>Aspergillus lentulus</i>	100	15 (ICU)

Table 2 (continued)

Study	Top Five Bacterial Microorganisms From Respiratory Tract Culture	Fungal Microorganisms From Respiratory Tract Culture	Empirical Antibi- otic Therapy (%)	Median Day to Secondary Pulmonary Infection Diagnosis From Admission
Nasir et al. [26]	25% <i>Pseudomonas aeruginosa</i> , 25% <i>Acinetobacter</i> spp., 25% <i>Stenotrophomonas maltophilia</i> , 12.5% <i>Klebsiella pneumoniae</i> , 12.5% MRSA	60% <i>Aspergillus flavus</i> , 20% <i>Aspergillus fumigatus</i> , 20% <i>Aspergillus terreus</i>	77.7	7
Roman-Montes et al. [73]	NR	54.5% <i>Aspergillus fumigatus</i> , 27.3% <i>Aspergillus</i> spp., 9.1% <i>Aspergillus flavus</i> , 9.1% <i>Aspergillus niger</i>	NR	9
Rouze et al. [13]	22.3% <i>Pseudomonas aeruginosa</i> , 18.8% <i>Enterobacter</i> spp., 11.5% <i>Klebsiella</i> spp., 9.4% MSSA, 8.4% <i>Escherichia coli</i>	NR	93.2	NR
Rutsaert et al. [15]	NR	83% <i>Aspergillus fumigatus</i> , 17% <i>Aspergillus Flavus</i>	NR	6
Segrelles-Calvo et al. [74]	NR	42.8% <i>Aspergillus fumigatus</i> , 28.6% <i>Aspergillus flavus</i> , 28.6% <i>Aspergillus niger</i>	NR	NR
Torrego et al. [11]	39% <i>Pseudomonas aeruginosa</i> , 11% <i>Enterobacter cloacae</i> , 11% <i>Enterococcus faecalis</i> , 11% <i>Klebsiella aerogenes</i> , 11% <i>Staphylococcus aureus</i>	NR	NR	<21
Van Arkel et al. [75]	NR	100% <i>Aspergillus fumigatus</i>	NR	12
Van Biesen et al. [16]	NR	71.4% <i>Aspergillus fumigatus</i> , 14.3% <i>Aspergillus flavus</i> , 14.3% <i>Aspergillus terreus</i>	NR	5 (ICU)
Wang et al. [57]	NR	NR	NR	NR
White et al. [34]	NR	91.6% <i>Aspergillus fumigatus</i> , 8.4% <i>Aspergillus versicolor</i>	98.0	8 (ICU)
Yang et al. [24]	25% ESBK <i>Klebsiella pneumoniae</i> , 25% <i>Carbapenem-resistant Klebsiella pneumoniae</i> , 25% MDR <i>Pseudomonas aeruginosa</i> , 25% <i>Serratia marcescens</i>	50% <i>Aspergillus fumigatus</i> , 50% <i>Aspergillus flavus</i>	94.2	NR
Zhou et al. [31]	NR	NR	94.8	17
Case Series/Reports	Top Five Bacterial Microorganisms From Respiratory Tract Culture	Fungal Microorganisms From Respiratory Tract Culture	Antibiotic/ Antifungal Therapy* (%)	Median Day to Secondary Pulmonary Infection Diagnosis From Admission
Abdalla et al. [76]	NR	50% <i>Aspergillus niger</i> , 50% <i>Aspergillus terreus</i>	100	12 (ICU)
Blanco et al. [19]	NR	100% <i>Pneumocystis jirovecii</i>	60.0	12
Falces-Romero et al. [77]	NR	87.5% <i>Aspergillus fumigatus</i> , 12.5% <i>A. nidulans</i>	80.0	17.1
Koehler et al. [78]	NR	100% <i>Aspergillus fumigatus</i>	NR	NR
Lahmer et al. [79]	NR	100% <i>Aspergillus fumigatus</i>	NR	5.5 (ICU)
Lescure et al. [80]	100% <i>Acinetobacter baumannii</i>	100% <i>Aspergillus fumigatus</i>	NR	NR
Sharifipour et al. [81]	89.4% <i>Acinetobacter baumannii</i> , 5.3% MSSA, 5.3% MRSA	NR	NR	NR



Table 2 (continued)

Case Series/Reports	Top Five Bacterial Microorganisms From Respiratory Tract Culture	Fungal Microorganisms From Respiratory Tract Culture	Antibiotic/Antifungal Therapy* (%)	Median Day to Secondary Pulmonary Infection Diagnosis From Admission
Antinori et al. [82]	NR	100% <i>Aspergillus fumigatus</i>	100	9
Blaize et al. [83]	NR	100% <i>Aspergillus fumigatus</i>	100	4
Duployez et al. [84]	100% Panton-Valentine Leukocidin (PVL) Methicillin-Sensitive <i>Staphylococcus aureus</i> (MSSA)	N/A	100	5
Fernandez et al. [85]	100% <i>Enterococcus faecalis</i> , <i>Acinetobacter baumannii</i>	100% <i>Aspergillus fumigatus</i>	100	10
Ghelfenstein et al. [86]	NR	100% <i>Aspergillus fumigatus</i>	100	6
Kelly et al. [21]	NR	100% <i>Pneumocystis jirovecii</i>	100	10
Mang et al. [20]	NR	100% <i>Pneumocystis jirovecii</i>	100	20
Meijer et al. [87]	NR	100% <i>Aspergillus fumigatus</i>	100	6
Menon et al. [18]	NR	100% <i>Pneumocystis jirovecii</i>	NR	NR
Mohamed et al. [47]	<i>Klebsiella Varicola</i>	100% <i>Aspergillus fumigatus</i>	100	7
Nasri et al. [88]	NR	100% <i>Aspergillus spp.</i>	100	9
Prattes et al. [89]	NR	100% <i>Aspergillus fumigatus</i>	100	3 (ICU)
Schein et al. [90]	NR	100% <i>Aspergillus spp.</i>	100	16
Sharma et al. [91]	NR	100% <i>Aspergillus fumigatus</i>	100	16

\*Among those with secondary infections

\*\*\*Percentage frequency of bacterial or fungal microorganisms not reported

NR not reported, *Spp.* Species

admission, the variability in time to diagnosis can be due to the inconsistency on when and a lack of information on why surveillance cultures are obtained.

Bronchoscopy may be a useful tool to obtain respiratory tract cultures of sufficient quantity to help diagnose and isolate microorganisms in secondary pulmonary infections while determining the antibiotic sensitivities in hospitalized COVID-19 patients. The routine use of bronchoscopy may even lead to over-diagnosis of secondary pulmonary infections from respiratory tract colonization. According to three observational studies, the incidence of secondary bacterial pulmonary infections was 15% and more when routine bronchoscopy with BAL was performed in critically ill COVID-19 patients requiring IMV [11–13]. Four observational studies reported that the incidence of secondary fungal pulmonary infections was 20% and more in critically ill COVID-19 patients when bronchoscopy with BAL was performed routinely post-intubation, in a serial fashion, or any change in clinical status due to atelectasis, new lung infiltrates on imaging, and thick secretions [14–16, 43]. Chang et al. described that respiratory tract cultures obtained from BAL have a higher positivity rate when compared to endotracheal aspirate and a greater tendency to detect different or second microorganisms as a cause of secondary pulmonary infections [12]. In the study by Torrego et al., the microbiology findings on BAL resulted in a change in antibiotic prescribed in 83% of critically ill COVID-19 patients requiring IMV. However, the bacterial microorganisms identified such as *Pseudomonas aeruginosa*, *Klebsiella* species, *Enterobacter cloacae*, and *Staphylococcus aureus* was similar to bacterial microorganisms in mechanically ventilated non-COVID-19 patients [11]. However, bronchoscopy is often avoided as it is an aerosol-generating procedure that will predispose healthcare workers and patients to a substantial risk of further transmitting COVID-19 infection. The use of bronchoscopy in COVID-19 patients has been recommended when current respiratory samples from sputum and endotracheal aspirates are negative, in which an alternate diagnosis provided by BAL would significantly impact clinical management [44]. Nevertheless, two recent single-center retrospective studies showed no increase in the risk of COVID-19 transmission to healthcare providers when bronchoscopy is routinely performed while adhering to the proper infection control protocol [12, 45].

The current knowledge of the risk factors for secondary pulmonary infections in SARS-CoV-2 is continuously evolving but remains poorly understood. Although it is becoming apparent that secondary pulmonary infections that occur in hospitalized COVID-19 patients can be associated with worse outcomes, it remains unclear if critically ill COVID-19 patients are at a greater likelihood of developing secondary pulmonary infections. COVID-19 infection will trigger innate and adaptive immune responses, including

local immune response, recruitment of macrophages and monocytes, the release of cytokines, and prime adaptive T- and B-cell in an effort to resolve underlying inflammation [46–49]. However, in some cases, a dysfunctional immune response occurs that renders COVID-19 patients vulnerable to secondary pulmonary infections. Lymphocyte count, specifically T-cells, is substantially decreased, whereas inflammatory mediators of interleukins- (IL-)2, IL-6, IL-8, IL-10, tumor necrosis factor-alpha (TNF-a), and interferon-gamma are markedly increased within a week from COVID-19 presentation before recovering to normal levels, two weeks later [30, 50–53]. This dysregulated immune response that is seen to a greater degree in those with severe COVID-19 infections has an immunosuppression stage following the proinflammatory phase characterized by a sustained and substantial reduction in peripheral lymphocyte count [48, 50, 54]. Similar immunological findings have been described in SARS-CoV patients during the 2003 epidemic and H1N1 influenza during the 2009 pandemic [53–55]. This state of lymphocytopenia-induced immunosuppression observed in many hospitalized COVID-19 patients may explain the time taken for secondary pulmonary infection diagnosis seen in studies included in our review [30–32, 56].

Furthermore, in a multi-center study involving 410 COVID-19 patients, secondary pulmonary infections were significantly associated with outcome severity. Critically ill patients had the highest percentage of secondary pulmonary infections (34.5%) compared to severely ill (8.3%) and moderately ill (3.9%) COVID-19 patients [32]. This high rate of secondary pulmonary infections occurs despite a majority of critically ill patients (92.9%) receiving antibiotics compared to 83.3% and 59.4% in the severely ill and moderately ill groups. Five observational studies reported that among critically ill COVID-19 patients, non-survivors/critically ill patients had a greater tendency to suffer from multi-organ dysfunction and develop secondary pulmonary infections despite up to 98% of them received antibiotics [24, 30, 31, 56, 57]. In all these studies, the degree of lymphocytopenia and corticosteroids administration was significantly higher in the critically ill/non-survivor group than in other groups. Furthermore, although the nadir CD4 + T-cell count was less than 200 cells/ $10^6$ L in the majority of case reports/series describing PJP among HIV patients co-infected with COVID-19 [19–21], a case report by Menon et al. described a hospitalized COVID-19 patient diagnosed with PJP despite the absence of HIV infection. Though her nadir CD4 + T-cell count was 291 cells/ $10^6$ L and she was on chronic oral budesonide for her ulcerative colitis, the improvement with trimethoprim-sulfamethoxazole supported the diagnosis of secondary PJP infection [18]. On the contrary, a retrospective study by Karmen-Tuohy et al. reported no increased incidence of secondary bacterial or even PJP pulmonary infections in HIV-positive COVID-19 patients who were

compliant with antiretroviral therapy, regardless of their CD4 + T cell count [58]. There is no single study to the current date, which has formally assessed lymphocytopenia as a risk factor for secondary pulmonary infections. Moreover, corticosteroids are frequently used in COVID-19 patients to prevent and treat cytokine storm and ARDS, which are suspected to be partly caused by dysregulated host immune response [50, 53, 59]. Recent studies assessing the use of corticosteroids in hospitalized COVID-19 patients demonstrated that a short course of corticosteroids over ten days has shown to be beneficial in the setting of hypoxic respiratory failure requiring oxygen therapy and mechanical ventilation requirement [60, 61]. However, previous studies have demonstrated that corticosteroids may inadvertently increase the mortality and secondary infections in influenza patients, and prolong viral shedding and induce lymphocytopenia in SARS-CoV patients by down-regulating the innate and adaptive immune system.[29, 54, 62, 63] Currently, there is no formal study to assess the risk of secondary pulmonary infections associated with corticosteroids administration in COVID-19 patients. However, in our review, although the majority of patients were receiving corticosteroids, the timing of administration, duration, and the dose of corticosteroids were not clearly described.

Differentiating viral from secondary bacterial and fungal pulmonary infections remains a challenge for clinicians. This diagnostic uncertainty has contributed to the overuse of antibiotics in patients with COVID-19 viral illness. Although the incidence of secondary bacterial pulmonary infections in COVID-19 patients is low, the reported use of empirical antibiotics is 60–100% among the observational studies included in our review (Table 2). These findings vastly differed when compared to patients with seasonal/pandemic influenza, in which the reported use of empirical antibiotics was 12–50% [36]. It is essential to consider how the frequent use of empirical antibiotic therapy could affect the prevalence of multidrug-resistant bacteria. The rising number of antibiotic use may predispose COVID-19 patients, especially those who are critically ill, to sepsis from secondary multidrug-resistant bacterial infections. An observational study during the SARS outbreak in 2003 demonstrated that MRSA acquisition identified on screening using nasal swabs drastically increased from 2.2 to 3.5 cases per 100 ICU admissions (pre-SARS and post-SARS period) to 25.3% per 100 ICU admissions (during SARS period), despite extensive infection control precautions [64]. This finding coincides with the increased use of broad-spectrum empiric antibiotics (4<sup>th</sup> generation cephalosporins, fluoroquinolones, aminoglycosides, and carbapenems) during the SARS period, in which MRSA was responsible for up to 48% of microorganisms isolated in patients with VAP. Furthermore, common bacterial microorganisms identified on post-mortem examination of SARS patients were *Pseudomonas aeruginosa*, *Klebsiella*

species, and *Staphylococcus aureus*, which are known for their high resistance to broad-spectrum antibiotics [65, 66]. In the studies that we reviewed, antibiotic sensitivities of microorganisms and treatment duration were not reported even though MDR microorganisms were observed. Based on the current microbiological data from our review, it remains imperative that empiric antibiotic therapy covers multidrug-resistant microorganisms such as MRSA and ESBL that are associated with a high fatality rate when concerns exist of possible secondary pulmonary infections in critically ill COVID-19 patients [39, 40]. Our review supports the notion of frequently obtaining surveillance cultures (from sputum, endotracheal aspirate, blood, and BAL if beneficial) and daily decision-making on antibiotic requirements to de-escalate and avoid prolonged therapy that will lead to the development of antibiotic resistance.

The considerable variability in the incidence of secondary bacterial (4.8–42.8%) and fungal (0.9–33.3%) pulmonary infections reported and time taken for the diagnosis can be due to several limitations across the various observational studies included in this review. (1) The majority of the studies examining the incidence of secondary pulmonary infections are of poor quality and limited by the lack of a clear definition of secondary infections versus co-infections. There is also an absence of a standardized definition with the heterogeneity of diagnostic criteria used to differentiate between true invasive pulmonary fungal infection from colonization.[33–35] (2) Moreover, secondary pulmonary infections observed are a bystander (minor secondary outcome) result or identified during subgroup analysis while assessing the many characteristics, risk factors, and outcomes of hospitalized COVID-19 patients [3, 24, 30–32, 56, 58]. (3) It is not uncommon for early and late secondary infections to be frequently clustered together in the currently available literature for COVID-19 patients that may lead to the under- or overestimation of the exact incidence of secondary pulmonary infections, depending on the duration of the study period, especially among the 78.6% retrospective studies included in our review [67]. (4) The wide range of incidence rates reported for secondary pulmonary infections might have been due to the differences in the patient population, severity of illness, diagnostic sampling, and frequency of surveillance cultures obtained across various observational studies from multiple different countries. The routine use of bronchoscopy with BAL in critically ill COVID-19 patients where many are intubated and requiring IMV may lead to the over-diagnosis of secondary pulmonary infections [14–16, 43]. (5) The restricted search methodology that is confined to English literature as we (authors) are not well-versed in other languages during this global pandemic likely contribute to the under-recognition of the true incidence of secondary pulmonary infections. (6) Furthermore, the high mortality rate associated with COVID-19 pneumonia may be an independent competing risk factor for the development of

late secondary infection, leading to an unintended underestimation of the actual risk in non-deceased COVID-19 patients [67]. (7) Lastly, the widespread use of empirical antibiotics, analgesics, and corticosteroids likely mask underlying symptoms of infections, and lead to the delay and also underdiagnosis of secondary pulmonary infections. This could be due to the lack of routine surveillance cultures obtained because of fear towards COVID-19 transmission to health care professionals with prolonged patient contact [68]. These explain the variable incidence rate and inability to effectively perform a meta-analysis to determine better the incidence, risk factor, prognostic marker, and secondary pulmonary infection outcome in COVID-19 patients.

## Conclusion

Our review on secondary pulmonary infections is limited by the lack of a clear definition of secondary infections versus co-infections, the inconsistency of the type microorganisms identified and time that surveillance cultures are obtained, the lack of information available on the associated antibiotic sensitivities of microorganisms, and duration of antibiotic treatment across various observational studies, small case reports and series, and variability in clinical characteristics reported in hospitalized COVID-19 patients. Additionally, with an observed strain being placed on the healthcare systems during the ongoing COVID-19 pandemic, there is a need for organized antimicrobial stewardship programs in the hospital to minimize the use of unnecessary empiric antibiotics and de-escalation of antibiotics when possible. As variation continues to exist on what constitutes a secondary infection (that we defined as infections occurring 48 h after admission) due to the lack of clear and consistent definition among many observational studies, we hope that a large, well-designed study can be performed in the future to accurately determine the incidence, microorganisms, risk factors, predictors, and outcomes of secondary pulmonary infections in hospitalized COVID-19 patients.

**Author contribution** All authors had access to the data and were involved in writing the manuscript.

**Funding** None.

## Declarations

**Conflict of interest** None.

## References

- Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med.* 2020;382:727–33.
- Guan W, Ni Z, Hu Y, Liang W, Ou C, He J, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med.* 2020;382:1708–20.
- Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan. *China JAMA.* 2020;323:1061–9.
- Assiri A, Al-Tawfiq JA, Al-Rabeeh AA, Al-Rabiah FA, Al-Hajjar S, Al-Barrak A, et al. Epidemiological, demographic, and clinical characteristics of 47 cases of Middle East respiratory syndrome coronavirus disease from Saudi Arabia: a descriptive study. *Lancet Infect Dis.* 2013;13:752–61.
- Tsang KW, Ho PL, Ooi GC, Yee WK, Wang T, Chan-Yeung M, et al. A cluster of cases of severe acute respiratory syndrome in Hong Kong. *N Engl J Med.* 2003;348:1977–85.
- Ksiazek TG, Erdman D, Goldsmith CS, Zaki SR, Peret T, Emery S, et al. A novel coronavirus associated with severe acute respiratory syndrome. *N Engl J Med.* 2003;348:1953–66.
- Lee N, Cameron P, Yung MY, Leung CB, Szeto CC (2003) A major outbreak of severe acute respiratory syndrome in Hong Kong. *The New England Journal of Medicine* 9.
- Seitz T, Hoepler W, Weseslindtner L, Aberle JH, Aberle SW, Puchhammer-Stoeckl E, et al. Successful management of the first reported case in Austria of COVID-19 with ARDS. *Infection.* 2020;48:647–51.
- Spiro JE, Sisovic S, Ockert B, Böcker W, Siebenbürger G. Secondary tension pneumothorax in a COVID-19 pneumonia patient: a case report. *Infection.* 2020;48:941–4.
- Zeng J-H, Liu Y-X, Yuan J, Wang F-X, Wu W-B, Li J-X, et al. First case of COVID-19 complicated with fulminant myocarditis: a case report and insights. *Infection.* 2020;48:773–7.
- Torrego A, Pajares V, Fernández-Arias C, Vera P, Mancebo J. Bronchoscopy in COVID-19 Patients with Invasive Mechanical Ventilation: A Center Experience. *Am J Respir Crit Care Med.* 2020 May 15;rccm.202004–0945LE.
- Chang SH, Jiang J, Kon ZN, Williams DM, Geraci TC, Smith DE, et al. Safety and Efficacy of Bronchoscopy in Critically Ill Patients With Coronavirus Disease 2019. *Chest.* 2020 Oct;S001236922034873X.
- On behalf of the coVAPid study Group, Rouzé A, Martin-Loeches I, Povoja P, Makris D, Artigas A, et al. Relationship between SARS-CoV-2 infection and the incidence of ventilator-associated lower respiratory tract infections: a European multicenter cohort study. *Intensive Care Med* [Internet]. 2021 Jan 3 [cited 2021 Jan 23]; <http://link.springer.com/https://doi.org/10.1007/s00134-020-06323-9>
- Bartoletti M, Pascale R, Cricca M, Rinaldi M, Maccaro A, Buscini L, et al. Epidemiology of invasive pulmonary aspergillosis among intubated patients with covid-19: a prospective study. *Clin Infect Dis.* 2020;28:ciaa1065.
- Rutsaert L, Steinfurt N, Van Hunsel T, Bomans P, Naesens R, Mertes H, et al. COVID-19-associated invasive pulmonary aspergillosis. *Ann Intensive Care.* 2020;10:71.
- Van Biesen S, Kwa D, Bosman RJ, Juffermans NP. Detection of invasive pulmonary aspergillosis in covid-19 with nondirected BAL. *Am J Respir Crit Care Med.* 2020;202:1171–3.
- Fekkar A, Lampros A, Mayaux J, Poignon C, Demeret S, Constantin J-M, et al. Occurrence of invasive pulmonary fungal infections in severe COVID-19 patients admitted to the ICU. *Am J Respir Crit Care Med.* 2020;203:307–17.
- Menon AA, Berg DD, Brea EJ, Deutsch AJ, Kidia KK, Thurber EG, et al. A Case of COVID-19 and *Pneumocystis jirovecii* Co-infection. *Am J Respir Crit Care Med.* 2020;rccm.202003–0766LE.

19. Blanco JL, Ambrosioni J, Garcia F, Martínez E, Soriano A, Mallolas J, et al. COVID-19 in patients with HIV: clinical case series. *Lancet HIV*. 2020;7:e314–6.
20. Mang S, Kaddu-Mulindwa D, Metz C, Becker A, Seiler F, Smola S, et al. Pneumocystis jirovecii pneumonia and severe acute respiratory syndrome coronavirus 2 coinfection in a patient with newly diagnosed HIV-1 infection. *Clin Infect Dis*. 2020;1:ciaa906.
21. Kelly S, Waters L, Cevik M, Collins S, Lewis J, Wu M-S, et al. Pneumocystis pneumonia, a COVID-19 mimic, reminds us of the importance of HIV testing in COVID-19. *Clin Med*. 2020;20:590–2.
22. Garcia-Vidal C, Sanjuan G, Moreno-García E, Puerta-Alcalde P, Garcia-Pouton N, Chumbita M, et al. Incidence of co-infections and superinfections in hospitalised patients with COVID-19: a retrospective cohort study. *Clinical Microbiology and Infection*. 2020 Jul;S1198743X2030450X.
23. Fu Y, Yang Q, Xu M, Kong H, Chen H, Fu Y, et al. Secondary bacterial infections in critical Ill patients with coronavirus disease 2019. *Open Forum Infect Dis*. 2020;7:220.
24. Yang X, Yu Y, Xu J, Shu H, Xia J, Liu H, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med*. 2020. [https://doi.org/10.1016/S2213-2600\(20\)30079-5](https://doi.org/10.1016/S2213-2600(20)30079-5).
25. Barrasa H, Rello J, Tejada S, Martín A, Balziskueta G, Vinuesa C, et al. SARS-CoV-2 in Spanish intensive care units: early experience with 15-day survival in Vitoria. *Anaesth Crit Care Pain Med*. 2020;39:553–61.
26. Nasir N, Farooqi J, Mahmood SF, Jabeen K. COVID-19-associated pulmonary aspergillosis (CAPA) in patients admitted with severe COVID-19 pneumonia: an observational study from Pakistan. *Mycoses*. 2020;63:766–70.
27. Rice TW, Rubinson L, Uyeki TM, Vaughn FL, John BB, Miller RR, et al. Critical illness from 2009 pandemic influenza A virus and bacterial coinfection in the United States. *Crit Care Med*. 2012;40:1487–98.
28. Schauwvlieghe AFAD, Rijnders BJA, Philips N, Verwijs R, Vanderbeke L, Van Tienen C, et al. Invasive aspergillosis in patients admitted to the intensive care unit with severe influenza: a retrospective cohort study. *Lancet Respir Med*. 2018;6:782–92.
29. Vanderbeke L, Spriet I, Breynaert C, Rijnders BJA, Verweij PE, Wauters J. Invasive pulmonary aspergillosis complicating severe influenza: epidemiology, diagnosis and treatment. *Curr Opin Infect Dis*. 2018;31:471–80.
30. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet*. 2020;395:497–506.
31. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *The Lancet*. 2020;395:1054–62.
32. Feng Y, Ling Y, Bai T, Xie Y, Huang J, Li J, et al. COVID-19 with different severities: a multicenter study of clinical features. *Am J Respir Crit Care Med*. 2020;201:1380–8.
33. Bassetti M, Giacobbe DR, Grecchi C, Rebuffi C, Zuccaro V, Scudeller L, et al. Performance of existing definitions and tests for the diagnosis of invasive aspergillosis in critically ill, adult patients: a systematic review with qualitative evidence synthesis. *J Infect*. 2020;81:131–46.
34. White PL, Dhillon R, Cordey A, Hughes H, Faggian F, Soni S, et al. A national strategy to diagnose COVID-19 associated invasive fungal disease in the ICU. *Clin Infect Dis*. 2020. <https://doi.org/10.1093/cid/ciaa1298>.
35. Machado M, Valerio M, Álvarez-Uría A, Olmedo M, Veintimilla C, Padilla B, et al. Invasive pulmonary aspergillosis in the COVID-19 era: An expected new entity. *Mycoses*. 2020 Nov 29;myc.13213.
36. Klein EY, Monteforte B, Gupta A, Jiang W, May L, Hsieh Y-H, et al. The frequency of influenza and bacterial coinfection: a systematic review and meta-analysis. *Influenza Other Respir Viruses*. 2016;10:394–403.
37. Vaillancourt M, Jorth P. The unrecognized threat of secondary bacterial infections with COVID-19. *mBio*. 2020;11:e01806–20.
38. Jones RN. Microbial etiologies of hospital-acquired bacterial pneumonia and ventilator-associated bacterial pneumonia. *Clin Infect Dis*. 2010;51:S81–7.
39. Bereket W, Hemalatha K, Getenet B, Wondwossen T, Solomon A, Zeynudin A, et al. Update on bacterial nosocomial infections. *Eur Rev Med Pharmacol Sci*. 2012;16:1039–44.
40. Cabrera CE, Gómez RF, Zuñiga AE, Corral RH, López B, Chávez M. Epidemiology of nosocomial bacteria resistant to antimicrobials. *Colombia Medica*. 2011;1:117–25.
41. Silvestri L, Sarginson RE, Hughes J, Milanese M, Gregori D, Van Saene HKF. Most nosocomial pneumonias are not due to nosocomial bacteria in ventilated patients: evaluation of the accuracy of the 48 h time cut-off using carriage as the gold standard. *Anaesth Intens Care*. 2002;30:275–82.
42. Lansbury L, Lim B, Baskaran V, Lim WS. Co-infections in people with COVID-19: a systematic review and meta-analysis. *J Infect*. 2020;81:266–75.
43. Alanio A, Dellièrè S, Fodil S, Bretagne S, Mégarbane B. Prevalence of putative invasive pulmonary aspergillosis in critically ill patients with COVID-19. *Lancet Respir Med*. 2020;8:e48–9.
44. Wahidi MM, Lamb C, Murgu S, Musani A, Shojaee S, Sachdeva A, et al. American Association for Bronchology and Interventional Pulmonology (AABIP) Statement on the use of bronchoscopy and respiratory specimen collection in patients with suspected or confirmed COVID-19 infection. *J Bronchol Int Pulmonol*. 2020;27:52–4.
45. Gao CA, Bailey JI, Walter JM, Coleman JM, Malsin ES, Argento AC, et al. Bronchoscopy on intubated COVID-19 patients is associated with low infectious risk to operators. *Annals ATS*. 2021. <https://doi.org/10.1513/AnnalsATS.202009-1225RL>.
46. Arastehfar A, Carvalho A, van de Veerdonk FL, Jenks JD, Koehler P, Krause R, et al. COVID-19 associated pulmonary aspergillosis (CAPA)—from immunology to treatment. *JoF*. 2020;6:91.
47. Mohamed A, Rogers TR, Talento AF. COVID-19 Associated invasive pulmonary aspergillosis: diagnostic and therapeutic challenges. *JoF*. 2020;6:115.
48. Thompson Iii GR, Cornely OA, Pappas PG, Patterson TF, Hoenigl M, Jenks JD, et al. Invasive aspergillosis as an under-recognized superinfection in COVID-19. *Open Forum Infect Dis*. 2020;7:242.
49. Qin C, Zhou L, Hu Z, Zhang S, Yang S, Tao Y, et al. Dysregulation of immune response in patients with coronavirus 2019 (COVID-19) in Wuhan. *China Clin Infect Dis*. 2020;71:762–8.
50. Li H, Liu L, Zhang D, Xu J, Dai H, Tang N, et al. SARS-CoV-2 and viral sepsis: observations and hypotheses. *The Lancet*. 2020;395:1517–20.
51. Liu J, Li S, Liu J, Liang B, Wang X, Wang H, et al. Longitudinal characteristics of lymphocyte responses and cytokine profiles in the peripheral blood of SARS-CoV-2 infected patients. *EBioMedicine*. 2020;55:102763.
52. Tian S, Xiong Y, Liu H, Niu L, Guo J, Liao M, et al. Pathological study of the 2019 novel coronavirus disease (COVID-19) through postmortem core biopsies. *Mod Pathol*. 2020;33:1007–14.
53. Lin L, Lu L, Cao W, Li T. Hypothesis for potential pathogenesis of SARS-CoV-2 infection—a review of immune changes in patients with viral pneumonia. *Emerg Microbes Infect*. 2020;9:727–32.
54. Gu J, Gong E, Zhang B, Zheng J, Gao Z, Zhong Y, et al. Multiple organ infection and the pathogenesis of SARS. *J Exp Med*. 2005;202:415–24.

55. Wong RSM. Haematological manifestations in patients with severe acute respiratory syndrome: retrospective analysis. *BMJ*. 2003;326:1358–62.
56. Du R-H, Liang L-R, Yang C-Q, Wang W, Cao T-Z, Li M, et al. Predictors of mortality for patients with COVID-19 pneumonia caused by SARS-CoV-2: a prospective cohort study. *Eur Respir J*. 2020;55:2000524.
57. Wang L, He W, Yu X, Hu D, Bao M, Liu H, et al. Coronavirus disease 2019 in elderly patients: characteristics and prognostic factors based on 4-week follow-up. *J Infect*. 2020;80:639–45.
58. Karmen-Tuohy S, Carlucci PM, Zervou FN, Zacharioudakis IM, Rebick G, Klein E, et al. Outcomes among HIV-positive patients hospitalized with COVID-19. *J Acquir Immune Defic Syndr*. 2020;85:6–10.
59. Berlin DA, Gulick RM, Martinez FJ. Severe Covid-19. *N Engl J Med*. 2020;383:2451–60.
60. Chopra A, Chieng HC, Austin A, Tiwari A, Mehta S, Nautiyal A, et al. Corticosteroid administration is associated with improved outcome in patients with severe acute respiratory syndrome coronavirus 2-related acute respiratory distress syndrome. *Crit Care Explor*. 2020;2:e0143.
61. The RECOVERY Collaborative Group. Dexamethasone in hospitalized patients with Covid-19—preliminary report. *N Engl J Med*. 2020;384:693–704.
62. Moreno G, Rodríguez A, Reyes LF, Gomez J, Sole-Violan J, Díaz E, et al. Corticosteroid treatment in critically ill patients with severe influenza pneumonia: a propensity score matching study. *Intensive Care Med*. 2018;44:1470–82.
63. Russell CD, Millar JE, Baillie JK. Clinical evidence does not support corticosteroid treatment for 2019-nCoV lung injury. *The Lancet*. 2020;395:473–5.
64. Yap FHY, Gomersall CD, Fung KSC, Ho P-L, Ho O-M, Lam PKN, et al. Increase in methicillin-resistant staphylococcus aureus acquisition rate and change in pathogen pattern associated with an outbreak of severe acute respiratory syndrome. *Clin Infect Dis*. 2004;39:511–6.
65. Franks TJ, Chong PY, Chui P, Galvin JR, Lourens RM, Reid AH, et al. Lung pathology of severe acute respiratory syndrome (SARS): a study of 8 autopsy cases from Singapore. *Hum Pathol*. 2003;34:743–8.
66. Tse GM-K. Pulmonary pathological features in coronavirus associated severe acute respiratory syndrome (SARS). *J Clin Pathol*. 2004;57:260–5.
67. Bassetti M, Kollef MH, Timsit J-F. Bacterial and fungal superinfections in critically ill patients with COVID-19. *Intensive Care Med*. 2020;46:2071–4.
68. Rawson TM, Moore LSP, Zhu N, Ranganathan N, Skolimowska K, Gilchrist M, et al. Bacterial and fungal co-infection in individuals with coronavirus: a rapid review to support COVID-19 antimicrobial prescribing. *Clinical Infectious Diseases*. 2020. <https://doi.org/10.1093/cid/ciaa530>.
69. Dupont D, Menotti J, Turc J, Miossec C, Wallet F, Richard J-C, et al. Pulmonary aspergillosis in critically ill patients with Coronavirus Disease 2019 COVID-19. *Med Mycol*. 2019. <https://doi.org/10.1093/mmy/myaa078>.
70. Gangneux J-P, Reizine F, Guegan H, Pinceaux K, Le Balch P, Prat E, et al. Is the COVID-19 pandemic a good time to include aspergillus molecular detection to categorize Aspergillosis in ICU patients? A monocentric experience. *JoF*. 2020;6:105.
71. Helleberg M, Steensen M, Arendrup MC. Invasive aspergillosis in patients with severe COVID-19 pneumonia. *Clin Microbiol Infect*. 2021;27:147–8.
72. Lamoth F, Glampedakis E, Boillat-Blanco N, Oddo M, Pagani J-L. Incidence of invasive pulmonary aspergillosis among critically ill COVID-19 patients. *Clin Microbiol Infect*. 2020;26:1706–8.
73. Roman-Montes Authors CM, Martinez-Gamboa A, Diaz-Lomeli P, Cervantes-Sanchez A, Rangel-Cordero A, Sifuentes-Osornio J, et al. Accuracy of galactomannan testing on tracheal aspirates in COVID-19-associated pulmonary aspergillosis. *Mycoses*. 2020. <https://doi.org/10.1111/myc.13216>.
74. Segrelles-Calvo G, Araújo GRS, Llopis-Pastor E, Carrillo J, Hernández-Hernández M, Rey L, et al. Prevalence of opportunistic invasive aspergillosis in COVID-19 patients with severe pneumonia. *Mycoses*. 2020. <https://doi.org/10.1111/myc.13219>.
75. van Arkel ALE, Rijpstra TA, Belderbos HNA, van Wijngaarden P, Verweij PE, Bentvelsen RG. COVID-19 associated pulmonary Aspergillosis. *Am J Respir Crit Care Med*. 2020;202:132–5.
76. Abdalla S, Almaslamani MA, Hashim SM, Ibrahim AS, Omrani AS. Fatal Coronavirus Disease 2019-associated pulmonary Aspergillosis; a report of two cases and review of the literature. *IDCases*. 2020;22:e00935.
77. Falces-Romero I, Ruiz-Bastián M, Díaz-Pollán B, Maseda E, García-Rodríguez J, SARS-CoV-2 Working Group. Isolation of *Aspergillus* spp. in respiratory samples of patients with COVID-19 in a Spanish Tertiary Care Hospital. *Mycoses*. 2020. <https://doi.org/10.1111/myc.13155>.
78. Koehler P, Cornely OA, Böttiger BW, Dusse F, Eichenauer DA, Fuchs F, et al. COVID-19 associated pulmonary aspergillosis. :7.
79. Lahmer T, Rasch S, Spinner C, Geisler F, Schmid RM, Huber W. Invasive pulmonary aspergillosis in severe coronavirus disease 2019 pneumonia. *Clin Microbiol Infect*. 2020;26:1428–9.
80. Lescuré F-X, Bouadma L, Nguyen D, Parisey M, Wicky P-H, Behillil S, et al. Clinical and virological data of the first cases of COVID-19 in Europe: a case series. *Lancet Infect Dis*. 2020;20:697–706.
81. Sharifipour E, Shams S, Esmkhani M, Khodadadi J, Fotouhi-Ardakani R, Koohpaei A, et al. Evaluation of bacterial co-infections of the respiratory tract in COVID-19 patients admitted to ICU. *BMC Infect Dis*. 2020;20:646.
82. Antinori S, Rech R, Galimberti L, Castelli A, Angeli E, Fossali T, et al. Invasive pulmonary aspergillosis complicating SARS-CoV-2 pneumonia: a diagnostic challenge. *Travel Med Infect Dis*. 2020;38:101752.
83. Blaize M, Mayaux J, Nabet C, Lampros A, Marcelin A-G, Thellier M, et al. Fatal invasive aspergillosis and coronavirus disease in an immunocompetent patient. *Emerg Infect Dis*. 2020;26:1636–7.
84. Duployez C, Le Guern R, Tinez C, Lejeune A-L, Robriquet L, Six S, et al. Panton-valentine leukocidin-secreting staphylococcus aureus pneumonia complicating COVID-19. *Emerging Infect Dis*. 2020;26:1939–41.
85. Fernandez NB, Caceres DH, Beer KD, Irrazabal C, Delgado G, Farias L, et al. Ventilator-associated pneumonia involving *Aspergillus flavus* in a patient with coronavirus disease (COVID-19) from Argentina. *Med Mycol Case Rep*. 2019. <https://doi.org/10.1016/j.mmcr.2020.07.001>.
86. Ghelfenstein-Ferreira T, Saade A, Alanio A, Bretagne S, Araujo de Castro R, Hamane S, et al. Recovery of a triazole-resistant *Aspergillus fumigatus* in respiratory specimen of COVID-19 patient in ICU—a case report. *Medical Mycology Case Reports*. 2020 Jul;S2211753920300439.
87. Meijer EFJ, Dofferhoff ASM, Hoiting O, Buil JB, Meis JF. Azole-resistant COVID-19-associated pulmonary aspergillosis in an immunocompetent host: a case report. *JoF*. 2020;6:79.

88. Nasri E. Fatal invasive pulmonary Aspergillosis in COVID-19 patient with acute myeloid leukemia in Iran. *Mycopathologia*. 2020;185:1077–84.
89. Prattes J, Valentin T, Hoenigl M, Talakic E, Reisinger AC, Eller P. (2020) Invasive pulmonary aspergillosis complicating COVID-19 in the ICU—a case report. *Med Mycol Case Rep*. May;S2211753920300300.
90. Schein F, Munoz-Pons H, Mahinc C, Grange R, Cathébras P, Flori P. Fatal aspergillosis complicating severe SARS-CoV-2 infection: a case report. *Journal de Mycologie Médicale*. 2020;30:101039.
91. Sharma A, Hofmeyr A, Bansal A, Thakkar D, Lam L, Harrington Z, et al. COVID-19 associated pulmonary aspergillosis (CAPA): an Australian case report. *Medical Mycology Case Reports*. 2020 Jun;S2211753920300397.