



Article

# Analysis of Blood Concentrations of Zinc, Germanium, and Lead and Relevant Environmental Factors in a Population Sample from Shandong Province, China

Long Li <sup>1,2</sup>, Guang Xu <sup>2</sup>, Hua Shao <sup>2,\*</sup>, Zhi-Hu Zhang <sup>2,\*</sup>, Xing-Fu Pan <sup>3</sup> and Jin-Ye Li <sup>4</sup>

<sup>1</sup> School of Medicine and Life Sciences, University of Jinan-Shandong Academy of Medical Sciences, Jinan 250062, China; tianya584526@163.com

<sup>2</sup> Shandong Academy Occupational Health and Occupational Medicine, 18877 Jingshi Road, Jinan 250062, China; sunshinefeng123@163.com

<sup>3</sup> Department of Occupational Diseases Control and Prevention, Fengtai Center for Disease Control and Prevention, Beijing 100071, China; cengcengc3@163.com

<sup>4</sup> Shandong Provincial Western Hospital, Jinan 250022, China; lijinyejn@163.com

\* Correspondence: chinashaohua5888@163.com (H.S.); zyhw@sina.com (Z.-H.Z.); Tel.: +86-82-595-888 (H.S.)

Academic Editor: Helena Solo-Gabriele

Received: 11 January 2017; Accepted: 17 February 2017; Published: 24 February 2017

**Abstract:** Trace elements, including zinc (Zn) and germanium (Ge), are essential for health; deficiency or excess levels of trace elements results is harmful. As a result of industrial and agricultural production, Pb widely exists in people's living environment. It is absorbed mainly through the respiratory and digestive tracts, producing systemic harm. Reference values for a normal, healthy population are necessary for health assessment, prevention and treatment of related diseases, and evaluation of occupational exposures. Reference ranges for the Chinese population have not been established. From March 2009 to February 2010; we collected data and blood samples ( $n = 1302$ ) from residents aged 6–60 years living in Shandong Province, China. We measured blood concentrations of Zn, Ge, and Pb using inductively coupled plasma mass spectrometry to determine reference ranges. Results were stratified by factors likely to affect the concentrations of these trace elements: sex, use of cosmetics or hair dye, age, alcohol intake, smoking habits, and consumption of fried food. The overall geometric mean (GM) concentrations (95% confidence interval) were 3.14 (3.08–3.20) mg/L for Zn, 19.9 (19.3–20.6)  $\mu\text{g/L}$  for Ge, and 24.1 (23.2–25.1)  $\mu\text{g/L}$  for Pb. Blood Zn concentrations were higher in women than in men ( $p < 0.001$ ), while the opposite was found for Pb ( $p < 0.001$ ) and sex did not influence Ge ( $p = 0.095$ ). Alcohol use was associated with higher blood concentrations of Zn ( $p = 0.002$ ), Ge ( $p = 0.002$ ), and Pb ( $p = 0.001$ ). The GM concentration of Zn was highest in 20–30-year-olds ( $p < 0.001$ ), while Pb concentrations were highest in 12–16-year-olds ( $p < 0.001$ ). Use of hair dye was associated with lower blood concentrations of Ge ( $p < 0.05$ ). GM blood concentrations of Pb differed significantly between those who consumed fried foods 1–2 times/month (18.7  $\mu\text{g/L}$ ), 1–2 times/week (20.9  $\mu\text{g/L}$ ), and every day (28.5  $\mu\text{g/L}$ ;  $p < 0.001$ ). Blood Pb concentrations were higher in subjects who used cosmetics ( $p < 0.05$ ), hair dye ( $p < 0.05$ ), and who smoked cigarettes ( $p < 0.001$ ) than in those who did not.

**Keywords:** reference values; blood; metal; trace element

## 1. Introduction

The development of mining and manufacturing industries has led to a rise in occupational and non-occupational metal poisoning, which has become a major public health problem. To assess and

monitor risk, reference values of potentially harmful metals-including trace elements required by the body-for a normal, healthy population are essential for comparison purposes [1,2]. Reference values enable assessment of population health, disease prevention and treatment [3], and evaluation of occupational exposures and environmental conditions [4].

Trace elements are essential to the human body, even though the daily demand for them is very low. They are involved in bodily physiology [5–7] and are important components of vitamins, hormones, and enzyme systems [8,9]. Insufficient or excess trace element concentrations in the body will result in harm [10].

The normal human body contains 2–4 g of zinc (Zn), one of the fourteen essential trace elements [11,12]. Biologically, Zn has three main functions: catalysis, adjustment, and structure. Zn, an enzyme constituent [11,13], also catalyzes enzymatic reactions and plays an important role in metabolism [14,15], tissue respiration [16,17] and regeneration [18,19], and growth [17,20]. Zn may also affect the effectiveness of hormone receptor and target organ responses; hormone production, storage, and secretion; and sexual development [21,22]. Zn stabilizes insulin structure [23,24], maintains normal dark adaptation capacity [25,26], and is a component of salivary proteins that promote the sense of taste and appetite [27,28]. It is involved in iron transport and transfer [29–31] and enhances the immune system [32–34]. Zn can inhibit lipid peroxidation and thiol oxidation of biomembranes, and along with copper-protein, catalase, and vitamin E, maintains cell structure [35–38]. Zn deficiency causes digestive disorders, delayed sexual maturity/development of secondary sexual characteristics resulting in stunted growth, skin disease, stomatitis, alopecia [29,30,39], and can impair immune function. Zn overdoses can cause metal fume fever, reduce serum high-density lipoprotein-cholesterol levels, and lead to iron deficiency anemia and copper deficiency [40,41].

Germanium (Ge), another essential trace element, is widely distributed in the body. It is a constituent of the amino acid guanidine and the enzymes cytochrome oxidase and carbonic anhydrase. It is also distributed in the brain cortex and is a component of cell walls, chromosomes, vesicles, lysosomes, and cytoplasmic matrix. Ge has anti-mutation, anti-cancer [42,43], anti-aging [44,45], anti-malarial [46,47], and anti-inflammatory [48,49] properties. It can stimulate hematopoiesis [50] and enhance immune function [51–53]. Intake is through consumption of water, food, or drugs [54]. Excessive Ge can damage the kidneys [48,55,56], nervous system [57–59], and lungs [60].

Lead (Pb), widely distributed in the atmosphere, soil, and food, enters the body via the respiratory and digestive systems and is deposited in bones. Unlike Zn and Ge, Pb is not beneficial to health. Pb can result in neurologic [61–63], skeletal, reproductive [64,65], hematopoietic [66,67], and urinary system [68,69] toxicity.


Considering their beneficial (Zn and Ge) and harmful (Pb) effects, periodic biological monitoring of these trace elements should be conducted. In Europe and the USA, such monitoring is generally performed at the national level. However, in Shandong Province, China, no previous biomonitoring studies have been conducted, and it is not known whether the geographical or social characteristics of this area make it nationally representative. Therefore, we aimed to measure the blood  $^{66}\text{Zn}$ ,  $^{72}\text{Ge}$ , and  $^{208}\text{Pb}$  concentrations of the general population in Shandong Province, stratified by sex, age, smoking, alcohol consumption, cosmetic and hair dye use, and fried food consumption, to form a basis for biological monitoring and scientific research.

## 2. Materials and Methods

### 2.1. Subject Selection

Shandong Province, located in the east of China, has a population of 92.82 million people and a surface area of 153,300 km<sup>2</sup>. Using cluster sampling, the study subjects were selected as follows: First, Shandong Province was divided into three socioeconomic levels. Second, we randomly sampled one city from each socioeconomic level. Last, we randomly selected 1302 subjects in each community. Inclusion criteria were as follows: (1) living in the local area for at least five years; (2) living in

areas without relevant industrial pollution; (3) no history of liver or kidney diseases, diabetes, hyperthyroidism, cancer, or other chronic diseases; (4) no acute infection; (5) no use of pharmaceutical preparations or dietary supplements containing trace elements within the past 3 months; and (6) age 6–60 years old. Selected areas are shown in Figure 1: 418 research subjects from Qingdao were included, 345 from Jinan, and 539 from Heze.



**Figure 1.** Location of the study population. Shandong Province, located in the east of China, has a population of 92.8 million people and a land area of 153,300 km<sup>2</sup>. We included 418 research participants from Qingdao, 345 from Jinan, and 539 from Heze. ☆: Sample collection areas.

All participants completed a questionnaire regarding personal information, lifestyle and eating habits, and medical history. All questionnaires were recovered and meet the requirements. A total of 1302 blood samples and questionnaires were collected from March 2009 to February 2010. Written informed consent was obtained from each subject. The study was conducted in accordance with the Declaration of Helsinki, and the study was approved by the Ethical Censorship Committee of the Shandong Academy of Medical Sciences (YKYL-2009066). Participants agreed to the use of their blood samples for this biological monitoring research.

## 2.2. Sample Preparation and Analysis

All samples were collected and processed in a clean environment. Blood samples (6 mL) were collected in vacutainers containing lithium heparin (BD, Bergen, NJ, USA), and were immediately transferred to 2 mL freezing tubes (Axygen, San Francisco, CA, USA) after thorough mixing. All samples were stored at  $-80^{\circ}\text{C}$  until analysis. Before analyzing, the samples were warmed to room temperature ( $23^{\circ}\text{C}$ ). As described in previous literature [70], 0.5 mL of blood was added to 4.5 mL of a diluent containing 0.01% (V/V) Triton-X-100 (Sigma Aldrich, Bergen, NJ, USA) and 0.5% ultrapure concentrated nitric acid (Merck, Darmstadt, Germany). Samples were vortexed in a table-top vortexer (Multi Reax [XWT-204], Heidolph, San Francisco, CA, USA). Concentrations of Zn, Ge, and Pb in the diluted samples were then quantified using inductively coupled plasma mass spectrometry (ICP-MS, Thermo Fisher, Waltham, MA, USA). Yttrium (Y) solution with concentration of 10  $\mu\text{g/L}$  was used as the internal standard. The 0.01% Triton-X-100 and 0.5% ultrapure by 10 determination will respond to signals corresponding to 3 times the standard deviation of the analyte concentration as the detection limit; the 10 times value was determined by a solvent blank, and the response signals corresponding to 10 times the standard deviation of the measured element concentrations were defined as a quantification limit. Since the method for sample processing calls for dilution by a factor of 10,

the limit of detection and limit of quantification were both multiplied by 10. This resulted in limits of detection for Zn, Ge, and Pb of 4.30, 0.18 and 0.28 µg/L, respectively.

### 2.3. Quality Control


Contamination in the pre-analytic phase during sample collection may lead to inaccurate measurements [71]. Therefore, to minimize contamination, we pre-tested the heparin vacutainers and frozen vials. We soaked 20 vacutainers and 20 vials in 1% (V/V) ultrapure nitric acid for one hour and then determined the metal concentrations in the soaking solution using ICP-MS. The concentrations of Zn, Pb, and Ge in these vacutainers and vials were lower than the respective detection limits. Sets of 30 samples were processed after determination of a single point standard solution, provided that the determination result was within the allowable range (deviation < 10%) using nickel, arsenic, molybdenum, and tungsten as reference elements. The ICP-MS measurement procedures were referenced to previous research [72]. Sample preparation and analysis were performed by investigators with professional training and ICP-MS operators with professional experience in occupational hygiene and chemical analysis. The quality of laboratory instruments and procedures was periodically checked to ensure the reproducibility and recovery of the assays; using spiked recovery experiments, recovery was in the range of 90.0% (Pb) to 112.4% (Zn).

### 2.4. Statistical Analysis

All analyses were performed using SPSS version 22.0 statistical package (SPSS, IBM, Chicago, IL, USA) and EpiData 3.1 (EpiData ISOC, Funen, Denmark). The construct validity of the scale was evaluated by principal component analysis (PCA). The distributions of continuous variables were shown by the Kolmogorov-Smirnov test to be non-normal. Therefore, metal concentrations were described in terms of the median and interquartile range (IQR), geometric mean (GM), and 95% confidence interval (95% CI) of the geometric mean. Univariate statistical analysis was performed using the rank sum test. Univariate statistical analyses of the effect of cosmetics, sex, alcohol intake, and hair dye on serum concentrations of trace elements were performed using the Wilcoxon test; the Kruskal-Wallis test was used to assess the effects of age, smoking, and consumption of fried foods on serum concentrations of trace elements. A  $p$ -value ( $p \leq 0.05$ ) was considered statistically significant.

## 3. Results

The Kaiser-Meyer-Olkin measure (KMO) value was 0.911, and the partial correlation is very weak; the Bartlett spherical test, rejected the original hypothesis of the unit correlation matrix ( $p = 0.0008$ ), is suitable for factor analysis. There are seven factors which characteristic value in the principal component analysis was greater than 1, and the cumulative contribution to the total variance of the rate of 73.2%. According to the maximum factor load corresponding to the original variables, the original variables are divided into seven categories, which are in good agreement with the seven factors of the scale design. Subjects (765 men and 537 women) were grouped by age as follows: 6–12 ( $n = 231$ ), 12–16 ( $n = 214$ ), 16–20 ( $n = 168$ ), 20–30 ( $n = 187$ ), 30–45 ( $n = 255$ ), and 45–60 ( $n = 247$ ) years. The utilization rate of cosmetics, hair dye and alcohol were 5.5%, 8.6% and 14.7%, respectively. Other demographic data are presented in Figure 2. Chemical concentrations were above the limit of detection for all three trace elements. As shown in Table 1, the GM concentration of blood Zn (BZn) was 3.14 mg/L (95% CI: 3.08–3.20 mg/L). Women had significantly higher levels of Zn than men (GM, 3.28 mg/L vs. 3.04 mg/L,  $p < 0.001$ ). Alcohol use was associated with slightly higher BZn concentrations (GM for drinkers = 3.39 mg/L; GM for non-drinkers = 3.09 mg/L;  $p = 0.002$ ). The GM concentration of BZn was highest in the 20–30-year-old age group ( $p < 0.001$ ).



**Figure 2.** Characteristics of the study population. Univariate analyses of associations between cosmetics, alcohol intake and hair dye with serum concentrations of trace elements were performed using the Wilcoxon test; the Kruskal-Wallis test was used to assess associations between smoking, and consumption of fried foods with serum concentrations of trace elements. N: no; Y: yes; F: former smoker; A: 1–2 times per month; B: 1–2 times per week; C: Every day.

**Table 1.** Blood  $^{66}\text{Zn}$  levels (mg/L), stratified by relevant categorical variables.

Items	<i>n</i>	P5	P25	P50	P75	P95	GM (95% CI)
Total population	1302	1.41	2.40	3.10	3.83	5.02	3.14 (3.08–3.20)
Sex ( $p < 0.001$ )							
Male	765	1.32	2.29	3.00	3.74	4.86	3.04 (2.97–3.12)
Female	537	1.55	2.51	3.26	3.99	5.13	3.28 (3.18–3.37)
Use of cosmetics ( $p = 0.991$ )							
No	1230	1.43	2.40	3.10	3.82	5.01	3.14 (3.08–3.20)
Yes	72	1.23	2.35	3.08	3.97	5.35	3.14 (2.88–3.41)
Use of hair dye ( $p = 0.068$ )							
No	1190	1.43	2.41	3.13	3.84	5.03	3.16 (3.09–3.22)
Yes	112	1.36	2.25	2.84	3.71	4.92	2.95 (2.76–3.14)
Age in years ( $p < 0.001$ )							
6–12	231	1.03	1.94	2.72	3.55	4.32	2.75 (2.61–2.88)
12–16	214	1.65	2.49	3.17	3.70	4.91	3.13 (3.00–3.26)
16–20	168	1.86	2.59	3.27	4.33	5.34	3.41 (3.24–3.58)
20–30	187	1.78	2.89	3.63	4.27	5.56	3.61 (3.45–3.77)
30–45	255	1.39	2.22	2.83	3.69	5.08	3.01 (2.88–3.15)
>45	247	1.52	2.38	3.05	3.81	4.77	3.10 (2.97–3.22)
Alcohol intake ( $p = 0.002$ )							
No	1110	1.40	2.39	3.08	3.75	4.93	3.09 (3.03–3.16)
Yes	192	1.72	2.50	3.32	4.19	5.54	3.39 (3.22–3.56)
Smoking ( $p = 0.001$ )							
No	1111	1.36	2.39	3.07	3.77	4.96	3.10 (3.04–3.16)
Yes	170	1.72	2.56	3.45	4.18	5.29	3.42 (3.25–3.59)
Former smoker	21	1.44	2.28	2.73	3.52	5.35	2.91 (2.49–3.34)
Consumption of fried foods ( $p = 0.088$ )							
1–2 times per month	842	1.40	2.39	3.06	3.76	4.91	3.10 (3.03–3.17)
1–2 times per week	380	1.49	2.43	3.19	3.87	5.19	3.17 (3.06–3.28)
Every day	80	1.46	2.30	3.33	4.48	6.11	3.42 (3.12–3.72)



The GM concentration of blood Ge (BGe) for the total sample was 19.9 µg/L (95% CI: 19.3–20.6 µg/L), as shown in Table 2. The GM of BGe among the subjects who used hair dye was 17.6 µg/L, which was significantly lower than that of non-users (GM = 20.2 µg/L,  $p < 0.05$ ). The GM concentration of BGe in the 16–20-year-old age group was significantly higher than in the other age groups ( $p < 0.001$ ). Alcohol consumption was associated with slightly increased BGe levels (GM for drinkers = 19.6 µg/L; GM for non-drinkers = 21.7 µg/L;  $p = 0.002$ ). A statistically significant difference in blood Ge concentration was observed according to consumption of fried foods: 1–2 times per month (GM = 18.7 µg/L), 1–2 times per week (GM = 20.9 µg/L), or every day (GM = 28.5 µg/L),  $p < 0.001$ .

**Table 2.** Blood  $^{72}\text{Ge}$  levels (µg/L), stratified by relevant categorical variables.

Items	<i>n</i>	P5	P25	P50	P75	P95	GM (95% CI)
Total population	1302	5.10	12.1	18.3	24.5	45.2	19.9 (19.3–20.6)
Sex ( $p = 0.095$ )							
Male	765	4.40	12.5	18.9	25.2	40.9	19.8 (19.0–20.6)
Female	537	6.13	11.9	17.0	23.0	53.0	20.2 (19.1–21.3)
Use of cosmetics ( $p = 0.544$ )							
No	1230	5.11	12.2	18.3	24.6	43.3	19.9 (19.3–20.6)
Yes	72	4.65	10.4	17.2	24.1	55.3	20.5 (17.1–23.9)
Use of hair dye ( $p = 0.016$ )							
No	1190	5.11	12.3	18.5	24.6	46.7	20.2 (19.5–20.8)
Yes	112	4.99	10.9	15.3	22.6	38.7	17.6 (15.7–19.5)
Age in years ( $p < 0.001$ )							
6–12	231	1.04	9.96	17.9	23.1	27.0	16.2 (15.1–17.3)
12–16	214	6.96	14.3	19.9	25.9	42.8	21.5 (20.0–23.0)
16–20	168	9.61	15.0	22.8	45.2	62.2	29.3 (26.5–32.0)
20–30	187	5.21	10.8	16.0	23.1	32.5	17.3 (16.0–18.5)
30–45	255	4.89	12.6	19.4	26.8	42.5	20.9 (19.5–22.3)
>45	247	6.48	11.1	15.3	20.7	34.0	16.9 (15.8–17.9)
Alcohol intake ( $p = 0.001$ )							
No	1110	4.88	11.8	17.8	24.0	46.9	19.6 (18.9–20.3)
Yes	192	6.44	13.8	20.8	28.0	41.5	21.7 (20.2–23.2)
Smoking ( $p = 0.148$ )							
No	1111	4.91	11.9	17.9	24.5	47.1	19.9 (19.2–20.6)
Yes	170	5.70	13.8	18.9	24.6	36.4	19.9 (18.5–21.3)
Former smoker	21	10.7	16.1	19.7	25.6	42.0	21.8 (17.9–25.6)
Consumption of fried foods ( $p < 0.001$ )							
1–2 times per month	842	4.92	11.9	17.6	23.9	37.6	18.7 (18.0–19.4)
1–2 times per week	380	4.93	12.2	18.6	24.9	53.1	20.9 (19.5–22.3)
Every day	80	7.31	16.1	23.6	40.4	63.4	28.5 (24.7–32.3)

The geometric mean blood concentration of Pb (BPb) was 24.1 (95% CI: 23.2–25.1 µg/L) (Table 3). Men had higher BPb (GM = 26.6 µg/L, 95% CI: 25.3–27.9 µg/L) than women (GM = 20.6 µg/L, 95% CI: 19.34–21.8 µg/L,  $p < 0.001$ ). The GM concentration of BPb was significantly higher in subjects who used cosmetics compared to those who did not (24.4 µg/L vs. 20.0 µg/L,  $p < 0.05$ ), and was also higher in subjects who did than did not use hair dye compared to those who did (24.5 µg/L vs. 19.6 µg/L,  $p < 0.05$ ). Alcohol intake was associated with elevated BPb (GM for drinkers = 28.3 µg/L; GM for non-drinkers = 23.4 µg/L;  $p = 0.002$ ). The GM concentration of BPb was highest in the 12–16-year-old age group ( $p < 0.001$ ). There was a statistically significant difference in BPb between smokers (GM = 29.1 µg/L, 95% CI: 26.3–32.0 µg/L) and non-smokers (GM = 23.1 µg/L, 95% CI: 22.1–24.1 µg/L,  $p < 0.001$ ).

**Table 3.** Blood  $^{208}\text{Pb}$  levels ( $\mu\text{g/L}$ ), stratified by relevant categorical variables.

Items	<i>n</i>	P5	P25	P50	P75	P95	GM (95% CI)
Total population	1302	3.00	12.5	20.9	32.2	56.0	24.1 (23.2–25.1)
Sex ( $p < 0.001$ )							
Male	765	4.48	14.0	22.9	35.2	58.9	26.6 (25.3–27.9)
Female	537	2.05	10.3	17.9	28.4	47.6	20.6 (19.3–21.8)
Use of cosmetics ( $p = 0.019$ )							
No	1230	1.41	6.86	17.8	28.8	53.8	20.0 (16.0–24.0)
Yes	72	3.69	12.7	21.1	32.4	56.3	24.4 (23.4–25.3)
Use of hair dye ( $p = 0.004$ )							
No	1190	0.12	9.12	17.9	27.1	49.0	19.6 (17.0–22.3)
Yes	112	3.66	12.7	21.2	32.8	56.4	24.5 (23.5–25.5)
Age in years ( $p < 0.001$ )							
6–12	231	1.80	10.8	21.2	36.1	56.9	25.0 (22.7–27.3)
12–16	214	7.48	18.0	25.6	35.6	53.4	27.7 (25.7–29.6)
16–20	168	4.96	11.2	19.3	28.0	57.5	23.2 (20.4–26.0)
20–30	187	5.33	14.9	22.3	34.3	56.5	26.9 (24.1–29.8)
30–45	255	0.07	7.40	18.0	29.4	56.1	20.8 (18.6–22.9)
>45	247	4.99	12.8	18.6	27.4	55.0	22.13 (20.2–24.0)
Alcohol intake ( $p = 0.001$ )							
No	1110	2.67	12.0	20.4	31.5	51.6	23.4 (22.4–24.4)
Yes	192	5.89	15.6	23.4	37.7	60.7	28.3 (25.6–31.0)
Smoking ( $p < 0.001$ )							
No	1111	2.62	11.9	19.9	31.5	51.6	23.1 (22.1–24.1)
Yes	170	5.81	16.7	24.6	36.0	69.3	29.1 (26.3–32.0)
Former smoker	21	5.20	22.0	25.2	42.6	153.0	37.9 (23.3–52.5)
Consumption of fried foods ( $p = 0.319$ )							
1–2 times per month	842	2.94	12.5	20.6	32.4	56.8	24.1 (22.9–25.2)
1–2 times per week	380	3.56	12.1	21.2	31.6	53.0	23.8 (22.1–21.2)
Every day	80	4.36	14.4	23.1	34.8	57.4	26.0 (22.5–29.5)

#### 4. Discussion

Zn, Ge, and Pb are present in food, water, soil, and elsewhere in the natural environment. Appropriate levels of trace elements are required to maintain the body healthy. Therefore, knowing the reference ranges for these metals is useful for evaluation of occupational hazard exposure and to evaluate prevention or treatment strategies for diseases caused by deficiency or excess of these elements. Our results show that levels of trace elements in the body were associated with dietary and environment factors. Reference value of trace elements in human blood have been measured in many areas including Europe and North America since the 1990s [2,37]. Blood concentrations reflect short-term changes [73] and are considered a sensitive indicator of trace element deficiency or excess. In the present study, concentrations of Zn, Ge, and Pb were measured in blood, as a reflection of the total body content of these trace elements.

As early as the 1990s, European and American countries began biological monitoring of Zn [74,75]. The reference range for Zn obtained in an Italian sample (GM = 6.42 mg/L) was similar to the ranges obtained in populations in Spain and in the Czech Republic [76–80], but was significantly higher than that shown in our data (GM = 3.14 mg/L). Another study from China found results similar to ours [78]. Rice is the main staple of Asians, but in current high-yielding rice varieties the supply of zinc is poor, as polishing and shelling cause a huge loss of zinc [81,82]. Other reason is possible that discrepancies

in levels of trace elements between different countries are partly due to environmental factors, but specific reasons need to be explored further.

The BZn of drinkers was slightly higher than that of non-drinkers. This finding is consistent with results from a study in Italy [76] and likely reflects a causal relationship. Alcohol contains a large amount of Zn, derived from the soil via absorption by plants, which is then released when beverages are packaged in metal containers [83,84]. The GM concentration of BZn was significantly higher in the 20–30-year-old age group than in any other age group studied. This may be because sexual development in this period requires an increased intake of Zn [21,22].

Biological monitoring of Ge is necessary, as this element is used in many drugs. To date, there has been no estimate of Ge concentration from a national sample. Results reported from Chengde City (18.3–92.5 µg/L) [85] were similar to those from our study (GM = 19.9 µg/L), whereas Ge serum levels of 290 µg/L have been reported in the USA [86]. The results of the present study showed that the BGe among the subjects who used hair dye was significantly lower than among those who did not. This may be because hair dye contains a high concentration of Pb, which inhibits the absorption of Ge, but the specific mechanism needs further study. As adolescence is a critical period of growth and development, the demand for trace elements in this period is greatly increased. Accordingly, the BGe in the 16–20-year old age group was significantly higher than in other age groups. Those who consumed alcohol had significantly higher blood Ge concentrations than non-drinkers, and that the greater the frequency of fried food consumption, the higher the level of blood Ge. The reasons underlying these findings require further study.

Because Pb is widespread and harmful biomonitoring studies have been conducted in many countries [87–90]. Data from the present study revealed GM concentration of BPb to be 24.1 µg/L, which is lower than that observed in Brazilian, Czech, Danish, Italian, and Spanish studies, but higher than in that in American, Korean, Canadian, and Australian studies [79]. There are reports that BPb is higher in men than in women, as was found in our study [77,78]. We found that BPb concentrations were significantly higher in subjects who used cosmetics than in those who did not, similar to the findings for hair dye users. This is consistent with the results of previous reports; the vast majority of cosmetics contain Pb, which is absorbed through the skin into the body [91–96]. In the present study, alcohol consumption was associated with higher BPb. Pb in the soil is absorbed into plants [97], and ingestion of alcohol facilitates the absorption of Pb by the body [98]. There was a statistically significant difference in BPb concentrations between smokers and non-smokers. Pb in cigarettes enters the body through the respiratory tract [99], and may act synergistically with risk factors associated with hypertension [100]. Existing literature supports an identical trend with drinkers and smokers [101].

## 5. Conclusions

This study provides data on blood concentrations of Zn, Pb, and Ge in a sample of the population living in Shandong Province, China. It provides valid and reliable reference data for establishing reference values for blood levels of these trace elements for the Chinese population. Alcohol consumption was associated with blood concentrations of Zn, Ge, and Pb, while cigarette smoking had no significant influence on BZn and BPb. Use of cosmetics and hair dye was associated with higher blood Pb concentrations. In addition, there was a positive association between the frequency of fried food consumption and blood Pb concentration. Further research is needed to determine the factors underlying the associations we observed between these variables and blood levels of trace elements.

**Acknowledgments:** This study was supported by the Institute for Occupational Health and Poison Control under the China Center for Disease Control and Prevention (CDC), Ministry of Health of the People's Republic of China Grant no.2006BAI06B02 and U.S. National Institute of Health/National Institute of Environmental Health Science ES017055. The authors gratefully acknowledge the technical assistance of He Ze Center for Disease Control and Prevention, Ji Nan Center for Disease Control and Prevention and Ri Zhao Center for Disease Control and Prevention.



**Author Contributions:** Hua Shao and Zhi-Hu Zhang conceived and designed the experiments; Zhi-Hu Zhang, Long Li and Guang Xu performed the experiments; Long Li, Xing-Fu Pan and Jin-Ye Li analyzed the data; Xing-Fu Pan contributed reagents/materials/analysis tools; Long Li wrote the paper.

**Conflicts of Interest:** This manuscript has not been published or presented elsewhere in part or in entirety and is not under consideration by another journal. There are no conflicts of interest to declare.

## References

1. Albertini, R.; Bird, M.; Doerr, N.; Needham, L.; Robison, S.; Sheldon, L.; Zenick, H. The use of biomonitoring data in exposure and human health risk assessments. *Environ. Health Perspect.* **2006**, *114*, 1755–1762. [[CrossRef](#)] [[PubMed](#)]
2. Minoia, C.; Pietra, R.; Sabbioni, E.; Ronchi, A.; Gatti, A.; Cavalleri, A.; Manzo, L. Trace element reference values in tissues from inhabitants of the European Community. III. The control of preanalytical factors in the biomonitoring of trace elements in biological fluids. *Sci. Total Environ.* **1992**, *120*, 63–79. [[CrossRef](#)]
3. Korvela, M.; Lind, A.L.; Wetterhall, M.; Gordh, T.; Andersson, M.; Pettersson, J. Quantification of 10 elements in human cerebrospinal fluid from chronic pain patients with and without spinal cord stimulation. *J. Trace Elements Med. Biol.* **2016**, *37*, 1–7. [[CrossRef](#)] [[PubMed](#)]
4. Bloise, A.; Barca, D.; Gualtieri, A.F.; Pollastri, S.; Belluso, E. Trace elements in hazardous mineral fibres. *Environ. Pollut.* **2016**, *216*, 314–323. [[CrossRef](#)] [[PubMed](#)]
5. Czarnek, K.; Terpiłowska, S.; Siwicki, A.K. Selected aspects of the action of cobalt ions in the human body. *Cent. Eur. J. Immunol.* **2015**, *40*, 236–242. [[CrossRef](#)] [[PubMed](#)]
6. Anderson, R.A. Essentiality of chromium for human nutrition and health. *Sci. Total Environ.* **1989**, *86*, 75–81. [[CrossRef](#)]
7. Van Bakel, M.M.; Printzen, G.; Wermuth, B.; Wiesmann, U.N. Antioxidant and thyroid hormone status in selenium deficient phenylketonuric and hyperphenylalaninemic patients. *Am. J. Clin. Nutr.* **2000**, *72*, 976–981. [[PubMed](#)]
8. Araya, M.; Pizarro, F.; Olivares, M.; Arredondo, M.; González, M.; Méndez, M. Understanding copper homeostasis in humans and copper effects on health. *Biol. Res.* **2006**, *39*, 183–187. [[CrossRef](#)] [[PubMed](#)]
9. Bonham, M.; O'Connor, J.M.; Hannigan, B.M.; Strain, J.J. The immune system as a physiological indicator of marginal copper status? *Br. J. Nutr.* **2002**, *87*, 393–403. [[CrossRef](#)] [[PubMed](#)]
10. Davis, C.D. Low dietary copper increases fecal free radical production, fecal water alkaline phosphatase activity and cytotoxicity in healthy men. *J. Nutr.* **2003**, *133*, 522–527. [[PubMed](#)]
11. Sujiwattarat, P.; Pongsanarakul, P.; Tamsiripong, Y.; Tamsiripong, T.; Thawornkuno, C.; Uno, Y.; Unajak, S.; Matsuda, Y.; Choowongkamon, K.; Srikulnath, K. Molecular cloning and characterization of Siamese crocodile (*Crocodylus siamensis*) copper, zinc superoxide dismutase (CSI-Cu, Zn-SOD) gene. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* **2016**, *191*, 187–195. [[CrossRef](#)] [[PubMed](#)]
12. Maret, W.; Sandstead, H.H. Zinc requirements and the risks and benefits of zinc supplementation. *J. Trace Elements Med. Biol.* **2006**, *20*, 3–18. [[CrossRef](#)] [[PubMed](#)]
13. Gmshinskii, I.V.; Munkhuu, B.; Mazo, V.K. Trace elements in human nutrition: Biological indices of zinc insufficiency. *Vopr. Pitan.* **2006**, *75*, 4–11. [[PubMed](#)]
14. Zheng, J.; Zhang, Y.; Xu, W.; Luo, Y.; Hao, J.; Shen, X.L.; Yang, X.; Li, X.; Huang, K. Zinc protects HepG2 cells against the oxidative damage and DNA damage induced by ochratoxin A. *Toxicol. Appl. Pharmacol.* **2013**, *268*, 123–131. [[CrossRef](#)] [[PubMed](#)]
15. Sharif, R.; Thomas, P.; Zalewski, P. The role of zinc in genomic stability. *Mutat. Res.* **2012**, *733*, 111–121. [[CrossRef](#)] [[PubMed](#)]
16. Gazaryan, I.G.; Krasnikov, B.F.; Ashby, G.A.; Thorneley, R.N.; Kristal, B.S.; Brown, A.M. Zinc is a potent inhibitor of thiol oxidoreductase activity and stimulates reactive oxygen species production by lipoamide dehydrogenase. *J. Biol. Chem.* **2002**, *277*, 10064–10072. [[CrossRef](#)] [[PubMed](#)]
17. Brown, A.M.; Kristal, B.S.; Efron, M.S.; Shestopalov, A.I.; Ullucci, P.A.; Sheu, K.F.; Blass, J.P.; Cooper, A.J. Zn<sup>2+</sup> inhibits alpha-ketoglutarate-stimulated mitochondrial respiration and the isolated alpha-ketoglutarate dehydrogenase complex. *J. Biol. Chem.* **2000**, *275*, 13441–13447. [[CrossRef](#)] [[PubMed](#)]
18. Qiao, Y.; Zhang, W.; Tian, P.; Meng, F.; Zhu, H.; Jiang, X.; Liu, X.; Chu, P.K. Stimulation of bone growth following zinc incorporation into biomaterials. *Biomaterials* **2014**, *35*, 6882–6897. [[CrossRef](#)] [[PubMed](#)]

19. Sharir, H.; Zinger, A.; Nevo, A.; Sekler, I.; Hershfinkel, M. Zinc released from injured cells is acting via the Zn<sup>2+</sup>-sensing receptor, ZnR, to trigger signaling leading to epithelial repair. *J. Biol. Chem.* **2010**, *285*, 26097–26106. [[CrossRef](#)] [[PubMed](#)]
20. Cao, J.; Gao, Z.; Yan, J.; Li, M.; Su, J.; Xu, J.; Yan, C.H. Evaluation of trace elements and their relationship with growth and development of young children. *Biol. Trace Element Res.* **2016**, *171*, 270–274. [[CrossRef](#)] [[PubMed](#)]
21. Chang, C.S.; Choi, J.B.; Kim, H.J.; Park, S.B. Correlation between serum testosterone level and concentrations of copper and zinc in hair tissue. *Biol. Trace Element Res.* **2011**, *144*, 264–271. [[CrossRef](#)] [[PubMed](#)]
22. Björndahl, L.; Kvist, U. A model for the importance of zinc in the dynamics of human sperm chromatin stabilization after ejaculation in relation to sperm DNA vulnerability. *Syst. Biol. Reprod. Med.* **2011**, *57*, 86–92. [[CrossRef](#)] [[PubMed](#)]
23. Ranasinghe, P.; Pigera, S.; Galappaththy, P.; Katulanda, P.; Constantine, G.R. Zinc and diabetes mellitus: Understanding molecular mechanisms and clinical implications. *Daru* **2015**, *23*, 44–56. [[CrossRef](#)] [[PubMed](#)]
24. Slepchenko, K.G.; Daniels, N.A.; Guo, A.; Li, Y.V. Autocrine effect of Zn<sup>2+</sup> on the glucose-stimulated insulin secretion. *Endocrine* **2015**, *50*, 110–122. [[CrossRef](#)] [[PubMed](#)]
25. Mochizuki, K.; Murase, H.; Imose, M.; Kawakami, H.; Sawada, A. Improvement of scotopic electroretinograms and night blindness with recovery of serum zinc levels. *Jpn. J. Ophthalmol.* **2006**, *50*, 532–536. [[CrossRef](#)] [[PubMed](#)]
26. Kraft, S.P.; Parker, J.A.; Matuk, Y.; Rao, A.V. The rat electroretinogram in combined zinc and vitamin A deficiency. *Investig. Ophthalmol. Vis. Sci.* **1987**, *28*, 975–984.
27. Komai, M.; Goto, T.; Suzuki, H.; Takeda, T.; Furukawa, Y. Zinc deficiency and taste dysfunction, contribution of carbonic anhydrase, a zinc-metalloenzyme, to normal taste sensation. *Biofactors* **2000**, *12*, 65–70. [[CrossRef](#)] [[PubMed](#)]
28. Fukasawa, T.; Orii, T.; Tanaka, M.; Suzuki, N.; Kanzaki, Y. Relation between drug-induced taste disorder and chelating behavior with zinc ion; statistical approach to the drug-induced taste disorder, part II. *Chem. Pharm. Bull.* **2008**, *56*, 1177–1180. [[CrossRef](#)] [[PubMed](#)]
29. Desai, V.; Gaurav, I.; Kumar, M.V.; Gaurav, I.; Sharma, R. Molecular analysis of trace elements in oral submucous fibrosis and future perspective. *Univers. Res. J. Dent.* **2014**, *4*, 26–35. [[CrossRef](#)]
30. Ray, J.G.; Ghosh, R.; Mallick, D.; Swain, N.; Gandhi, P.; Ram, S.S.; Selvaraj, S.; Rathore, A.; Mathummal, S.; Chakraborty, A. Correlation of trace elemental profiles in blood samples of Indian patients with leukoplakia and oral submucous fibrosis. *Biol. Trace Element Res.* **2011**, *144*, 295–305. [[CrossRef](#)] [[PubMed](#)]
31. Hennigar, S.R.; Kelley, A.M.; McClung, J.P. Metallothionein and zinc transporter expression in circulating human blood cells as biomarkers of Zinc Status: A systematic review. *Adv. Nutr.* **2016**, *7*, 735–746. [[CrossRef](#)] [[PubMed](#)]
32. Tomiyama, K.; Arakawa, Y. Zinc and tin-induced apoptotic mechanisms in immune system and cranial nerve system. *Nihon Rinsho* **2016**, *74*, 1111–1119. [[PubMed](#)]
33. Maywald, M.; Rink, L. Zinc supplementation induces CD4<sup>+</sup>CD25<sup>+</sup>Foxp3<sup>+</sup> antigen-specific regulatory T cells and suppresses IFN- $\gamma$  production by upregulation of Foxp3 and KLF-10 and downregulation of IRF-1. *Eur. J. Nutr.* **2016**. [[CrossRef](#)] [[PubMed](#)]
34. Beck, F.W.; Prasad, A.S.; Kaplan, J.; Fitzgerald, J.T.; Brewer, G.J. Changes in cytokine production and T cell subpopulations in experimentally induced zinc-deficient humans. *Am. J. Physiol.* **1997**, *272*, 1002–1007.
35. Della Rovere, F.; Granata, A.; Pavia, R.; Tomaino, A.; Zirilli, A.; Monaco, F.; Familiari, D.; La Rocca, A.; Ientile, R.; Mondello, B.; et al. Vitamins A, E, microelements and membrane lipid peroxidation in patients with neoplastic disease treated with calcium antagonists and antagonists of receptors H2. *Anticancer Res.* **2004**, *24*, 1449–1453. [[PubMed](#)]
36. Chen, J.F.; Shi, Q.J.; Zheng, S.Y. Experimental research of protective and therapeutic effects of zinc and vitamin E on mouse liver radiational damage. *Hunan Yi Ke Da Xue Xue Bao* **2001**, *26*, 207–210. [[PubMed](#)]
37. Haberal, M.; Mavi, V.; Oner, G. The stabilizing effect of vitamin E, selenium and zinc on leucocyte membrane permeability: A study in vitro. *Burns Incl. Therm. Inj.* **1987**, *13*, 118–122. [[CrossRef](#)]
38. Cai, C.; Lin, P.; Zhu, H.; Ko, J.K.; Hwang, M.; Tan, T.; Pan, Z.; Korichneva, I.; Ma, J. Zinc binding to MG53 protein facilitates repair of injury to cell membranes. *J. Biol. Chem.* **2015**, *290*, 13830–13839. [[CrossRef](#)] [[PubMed](#)]
39. Yagi, T.; Asakawa, A.; Ueda, H.; Ikeda, S.; Miyawaki, S.; Inui, A. The role of zinc in the treatment of taste disorders. *Recent Pat. Food Nutr. Agric.* **2013**, *5*, 44–51. [[CrossRef](#)] [[PubMed](#)]

40. Shweta, G.; Prantesh, J.; Shashvat, S. Isolated zinc deficiency causing severe microcytosis and sideroblastic anemia. *Turk. J. Haematol.* **2014**, *31*, 339–340. [[CrossRef](#)] [[PubMed](#)]
41. Motooka, R.; Yamamoto, S. Copper deficiency myelopathy probably caused by long-lasting daily excessive intake of zinc. *Rinsho Shinkeigaku* **2016**, *56*, 690–693. [[CrossRef](#)] [[PubMed](#)]
42. Yang, F.; Jin, H.; Pi, J.; Jiang, J.H.; Liu, L.; Bai, H.H.; Yang, P.H.; Cai, J.Y. Anti-tumor activity evaluation of novel chrysin-organogermanium (IV) complex in MCF-7 cells. *Bioorg. Med. Chem. Lett.* **2013**, *23*, 5544–5551. [[CrossRef](#)] [[PubMed](#)]
43. Jao, S.W.; Lee, W.; Ho, Y.S. Effect of germanium on 1,2-dimethylhydrazine-induced intestinal cancer in rats. *Dis. Colon Rectum* **1990**, *33*, 99–104. [[CrossRef](#)] [[PubMed](#)]
44. Yang, M.K.; Kim, Y.G. Protective role of germanium-132 against paraquat-induced oxidative stress in the livers of senescence-accelerated mice. *J. Toxicol. Environ. Health A* **1999**, *58*, 289–297. [[PubMed](#)]
45. Wu, Z.; Chen, X.; Yang, K.; Xia, T. Studies on the hydroxyl free radical-scavenging effect of combined selenium and germanium. *Wei Sheng Yan Jiu* **2001**, *30*, 208–210.
46. Mrema, J.E.; Slavik, M.; Davis, J. Spirogermanium: A new drug with antimalarial activity against chloroquine-resistant *Plasmodium falciparum*. *Int. J. Clin. Pharmacol. Ther. Toxicol.* **1983**, *21*, 167–171. [[PubMed](#)]
47. Goodman, S. Therapeutic effects of organic germanium. *Med. Hypotheses* **1988**, *26*, 207–215. [[CrossRef](#)]
48. DiMartino, M.J.; Lee, J.C.; Badger, A.M.; Muirhead, K.A.; Mirabelli, C.K.; Hanna, N. Antiarthritic and immunoregulatory activity of spirogermanium. *J. Pharmacol. Exp. Ther.* **1986**, *236*, 103–110. [[PubMed](#)]
49. Lee, J.H.; Kim, K.W.; Yoon, M.Y.; Lee, J.Y.; Kim, C.J.; Sim, S.S. Anti-inflammatory effect of germanium-concentrated yeast against paw oedema is related to the inhibition of arachidonic acid release and prostaglandin E<sub>2</sub> production in RBL 2H3 cells. *Auton. Autacoid Pharmacol.* **2005**, *25*, 129–134. [[CrossRef](#)] [[PubMed](#)]
50. Via, S. The role of trace elements in hematopoiesis. *Terapevticheskii Arkhiv* **1963**, *35*, 3–14.
51. Badger, A.M.; Mirabelli, C.K.; DiMartino, M. Generation of suppressor cells in normal rats by treatment with spirogermanium, a novel heterocyclic anticancer drug. *Immunopharmacology* **1985**, *10*, 201–207. [[CrossRef](#)]
52. Nakamura, T.; Takeda, T.; Tokuji, Y. The oral intake of organic germanium, Ge-132, elevates  $\alpha$ -Tocopherol levels in the Plas-ma and modulates hepatic gene expression profiles to promote immune activation in mice. *Int. J. Vitam. Nutr. Res.* **2014**, *84*, 183–195. [[CrossRef](#)] [[PubMed](#)]
53. Hirayama, C.; Suzuki, H.; Ito, M.; Okumura, M.; Oda, T. Propagermanium: A nonspecific immune modulator for chronic hepatitis B. *J. Gastroenterol.* **2003**, *38*, 525–532. [[PubMed](#)]
54. Li, Y.; Ren-Lu, H. Germanium and human healthy. *Stud. Trace Elements Health* **2005**, *22*, 60–61.
55. Okada, K.; Okagawa, K.; Kawakami, K.; Kuroda, Y.; Morizumi, K.; Sato, H.; Morita, H.; Shimomura, S.; Saito, S. Renal failure caused by long-term use of a germanium preparation as an elixir. *Clin. Nephrol.* **1989**, *31*, 219–224. [[PubMed](#)]
56. Krapf, R.; Schaffner, T.; Iten, P.X. Abuse of germanium associated with fatal lactic acidosis. *Nephron* **1992**, *62*, 351–356. [[CrossRef](#)] [[PubMed](#)]
57. Kamijo, M.; Yagihashi, S.; Kida, K.; Narita, S.; Nakata, F. An autopsy case of chronic germanium intoxication presenting peripheral neuropathy, spinal ataxia, and chronic renal failure. *Rinsho Shinkeigaku* **1991**, *31*, 191–196. [[PubMed](#)]
58. Schauss, A.G. Nephrotoxicity and neurotoxicity in humans from organogermanium compounds and germanium dioxide. *Biol. Trace Element Res.* **1991**, *29*, 267–280. [[CrossRef](#)]
59. Kim, K.M.; Lim, C.S.; Kim, S.; Kim, S.H.; Park, J.H.; Ahn, C.; Han, J.S.; Lee, J.S. Nephropathy and neuropathy induced by a germanium-containing compound. *Nephrol. Dial. Transplant.* **1998**, *13*, 3218–3219. [[CrossRef](#)] [[PubMed](#)]
60. Dixon, C.; Hagemester, F.; Legha, S.; Bodey, G. Pulmonary toxicity associated with spirogermanium. *Cancer Treat Rep.* **1984**, *68*, 907–908. [[PubMed](#)]
61. Stewart, W.F.; Schwartz, B.S. Effects of lead on the adult brain: A 15-year exploration. *Am. J. Ind. Med.* **2007**, *50*, 729–739. [[CrossRef](#)] [[PubMed](#)]
62. Lanphear, B.P.; Hornung, R.; Khoury, J.; Yolton, K.; Baghurst, P.; Bellinger, D.C.; Canfield, R.L.; Dietrich, K.N.; Bornschein, R.; Greene, T.; et al. Low-level environmental lead exposure and children’s intellectual function: An international pooled analysis. *Environ. Health Perspect.* **2005**, *113*, 894–899. [[CrossRef](#)] [[PubMed](#)]

63. Kupraszewicz, E.; Brzóška, M.M. Excessive ethanol consumption under exposure to lead intensifies disorders in bone metabolism: A study in a rat model. *Chem. Biol. Interact.* **2013**, *203*, 486–501. [[CrossRef](#)] [[PubMed](#)]
64. Hernández-Ochoa, I.; García-Vargas, G.; López-Carrillo, L.; Rubio-Andrade, M.; Morán-Martínez, J.; Cebrián, M.E.; Quintanilla-Vega, B. Low lead environmental exposure alters semen quality and sperm chromatin condensation in northern Mexico. *Reprod. Toxicol.* **2005**, *20*, 221–228. [[CrossRef](#)] [[PubMed](#)]
65. Batra, N.; Nehru, B.; Bansal, M.P. Reproductive potential of male Portan rats exposed to various levels of lead with regard to zinc status. *Br. J. Nutr.* **2004**, *91*, 387–391. [[CrossRef](#)] [[PubMed](#)]
66. Molina, R.M.; Phattanarudee, S.; Kim, J.; Thompson, K.; Wessling-Resnick, M.; Maher, T.J.; Brain, J.D. Ingestion of Mn and Pb by rats during and after pregnancy alters iron metabolism and behavior in off spring. *Neurotoxicology* **2011**, *32*, 413–422. [[CrossRef](#)] [[PubMed](#)]
67. Ermentrout, R.M.; Layon, M.E.; Ackley, C.J.; Venkatesan, P.; Lowrey, C.H. The effects of lead and cadmium on GATA-1 regulated erythroid gene expression. *Blood Cells Mol. Dis.* **2006**, *37*, 164–172. [[CrossRef](#)] [[PubMed](#)]
68. He, X.Y.; Yuan, L.Y.; Li, Y.T.; Li, M.; Chen, Y.; Yuan, H.; Wu, J.; Guo, C.Z.; Li, J. Cytotoxic responses and apoptosis in rat kidney epithelial cells exposed to lead. *Biomed. Environ. Sci.* **2016**, *29*, 529–533. [[PubMed](#)]
69. Wilk, A.; Kalisińska, E.; Kosik-Bogacka, D.; Romanowski, M.; Róžański, J.; Ciechanowski, K.; Słojewski, M.; Łanocha-Arendarczyk, N. Cadmium, lead and mercury concentrations in pathologically altered human kidneys. *Environ. Geochem. Health* **2016**. [[CrossRef](#)] [[PubMed](#)]
70. Cornelis, R.; Sabbioni, E.; Vander Venne, M.T. Trace element reference values in tissues from inhabitants of the European Community. *Sci. Total Environ.* **1994**, *158*, 191–226. [[CrossRef](#)]
71. Wheal, M.S.; DeCourcy-Ireland, E.; Bogard, J.R.; Thilsted, S.H.; Stangoulis, J.C. Measurement of haem and total iron in fish, shrimp and prawn using ICP-MS: Implications for dietary iron intake calculations. *Food Chem.* **2016**, *201*, 222–229. [[CrossRef](#)] [[PubMed](#)]
72. Feillet-Coudray, C.; Meunier, N.; Rambeau, M.; Brandolini-Bunlon, M.; Tressol, J.C.; Andriollo, M.; Mazur, A.; Cashman, K.D.; Coudray, C. Long-term moderate zinc supplementation increases exchangeable zinc pool masses in late-middle-aged men: The zenith study. *Am. J. Clin. Nutr.* **2005**, *82*, 103–110. [[PubMed](#)]
73. Wessells, K.R.; Jorgensen, J.M.; Hess, S.Y.; Woodhouse, L.R.; Peerson, J.M.; Brown, K.H. Plasma zinc concentration responds rapidly to the initiation and discontinuation of short-term zinc supplementation in healthy men. *J. Nutr.* **2010**, *140*, 2128–2133. [[CrossRef](#)] [[PubMed](#)]
74. Khandekar, R.N.; Raghunath, R.; Mishra, U.C. Levels of lead, cadmium, zinc and copper in the blood of an urban population. *Sci. Total Environ.* **1987**, *66*, 185–191. [[CrossRef](#)]
75. Subramanian, K.S.; Meranger, J.C. Blood levels of cadmium, copper, lead and zinc in children in a British Columbia community. *Sci. Total Environ.* **1983**, *30*, 231–244. [[CrossRef](#)]
76. Ding, C.; Zhu, C.; Liu, D.; Dong, M.; Zhang, A.H.; Pan, Y.J.; Yan, H.F. Inductively coupled plasma mass spectrometry for the simultaneous determination of thirty metals and metalloids elements in blood samples. *Chin. J. Prev. Med.* **2012**, *46*, 745–749.
77. Bocca, B.; Madeddu, R.; Asara, Y.; Tolu, P.; Marchal, J.A.; Forte, G. Assessment of reference ranges for blood Cu, Mn, Se and Zn in a selected Italian population. *J. Trace Element Med. Biol.* **2011**, *25*, 19–26. [[CrossRef](#)] [[PubMed](#)]
78. Zhang, L.L.; Lu, L.; Pan, Y.J.; Ding, C.G.; Xu, D.Y.; Huang, C.F.; Pan, X.F.; Zheng, W. Baseline blood levels of manganese, lead, cadmium, copper, and zinc in residents of Beijing suburb. *Environ. Res.* **2015**, *140*, 10–17. [[CrossRef](#)] [[PubMed](#)]
79. Bazzi, A.; Nriagu, J.O.; Linder, A.M. Determination of toxic and essential elements in children's blood with inductively coupled plasma-mass spectrometry. *J. Environ. Monit.* **2008**, *10*, 1226–1232. [[CrossRef](#)] [[PubMed](#)]
80. Moreno, M.A.; Marin, C.; Vinagre, F.; Ostapczuk, P. Trace element levels in whole blood samples from residents of the city Badajoz, Spain. *Sci. Total Environ.* **1999**, *229*, 209–215. [[CrossRef](#)]
81. Joy, E.J.; Louise Ander, E.; Broadley, M.R.; Young, S.D.; Chilimba, A.D.; Hamilton, E.M.; Watts, M.J. Elemental composition of Malawian rice. *Environ. Geochem. Health* **2016**. [[CrossRef](#)] [[PubMed](#)]
82. Swamy, B.P.; Rahman, M.A.; Inabangan-Asilo, M.A.; Amparado, A.; Manito, C.; Chadha-Mohanty, P.; Reinke, R.; Slamet-Loedin, I.H. Advances in breeding for high grain Zinc in Rice. *Rice* **2016**, *9*, 49–64. [[CrossRef](#)] [[PubMed](#)]
83. Gao, Y.; Huang, A.; Li, H. Investigation on the content of zinc, copper and chromium in the beverage industry. *Stud. Trace Elements Health* **2004**, *21*, 41–46.



84. Fan, K.; Wang, X.; Tian, W. Six kinds of wine depends on the ICP-AES method in the determination of heavy metals. *Chin. Public Health* **2000**, *16*, 639.
85. Wei, J. *Investigation on 19 Trace Elements in Whole Blood among General Population in Chengde*; Hebei Medical University: Shijiazhuang, China, 2006.
86. Schroeder, H.A.; Balassa, J.J. Abnormal trace metals in man: Germanium. *J. Chronic Dis.* **1967**, *20*, 211–224. [[CrossRef](#)]
87. Schultze, B.; Lind, P.M.; Larsson, A.; Lind, L. Whole blood and serum concentrations of metals in a Swedish population-based sample. *Scand. J. Clin. Lab Investig.* **2014**, *74*, 143–148. [[CrossRef](#)] [[PubMed](#)]
88. Ikeda, M.; Zhang, Z.W.; Shimbo, S.; Watanabe, T.; Nakatsuka, H.; Moon, C.S.; Matsuda-Inoguchi, N.; Higashikawa, K. Urban population exposure to lead and cadmium in east and south-east Asia. *Sci. Total Environ.* **2000**, *249*, 373–384. [[CrossRef](#)]
89. Faro, A.R.; Pinto Wde, J.; Ferreira, A.P.; Barbosa Jr, F.; Souza, V.C.; Fujimoto, D.E.; Koifman, R.J.; Koifman, S. Serum cadmium levels in a sample of blood donors in the Western Amazon, Brazil, 2010–2011. *Cad. Saude Publica* **2014**, *30*, 403–414. [[CrossRef](#)] [[PubMed](#)]
90. Farzin, L.; Amiri, M.; Shams, H.; Farzin, L.; Amiri, M.; Shams, H.; Faghih, M.A.A.; Moassesi, M.E. Blood levels of lead, cadmium, and mercury in residents of Tehran. *Biol. Trace Element Res.* **2008**, *123*, 14–26. [[CrossRef](#)] [[PubMed](#)]
91. Nduka, J.K.; Orisakwe, O.E.; Ukaebgu, L.D.; Sokaibe, C.; Udowelle, N.A. Human health risk assessment of heavy metals in cosmetics in Nigeria. *J. Cosmet. Sci.* **2015**, *66*, 233–246. [[PubMed](#)]
92. Zakaria, A.; Ho, Y.B. Heavy metals contamination in lipsticks and their associated health risks to lipstick consumers. *Regul. Toxicol. Pharmacol.* **2015**, *73*, 191–195. [[CrossRef](#)] [[PubMed](#)]
93. Corbett, J.F. Cadmium, lead and nickel in hair care products in Turkey. *J. Cosmet. Sci.* **2015**, *66*, 65. [[PubMed](#)]
94. Borowska, S.; Brzóska, M.M. Metals in cosmetics: Implications for human health. *J. Appl. Toxicol.* **2015**, *35*, 551–572. [[CrossRef](#)]
95. Ozbek, N.; Akman, S. Determination of lead, cadmium and nickel in hennas and other hair dyes sold in Turkey. *Regul. Toxicol. Pharmacol.* **2016**, *79*, 49–53. [[CrossRef](#)] [[PubMed](#)]
96. Kaličanin, B.; Velimirović, D. A Study of the Possible Harmful Effects of Cosmetic Beauty Products on Human Health. *Biol. Trace Element Res.* **2016**, *170*, 476–484. [[CrossRef](#)] [[PubMed](#)]
97. Kristensen, L.J.; Taylor, M.P.; Evans, A.J. Reply to Gulson’s comments on “Tracing changes in atmospheric sources of lead contamination using lead isotopic compositions in Australian redwine”. *Chemosphere* **2016**, *165*, 579–584. [[CrossRef](#)] [[PubMed](#)]
98. Kumar, V.; Tripathi, V.K.; Jahan, S.; Agrawal, M.; Pandey, A.; Khanna, V.K.; Pant, A.B. Lead intoxication synergies of the ethanol-induced toxic responses in neuronal cells—PC12. *Mol. Neurobiol.* **2015**, *52*, 1504–1520. [[CrossRef](#)] [[PubMed](#)]
99. Taroni, M.; Zagà, V.; Bartolomei, P.; Gattavecchia, E.; Pacifici, R.; Zuccaro, P.; Esposito, M. 210Pb and 210Po concentrations in Italian cigarettes and effective dose evaluation. *Health Phys.* **2014**, *107*, 195–199. [[CrossRef](#)] [[PubMed](#)]
100. Afridi, H.I.; Talpur, F.N.; Kazi, T.G.; Brabazon, D. Estimation of aluminum, arsenic, lead and nickel status in the samples of different cigarettes and their effect on human health of Irish smoker hypertensive consumers. *Clin. Lab.* **2015**, *61*, 1147–1156. [[CrossRef](#)] [[PubMed](#)]
101. Forte, G.; Madeddu, R.; Tolu, P.; Asara, Y.; Marchal, J.A.; Bocca, B. Reference intervals for blood Cd and Pb in the general population of Sardinia (Italy). *Int. J. Hyg. Environ. Health* **2011**, *214*, 102–109. [[CrossRef](#)] [[PubMed](#)]

