

Nucleophilicities and Nucleofugalities of Thio- and Selenoethers

Biplab Maji,^[a, b] Xin-Hua Duan,^[a, c] Patrick M. Jüstel,^[a] Peter A. Byrne,^[a, d] Armin R. Ofial,^[a] and Herbert Mayr*^[a]


Dedicated to Professor Peter Klüfers on the occasion of his 70th birthday

Abstract: Rate constants for the reactions of dialkyl chalcogenides with laser flash photolytically generated benzhydrylium ions have been measured photometrically to integrate them into the comprehensive benzhydrylium-based nucleophilicity scale. Combining these rate constants with the previously reported equilibrium constants for the same reactions provided the corresponding Marcus intrinsic barriers and

made it possible to quantify the leaving group abilities (nucleofugalities) of dialkyl sulfides and dimethyl selenide. Due to the low intrinsic barriers, dialkyl chalcogenides are fairly strong nucleophiles (comparable to pyridine and *N*-methylimidazole) as well as good nucleofuges; this makes them useful group-transfer reagents.

Introduction

Dialkyl chalcogenides are known to act as nucleophiles in a variety of reactions.^[1] The amino acid L-methionine, for example, reacts as a sulfur-centred nucleophile at the 5'-position of adenosyl triphosphate (ATP) to form S-adenosyl methionine (Scheme 1), which functions as a methylating agent in living organisms.^[2] In several hydrolases, the chalcogen-containing amino acids serine, cysteine, and selenocysteine are essential, and their catalytic activities rely on the nucleophilic properties of the chalcogen atom.^[3] Dialkyl chalcogenides have successfully been employed as nucleophilic organocatalysts in various chalcogenide-ylide mediated reactions, for example, epoxida-



Scheme 1. Participation of S-centred nucleophiles in biological systems.

tions, aziridinations, cyclopropanations, and olefinations.^[4] Chalcogenides were also used in combination with Lewis acids as organocatalysts in Morita-Baylis-Hillman reactions.^[5]

Arnett has shown that the relative Brønsted basicities of chalcogenides are strongly dependent on the nature of the solvent.^[6] While in aqueous solution, dimethyl sulfide ($pK_{\text{aH}} = -6.95$) is a much weaker base than dimethyl ether ($pK_{\text{aH}} = -2.52$), the heat of protonation is the same for both compounds in HSO_3F . In the gas phase, the proton affinity of dimethyl sulfide ($197 \text{ kcal mol}^{-1}$) is significantly higher than that of dimethyl ether ($186 \text{ kcal mol}^{-1}$).^[6] Similar pK_{aH} values for cyclic sulfides have been measured by Scorrano.^[7] Because of the well-known limitations of the correlations between the reactivities of different types of nucleophiles and the corresponding Brønsted basicities,^[8] we have previously used benzhydrylium ions as reference electrophiles and reference Lewis acids for the construction of comprehensive nucleophilicity^[9] and Lewis basicity scales.^[10] Variation of the *p*- and *m*-substituents of the benzhydrylium ions allowed a wide variation of their electrophilicities and Lewis acidities while the steric demand in the vicinity of the reaction centre is kept constant. During earlier work, we have demonstrated that the rates of the reactions of η -, π -, and σ -nucleophiles with carbocations and Michael

[a] Prof. Dr. B. Maji, Prof. Dr. X.-H. Duan, P. M. Jüstel, Dr. P. A. Byrne, Dr. A. R. Ofial, Prof. Dr. H. Mayr
Department Chemie
Ludwig-Maximilians-Universität München
Butenandstr. 5–13, 81377 München (Germany)
E-mail: Herbert.Mayr@cup.uni-muenchen.de


[b] Prof. Dr. B. Maji
Department of Chemical Sciences
Indian Institute of Science Education and Research Kolkata
Mohanpur 741246 (India)

[c] Prof. Dr. X.-H. Duan
Department of Chemistry, School of Chemistry
Xi'an Key Laboratory of Sustainable Energy Material Chemistry
Xi'an Jiaotong University
No. 28, Xianning West Road, Xi'an 710049 (P. R. China)

[d] Dr. P. A. Byrne
School of Chemistry, University College Cork
College Road, Cork (Ireland)




Supporting information for this article is available on the WWW under <https://doi.org/10.1002/chem.202100977>

© 2021 The Authors. Chemistry - A European Journal published by Wiley-VCH GmbH. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.



Scheme 2. Dialkyl chalcogenides 1–3 used in this study.

Table 1. Electrophilicity (E), electrofugality (E_f), and Lewis acidity (LA) parameters of the benzhydrylium ions used in this work.

						
	X	Y	Abbreviations	$E^{[a]}$	$E_f^{[b]}$	$LA^{[c]}$
4a	OPh	H	(pop)(Ph)CH ⁺	2.90	−3.52	4.42
4b	OMe	H	(ani)(Ph)CH ⁺	2.11	−2.09	3.10
4c	OMe	Me	(ani)(tol)CH ⁺	1.48	−1.32	2.00
4d	OMe	OPh	(ani)(pop)CH ⁺	0.61	−0.86	0.90
4e	OMe	OMe	(ani) ₂ CH ⁺	0	0	0
4f			(fur)(ani)CH ⁺	−0.81	0.61	(−1.11) ^[d]
4g			(fur) ₂ CH ⁺	−1.36	1.07	−1.29

[a] Electrophilicity parameters E for benzhydrylium ions from refs. [9] and [11]. [b] Electrofugality parameters E_f for benzhydrylium ions from ref. [12]. [c] Lewis acidity LA of benzhydrylium ions in dichloromethane, from ref. [10b]. [d] Interpolated Lewis acidity LA , see ref. [10b].

acceptors can be described by the linear-free-energy relationship [Eq. (1)].^[9]

$$\log k(20^\circ\text{C}) = s_N(N + E) \quad (1)$$

In Equation (1), nucleophiles are characterized by the solvent-dependent nucleophilicity parameter N and the susceptibility parameter s_N , while electrophiles are characterized by the solvent-independent electrophilicity parameter E .^[9] As explained in detail previously,^[9b] the unconventional expression of the linear free energy relationship (1) avoids far-reaching extrapolations by defining nucleophilicity N as the negative intercept on the abscissa in $\log k$ versus E plots. Based on Equation (1), the nucleophilicities of numerous n -, π -, and σ -nucleophiles, including amines, pyridines, imidazoles, imidazolines, and phosphanes, have been quantified.^[9d] In this work, we set out to determine the nucleophilicities of dialkyl chalcogenides 1–3 (Scheme 2) using the benzhydrylium ions 4 depicted in Table 1 as reference electrophiles.


Results and Discussion

Kinetics of the reactions of dialkyl chalcogenides 1–3 with benzhydrylium ions 4

To characterize the sulfonium ions 4-SR₂ generated by the reactions of the thioethers 2a–d with benzhydrylium ions, benzhydrylium triflate 4e-OTf (generated by mixing 4e-Cl with trimethylsilyl triflate) was combined with 2a–d in dichloromethane. As specified by the NMR spectroscopic analysis of the crude products (see the Supporting Information) the corresponding dialkylbenzhydrylsulfonium triflates (4-SR₂)⁺TfO[−] were formed exclusively.

Previous work showed that due to the low Lewis basicities of 1–3 their combinations with benzhydrylium ions, which are better stabilized than 4g (Lewis acidity $LA < -2$), do not lead to adduct formation in 2 mM dichloromethane solutions.^[10a] On the other side, reactions of thio- and selenoethers with 4g and less stabilized benzhydrylium ions (4a–f, $LA > -1$) were so fast that conventional UV/Vis spectroscopy, even when combined with stopped-flow techniques, was not suitable to follow the decay of the absorbances of the benzhydrylium ions 4 because the reactions are faster than the mixing time in the stopped-flow instrument. For that reason, laser-flash photolytic techniques as described previously^[13] were employed to study the kinetics of the reactions of diethyl ether (1) and the thio- and selenoethers 2 and 3 with the benzhydrylium ions 4 (Scheme 3).

The benzhydrylium ions 4 were generated by laser flash irradiation (7 ns pulse, 266 nm, 40–60 mJ/pulse) of the benzhydryltriphenylphosphonium tetrafluoroborates 4-PPh₃BF₄ in the presence of the nucleophiles 2 or 3 in acetonitrile or dichloromethane at 20 °C. The intermediate benzhydrylium ions 4 were identified by their UV/Vis spectra.^[13] The rates of the combina-



Scheme 3. Generation of benzhydrylium ions 4 by laser-flash irradiation of the precursor phosphonium tetrafluoroborates 4-PPh₃BF₄ and their combination with dialkyl chalcogenides (at 20 °C in MeCN or CH₂Cl₂).

tion reactions were followed by monitoring the decay of the absorbances of **4** at or close to their absorption maxima.

Quantitative kinetic measurements were not carried out with diethyl ether (**1**) as the addition of two equivalents of **1** to a 0.035 M solution of (ani)₂CH⁺ BF₄⁻ (**4e**) in CH₂Cl₂ at ambient temperature neither resulted in decolorization of the solution nor in a change of the ¹H NMR spectrum of **4e** within 10 minutes. After 1 h, about one-third of **4e** was reduced with formation of bis(4-methoxyphenyl)methane, indicating that hydride transfer will disturb the kinetics of Lewis adduct formation with more reactive benzhydrylium ions. This observation is in line with Penczek's thorough investigation of the interactions of ethers and acetals with Ph₃C⁺, which has an electrophilicity similar to **4e**.^[9d] For the coordination of Ph₃C⁺ with Et₂O (**1**) at 25 °C an equilibrium constant $K=0.23\text{ M}^{-1}$,^[14] and for the hydride transfer from Et₂O to Ph₃C⁺ a second-order rate constant of $3\times 10^{-4}\text{ M}^{-1}\text{ s}^{-1}$ was determined.^[15]

All reactions of the benzhydrylium ions **4** with the thioethers **2a–d** and dimethyl selenide **3** reported herein were well behaved and gave rise to mono-exponential decays of the absorbances of **4** as shown in Figure 1 for the reaction of **4d** with **2b**. Because of the low concentrations of PPh₃ generated

Table 2. Second-order rate constants (k) for the reactions of the thio- and selenoethers **2** and **3** with benzhydrylium ions **4** in CH₂Cl₂ at 20 °C.

Nucleophile	N / s_N ^[a]	Ar ₂ CH ⁺	4	k [M ⁻¹ s ⁻¹]
Me ₂ S (2a)	12.32, 0.72	(fur) ₂ CH ⁺	4g	8.10×10^7
		(fur)(ani)CH ⁺	4f	2.02×10^8
		(ani) ₂ CH ⁺	4e	5.14×10^8 ^[b]
		(ani)(pop)CH ⁺	4d	8.80×10^8 ^[b]
2a in MeCN	(12.7, 0.72) ^[c]	(fur) ₂ CH ⁺	4g	1.55×10^8
		(fur)(ani)CH ⁺	4f	3.29×10^8 ^[b]
		(ani) ₂ CH ⁺	4e	7.46×10^8 ^[b]
		(ani)(pop)CH ⁺	4d	1.22×10^9 ^[b]
<i>n</i> Bu ₂ S (2b)	11.86, 0.74	(ani)(Ph)CH ⁺	4b	2.87×10^9 ^[b]
		(fur) ₂ CH ⁺	4g	5.13×10^7
		(fur)(ani)CH ⁺	4f	1.30×10^8
		(ani) ₂ CH ⁺	4e	2.73×10^8 ^[b]
THT (2c)	(13.1, 0.72) ^[c]	(ani)(pop)CH ⁺	4d	4.61×10^8 ^[b]
		(fur) ₂ CH ⁺	4g	2.88×10^8 ^[b]
		(fur)(ani)CH ⁺	4f	3.85×10^8 ^[b]
		(ani) ₂ CH ⁺	4e	8.15×10^8 ^[b]
2c in MeCN	(13.3, 0.72) ^[c]	(ani)(pop)CH ⁺	4d	1.22×10^9 ^[b]
		(fur) ₂ CH ⁺	4g	3.99×10^8 ^[b]
		(fur)(ani)CH ⁺	4f	6.89×10^8 ^[b]
		(ani) ₂ CH ⁺	4e	1.65×10^9 ^[b]
THTP (2d)	11.94, 0.75	(ani)(pop)CH ⁺	4d	1.68×10^9 ^[b]
		(ani)(Ph)CH ⁺	4b	4.97×10^9 ^[b]
		(pop)(Ph)CH ⁺	4a	5.17×10^9 ^[b]
		(fur) ₂ CH ⁺	4g	8.37×10^7
Me ₂ Se (3)	(12.6, 0.72) ^[c]	(fur)(ani)CH ⁺	4f	2.16×10^8
		(ani) ₂ CH ⁺	4e	5.57×10^8 ^[b]
		(ani)(pop)CH ⁺	4d	7.34×10^8 ^[b]
		(fur)(ani)CH ⁺	4f	3.16×10^8 ^[b]
		(ani) ₂ CH ⁺	4e	4.90×10^8 ^[b]
		(ani)(pop)CH ⁺	4d	1.08×10^9 ^[b]
		(ani)(tol)CH ⁺	4c	2.06×10^9 ^[b]

[a] Nucleophile-specific parameters N and s_N according to Equation (1). [b] Because of the proximity of the diffusion limit, not used for the calculation of N and s_N . [c] As the available rate constants are close to the diffusion limit, they are not used for the calculation of N and s_N ; for that reason, $s_N=0.72$ was assumed to be the same as for structurally analogous nucleophiles and combined with the smallest rate constant of the series to obtain an estimate for N .




Figure 1. a) Kinetics of the reaction of **2b** ($[2b]_0 = 1.68\text{ mM}$) with **4d** at 20 °C in CH₂Cl₂ were monitored b) by the exponential decay of the absorbance of **4d** at 516 nm during the course of the reaction (**4d** generated by laser flash photolysis of **4d**-PPh₃BF₄). c) The second-order rate constant $k = 4.61\times 10^8\text{ M}^{-1}\text{ s}^{-1}$ corresponds to the slope of the linear correlation of k_{obs} with $[2b]$.

by photolytic cleavage, recombination of PPh₃ with the benzhydrylium ions did not compete with the reactions of the chalcogenides. First-order rate constants k_{obs} [s⁻¹] were obtained by fitting the mono-exponential function $A_t = A_0 e^{-k_{\text{obs}} t} + C$ to the decays of the absorbances. Plots of k_{obs} versus the concentrations of the nucleophiles were linear, indicating second-order rate laws, and the second-order rate constants k [M⁻¹ s⁻¹] listed in Table 2 were derived from the slopes of such plots.

Determination of nucleophile-specific parameters N and s_N

As previously reported for numerous reactions of benzhydrylium ions with different families of nucleophiles, plots of $\log k$ versus electrophilicity E are generally linear up to rate constants of approximately $2\times 10^8\text{ M}^{-1}\text{ s}^{-1}$ and flatten as the diffusion limit

is approached.^[11] The linear parts of these correlations, which follow Equation (1), have been used to derive N from the intercepts on the abscissa and s_N as the slopes.^[9]

Analogously, Figure 2 shows a flattening of $\log k$ versus E plots when the second-order rate constants k for the reactions of dialkyl sulfides **2** with benzhydrylium ions **4** exceed $2 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$. Only the reactions of the least electrophilic benzhydrylium ions **4f** and **4g** with the thioethers **2a**, **2b**, and **2d** proceed with rate constants smaller than $2 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$ and can be used to evaluate N and s_N for these nucleophiles in the conventional way as illustrated in Figure 2. Since all rate constants determined for the stronger nucleophiles **2c** and **3** are greater than $2 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$, a different way for characterizing the nucleophilic reactivities of **2c** and **3** was needed.

Reactions with less electrophilic carbocations cannot be used for this purpose because such carbocations do not give adducts at 20 °C due to the lack of a thermodynamic driving force. As specified in the footnotes of Table 2, approximate N values for the more nucleophilic Lewis bases **2c** and **3** (and for **2a** in acetonitrile) were calculated from the rate constants of the slowest reactions of these chalcogenides by applying




Figure 2. Plot of $\log k$ versus the electrophilicity parameter E for the reactions of dimethyl sulfide (**2a**), dibutyl sulfide (**2b**), and tetrahydrothiophene (THT, **2c**) with benzhydrylium ions **4** in CH_2Cl_2 at 20 °C.

Equation (1) with an assumed $s_N=0.72$, the susceptibility determined for the structurally related dimethyl sulfide **2a**. Similar values of s_N have also been obtained for **2b** and **2d**.

Intrinsic barriers

For some of the reactions listed in Table 2, equilibrium constants have previously been determined^[10a] (Table 3). According to the Marcus Equation (2), the activation energy ΔG^\ddagger of a reaction can be expressed by the combination of the reaction Gibbs energy $\Delta_r G^\circ$ with the intrinsic barrier ΔG_0^\ddagger , which corresponds to the activation Gibbs energy for a reaction with $\Delta_r G^\circ = 0$.^[16]

$$\Delta G^\ddagger = \Delta G_0^\ddagger + 0.5 \Delta_r G^\circ + (\Delta_r G^\circ)^2 / (16 \Delta G_0^\ddagger) \quad (2)$$

Substitution of $\Delta_r G^\circ$ and ΔG^\ddagger into Equation (2) yields the intrinsic barriers ΔG_0^\ddagger . Values of ΔG_0^\ddagger determined for the reactions of nucleophiles **2** and **3** with benzhydrylium ions (**4**) are listed in Table 3. Zhu has criticized Marcus' derivation of Equation (2) and developed an alternative expression, which differs from Equation (2) by the absence of the last term.^[17] Since the intrinsic barriers calculated by Zhu's equation differ only insignificantly from those calculated by Equation (2) (+0.4 to 1.4 kJ mol^{-1}), we only list the Marcus intrinsic barriers in Table 3 in order to retain comparability with the majority of published intrinsic barriers.

In previous work, we showed that within a reaction series (one nucleophile with different benzhydrylium ions) Marcus intrinsic barriers ΔG_0^\ddagger decrease with increasing reactivity of the benzhydrylium ions for nucleophiles with $s_N > 0.67$, and increase for nucleophiles with $s_N < 0.67$.^[18] Accordingly, the intrinsic barriers are almost constant within the different reaction series of Table 3, because the corresponding susceptibility parameters ($0.72 < s_N < 0.75$) are close to 0.67.

Intrinsic barriers below 35 kJ mol^{-1} , as derived for the reactions of benzhydrylium ions with thio- and selenoethers (Table 3), are the smallest ones we have observed so far. These

Table 3. Equilibrium constants K , Gibbs reaction energies $\Delta_r G^\circ$, Gibbs activation energies ΔG^\ddagger , and Marcus intrinsic barriers ΔG_0^\ddagger for the reactions of the thio- and selenoethers **2** and **3** with benzhydrylium ions (Ar_2CH^+ , **4**) in CH_2Cl_2 at 20 °C.

Nucleophile (Lewis base)	Ar_2CH^+	4	$K^{[a]}$ [M^{-1}]	$\Delta_r G^\circ^{[a]}$ [kJ mol^{-1}]	$\Delta G^\ddagger^{[b]}$ [kJ mol^{-1}]	$\Delta G_0^\ddagger^{[c]}$ [kJ mol^{-1}]
Me ₂ S (2a)	(fur) ₂ CH ⁺	4g	2.16×10^2	-13.1	27.4	33.6
	(ani) ₂ CH ⁺	4e	8.91×10^3	-22.2	22.9	33.0
	(ani)(pop)CH ⁺	4d	6.67×10^4	-27.1	21.6	33.7
nBu ₂ S (2b)	(fur) ₂ CH ⁺	4g	1.52×10^2	-12.2	28.5	34.3
	(ani) ₂ CH ⁺	4e	6.15×10^3	-21.3	24.4	34.2
	(ani)(pop)CH ⁺	4d	3.61×10^4	-25.6	23.1	34.7
THT (2c)	(fur) ₂ CH ⁺	4g	7.94×10^2	-16.3	24.3	31.9
	(ani) ₂ CH ⁺	4e	1.22×10^4	-22.9	21.7	32.2
	(ani)(pop)CH ⁺	4d	6.43×10^4	-27.0	20.8	32.9
THTP (2d)	(fur) ₂ CH ⁺	4g	8.86×10^2	-16.5	27.3	35.1
	(ani) ₂ CH ⁺	4e	3.04×10^4	-25.2	22.7	34.1
	(ani)(pop)CH ⁺	4d	1.05×10^5	-28.2	22.0	34.7
Me ₂ Se (3)	(ani) ₂ CH ⁺	4e	9.50×10^2	-16.7	23.0	30.8
	(ani)(pop)CH ⁺	4d	7.65×10^3	-21.8	21.0	31.0

[a] From ref. [10a]. [b] Calculated by applying the Eyring equation on the rate constants in Table 2. [c] Calculated by using Marcus Equation (2).

low intrinsic barriers account for the fact that only a small number of benzhydrylium ions undergo activation-controlled reactions with **2** and **3**. The reactions of **2** and **3** with slightly more reactive carbenium ions occur under diffusion control, while the reactions of **2** and **3** with slightly less reactive carbenium ions do not occur at all (i.e., no products are formed).

Intrinsic barriers ΔG_0^\ddagger , that is, the barriers for reactions with reaction Gibbs energies $\Delta_r G^\circ = 0$, are linked to the reorganization energies λ by the relationship $\Delta G_0^\ddagger = \lambda/4$. Since little structural reorganization is required when electrophiles attack at the lone electron pair of sp^3 -hybridized atoms, dialkyl chalcogenides as well as tertiary amines (for DABCO and quinuclidine, $\Delta G_0^\ddagger \approx 40 \text{ kJ mol}^{-1}$)^[10b] react via low intrinsic barriers. In previous work, the kinetically controlled S attack at thiocyanate ions has also been assigned to the lower reorganization energy for this site of attack ($\Delta G_0^\ddagger = 35\text{--}38 \text{ kJ mol}^{-1}$ for the S attack at SCN^-).^[19]

Nucleofugalities

In analogy to Equation (1), Equation (3) has been used for the construction of a comprehensive nucleofugality scale,^[12] which allows one to calculate the rates of heterolytic cleavages k_{rev} [s^{-1}] of R–X from the electrofugality parameter E_f of R^+ and the solvent-specific nucleofuge parameters N_f and s_f of X or X^- .

$$\log k_{\text{rev}}(25^\circ\text{C}) = s_f(E_f + N_f) \quad (3)$$

From the rate constants k of the reactions of the benzhydrylium ions **4** with the thio- and selenoethers **2** and **3** determined in this work (Table 2) and the previously reported equilibrium constants K ,^[10a] the heterolysis rate constants k_{rev} [s^{-1}] of the corresponding trialkyl sulfonium ions have now been calculated as the ratios k/K at 20°C in CH_2Cl_2 (Table 4, column 5). Plots of $\log k_{\text{rev}}$ against the known electrofugality parameters E_f of benzhydrylium ions,^[12] resulted in linear




Figure 3. Plot of the reverse rate constants ($\log k_{\text{rev}}$) for the reactions of nucleophiles **2** and **3** with benzhydrylium ions **4** in CH_2Cl_2 at 20°C against the electrofugality parameters E_f of **4** (see the Supporting Information for analogous $\log k_{\text{rev}}$ vs. E_f plots for **2b** and **2c**).

correlations (Figure 3) from which the nucleofuge-specific parameters s_f and N_f for **2** and **3** were derived (Table 4).

Jurić, Denegri, and Kronja recently used Equation (3) to derive the nucleofugalities of dimethyl sulfide (**2a**) and THT (**2c**) from the solvolysis rate constants k_{rev} [s^{-1}] of the corresponding benzhydryl sulfonium ions in pure and aqueous ethanol and methanol.^[20,21] By using their N_f and s_f parameters^[20,21] to calculate k_{rev} for the corresponding sulfonium ions in ethanol at 25°C by Equation (3) one arrives at values for k_{rev} that are 2 to 3 orders of magnitude smaller than those calculated from the k/K ratios in dichloromethane at 20°C (two right columns of Table 4). This large difference cannot be due to the use of different solvents used in both studies because a variation of

Table 4. Reverse rate constants (k_{rev}) for the reactions of nucleophiles **2** and **3** with benzhydrylium ions (Ar_2CH^+ , **4**) and the resulting nucleofuge-specific parameters (N_f and s_f) in CH_2Cl_2 at 20°C .

Nucleofuge	N_f, s_f [a]	Ar_2CH^+	E_f [b]	k_{rev} [c] [s^{-1}]	k_{rev} [d] [s^{-1}]
Me ₂ S (2a)	6.33, 0.75	(fur) ₂ CH ⁺	4g	3.75×10^5	1.04×10^3
		(ani) ₂ CH ⁺	4e	5.77×10^4	1.25×10^2
		(ani)(pop)CH ⁺	4d	1.32×10^4	2.28×10^1
<i>n</i> Bu ₂ S (2b)	6.36, 0.74	(fur) ₂ CH ⁺	4g	3.38×10^5	
		(ani) ₂ CH ⁺	4e	4.44×10^4	
		(ani)(pop)CH ⁺	4d	1.28×10^4	
THT (2c)	7.26, 0.66	(fur) ₂ CH ⁺	4g	3.63×10^5	1.38×10^3
		(ani) ₂ CH ⁺	4e	6.68×10^4	1.66×10^2
		(ani)(pop)CH ⁺	4d	1.90×10^4	3.01×10^1
THTP (2d)	7.33, 0.59	(fur) ₂ CH ⁺	4g	9.45×10^4	
		(ani) ₂ CH ⁺	4e	1.83×10^4	
		(ani)(pop)CH ⁺	4d	6.99×10^3	
Me ₂ Se (3)	8.72, 0.66	(ani) ₂ CH ⁺	4e	5.16×10^5	
		(ani)(pop)CH ⁺	4d	1.41×10^5	
		(ani)(pop)CH ⁺	4d		

[a] From Equation (3). [b] From ref. [12]. [c] Calculated by the relation $k_{\text{rev}} = k/K$ in Tables 2 and 3. [d] Extrapolated by Equation (3) from E_f in Table 4 and solvolysis rate constants in ethanol at 25°C (from ref. [20]).

the solvent has been reported to have only a minor effect on the nucleofugality of neutral leaving groups.^[20–22]

In order to resolve the discrepancy of k_{rev} determined in this and earlier work^[20,21] we performed dynamic NMR (DNMR) studies in analogy to previously reported DNMR investigations of trialkyl sulfonium ions.^[23] The *S*-methyl groups of a mixture of **4g-SMe₂** and Me₂S (**2a**, marked red in Figure 4) in CD₂Cl₂ at –80 °C resonated separately at δ 2.78 for **4g-SMe₂** and δ 2.02 for **2a** (red) in the 400 MHz ¹H NMR spectrum (Figure 4). Coalescence of these *S*-methyl resonances was observed when the temperature was gradually raised (Figure 4b). Because the concentration of Me₂S arising from heterolytic cleavage of the sulfonium ion **4g-SMe₂** (black in Figure 4) is low compared to the concentration of extra Me₂S in solution (red in Figure 4), benzhydrylium ion **4g** reacts preferentially with **2a** from the solution and gives rise to the observed dynamic NMR phenomenon. Line shape analysis (LSA) for ¹H NMR spectra in the temperature range from –45 to –25 °C furnished the corresponding exchange rate constants $k_{\text{rev}}(T)$ and Eyring activation parameters (Supporting Information), which allowed us to extrapolate a first-order rate constant of $1.3 \times 10^5 \text{ s}^{-1}$ for the Me₂S exchange at 20 °C. Non-coordinated **4g** is not detectable in the NMR spectrum, in accord with the equilibrium constant $K = 216 \text{ M}^{-1}$ for the Lewis adduct formation from **4g** and **2a** (Table 3). The (pseudo)-first-order rate constant for the reaction of **4g** with **2a** can be calculated as $4.3 \times 10^6 \text{ s}^{-1}$ ($= 8.1 \times 10^7 \text{ M}^{-1} \text{ s}^{-1} \times 0.053 \text{ M}$) under these conditions. Since this value is 33 times greater than the exchange rate constant determined by DNMR, we can conclude that the latter (k_{rev} in Figure 4) corresponds to the heterolytic cleavage of the carbon-sulfur bond in the sulfonium ion **4g-SMe₂**.

The value of k_{rev} for **4g-SMe₂** determined by DNMR is three times smaller than that derived as k/K from experimentally measured equilibrium and forward rate constants (k_{rev} in CH₂Cl₂, Table 4), an acceptable agreement in view of the completely different ways of derivation. The analogous DNMR investigation of (ani)₂CH-SMe₂⁺ (**4e-SMe₂**) in CD₂Cl₂ in the temperature range from –50 to 0 °C resulted in an extrapolated $k_{\text{rev}}(20 \text{ °C}) = 2.6 \times 10^4 \text{ s}^{-1}$, two times smaller than k_{rev} derived from the k/K ratio.

After confirming the correct order of magnitude of k_{rev} derived from k/K for alkoxy-substituted benzhydryldimeth-

ylsulfonium ions, let us return to the question, why the k_{rev} values in Table 4, column 5 are so much larger than those calculated by Equation (3) from the ethanolysis rate constants of the Zagreb group (Table 4, column 6).

The ethanolysis rate constants in refs^[20,21] refer to benzhydryldimethylsulfonium ions composed of the parent and halogen-substituted benzhydrylium ions, all of which are highly electrophilic ($E \geq 5.20$).^[9] Substitution of this E value and the nucleophilicity parameters of **2a** and **2c** (Table 2) into Equation (1) shows that all benzhydrylium ions generated in the ethanolysis studies undergo diffusion-controlled reactions with the thioethers **2** ($k^{\text{eq}(1)} > 10^{10} \text{ M}^{-1} \text{ s}^{-1}$). In contrast, the reactions of the thioethers **2** with the alkoxy-substituted benzhydrylium ions **4d–g** investigated in this work are activation controlled (Table 2). As a consequence of the principle of microscopic reversibility, the transition states of the heterolyses of the parent and halogen-substituted benzhydryl sulfonium ions correspond to the carbocations (Figure 5a), while those of the alkoxy-substituted analogues only resemble the carbocations (Figure 5b).

As linear free energy relationships break down when a change from activation-control to diffusion-control^[10b,c] is involved, one can explain why the nucleofuge-specific parameters N_f and s_f derived from ethanolysis rate constants (Figure 5a) are not applicable to the heterolysis rates of the alkoxy-substituted benzhydryl sulfonium ions (Figure 5b).

Relationships between structures and reactivities

The observation that diethyl ether (**1**) does not form adducts ($K < 10$) with benzhydrylium ions of $LA < 1$ (Table 1) at 20 °C shows that ethers are weaker Lewis bases towards carbenium ions than structurally related sulfides. This ordering is in analogy with the relative proton affinities in the gas phase (Me₂S 197, Me₂O 186 kcal mol^{–1}), but not with the relative Brønsted basicities in an aqueous solution [$\text{p}K_{\text{aH}}(\text{Me}_2\text{O}) = -2.52$ vs. $\text{p}K_{\text{aH}}(\text{Me}_2\text{S}) = -6.95$].^[6]

Scheme 4 shows that Me₂S (**2a**) is a slightly stronger nucleophile and stronger Lewis base than *n*Bu₂S (**2b**), with the consequence that the nucleofugalities (k_{rev}) of these two




Figure 4. a) Dimethylsulfide exchange between **4g-SMe₂** and free **2a** in CD₂Cl₂. b) Temperature-dependent proton resonances for the *S*-methyl groups evaluated by line-shape analysis (DNMR6 algorithm) for the determination of exchange rate constants $k_{\text{rev}}(T)$.






Figure 5. Gibbs energy [kJ mol^{-1}] profiles for the heterolysis of benzhydryldimethylsulfonium ions a) **4h-SMe₂** and b) **4e-SMe₂**.



Scheme 4. Rate and equilibrium constants for the reactions of thioethers **2** with the benzhydrylium ion **4g** in CH_2Cl_2 at 20°C (with $k_{\text{rev}} = k/K$).



Scheme 5. Rate and equilibrium constants for the reactions of dialkyl chalcogenides with the benzhydrylium ion **4e** in CH_2Cl_2 at 20°C (with $k_{\text{rev}} = k/K$).

sulfides are almost identical. Interestingly, dimethyl sulfide (**2a**) and tetrahydrothiopyran (**2d**) have the same nucleophilic reactivity, though the cyclic sulfide **2d** is a four-times stronger Lewis base. The higher intrinsic barriers in reactions of **2d** (Table 3), which account for this ranking, also account for the fact that despite the comparable Lewis basicities of the cyclic sulfides **2c** and **2d**, tetrahydrothiopyran (**2d**) reacts 3-times more slowly than tetrahydrothiophene (**2c**) with **4g**.

Due to the low Lewis basicity of dimethylselenide (**3**), its reaction with $(\text{fur})_2\text{CH}^+$ (**4g**) proceeds only with a very low degree of conversion. For that reason, its nucleophilic reactivity




Figure 6. Scope of reactions of thio- and selenoethers with benzhydrylium ions of different electrophilicity E .

could not be included in the comparisons of Scheme 4. On the other hand, Scheme 5 shows that the more Lewis acidic benzhydrylium ion **4e** can be used as a reference for the comparison of dimethyl selenide (**3**) with dimethyl sulfide (**2a**).

Though dimethyl sulfide (**2a**) is a tenfold stronger Lewis base towards benzhydrylium ions than dimethyl selenide (**3**), according to Table 3, both compounds have equal nucleophilicities, indicating a higher intrinsic barrier for the reaction of **2a**. As a consequence, selenoether **3** is a tenfold better nucleofuge than thioether **2a** (Scheme 5).

Conclusion

Dialkyl sulfides **2a–d** and dimethyl selenide **3** react with benzhydrylium ions (which can be considered representative carbenium ions and π -electrophiles in general) with unusually low intrinsic barriers. For this reason, these thio- and selenoethers undergo diffusion-controlled reactions with almost all carbenium ions that have sufficiently high Lewis acidities for the reactions to be exergonic. Thus, only a small number of carbenium ions undergo activation-controlled reactions with these dialkyl chalcogenides, providing rate constants that can be used to calculate the corresponding nucleophilicity parameters N and s_N from Equation 1 (Figure 6).

Because of the similarities of their susceptibilities s_N , the relative reactivities of the nucleophiles depicted in Figure 7 exhibit only a minor dependence on the nature of their electrophilic reaction partners. For that reason, their relative nucleophilic reactivities can be approximated by the nucleophilicity parameters N . Several entries in Figure 7 show that the nucleophilicities N and Lewis basicities LB depend only slightly on the solvent (CH_2Cl_2 vs. MeCN) due to the fact that the combination of a cation and a neutral reactant yields another cation. For that reason, we shall not specifically mention solvent effects in the following discussion.

The thioethers **2** and the selenoether **3** cover a very small range of both the nucleophilicity and the Lewis basicity scales in Figure 7. Although they are by far the weakest Lewis bases in this ranking, their nucleophilicities are comparable to those of several commonly used organocatalysts. Thus, dialkyl chalcogenides **2** and **3** have a similar nucleophilicity to pyridine and *N*-methylimidazole, though the Lewis basicities of the latter

derived from equilibrium constants of their reactions with benzhydrylium ions are 8 to 11 orders of magnitude greater. Analogously, the nucleophilic reactivities of the chalcogenides are only 10 to 1000 times smaller than those of PPh_3 and PBu_3 , whereas the corresponding Lewis basicities differ by more than 10 orders of magnitude. Both comparisons reflect the very low intrinsic barriers of the reactions of the chalcogenides.

What is the reason for the low Marcus intrinsic barriers of these reactions of chalcogenides which are responsible for their unique reactivities? Hoz has reported that the barriers of the identity reactions in Equation (4) (i.e., the intrinsic barriers) decrease, the further right X is in the periodic table,^[24] which can be explained by increasing electronegativity of X .



That the intrinsic barriers remain almost constant as one goes down a particular group in the periodic table was




Figure 7. a) Nucleophilicity parameters N/s_N and b) Lewis-basicity parameters LB of chalcogenides and pnictogenides in acetonitrile and dichloromethane.

explained by Arnaut and colleagues by decreasing C–X force constants associated with flattening of the intersecting parabola in the Marcus model,^[25a] which is almost compensated for by increasing the C–X bond length. More recent work showed that in analogous identity reactions with neutral nucleophiles at benzyl derivatives, intrinsic barriers increase as one goes down the periodic table, but again, the intrinsic barrier for SMe_2 exchange is lower than that for NMe_3 exchange.^[25b] As the relative nucleophilicities toward C_{sp^2} centres have been reported to be linearly correlated with relative nucleophilicities toward C_{sp^3} centres,^[26] the same reason appears to account for the fact that the intrinsic barriers for the reactions of S-nucleophiles with carbenium ions are lower than those for N- and P-nucleophiles.

The astonishing result that dialkyl sulfides and selenides are as nucleophilic as the much stronger (Brønsted and Lewis) bases pyridine and imidazole is not only due to the exceptionally low intrinsic barriers for the reactions of the chalcogenides but also due to the particularly high intrinsic barriers for the reactions of the N-heteroarenes. As intrinsic barriers are associated with the degree of reorganization,^[16,27] the nuclear movements accompanying the reorganization of the aromatic π -system during the electrophilic attack at the nitrogen of the heteroarenes^[28] account for the fact that pyridines and imidazoles react via higher intrinsic barriers than alkylamines with the consequence that N-heteroarenes and alkylamines differ much more in nucleophilicity than in basicity (Figure 7).

The exceptionally low intrinsic barriers of the reactions of dialkyl sulfides and selenides, which are responsible for their high nucleophilicities despite their low Lewis basicities, also account for their high nucleofugalities and thus for their suitability as group-transfer reagents.

Acknowledgements

We thank the Department Chemie (LMU München) for financial support. We are grateful to the Alexander von Humboldt Foundation and the Carl Friedrich von Siemens Foundation for provision to P.A.B. of a Humboldt Research Fellowship for Postdoctoral Researchers. Open access funding enabled and organized by Projekt DEAL.

Conflict of Interest

The authors declare no conflict of interest.

Keywords: kinetics · Lewis bases · linear free energy relationships · thermodynamics · thioethers

- [1] a) L. A. Wessjohann, M. C. N. Brauer, K. Brand in *Enantioselective Organocatalyzed Reactions I* (Ed.: R. Mahrwald), Springer, Dordrecht, 2011, pp 209–314; b) Y. Nishibayashi, S. Uemura in *Organoselenium Chemistry* (Ed.: T. Wirth), Springer, Berlin, 2000, pp 235–255; c) *Handbook of Chalcogen Chemistry: New Perspectives in Sulfur, Selenium and Tellurium*

- (Ed.: F. A. Devillanova), RSC Publishing, Cambridge, 2007; d) Y. Tang, X.-L. Sun, *Science of Synthesis, Vol. 39* Thieme, Stuttgart, 2008, pp 501–541. [2] a) G. L. Cantoni, *J. Am. Chem. Soc.* 1952, 74, 2942–2943; b) W. Struck, M. L. Thompson, L. S. Wong, J. Micklefield, *ChemBioChem* 2012, 13, 2642–2655; c) J. Deen, C. Vranken, V. Leen, R. K. Neely, K. P. F. Janssen, J. Hofkens, *Angew. Chem. Int. Ed.* 2017, 56, 5182–5200; *Angew. Chem.* 2017, 129, 5266–5285. [3] a) W. Appel, *Clinical Biochemistry* 1986, 19, 317–322; b) H.-H. Otto, T. Schirmeister, *Chem. Rev.* 1997, 97, 133–172; c) T. Stadtman, *J. Biol. Chem.* 1991, 266, 16257–16260. [4] a) R. Okazaki, N. Tokitoh, V. K. Aggarwal, C. L. Winn, *Encyclopedia of Reagents for Organic Synthesis*, Wiley, New York, 2006, DOI: 10.1002/047084289X.rd372.pub2; b) A.-H. Li, L.-X. Dai, V. K. Aggarwal, *Chem. Rev.* 1997, 97, 2341–2372; c) E. M. McGarrigle, E. L. Myers, O. Illa, M. A. Shaw, S. L. Riches, V. K. Aggarwal, *Chem. Rev.* 2007, 107, 5841–5883. [5] a) D. Basavaiah, A. J. Rao, T. Satyanarayana, *Chem. Rev.* 2003, 103, 811–892; b) D. Basavaiah, B. S. Reddy, S. S. Badsara, *Chem. Rev.* 2010, 110, 5447–5674; c) D. Basavaiah, R. T. Naganaboina, *New J. Chem.* 2018, 42, 14036–14066. [6] E. M. Arnett, *Acc. Chem. Res.* 1973, 6, 404–409. [7] R. Curci, F. Di Furia, A. Levi, V. Lucchini, G. Scorrano, *J. Chem. Soc. Perkin Trans. 2* 1975, 341–344. [8] a) J. E. Leffler, E. Grunwald, *Rates and Equilibria of Organic Reactions*, Wiley, New York, 1963; b) A. Williams, *Free Energy Relationships in Organic and Bio-Organic Chemistry*, The Royal Society of Chemistry, Cambridge, 2003; c) F. G. Bordwell, T. A. Cripe, D. L. Hughes in *Nucleophilicity* (Eds.: J. M. Harris, S. P. McManus), American Chemical Society, Chicago, 1987, pp 137–153; d) R. G. Pearson, H. R. Sobel, J. Songstad, *J. Am. Chem. Soc.* 1968, 90, 319–326; e) J. Hine, R. D. Weimar Jr., *J. Am. Chem. Soc.* 1965, 87, 3387–3396. [9] a) H. Mayr, T. Bug, M. F. Gotta, N. Hering, B. Irrgang, B. Janker, B. Kempf, R. Loos, A. R. Ofial, G. Remennikov, H. Schimmel, *J. Am. Chem. Soc.* 2001, 123, 9500–9512; b) H. Mayr, *Angew. Chem. Int. Ed.* 2011, 50, 3612–3618; *Angew. Chem.* 2011, 123, 3692–3698; c) H. Mayr, *Tetrahedron* 2015, 71, 5095–5111; d) A database of reactivity parameters E , N , and s_N is freely accessible through the webpage www.cup.lmu.de/oc/mayr/DBintro.html (accessed on March 2021). [10] a) H. Mayr, J. Ammer, M. Baidya, B. Maji, T. A. Nigst, A. R. Ofial, T. Singer, *J. Am. Chem. Soc.* 2015, 137, 2580–2599; b) H. Mayr, A. R. Ofial, *Acc. Chem. Res.* 2016, 49, 952–965; c) H. Mayr, A. R. Ofial, *Pure Appl. Chem.* 2017, 89, 729–744. [11] J. Ammer, C. Nolte, H. Mayr, *J. Am. Chem. Soc.* 2012, 134, 13902–13911. [12] N. Streidl, B. Denegri, O. Kronja, H. Mayr, *Acc. Chem. Res.* 2010, 43, 1537–1549. [13] a) J. Ammer, C. F. Sailer, E. Riedle, H. Mayr, *J. Am. Chem. Soc.* 2012, 134, 11481–11494; b) J. Ammer, H. Mayr, *J. Phys. Org. Chem.* 2013, 26, 956–969. [14] a) S. Słomkowski, S. Penczek, *J. Chem. Soc. Perkin Trans. 2* 1974, 1718–1722; b) S. Penczek, R. Szymanski, *Polym. J.* 1980, 12, 617–628. [15] S. Penczek, *Makromol. Chem.* 1974, 175, 1217–1252. [16] a) R. A. Marcus, *Annu. Rev. Phys. Chem.* 1964, 15, 155–196; b) W. J. Albery, *Annu. Rev. Phys. Chem.* 1980, 31, 227–263; c) R. A. Marcus, *Pure Appl. Chem.* 1997, 69, 13–29. [17] G.-B. Shen, K. Xia, X.-T. Li, J.-L. Li, Y.-H. Fu, L. Yuan, X.-Q. Zhu, *J. Phys. Chem. A* 2016, 120, 1779–1799. [18] C. Schindele, K. N. Houk, H. Mayr, *J. Am. Chem. Soc.* 2002, 124, 11208–11214. [19] R. Loos, S. Kobayashi, H. Mayr, *J. Am. Chem. Soc.* 2003, 125, 14126–14132. [20] a) S. Jurić, B. Denegri, O. Kronja, *J. Org. Chem.* 2010, 75, 3851–3854; b) S. Jurić, B. Denegri, O. Kronja, *J. Phys. Org. Chem.* 2012, 25, 147–152. [21] B. Denegri, M. Matić, O. Kronja, *Synthesis* 2017, 49, 3422–3432. [22] D. N. Kevill, S. W. Anderson, N. H. Ismail, *J. Org. Chem.* 1996, 61, 7256–7262. [23] a) D. Darwish, G. Tourigny, *J. Am. Chem. Soc.* 1966, 88, 4303–4304; b) M. Nakamura, H. Kihara, N. Nakamura, M. Oki, *Org. Magn. Reson.* 1979, 12, 702–707; c) D. Darwish, S. H. Hui, R. Tomilson, *J. Am. Chem. Soc.* 1968, 90, 5631–5632. [24] a) S. Hoz, H. Basch, J. L. Wolk, T. Hoz, E. Rozental, *J. Am. Chem. Soc.* 1999, 121, 7724–7725; b) R. Yi, H. Basch, S. Hoz, *J. Org. Chem.* 2002, 67, 5891–5895. [25] a) L. Arnaut, A. Pais, S. Formosinho, *J. Mol. Struct.* 2001, 563, 1–17; b) R. Robiette, T. Trieu-Van, V. K. Aggarwal, J. N. Harvey, *J. Am. Chem. Soc.* 2016, 138, 734–737.

- [26] a) J. W. Bunting, J. M. Mason, C. K. M. Heo, *J. Chem. Soc. Perkin Trans. 2* **1994**, 2291–2300; b) J. P. Richard, M. M. Toteva, J. Crueiras, *J. Am. Chem. Soc.* **2000**, *122*, 1664–1674; c) T. B. Phan, M. Breugst, H. Mayr, *Angew. Chem. Int. Ed.* **2006**, *45*, 3869–3874; *Angew. Chem.* **2006**, *118*, 3954–3959; d) P. M. Jüstel, C. D. Pignot, A. R. Ofial, *J. Org. Chem.* **2021**, *86*, 5965–5972.
- [27] J. Hine, *Adv. Phys. Org. Chem.* **1977**, *15*, 1–61.
- [28] C. F. Bernasconi, *Adv. Phys. Org. Chem* **1992**, *27*, 119–238.

Manuscript received: March 17, 2021
Accepted manuscript online: May 18, 2021
Version of record online: June 30, 2021