



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

# Journal Pre-proof

Near Hanging: evaluation and management

Jon Dorfman

PII: S0012-3692(22)04065-X

DOI: <https://doi.org/10.1016/j.chest.2022.11.004>

Reference: CHEST 5364

To appear in: *CHEST*

Received Date: 29 October 2022

Accepted Date: 3 November 2022

Please cite this article as: Dorfman J, Near Hanging: evaluation and management, *CHEST* (2022), doi: <https://doi.org/10.1016/j.chest.2022.11.004>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Copyright © 2022 Published by Elsevier Inc under license from the American College of Chest Physicians.



Title: Near Hanging: evaluation and management

Abstract: 210 words

Text: 2947 words

Jon Dorfman  
Associate Professor of Surgery  
Department of Surgery  
Division of Trauma and Surgical Critical Care  
UMass Memorial  
55 Lake Avenue  
Worcester, MA 01655  
Jon.Dorfman@umassmemorial.org

No conflicts of interest exist.

Acknowledgments

Thank you to Timothy Emhoff, MD for manuscript review and insights.

Key Words: Suicide, Near hanging, Hangman's fracture, Blunt cerebrovascular injury, Laryngeal fracture

Abbreviations List: Blunt cerebrovascular injury (BCVI), Targeted Temperature Management (TTM)

## Abstract

Prior to the COVID-19 pandemic, the incidence of self-harm was already on the rise. Hanging/suffocation accounted for 50% of the increase in suicide attempts and remains the second leading cause of death from self-harm in the United States. Studies on the management of near hanging patients are lacking and most published literature is retrospective. Following airway and circulation assessment, clinical examination and imaging, namely CT angiography, remain the standard to identify the injuries associated with near hanging - cervical spine fracture, blunt cerebrovascular injury, laryngeal injury, injury to the trachea and oropharynx. These injuries, however, are uncommon, and each occur in less than 5% of patients in most series. In a large series of critically ill near hanging patients, over 50% survived to hospital discharge, however cardiac arrest predicted a poor outcome. The management of asphyxia related arrest remains controversial. Targeted Temperature Management (TTM) has only been studied in a single large multicenter trial which was retrospective. Given the significant selection bias of TTM treatment for the most ill patients, no firm recommendations can be made. Finally, for survivors, the underlying mental health issues must be addressed to avoid recurrent suicide attempts. Thirty percent of patients in a large near hanging series were admitted for their second suicide attempt.

## Introduction

Self-injury by hanging, otherwise described as “near hanging” if the patient survives to arrival at the emergency department is a relatively uncommon trauma mechanism. A retrospective 22 year review at 31 European medical centers included only 886 intensive care unit admissions.<sup>1</sup> At

one of the busiest trauma centers in the North America, 71 cases occurred over an 8-year period.<sup>2</sup> Near hanging is not a mechanism explicitly included in published trauma center triage<sup>3</sup> but may be included by institutional or regional level trauma triage criteria. However, this mechanism meets the criteria of other traumatically injured patients - urgent, potentially life threatening, multisystem injuries requiring an on-site team to make critical decisions and formulate a timely treatment plan which may involve multiple services in a critical care environment.

Suicide attempts and mortality have been increasing in the United States<sup>4</sup> with suicide surpassing motor vehicle crash mortalities by 2010.<sup>5</sup> In terms of modality, “hanging/suffocation” accounted for more than 50% of the increase in suicides in the period 2000-2010<sup>5</sup> and is the second leading mechanism of self-harm in the United States.<sup>6</sup> This phenomenon is not unique to the United States but may be seen worldwide.<sup>7</sup> Furthermore, suicidal ideation and mental illness have experienced a dramatic increase during the past two years of the COVID-19 pandemic. The increase in mental health issues and suicides has occurred in pediatrics<sup>8,9</sup> as well as adults<sup>10</sup>. Given the prevalence and increase of this problem, it is essential that all healthcare practitioners understand the evaluation and management of the near hanging patient population.

### Initial Assessment

Management of the near hanging patient begins with Advanced Trauma Life Support primary survey with assessment of Airway (A), Breathing (B) and Circulation (C). A patent airway can be rapidly assessed by asking the patient their name. Patients with poor mental status (Glasgow

Coma Scale score of less than 9), signs of airway compromise or injury should be intubated carefully with the concern that there could be direct airway injury (such as laryngeal fracture or tracheal trauma). Clinical examination findings that suggest airway injury would include a high-pitched voice, hoarse voice, inability to manage secretions as well as subcutaneous emphysema. In a series of near hanging patients, the risk of laryngeal injury, however, is less than 5%.<sup>11</sup> The risk of oropharyngeal is even lower. If intubation is required, it should occur with cervical spine precautions with the assumption that a cervical spine fracture and instability is present. Either the cervical collar should remain in place during the procedure, or a second practitioner should hold the head in neutral position. A glide scope may facilitate intubation. The intubating practitioner, however, should choose their most facile method for intubation but be prepared to change their approach if securing the airway is challenging, including intubation over a bougie or even a surgical airway.

Once the airway is assessed to be patent or is secured with a cuffed endotracheal tube, respiratory status and circulation are then evaluated. Adjunctive testing with chest wall ultrasound and chest radiograph can be used to determine the presence of a pneumothorax. Once the primary survey is completed, the secondary survey is performed. The secondary survey is a head-to-toe physical examination looking for other signs of injury. Findings on secondary survey in the near hanging patient may include ligature mark on the neck as well as petechial hemorrhages on the face.<sup>12</sup>

A history should be obtained from the patient or the prehospital ambulance crew to determine whether the near hanging was accidental or intentional. Intentional injury may be complex

suicide with more than one suicide modality utilized, for example overdose ingestion and hanging.<sup>13</sup> These suicide methods can be chosen simultaneously or one after the other if the first method was unsuccessful. (Please note this differs from complicated suicide which occurs due to unintended trauma from a suicide attempt that goes awry.) Complex suicide occurs in less than 5% of cases with most published literature case reports or small case series. Signs of penetrating trauma such as a stab wound should be sought on secondary survey by examining the torso and extremities. If awake and cooperative, the patient should be questioned about access to medications and whether any medications were taken in excess. A postmortem single city series of hanging victims revealed nearly 70% had positive serum toxicology for licit and illicit drugs.<sup>14</sup> With our toxicology service, we have developed our own institutional approach for the assessment of overdose. Our guidelines include obtaining a serum salicylate level, serum acetaminophen level and EKG to determine the QTc. Salicylate<sup>15</sup> and acetaminophen overdose<sup>16</sup> treatment is based on serum level. For many other medications including antipsychotics, the treatment of overdose is supportive care as well as bicarbonate for prolonged QTc.<sup>17</sup>

Additional work up for the hanging patient typically includes at a CT angiogram of the neck to determine the presence of associated injuries (Table 1) which will be addressed separately below.

#### Cervical spine fracture

Injury to the cervical spine is assumed to be present until proven otherwise, although surprisingly, fractures are uncommon in modern series.<sup>2</sup> A cervical collar should remain in place until the work up is completed. The prehospital extrication collar may remain in place but should be changed to a padded collar within hours to prevent skin breakdown. Even with a

padded collar, skin break down<sup>18,19</sup> can occur which would meet the Agency for Healthcare Research and Quality Patient Safety Indicator 3 for hospital acquired pressure ulcer.<sup>20</sup> Many different commercial padded collars exist with institutional preference determining which one is available for use. Previously, imaging could include either plain film radiography of the cervical spine or CT scans in the alert and stable patient.<sup>21</sup> However, plain film radiography has been shown to be inadequate in adults, with a sensitivity of only 50% for detection of cervical spine fractures.<sup>18</sup> For this reason as well as the desire to rule out blunt cerebrovascular injury (BCVI) with angiography, imaging of the cervical spine should be performed with CT scan. In the obtunded patient, controversy remains whether CT scan alone is adequate for cervical spine clearance and removal of the cervical collar.<sup>22</sup>

The classic cervical spine fracture is the Hangman's Fracture. The Hangman's Fracture pattern describes bilateral C2 pars interarticularis fractures, a pattern which account for up to 20% of all cervical spine fractures and occurs more frequently from other mechanisms such as motor vehicle crashes. A Hangman's Fracture is typically managed nonoperatively with a rigid orthosis cervical collar and cervical spine precautions. In a single center series, a review of 10 years of near hanging patients found 5% of patients sustained a cervical spine fracture. The type of spine fracture, however, was not described.<sup>11</sup> A retrospective multicenter trial reported 20 (2.9%) cervical spine injuries in 692 patients but also did not report the specific pattern of cervical spine fracture identified.<sup>24</sup>

The Hangman's Fracture obtained its name from judicial hangings which, due to the height and force at which they occur, lead to an extension injury- the head snaps back and the chin moves upward. Judicial hangings are considered complete hangings as the height from the fall exceeds



the patient's height. All the person's weight is transmitted to the neck as the feet remain suspended above the ground. Classically the knot of the rope must be under the chin to cause head extension.<sup>12</sup> Most suicide attempts are incomplete hangings with the height of the fall less than the height of the person and, therefore, do not occur with the same force. Also, the knot is often behind the person's head in suicide and not under the chin, and no head extension occurs. Death is usually by asphyxiation: loss of the airway and cerebral circulation with subsequent cardiac arrest. Patients who recover vital signs usually meet brain death criteria (which vary by the state and country).<sup>23</sup>

Cervical spine precautions entail wearing the cervical at all times- in bed when sleeping and even in the shower. The padding is changed with an assistant when soiled or wet. Please note that cervical spine precautions do not require bedrest or head of bed restrictions. A spine surgery consultation should be obtained for any cervical spine fracture that is identified. In the awake and cooperative patient, a neurological examination should be performed to assess for spinal cord injury which is reported by some series as occurring in 1% of patients.<sup>24</sup> Spinal shock maybe noted on physical examination. In patients with neurological deficits, this is the clinical loss of motor function, sensation and reflexes. For spine fractures that lead to a high spinal cord injury (above T6) as would occur from near hanging, neurogenic shock may also be present. Neurogenic shock is the loss of the sympathetic innervation. These patients have hypotension with inappropriate bradycardia (e.g., a systolic blood pressure of 60 mmHg with heart rate of 80 bpm) and have brisk capillary refill revealing a lack of vasomotor tone on clinical examination. After administration of up to 2 liters of Lactated Ringer's crystalloid, inotropes and/or vasopressors may be administered. Dopamine, norepinephrine or phenylephrine may be used if

there is no sign of hemorrhage. Phenylephrine has only alpha receptor effects and may improve vasoconstriction and subsequently the patient's blood pressure at the expense of worsening the bradycardia, however.

### Blunt Cerebrovascular Injury

Injury to the carotid or vertebral arteries of the neck is termed blunt cerebrovascular injury (BCVI). Many assume these injuries occur from direct vascular perforation from a bone fragment, however the pathophysiology relates to shear injury. Therefore, they can occur without an identified cervical spine fracture. Whom to screen for BCVI remains highly debated. Most recently some advocate for screening all blunt trauma patients.<sup>25</sup> Risk factors have been identified for BCVI, one of which is hanging with anoxic brain injury.<sup>26,27</sup> An argument for select screening of near hanging patient has also been recently made,<sup>28,29</sup> however, these are all small retrospective studies, and no strong conclusion can be formed. Arterial vascular injuries are graded on a scale of 1-5 with low grade injuries scoring a 1 and the most severe injuries scored as a 5. Low grade injuries involve the intima and high-grade injuries lead to occlusion or transection.<sup>30</sup> The feared complication from BCVI is stroke which can lead to significant disability or death within 24 hours of the traumatic event.<sup>30</sup> CT angiography is the accepted gold standard for diagnosis.<sup>25</sup> Management of BCVI also remains controversial. Fortunately, most injuries are low grade intimal injuries. Studies comparing aspirin, clopidogrel and anticoagulation<sup>31</sup> did not show a significant difference in outcome between these therapies. Therefore, aspirin is the most common treatment for low grade injuries given the risk of hemorrhage in many trauma patients. For higher grade injuries, interventional angiographic

procedures may be required. In a retrospective multicenter trial of near hanging patients, 6.1% of patients had BCVI detected.<sup>24</sup>

### Traumatic Brain Injury

In regards to including neuroimaging of the brain, CT scans may be performed at the same time as the cervical spine imaging. Findings at this early time point may include edema and loss of gray white matter differentiation as well as intracranial hemorrhage. In a large retrospective multicenter trial where MRI and brain CT findings were reported, only 3 of 886 had an intracranial hemorrhage.<sup>1</sup> Limited data suggests that CT imaging of the head does not change management, particularly if the patient has a normal mental status (Glasgow Coma Scale 15).<sup>33</sup> Less than 3% (4 patients) of normal mental status patients had any intracranial finding in their small series. Only one patient had a traumatic intracranial hemorrhage, but this patient had also been assaulted. In their review of the National Trauma Databank, a similar correlation was noted- isolated near hanging patients who had no additional traumatic mechanism and a normal mental status (GCS 15) had no intracranial injuries.<sup>33</sup>

### Laryngeal injury

Laryngeal injuries include fractures of the hyoid bone, thyroid, or cricoid cartilage. Hyoid bone and thyroid cartilage fractures overall are uncommon from trauma and therefore limited literature is available. Cricoid cartilage injuries are even less frequent. Symptoms may include changes in voice, dysphagia, pain with coughing or shortness of breath. Pain and crepitus may be noted on palpation.<sup>32,34</sup>

Laryngeal fracture occurs at variable rates in the near hanging literature, however, this mechanism places the patient at a higher risk than other traumatically injured patients. The range

of injury is from 5%<sup>11</sup> to nearly 40% of near hanging patients. The most common site of injury was the superior horn of the thyroid cartilage.<sup>35</sup> Unsurprisingly, autopsy series show a higher rate of injury, however, the published rate in autopsy series are just as variable. The hyoid and thyroid cartilage were fractured in 15 to 72.5 % and 9 to 43% of cases of hanging respectively.<sup>36,37,38</sup> . These published studies were based on CT scan or autopsy and include less than 200 patients each.

One proposed system to classify injuries is based on clinical symptoms and their extent, as well as diagnostic findings.<sup>39</sup> Hyoid fracture management is typically nonoperative. Nonoperative management may include dietary restrictions (liquid or soft diet), placement of a feeding tube and pain control. Operative strategies are not limited to open reduction and fixation to repair the hyoid but can include the removal of the injured portion of hyoid as well as tracheostomy if respiratory distress occurs.<sup>32</sup> Laryngeal fracture management is dependent upon the respiratory status as well. Operative management with tracheostomy was more common in a small series in which near hanging was not a common mechanism. All patients requiring a tracheostomy were eventually successful decannulated. All comminuted laryngeal fractures were repaired and most of these patients did not require a tracheostomy.<sup>34</sup>

#### Post Obstructive Pulmonary Edema

Post obstructive pulmonary edema occurs when patients breath against an occluded airway. Airway occlusion occurs from the noose and the patient creates significant negative intrathoracic pressure with their respiratory effort. Negative inspiratory pressures as high as -140 cm H<sub>2</sub>O have been measured.<sup>40</sup> While the pathophysiology has not been completely elucidated, fluid rapidly accumulates in the alveolar interstitial space and then within the alveolus.<sup>41</sup> The

diagnosis is made based on hypoxia, frothy secretions and chest radiograph opacification. The treatment is with oxygen supplementation, relieving the airway occlusion, diuresis to achieve a net negative fluid balance if the patient's cardiovascular status permits and positive pressure ventilation. Rapid resolution should occur within 48 hours.<sup>40</sup>

### Cardiac Arrest

The near hanging victim may experience cardiac arrest. The etiology is typically asphyxiation rather than cardiac in origin. Other etiologies may include high spinal cord injury with involvement of the medulla oblongata and pons, the respiratory centers of the brain. Cardiac arrest portends a poor outcome. A 5-year evaluation of 231 near hanging patients admitted to an ICU found 45% had had a cardiac arrest. Ninety five percent of the cardiac arrest patients did not survive to hospital discharge compared to a 1.6% mortality in the non-cardiac arrest group.<sup>42</sup> Similar findings were found in a large multicenter European study of ICU patients. Four hundred and fifty of 886 (50%) experienced a cardiac arrest. Ninety five percent of these patients did not survive to hospital discharge. In their study, cardiac arrest was the single largest predictor of mortality.<sup>1</sup> How to manage the arrest after return of spontaneous circulation occurs remains controversial. The literature on targeted temperature management (TTM) relates to cardiac arrest and not asphyxia.<sup>43,44</sup> Prospective studies specifically examining TTM for respiratory arrest patient are lacking. A large retrospective multicenter trial examined outcomes in near hanging patients with arrest. Asphyxia arrest patients who received TTM had worse outcomes (mortality, neurologic outcome) than patients who did not, however, the patients selected for TTM had higher severity of illness.<sup>24</sup> As this study was retrospective, the time from return of spontaneous circulation to TTM initiation was missing in majority of patients and there was variability in the target temperature chosen.<sup>24</sup> The variability in targeted temperature is not

surprising as the study included patients from 2005 to 2015 and the recommendation for TTM have changed over time.<sup>45</sup> In the previously discussed multicenter European study, TTM use increased over time from 23% in 1992 to 67% by 2014.<sup>1</sup> No strong recommendations exist for TTM in the near hanging patient and prospective studies will be required to determine which patients may benefit.

#### Psychiatric Evaluation

Not to be forgotten, treatment of the patient's underlying psychiatric condition is key. A pre-existing psychiatric diagnosis is present in up to 67% of near hanging patients.<sup>1,42</sup> In a study based in Northern Ireland, the risk of self-harm after a suicide attempt was 41%.<sup>46</sup> When examining the risk of successful second suicide by initial method, hanging, strangulation, suffocation is an important high risk factor. Over 50% of near hanging patients successfully committed suicide on their second attempt in a national Swedish study.<sup>47</sup> This study had up to a 31 year follow up period, however, most recurrent suicide attempts occurred within a year of the first attempt. The initial attempt occurred from 1972-1982 and hanging, strangulation, or suffocation accounted for only 1% of all suicide attempts. This study may not be applicable given the difference in suicide methods today. A more contemporary study, from 2000-2013, also found that the risk of suicide was highest in the year after the first episode of self-harm.

Near hanging was again noted as a high-risk factor.<sup>48</sup>

#### Conclusion

Near hanging is a mechanism many practitioners will encounter in their careers. The literature on this entity has limited published data which is mostly retrospective. Institutions should develop their own guidelines based on their available resources and best practices based on the literature to ensure an appropriate evaluation, identify cervical injuries, as well as other possible

concomitant suicide mechanisms. Psychiatric evaluation and treatment for survivors is essential to reduce the risk of another self-harm attempt.

#### Acknowledgments

Thank you to Timothy Emhoff, MD for manuscript review and insights.

#### References.

1. de Charentenay L, Schnell G, Pichon N, Schenck M, Cronier P, Perbet S et al. Outcomes in 886 critically ill patients after near-hanging injury. *Chest*. 2020;158(6): 2404-2413.
2. Schellenberg M, Inaba K, Warriner Z et al. Near hangings: epidemiology, injuries, and investigations. *Journal of trauma and acute care surgery*. 2019;86(3):454-7.
3. Sasser SM, Hunt RC, Faul M et al. Guidelines for field triage of injured patients: recommendations of the National Expert Panel on Field Triage, 2011. *Morbidity and Mortality Weekly Report: Recommendations and Reports*. 2012;61(1):1-20.
4. Wang J, Sumner SA, Simon TR et al. Trends in the incidence and lethality of suicidal acts in the United States, 2006 to 2015. *JAMA psychiatry*. 2020;77(7):684-93.
5. Baker SP, Hu G, Wilcox HC, Baker TD. Increase in suicide by hanging/suffocation in the US, 2000–2010. *American journal of preventive medicine*. 2013;44(2):146-9.
6. Centers for Disease Control and Prevention, National Center for Injury Prevention and Control. Web-based injury statistics query and reporting system (WISQARS), fatal injury data, leading causes of death. Available at: <http://www.cdc.gov/ncipc/wisqars>. Accessed July 22, 2022

7. Alqahtani S, Nehme Z, Williams B, Bernard S, Smith K. Temporal trends in the incidence, characteristics, and outcomes of hanging-related out-of-hospital cardiac arrest. *Prehospital Emergency Care*. 2020;24(3):369-77.
8. Charpignon M, Ontiveros J, Sundaresan S, et al. Evaluation of Suicides Among US Adolescents During the COVID-19 Pandemic. *JAMA Pediatr*. Published online April 25, 2022. doi:10.1001/jamapediatrics.2022.0515
9. McCluskey CK, Klein MJ, Steward SC, Rotta AT. Suicide and self-harm in children and adolescents admitted to PICUs in the United States. *Pediatric critical care medicine*. 2022;23(1):e66-70.
10. Czeisler MÉ, Lane RI, Petrosky E et al. Mental health, substance use, and suicidal ideation during the COVID-19 pandemic—United States, June 24–30, 2020. *Morbidity and Mortality Weekly Report*. 2020;69(32):1049.
11. Salim A, Martin M, Sangthong B, Brown C, Rhee P, Demetriades D. Near-hanging injuries: a 10-year experience. *Injury*. 2006;37(5):435-9.
12. Hua A, Shah KH, Garg M, Legome E, Ufberg J. A Hanging and Its Complications. *J Emerg Med*. 2016; 51(6):691-696.
13. Töro K, Pollak S. Complex suicide versus complicated suicide. *Forensic Sci Int*. 2009;184(1-3):6-9
14. San Nicolas AC, Lemos NP. Toxicology findings in cases of hanging in the City and County of San Francisco over the 3-year period from 2011 to 2013. *Forensic Sci Int*. 2015;255:146-55. .
15. Palmer BF, Clegg DJ. Salicylate Toxicity. *N Engl J Med*. 2020;382(26):2544-2555.
16. Gosselin S, Hoffman RS, Juurlink DN, Whyte I, Yarema M, Caro J. Treating acetaminophen overdose: thresholds, costs and uncertainties. *Clin Toxicol*. 2013;51(3):130-3.



17. Pourmand A, Mazer-Amirshahi M, Chistov S, Sabha Y, Vukomanovic D, Almulhim M. Emergency department approach to QTc prolongation. *Am J Emerg Med.* 2017; 35(12):1928-1933.
18. Como JJ, Diaz JJ, Dunham CM et al. Practice management guidelines for identification of cervical spine injuries following trauma: update from the eastern association for the surgery of trauma practice management guidelines committee. *J Trauma.* 2009;67(3):651-9.
19. Ham HW, Schoonhoven LL, Galer AA, Shortridge-Baggett LL. Cervical collar-related pressure ulcers in trauma patients in intensive care unit. *J Trauma Nurs.* 2014;21(3):94-102.
20. Calianno C. Pressure ulcers in acute care: a quality issue. *Nurs Manage.* 2007;38(5):42-51.
21. Stiell IG, Wells GA, Vandemheen KL et al. The Canadian C-spine rule for radiography in alert and stable trauma patients. *JAMA.* 2001;286(15):1841-8.
22. Patel MB, Humble SS, Cullinane DC et al. Cervical spine collar clearance in the obtunded adult blunt trauma patient: a systematic review and practice management guideline from the Eastern Association for the Surgery of Trauma. *J Trauma Acute Care Surg.* 2015;78(2):430-41
23. Greer DM, Shemie SD, Lewis A, et al. Determination of Brain Death/Death by Neurologic Criteria: The World Brain Death Project. *JAMA.* 2020;324(11):1078–1097.
24. Hsu CH, Haac BE, Drake M et al. EAST Multicenter Trial on targeted temperature management for hanging-induced cardiac arrest. *J Trauma Acute Care Surg.* 2018;85(1):37-47.
25. Leichtle SW, Banerjee D, Schrader R et al. Blunt cerebrovascular injury: The case for universal screening. *J Trauma Acute Care Surg.* 2020;89(5):880-886.
26. Biffi WL, Cothren CC, Moore EE et al. Western Trauma Association critical decisions in trauma: screening for and treatment of blunt cerebrovascular injuries. *J Trauma.* 2009;67(6):1150-3.

27. Bromberg WJ, Collier BC, Diebel LN, et al. Blunt cerebrovascular injury practice management guidelines: the Eastern Association for the Surgery of Trauma. *J Trauma*. 2010;68(2):471-7.
28. Berke DM, Helmer SD, Reyes J, Haan JM. Injury Patterns in Near-Hanging Patients: How Much Workup Is Really Needed? *Am Surg*. 2019 May 1;85(5):549-555.
29. Ribaute C, Darcourt J, Patsoura S et al. Should CT angiography of the supra-aortic arteries be performed systematically following attempted suicide by hanging? *J Neuroradiol*. 2021;48(4):271-276.
30. Biffi WL, Moore EE, Offner PJ, Brega KE, Franciose RJ, Burch JM. Blunt carotid arterial injuries: implications of a new grading scale. *J Trauma*. 1999;47(5):845-53.
31. Cothren CC, Biffi WL, Moore EE, Kashuk JL, Johnson JL. Treatment for blunt cerebrovascular injuries: equivalence of anticoagulation and antiplatelet agents. *Arch Surg*. 2009;144(7):685-90.
32. Ramchand T, Choudhry OJ, Shukla PA, Tomovic S, Kuperan AB, Eloy JA. Management of hyoid bone fractures: a systematic review. *Otolaryngol Head Neck Surg*. 2012;147(2):204-8.
33. Bordia R, Freeman C, Kou H, Culhane J. Do we need neuroimaging in every case of near-hanging?: experience from a level 1 trauma center and analysis of the National Trauma Data Bank. *Emergency radiology*. 2022; 29(1): 49-57.
34. Jalisi S, Zoccoli M. Management of laryngeal fractures--a 10-year experience. *J Voice*. 2011;25(4):473-9.
35. Fukumoto W, Mitani H, Kuno Y et al. Incidence and factor analysis of laryngo-hyoid fractures in hanging individuals-computed tomography study. *Eur Radiol*. 2021;31(10):7827-7833.

36. Sharma BR, Harish D, Sharma A, Sharma S, Singh H. Injuries to neck structures in deaths due to constriction of neck, with a special reference to hanging. *J Forensic Leg Med*. 2008;15(5):298-305.
37. Nikolic S, Micic J, Atanasijevic T, Djokic V, Djonic D. Analysis of neck injuries in hanging. *Am J Forensic Med Pathol*. 2003;24(2):179-82.
38. Zátopková L, Janík M, Urbanová P, Mottlová J, Hejna P. Laryngo-hyoid fractures in suicidal hanging: A prospective autopsy study with an updated review and critical appraisal. *Forensic Sci Int*. 2018;290:70-84.
39. Bell RB, Verschueren DS, Dierks EJ. Management of laryngeal trauma. *Oral Maxillofac Surg Clin North Am*. 2008;20(3):415-30.
40. Bhattacharya M, Kallet RH, Ware LB, Matthay MA. Negative-Pressure Pulmonary Edema. *Chest*. 2016;150(4):927-933.
41. Lemyze, M, and Mallat J. Understanding negative pressure pulmonary edema. *Intensive care medicine*. 2014;40(8): 1140-1143.
42. Gantois G, Parmentier-Decrucq E, Duburcq T, Favory R, Mathieu D, Poissy J. Prognosis at 6 and 12 months after self-attempted hanging. *The American Journal of Emergency Medicine*. 2017; 35(11):1672-1676.
43. Dankiewicz J, Cronberg T, Lilja G et al. Targeted hypothermia versus targeted Normothermia after out-of-hospital cardiac arrest (TTM2): A randomized clinical trial-Rationale and design. *Am Heart J*. 2019;217:23-31
44. Granfeldt A, Holmberg MJ, Nolan JP, Soar J, Andersen LW; International Liaison Committee on Resuscitation (ILCOR) Advanced Life Support Task Force. Targeted temperature

management in adult cardiac arrest: Systematic review and meta-analysis. *Resuscitation*. 2021;167:160-172.

45. Nielsen N, Wetterslev J, Cronberg T et al; TTM Trial Investigators. Targeted temperature management at 33°C versus 36°C after cardiac arrest. *N Engl J Med*. 2013;369(23):2197-206.

46. Griffin E, Bonner B, O'Hagan D, Kavalidou K, Corcoran P. Hospital-presenting self-harm and ideation: Comparison of incidence, profile and risk of repetition. *General hospital psychiatry*. 2019; 61: 76-81.

47. Runeson B, Tidemalm D, Dahlin M, Lichtenstein P, Långström N. Method of attempted suicide as predictor of subsequent successful suicide: national long term cohort study. *BMJ* 2010;340:c3222.

48. Geulayov G, Casey D, Bale L, Brand F, Clements C, Farooq B et al. Suicide following presentation to hospital for non-fatal self-harm in the Multicentre Study of Self-harm: a long-term follow-up study. *The Lancet Psychiatry*. 2019; 6(12): 1021-1030.

Table 1. Injuries from near hanging <sup>1, 10, 22, 26</sup>

Injury	
Anoxic brain injury	8% - 35.7%
Cervical spine fracture	2% - 6%
Laryngeal fracture	1.4% - 5.3%
Tracheal injury	Up to 1.6%
Pharyngeal laceration	Up to 1.6%
Blunt cerebrovascular injury	1.6% - 6.1%

Injuries and the rates of occurrence from near hanging.

**Declaration of interests**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Journal Pre-proof