



STUDY PROTOCOL

REVISED

Facilitators and barriers to stakeholder engagement in advance care planning for older adults in community settings: a hybrid systematic review protocol [version 2; peer review: 2 approved]

Monika Pilch ¹, Victoria Lunt ², Peter May ^{1,3}, David Mockler ⁴, Stephen Thomas ¹, Frank Doyle ⁵

¹Centre for Health Policy and Management, School of Medicine, Trinity College Dublin, University of Dublin, Dublin, D2, Ireland

²Beaumont Hospital and St Luke's Radiation Oncology Centre at Beaumont Hospital, Beaumont Hospital, Beaumont, Dublin, D9, Ireland

³The Irish Longitudinal study on Ageing (TILDA), Trinity College Dublin, Dublin, D2, Ireland

⁴The Library of Trinity College Dublin, Trinity College Dublin, University of Dublin, Dublin, D2, Ireland

⁵Division of Population Health Sciences, Royal College of Surgeons in Ireland, Dublin, D2, Ireland

v2 First published: 23 Jun 2020, 3:38
<https://doi.org/10.12688/hrbopenres.13082.1>

Latest published: 11 Jun 2021, 3:38
<https://doi.org/10.12688/hrbopenres.13082.2>

Abstract

Background: Poor stakeholder engagement in advance care planning (ACP) poses national and international challenges, preventing maximisation of its potential benefits. Conceptualisation of advance care planning as a health behaviour highlights the need to design innovative, evidence-based strategies that will facilitate meaningful end-of-life care decision-making.

Aim: To review systematically and synthesise quantitative and qualitative evidence on barriers and facilitators to stakeholders' engagement in ACP for older adults (≥ 50 years old) in a community setting.

Methods: A hybrid systematic review will be conducted, identifying studies for consideration in two phases. First, databases will be searched from inception to identify relevant prior systematic reviews, and assess all studies included in those reviews against eligibility criteria (Phase 1). Second, databases will be searched systematically for individual studies falling outside the timeframe of those reviews (Phase 2). A modified SPIDER framework informed eligibility criteria. A study will be considered if it (a) included relevant adult stakeholders; (b) explored engagement in ACP among older adults (≥ 50 years old); (c) employed any type of design; (d) identified enablers and/or barriers to events specified in the Organising Framework of ACP Outcomes; (e) used either quantitative, qualitative, or mixed methods methodology; and (f) evaluated phenomena of interest in a community setting (e.g.,

Open Peer Review

Reviewer Status

	Invited Reviewers	
	1	2
version 2 (revision) 11 Jun 2021		 report
version 1 23 Jun 2020	 report	 report

1. **Jessica E. Simon** , University of Calgary, Calgary, Canada

2. **Susan Fiona Lee** , Monash University, Frankston, Australia

Any reports and responses or comments on the article can be found at the end of the article.

primary care or community healthcare centres). Screening, selection, bias assessment, and data extraction will be completed independently by two reviewers. Integrated methodologies will be employed and quantitative and qualitative data will be combined into a single mixed method synthesis. The Behaviour Change Wheel will be used as an overarching analytical framework and to facilitate interpretation of findings. The Joanna Briggs Institute (JBI) Reviewers' Manual and PRISMA-P guidelines have been used to inform this protocol development.

Registration: This protocol has been submitted for registration on PROSPERO, registration number CRD42020189568 and is awaiting review.

Keywords

Advance care planning, community, behaviour change, systematic review



This article is included in the [TILDA](#) gateway.

Corresponding author: Monika Pilch (pilchm@tcd.ie)

Author roles: **Pilch M:** Conceptualization, Data Curation, Formal Analysis, Funding Acquisition, Investigation, Methodology, Project Administration, Writing – Original Draft Preparation, Writing – Review & Editing; **Lunt V:** Data Curation, Formal Analysis, Investigation, Writing – Review & Editing; **May P:** Conceptualization, Investigation, Methodology, Supervision, Writing – Review & Editing; **Mockler D:** Data Curation, Investigation, Methodology, Software, Writing – Review & Editing; **Thomas S:** Conceptualization, Funding Acquisition, Investigation, Resources, Supervision, Writing – Review & Editing; **Doyle F:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Supervision, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: Health Research Board [SPHeRE-2018-1]. The Health Research Board (HRB) supports excellent research that improves people's health, patient care and health service delivery. HRB aims to ensure that new knowledge is created and then used in policy and practice. In doing so, HRB supports health system innovation and create new enterprise opportunities.
The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2021 Pilch M *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Pilch M, Lunt V, May P *et al.* **Facilitators and barriers to stakeholder engagement in advance care planning for older adults in community settings: a hybrid systematic review protocol [version 2; peer review: 2 approved]** HRB Open Research 2021, 3:38 <https://doi.org/10.12688/hrbopenres.13082.2>

First published: 23 Jun 2020, 3:38 <https://doi.org/10.12688/hrbopenres.13082.1>

REVISED Amendments from Version 1

We clarified that few reviews have synthesised evidence to identify patterns across older adults, irrespective of their health status or condition, while taking stakeholder perspective and level of influence into account (p. 4). We specified that “grey literature” will not be considered for inclusion in this review (p. 6). We indicated that if a study evaluated an intervention, we will also extract the name of the strategy and identify COM-B factors addressed in that intervention” (p. 8). We specified a timeframe for the completion of Phase 1 of the search (June 2020) (p. 6). We narrowed down the number of items to be extracted from eligible systematic reviews (p. 8). Minor spelling and grammar corrections were made on pages 3, 4, and 8.

Any further responses from the reviewers can be found at the end of the article

Introduction

People living and dying with serious illness are a societal and policy priority worldwide (Centeno & Arias-Casais, 2019). As prognosis worsens and the end-of-life phase approaches, many people with terminal illness continue to receive aggressive, high-intensity treatment; e.g., chemotherapy in advanced cancer, mechanical ventilation in respiratory failure, and dialysis in chronic kidney disease. These treatments are often instigated and continued without checking patient preferences (Kelley *et al.*, 2010), and as a result of poor physician understanding of those preferences (Downey *et al.*, 2013). This may result in failure to address quality of life domains, such as pain and symptom management, or psychosocial and spiritual supports; often with limited survival benefit (Earle *et al.*, 2008). They may result in longer hospital stays at a time when many people prefer to be at home or in hospice, and this time in hospital reduces remaining quality of life and time with loved ones (Davison, 2010; Gomes *et al.*, 2012).

Advance care planning (ACP) provides mechanisms through which individual preferences can be meaningfully communicated and supported. A broad definition of ACP indicates that communication and decision-making processes regarding future healthcare wishes should include identification of patients’ goals of care, values, preferences, and priorities; and not be limited to mere completion of advance care directives. It also emphasises the importance of involving all relevant stakeholders and all domains of care; including biological, psychological, social, and spiritual (Rietjens *et al.*, 2017). Although older people who engaged in ACP are less likely to be admitted to intensive care units (ICUs), acute care settings (e.g., ICUs or emergency departments) are not ideal contexts for the initiation of end-of-life conversations or decision-making (Khandelwal *et al.*, 2015). Similarly, individuals presenting to an emergency department from residential aged care settings are more likely to have an advance care plan than community dwellers (Street *et al.*, 2014). Therefore, opportunities to engage individuals in ACP at an early stage and facilitate ongoing conversations regarding future healthcare needs in a community, non-residential setting (e.g., primary care or community healthcare centres) needs to be explored to bridge this gap.

ACP is associated with increased quality of life (Garrido *et al.*, 2015); a reduction in unwanted admissions to hospitals; care consistent with patients’ goals (Jimenez *et al.*, 2018); improved quality of end-of-life care, increased use of hospice services and reduced hospital deaths (Bischoff *et al.*, 2013); compliance with patients’ end-of-life wishes (Brinkman-Stoppelenburg *et al.*, 2014; Silveira *et al.*, 2010); decreased caregiver burden (Balein, 2009; Earle *et al.*, 2008); better bereavement processes and more positive psychological outcomes for family members (Wright *et al.*, 2008); reduced health care spending (Gidwani-Marszowski *et al.*, 2019; Zhang *et al.*, 2009); and cost-effective strategies for facilitating patients’ choice (Nguyen *et al.*, 2017). However, low engagement in ACP poses national and international challenges, preventing maximisation of these potential benefits.

Community-based interventions have the potential to overcome many of the barriers to ACP, to include the introduction of a more person-centered, holistic, inclusive, population-based, and integrated care approach. The biggest challenge to ACP implementation is the identification of the right time to initiate these sensitive end-of-life conversations. Diagnosis is often associated with increased psychological distress, which might impact patients’ readiness, mental ability to comprehend novel medical concepts, and decision-making capacity (Pinquart & Duberstein, 2010). Although clinical guidelines specify that ACP should commence when there is evidence of a life-limiting advanced progressive illness (Clayton *et al.*, 2007), research findings and clinical practice suggest that this might be too late (Jimenez *et al.*, 2018). The preparation for end-of-life conversations is a process that could begin early in the life course, e.g., when individuals reach their 50th year of life, and irrespective of health status (Howard *et al.*, 2018). As this age group is at an increased risk of presenting with a chronic illness (Mullaney *et al.*, 2016), end-of-life issues are likely to be personally relevant to them (Simon *et al.*, 2015). As readiness for ACP engagement is linked to the concept of preparedness, the interventions introduced at an *early* stage, in a community, could prepare individuals for patient-healthcare practitioner end-of-life conversations in specialised palliative care settings, and take place when individuals are ageing but still healthy (Lewis *et al.*, 2016).

Irrespective of supportive legal frameworks and relevant policies (Kelly, 2017), patient engagement in ACP is poor (Brinkman-Stoppelenburg *et al.*, 2014; Jimenez *et al.*, 2018). Effectiveness studies suggest that an *early* and *phased process* of preparation for the last stage of life, incorporating: 1) longitudinal and repeated ACP discussions (Prendergast, 2001); 2) customised to patients’ needs, circumstances, and readiness (Fried *et al.*, 2009; Simon *et al.*, 2015); 3) involving digital decision aids (Austin *et al.*, 2015; Ostherr *et al.*, 2016); 4) targeting multiple stakeholders; and 5) taking context into account (Jimenez *et al.*, 2018) would increase *informed* and *meaningful* participation in ACP, as well as optimise end-of-life outcomes.

There is a growing recognition of engaging in ACP as a *health behaviour* (Fried *et al.*, 2009; Fried *et al.*, 2010; Sudore *et al.*, 2008). This behavioural conceptualisation allows a view of ACP

as subject to health behaviour change interventions (Fried *et al.*, 2012; Fried *et al.*, 2016; Fried *et al.*, 2018; Sudore *et al.*, 2017; Van Scoy *et al.*, 2017). To date, most approaches to behaviour change have focused on individuals, and although this level is crucial when highly sensitive ACP conversations are considered, approaches that ignore contextual influences are open to criticism (Lin *et al.*, 2019). The Organising Framework of ACP Outcomes suggests that readiness for ACP engagement is a complex construct, which needs to be systematically assessed with regards to all relevant stakeholders. Therefore, a comprehensive systematic synthesis of findings, across all aspects of the framework, is essential to informing efforts in behaviour change. This proposed conceptualisation of ACP supports the use of the Behaviour Change Wheel (BCW) framework (Michie *et al.*, 2011), which is an encapsulating model, and increasingly employed in ACP empirical investigations (e.g., Biondo *et al.*, 2019; Peck *et al.*, 2018; Tam-Tham *et al.*, 2016).

Given the need to increase meaningful ACP engagement, the design of innovative interventions; targeting complex interactions between facilitators, barriers, and stakeholders, is required. There is a consensus that the development of ACP interventions should be based on a careful and critical synthesis of available evidence and the understanding of barriers and enablers to engagement (Risk *et al.*, 2019). However, existing systematic reviews tend to focus on acute and specialised contexts (Gilissen *et al.*, 2019), one group of stakeholders (De Vleminck *et al.*, 2013), and/or specific clinical conditions (Tilburgs *et al.*, 2018; Van Der Steen *et al.*, 2014). Although several studies have been conducted to explore factors associated with the uptake of ACP among older adults in community settings; and these have been synthesised in previous systematic reviews (e.g., De Vleminck *et al.*, 2013; Ramsaroop *et al.*, 2007; Risk *et al.*, 2019; Solis *et al.*, 2018), few studies have synthesised available evidence to identify patterns across older adults, irrespective of their health status or condition, while taking stakeholder perspective and level of influence into account. A synthesis utilising the BCW framework is necessary to identify influences within the contextual layers of health behaviour among older adults living in a community, while highlighting individual, interpersonal, provider, and system levels of influence.

Objectives and review questions

The overarching aim of the study will be to systematically review and synthesise quantitative and qualitative evidence on barriers and facilitators to stakeholders' engagement in ACP for older adults (≥ 50 years old) in a community setting.

The synthesis will be guided by the following review questions (RQs):

- RQ1: What individual-level, service-level, and system-level factors facilitate and/or impede stakeholders' ACP engagement for older adults in a community setting?
- RQ2: When categorised into the COM-B ('capability', 'opportunity', 'motivation' dimensions of behaviour) model, which of the identified barriers and facilitators are grouped into the categories of (a) Capability, (b) Motivation, and (c) Opportunity?

Methods

Design

As several systematic reviews have been published in this area, a hybrid systematic review will be completed, adopting the methodology of Doyle *et al.* (2019). In a hybrid review approach, potentially relevant studies are searched for in two phases (Doyle *et al.*, 2019). First, a systematic search for systematic reviews will be completed and eligible primary studies included in those reviews will be identified and extracted (Phase 1). Second, this process will be supplemented with an updated systematic review of more recently published individual studies (Phase 2). When available, relevant data will be extracted from the eligible systematic reviews. If not available, two reviewers will independently extract data from the primary studies.

While no prior systematic review has examined the specific research questions, multiple prior reviews have addressed questions that partially overlap. The advantage of the hybrid method is, therefore, that it leverages the efforts of multiple prior studies and minimises duplication of effort with those same teams. Derived results will nevertheless represent a distinctive contribution to knowledge in specifically addressing, for the first time, the barriers and facilitators for ACP for older people (≥ 50 years old) in a community setting; which is important for the reasons detailed elsewhere in this protocol.

A mixed-method approach will be used and quantitative, qualitative, and mixed-methods reviews will be included. The BCW will be used as an overarching analytical framework (Michie *et al.*, 2011). The Joanna Briggs Institute (JBI) Reviewers' Manual (Lizarondo *et al.*, 2019) and PRISMA-P guidelines (Moher *et al.*, 2015) have been used to inform this protocol development (see *Reporting guidelines*, Pilch, 2020). This protocol has been submitted for registration on PROSPERO, registration number: CRD42020189568.

Eligibility criteria

The modified SPIDER reporting framework (Cooke *et al.*, 2012) informed eligibility criteria, which were specified for the two stages of the search: the search for systematic reviews (Phase 1) and the supplementary search for primary studies (Phase 2). A "setting" ("s") eligibility criterion was added to incorporate a contextual variable (resulting in SPIDER"s" framework).

SPIDERS reporting framework

A review or a primary study will be deemed eligible if it: (a) included adult stakeholders (relevant older adults ≥ 50 years old, their significant others, healthcare professionals, non-medical peers, service managers, and/or policy makers) (**Sample**); (b) explored engagement in ACP among older adults (≥ 50 years old) (**Phenomena of Interest**); (c) employed any type of design (**Design**); (d) identified enablers and/or barriers to events specified in the Organising Framework of ACP Outcomes (Sudore *et al.*, 2018) (**Evaluation**); (e) used either quantitative, qualitative, or mixed methods methodology (**Research Type**); and (f) evaluated phenomena of interest (or included primary studies that aimed to do so) in a community setting (**Setting**). Only reviews and primary studies reported in English and published in peer-reviewed journals will be

considered for inclusion. The necessary differences in the approach taken when identifying systematic reviews (Phase 1) or the supplementary primary studies (Phase 2), are presented in [Table 1](#). The specification of the inclusion and exclusion criteria is presented below.

Sample. In the context of this review, the eligible population includes adult stakeholders who have the potential to influence ACP engagement among older adults in a community setting. They include older adults themselves, their significant others, healthcare professionals, non-medical peers, service managers,

and/or policy makers. A study will be deemed eligible if it collected data related to the population of interest; adults in community settings, aged ≥ 50 years old. This population has been chosen as this age group is at increased risk of presenting with a chronic illness, therefore, end-of-life issues are likely more relevant to them ([Mullaney et al., 2016](#)). There will be no limits on the health status and/or type of a condition or diagnosis as capturing the diversity of perspectives of various groups and the identification of common patterns across diagnoses, as well as healthy participants, is an ultimate goal of this review.

Table 1. Inclusion and exclusion criteria.

SPIDER(s)	Search for Systematic Reviews (Phase 1)	Supplementary Search for Primary Studies (Phase 2)
Sample	<p>Stakeholder: Relevant older adults, their significant others, healthcare professionals, non-medical peers, service managers, and/or policy makers.</p> <ul style="list-style-type: none"> Population of interest: those in a community setting and >50 years old. 	<p>Stakeholder: Relevant older adults, their significant others, healthcare professionals, non-medical peers, service managers, and/or policy makers.</p> <ul style="list-style-type: none"> Population of interest: those in the community setting and >50 years old.
Phenomena of Interest	<p>Events or actions associated with the stakeholders' engagement, uptake, utilisation, implementation, and/or initiation of ACP (e.g., communication, decision making, documentation completion, etc.), as informed by the Organising Framework of ACP Outcomes (Sudore et al., 2018).</p>	<p>Events or actions associated with the stakeholders' engagement, uptake, utilisation, implementation, and/or initiation of ACP (e.g., communication, decision making, documentation completion, etc.), as informed by the Organising Framework of ACP Outcomes (Sudore et al., 2018).</p>
Design	<p>Systematic reviews that employed all types of review methodology.</p> <p>Exclusion: protocols, scoping reviews, narrative reviews, overviews of reviews, editorials, comments, and expert opinion.</p>	<p>Empirical studies</p> <ul style="list-style-type: none"> Descriptive studies as well as experimental and observational study designs that explored the link between predictor, mediator, and/or moderator variables and relevant ACP events and outcomes. All qualitative approaches and designs.
Evaluation	<p>Facilitators and/or barriers to events or actions specified in the Organising Framework of ACP Outcomes; and identified at the level of an individual, a service, and/or a system.</p> <ul style="list-style-type: none"> A barrier: any factor that has been identified as an obstacle, impediment, deterrent, hindrance, or difficulty in a stakeholder's engagement, uptake, utilisation, implementation, and or initiation in ACP activities/events. A facilitator: any enabler or factor that has been shown to increase the chances of a stakeholder's engagement, uptake, utilisation, implementation, and or initiation in ACP activities/events. 	<p>Facilitators and/or barriers to events or actions specified in the Organising Framework of ACP Outcomes; and identified at the level of an individual, a service, and/or a system.</p> <ul style="list-style-type: none"> A barrier: any factor that has been identified as an obstacle, impediment, deterrent, hindrance, or difficulty in a stakeholder's engagement, uptake, utilisation, implementation, and or initiation in ACP activities/events. A facilitator: any enabler or factor that has been shown to increase the chances of a stakeholder's engagement, uptake, utilisation, implementation, and or initiation in ACP activities/events.
Research Type	All types of quantitative, qualitative, and mixed-methods review methodologies will be included in the review.	Quantitative, qualitative, and mixed-method approaches employed in the primary studies will be considered for inclusion.
Setting	<p>Reviews that included studies that meet the inclusion criteria for the primary studies.</p> <p>If evidence from multiple settings were included, a review will be included only if data relating to community settings can be extracted separately.</p> <p>Exclusion: Studies conducted in the hospital (e.g., ICU or an emergency department), ambulatory setting, outpatient or hospital clinics, and/or long term care facilities (e.g., residential care, to include a nursing home or a hospice) will be excluded.</p>	<p>Studies conducted in, or related to, all kind of community-based centres and/or primary care. Eligibility determined by the setting of receiving care. Studies from both rural and urban settings will be included.</p> <p>Exclusion: Studies conducted in the hospital (e.g., ICU or an emergency department), ambulatory setting, outpatient or hospital clinics, and/or long term care facilities (e.g., residential care, to include a nursing home or a hospice) will be excluded.</p>

ACP, advance care planning; ICU, intensive care unit.

Phenomena of interest. The focus of this review will be on engagement (or not) in ACP among older adults in a community setting. The conceptualisation of the phenomena of interest was informed by the Organising Framework of ACP Outcomes (Sudore *et al.*, 2018) and involves events or actions associated with the stakeholders' engagement, uptake, utilisation, implementation, and/or initiation of ACP. Therefore, in the context of this review, the relevant events and/or outcomes include, but are not limited to: (a) documentation of values, goals, and/or preferences (e.g., completion of the living will and/or advance directives); (b) the choice and/or documentation of a decision-maker; (c) communications about the goals of care, to include quality vs quantity of life (with family and/or health-care providers); and (d) communications about life-sustaining treatments (with family and/or health-care providers).

Design. Empirical studies that employed all types of methodology will be included in this review. In the first stage of the search (Phase 1), only systematic reviews will be selected and protocols, scoping reviews, narrative reviews, overviews of reviews, editorials, comments, and expert opinion will be excluded. When selecting primary studies, descriptive, experimental, and observational study designs that explored links between variables identified as facilitators or barriers and relevant ACP events/outcomes will be considered for the quantitative component of the review. An inclusive approach will be taken and the qualitative component of the review will not be limited by a specific research design.

Evaluation. Facilitators and/or barriers to events specified in the Organising Framework of ACP Outcomes (Sudore *et al.*, 2018) will be the core evaluation in the context of this study. Therefore, reviews and/or primary studies exploring individual, service, and system level factors that facilitated or impeded stakeholders' engagement, uptake, utilisation, implementation, and/or initiation in ACP activities/events (communication, decision making, documentation completion, etc.) will be included. For the purposes of this review, a barrier has been broadly defined as any factor that has been identified as an obstacle, impediment, deterrent, hindrance, or difficulty. A facilitator is defined as any kind of an enabler or factor that has been shown to increase the chances of ACP engagement. Following the model proposed by Sudore *et al.* (2018), studies that explored relevant factors in relation to individuals (older adults, significant others, healthcare professional, service managers, and/or policy makers), communities (public health, community incentives, legal support, policy and media), and healthcare system (documentation, training, facilitators, and palliative care) will be included.

When selecting primary studies, those that explored predictors, moderators, and/or mediators of ACP engagement will be included in the quantitative component of the review. The quantitative component will also consider studies that identified factors (e.g., active ingredients) contributing to the effectiveness of interventions, tools, and/or strategies designed to increase engagement in ACP activities or events. The qualitative component of the review will consider studies that explored

perspectives, views, opinions, and/or experiences of relevant stakeholders in relation to the phenomena of interest.

Research type. All types of quantitative, qualitative, and mixed-methods review methodologies will be included in the first stage of the review (Phase 1). Quantitative, qualitative, and mixed-method approaches employed in the primary studies will be considered for inclusion (Phase 1 & 2). Although we acknowledge the importance of grey literature, due to envisaged amount of existing evidence on the topic, grey literature will not be considered in this review.

Setting. A review will be eligible only if it included studies conducted in community, non-residential settings. In the selection of primary studies, the setting of receiving care will determine whether a study is eligible. Therefore, a study will be included if it recruited patients from all kind of community-based centres and/or primary care. Studies that took place in the hospital (e.g., ICU or an emergency department), outpatient or ambulatory settings, hospital clinics, and/or long term care facilities (e.g., residential care, to include a nursing home or a hospice) will be excluded. This decision has been made as the aim is to explore opportunities to facilitate the initiation of end-of-life conversations at an early stage, when individuals are relatively well. Although it is acknowledged that hospital-based services may have this function for some individuals, and in certain contexts, the aim of this synthesis is to enhance an understanding of patterns of ACP engagement outside of long-term and acute care environments.

Search strategy

As a hybrid systematic review will be conducted (Doyle *et al.*, 2019), a two-stage search strategy has been developed in collaboration with a medical librarian (DM). A comprehensive search for systematic reviews will be conducted in each database from its inception (Phase 1), and this process will be completed in June 2020. The eligible reviews will be used to identify and extract relevant primary studies. This data will be supplemented by an updated search for more recent studies; falling outside the timeframe of those reviews (Phase 2). The search strategy will also include screening of relevant overviews of reviews (in Phase 1), "backward and forward" citation search (Phase 1 & 2), and reference list screening (Phase 1 & 2).

Electronic searches. All searches will be completed by the first reviewer (MP). The search for systematic reviews (Phase 1) will include comprehensive searching of electronic databases from inception, and will include MEDLINE (Ovid), EMBASE, PsycINFO, the Cochrane Library (CENTRAL), CINAHL, and Epistemonikos. A sample search strategy for EMBASE in Phase 1 of the search is included in Appendix B (see *Extended data*, (Pilch, 2020)).

It is estimated that the time limit for the supplementary identification of primary studies (Phase 2) will be the last three years; however, it will be determined and specified after completing Phase 1 of the search. The supplementary search will include

comprehensive searching of the same databases, with the addition of the Web of Science. Epistemionikos will not be searched in Phase 2 as it is likely to identify systematic reviews and primary studies already included in the identified reviews (El-Khayat, 2017). A sample search strategy for EMBASE in Phase 2 of the search is included in Appendix C (see *Extended data*, (Pilch, 2020)). The search strategy will be adapted to each database interface.

The search terms used are specified in Table 2. In searching for systematic reviews, terms relating to the phenomena of interest, design, evaluation, and setting will be utilised in the database searches, combined with ‘AND’. In the supplementary search for primary studies, the search terms relating to the design category will not be used.

Screening and selection of primary studies

The two-stage process of primary studies identification is illustrated in Figure 1. Upon completion of the first stage of the search (Phase 1), all references will be downloaded into Endnote. Duplicates will be removed using the automatic function of the software. Then, the references will be exported to Covidence. Titles and abstracts will be independently screened by the first and second reviewer (MP & VL) and those clearly not relevant to the review will be eliminated. Full-text copies of reports will be retrieved for all papers passing title/abstract review. Full texts will be screened by MP and VL and those clearly not relevant to the review will be eliminated. Conflicts will be resolved through discussion and consensus with other reviewers (PM, FD, or ST). A database of eligible *systematic reviews* will be created at the end of this process.

The eligible primary studies will be independently extracted from the selected systematic reviews by MP and VL. Eligibility of primary studies will be determined by screening the summary tables included in the systematic reviews (if available) or by reviewing the full report (if summary tables are not available or do not include relevant information). A database of the eligible primary studies will be created at this stage (Database A in Figure 1).

Upon completion of the second stage of the search (Phase 2), all references will be downloaded into Endnote and duplicates removed. The systematic processes described above will be applied to the supplementary search and will include: screening of titles and abstracts and elimination of non-relevant studies (MP & VL); retrieval of full-text copies of reports for all papers passing title/abstract review (MP & VL); screening of the full-texts and elimination of those clearly not relevant to the review (MP & VL). Conflicts will be resolved through discussion and consensus with other reviewers (PM, FD, or ST). A database of primary studies identified during the supplementary search will be created at the end of this process (Database B in Figure 1). Both databases (Database A & Database B) will be merged.

Multiple reports of the same study, if identified, will be treated as a single study. The details of both screening and selection processes (one for systematic reviews and one for primary studies), with the indication of included and excluded studies at each stage, will be documented and demonstrated on the PRISMA flow diagrams. Reasons for rejection at the full-text stage will be recorded.

Table 2. Search Terms.

	Search Terms
Phenomena of Interest	‘advance* care plan*’ OR ‘patient advance* care plan*’ OR ‘advance* health care plan*’ OR ‘advance* healthcare plan*’ OR ‘anticipat* care’ OR ‘anticipat* healthcare’ OR ‘anticipat* health care’ OR ‘advance* directive*’ OR ‘healthcare directive*’ OR ‘advance* care statement*’ OR ‘living will’ OR ‘resuscitation order*’ OR ‘do not resuscitate’ OR ‘DNR’ OR ‘DNAR’ OR ‘ADRT’ OR ‘right to die’ OR ‘right-to-die’ OR ‘do not hospitali?e’ OR ‘prefer* place* of care’ OR ‘prefer* place* of death’ OR ‘end of life prefer*’ OR ‘end-of-life prefer*’ OR ‘EOL prefer*’ OR ‘end of life plan’ OR ‘end-of-life plan’ OR ‘EOL plan’ OR ‘end of life choice’ OR ‘end-of-life choice’ OR ‘EOL choice’ OR ‘end of life decision*’ OR ‘end-of-life decision*’ OR ‘EOL decision*’ OR ‘end of life communication*’ OR ‘end-of-life communication*’ OR ‘EOL communication*’ OR ‘end of life conversation*’ OR ‘end-of-life conversation*’ OR ‘EOL conversation*’ OR ‘end of life care goal*’ OR ‘end-of-life care goal*’ OR ‘EOL care goal*’ OR ‘end of life document*’ OR ‘end-of-life document*’ OR ‘EOL document*’ OR ‘end of life directive*’ OR ‘end-of-life directive*’ OR ‘EOL directive*’ OR ‘disease-specific advance* plan*’ OR ‘chronic illness advance* plan*’ OR ‘progressive illness advance* plan*’ OR ‘patient advance* plan*’ OR ‘patient advance* directive*’ OR ‘patient advance* statement*’ OR ‘patient advance* decision*’ OR ‘prefer* place* of care’ OR ‘prefer* place* of death’ OR ‘patient care plan*’ OR ‘anticipatory care’ OR ‘anticipatory plan’ OR ‘health care prox*’ OR ‘healthcare prox*’ OR ‘power of attorney’ OR ‘surrogate decision-maker’ OR ‘surrogate decision maker’ OR ‘decision making representative*’ OR ‘decision-making representative*’
Design	‘review’ OR ‘systematic review’ OR ‘synthesis’ OR ‘meta-synthesis’ OR ‘meta-analysis’
Evaluation	barrier* OR facilitator* OR impediment* OR obstacle* OR deterrent* OR difficult* OR enabler* OR promot* OR benefit* OR burden*
Setting	‘community care’ OR ‘community health care’ OR ‘community healthcare’ OR ‘primary care’ OR ‘primary health care’ OR ‘primary healthcare’ OR ‘general practice’ OR ‘general practitioner’ OR ‘family practice’ OR ‘community-dwelling person’ OR ‘community-dweller’ OR ‘home care’




Figure 1. A two-stage process of primary studies identification. SRs, systematic reviews.

Assessment of risk of bias in included studies

As the primary studies will be extracted from the eligible systematic reviews, the quality assessment of those studies will also be extracted (if available). If not available, the risk of bias assessment will be completed by MP and VL. The Crowe Critical Appraisal Tool (CCAT); a validated general critical appraisal instrument that can be used across a wide range of research designs, will be used (Crowe *et al.*, 2012). A similar procedure will be followed if reviews disagree on the assessment of bias of the included primary studies. The quality of newly identified primary studies (not included in the eligible systematic reviews) will also be assessed with the CCAT tool. Other reviewers (PM, FD, or ST) will be involved, if necessary. The appraisal checklist will be computerised and the electronic version will be used to facilitate critical assessment process. The tool will be used to identify bias but not to exclude studies.

Data extraction and management

Two data extraction forms will be designed to ensure a focused and systematic approach to data collection. The form for the extraction of data from systematic reviews will be informed by the JBI instrument for the conduct of umbrella reviews (Aromataris *et al.*, 2017). The JBI Mixed Methods data extraction form will be used to develop a data extraction form for individual studies (Lizarondo *et al.*, 2019). Each form will be piloted on a sample of three studies (reviews or individual studies, respectively) and adapted, if necessary. When finalised, the forms will be computerised and the electronic versions will be used to facilitate data extraction processes.

The following information will be extracted from the reviews: (a) review details (authors, year of publication, aims and objectives); (b) search details (sources searched, range of included studies, number of studies included in the review); (c) details of quality appraisal (appraisal tool used and appraisal rating), if available.

The following information will be extracted from the individual studies: (a) study details (authors, year of publication, journal, title, geographical area of the study, overarching conceptual frameworks); (b) methodology (study design, aims and objectives, context/settings, participant characteristics, description of phenomena of interest); (c) details of analysis/ analytical approach; (d) details of evaluation (outcomes or findings of significance to the review objectives, themes or subthemes and associated illustration), (e) authors' conclusions. If a study evaluated an intervention, we will also extract the name of the intervention used. If relevant information is reported in a primary study, we will identify COM-B factors addressed in that intervention". If available, data relevant to the review questions will be independently extracted from the eligible systematic reviews by the first and the second reviewer (MP & VL). If not available, a standard procedure will be followed to extract data from the eligible primary studies. The same process of data extraction will be repeated for the primary studies identified through the supplementary search (Phase 2). Conflicts will be resolved through discussion and consensus. Other reviewers (PM, FD, or ST) will be involved if there is a disagreement. To obtain information on missing data, authors will be contacted.

Analytical approach and integration of findings

Integrated methodologies will be employed and quantitative and qualitative data will be combined into a single mixed research synthesis (Sandelowski *et al.*, 2006). Data will be integrated by translating qualitative data into numerical format. This converted data will be presented along with quantitative data in a summary table (Lizarondo *et al.*, 2019). Then, following the similar approach taken elsewhere (Sharpe *et al.*, 2018), the findings will be mapped across the COM-B categories (see Figure 2). Specifically, MP and VL will record the frequency of reported facilitators and barriers and group the findings into the relevant categories of the COM-B model. Then, the BCW (Michie *et al.*, 2011) will be utilised as a broader analytical framework to differentiate individual-level, service-level, and




Figure 2. The COM-B system (adapted from: [Michie et al., 2011](#), under a [CC BY 2.0](#) license).

system-level factors that facilitate and/or impede stakeholders' engagement in ACP. Other reviewers (PM, FD, & ST) will be asked to provide a critical evaluation of the analytical process and the findings.

Dissemination

We will apply a peer-reviewed knowledge transfer and exchange model ([Payne et al., 2019](#)) and use multi-channel dissemination to reach different audiences and all key stakeholders. A systematic review article will be submitted for peer-reviewed publication.

Study status

Preliminary searches have been completed.

Discussion

Given the low engagement in ACP and the known challenges to introducing ACP in the last stage of illness trajectories, which usually occurs in acute healthcare settings, there is a need to explore salient factors facilitating ACP engagement in a community context. These settings are likely to offer space and time for ACP conversations, allowing their integration into standard care. This review will be the first to adopt a hybrid review methodology to present cumulative evidence on facilitators and barriers to ACP among older adults in a community setting, as perceived from the perspective of various stakeholders. By uniquely adopting the BCW framework, it will provide insights into different levels of influence (including individual, service-based, and systemic), and with the view of informing behaviour change strategies. This comprehensive approach will aim to enhance the understanding of modifiable and non-modifiable factors that may facilitate or impede early key stakeholders' engagement in ACP for older adults, irrespective of their health status.

The adoption of a specific methodology, combining a systematic search for systematic reviews with systematic supplementary searches for original studies, will allow a cumulative and time-effective approach. The strength of the hybrid approach is that it leverages the work undertaken in previous systematic reviews, e.g., by utilising the reported outcomes of comprehensive searches and selection processes ([Doyle et al., 2019](#)).

Given the amount of existing evidence, this methodology will prevent the repetition of already completed and reported processes, facilitate more appropriate management of resources, and allow the application of sophisticated analytical frameworks to cumulative evidence.

The findings of this review may inform the development of an innovative and evidence-based interventions for ACP. The Medical Research Council Guidelines highlight the need to identify strong evidence base and model processes and outcomes to inform the development of complex healthcare interventions ([Craig et al., 2019](#)). As different settings may influence engagement in ACP ([Lewis et al., 2016](#)), the need to keep the implementation context in mind when designing and assessing interventions has been highlighted ([Edmondson et al., 2001](#); [Jimenez et al., 2018](#)). A careful and critical synthesis of available evidence, enhancing the understanding of barriers and enablers of ACP engagement in a community setting, can facilitate the development of relevant ACP interventions ([Risk et al., 2019](#)). While suggesting direction for further empirical investigations, the findings will also allow to derive conclusions and recommendations for clinical and policy decision making.

Data availability

Underlying data

No underlying data are associated with this article.

Extended data

Open Science Framework: Facilitators and barriers to stakeholder engagement in Advance Care Planning for older adults in community settings: A hybrid systematic review protocol. <https://doi.org/10.17605/OSF.IO/6TS34> ([Pilch, 2020](#)).

Reporting guidelines

Open Science Framework: PRISMA-P checklist for 'Facilitators and barriers to stakeholder engagement in Advance Care Planning for older adults in community settings: A hybrid systematic review protocol.' <https://doi.org/10.17605/OSF.IO/6TS34> ([Pilch, 2020](#)).

Data are available under the terms of the [Creative Commons Attribution 4.0 International license](#) (CC-BY 4.0).

References

- Aromataris E, Fernandez R, Godfrey C, et al.: **Chapter 10: Umbrella Reviews.** In: E. Aromataris & Z. Munn (Eds.), *Joanna Briggs Institute Reviewer's Manual*. The Joanna Briggs Institute. 2017.
- Austin CA, Mohottige D, Sudore RL, et al.: **Tools to promote shared decision making in serious illness: A systematic review.** *JAMA Intern Med.* 2015; **175**(7): 1213–1221.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Balein GN: **Factors Associated with Good Death.** *Asian Bioeth Rev.* 2009; **1**(3): 252–258.
[Reference Source](#)
- Biondo PD, King S, Minhas B, et al.: **How to increase public participation in advance care planning: Findings from a World Café to elicit community group perspectives.** *BMC Public Health.* 2019; **19**(1): 679.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bischoff KE, Sudore R, Miao Y, et al.: **Advance Care Planning and the Quality of End-of-Life Care in Older Adults.** *J Am Geriatr Soc.* 2013; **61**(2): 209–214.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Brinkman-Stoppelenburg A, Rietjens JAC, van der Heide A: **The effects of advance care planning on end-of-life care: A systematic review.** *Palliat Med.* 2014; **28**(8): 1000–1025.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Centeno C, Arias-Casais N: **Global palliative care: from need to action.** *Lancet Glob Health.* 2019; **7**(7): e815–e816.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Clayton JM, Hancock KM, Butow PN, et al.: **Clinical practice guidelines for communicating prognosis and end-of-life issues with adults in the advanced stages of a life-limiting illness, and their caregivers.** *Med J Aust.* 2007; **186**(S12): S77–S105.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Cooke A, Smith D, Booth A: **Beyond PICO: The SPIDER Tool for Qualitative Evidence Synthesis.** *Qual Health Res.* 2012; **22**(10): 1435–1443.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Craig P, Dieppe P, Macintyre S, et al.: **Developing and evaluating complex interventions: Following considerable development in the field since 2006, MRC and NIHR have jointly commissioned an update of this guidance to be published in 2019.** *Med Res Coun.* 2019; 1–39.
[Reference Source](#)
- Crowe M, Sheppard L, Campbell A: **Reliability analysis for a proposed critical appraisal tool demonstrated value for diverse research designs.** *J Clin Epidemiol.* 2012; **65**(4): 375–383.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Davison SN: **End-of-life care preferences and needs: Perceptions of patients with chronic kidney disease.** *Clin J Am Soc Nephrol.* 2010; **5**(2): 195–204.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- De Vlemingck A, Houttekier D, Pardon K, et al.: **Barriers and facilitators for general practitioners to engage in advance care planning: A systematic review.** *Scand J Prim Health Care.* 2013; **31**(4): 215–226.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Downey L, Au DH, Curtis JR, et al.: **Life-sustaining treatment preferences: Matches and mismatches between patients' preferences and clinicians' perceptions.** *J Pain Symptom Manage.* 2013; **46**(1): 9–19.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Doyle F, Freedland K, Carney R, et al.: **Network meta-analysis of randomised trials of pharmacological, psychotherapeutic, exercise and collaborative care interventions for depressive symptoms in patients with coronary artery disease: Hybrid systematic review of systematic reviews protocol.** *Syst Rev.* 2019; **8**(1): 71.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Earle CC, Landrum MB, Souza JM, et al.: **Aggressiveness of cancer care near the end of life: Is it a quality-of-care issue?** *J Clin Oncol.* 2008; **26**(23): 3860–3866.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Edmondson AC, Bohmer RM, Pisano GP: **Disrupted routines: Team learning and new technology implementation in hospitals.** *Adm Sci Q.* 2001; **46**(4): 685–716.
[Publisher Full Text](#)
- El-Khayat YM: **Epistemonikos.** *J Med Libr Assoc.* 2017; **105**(4): 431–432.
[Publisher Full Text](#) | [Free Full Text](#)
- Fried TR, Bullock K, Iannone L, et al.: **Understanding advance care planning as a process of health behavior change.** *J Am Geriatr Soc.* 2009; **57**(9): 1547–1555.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Fried TR, Redding CA, Martino S, et al.: **Increasing engagement in advance care planning using a behaviour change model: Study protocol for the STAMP randomised controlled trials.** *BMJ Open.* 2018; **8**(8): e025340.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Fried TR, Redding CA, Robbins ML, et al.: **Development of Personalized Health Messages to Promote Engagement in Advance Care Planning.** *J Am Geriatr Soc.* 2016; **64**(2): 359–364.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Fried TR, Redding CA, Robbins ML, et al.: **Stages of change for the component behaviors of advance care planning.** *J Am Geriatr Soc.* 2010; **58**(12): 2329–2336.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Fried TR, Redding CA, Robbins ML, et al.: **Promoting advance care planning as health behavior change: Development of scales to assess Decisional Balance, Medical and Religious Beliefs, and Processes of Change.** *Patient Educ Couns.* 2012; **86**(1): 25–32.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Garrido MM, Balboni TA, Maciejewski PK, et al.: **Quality of life and cost of care at the end of life: The role of advance directives.** *J Pain Symptom Manage.* 2015; **49**(5): 828–835.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gidwani-Marszowski R, Asch SM, Mor V, et al.: **Health System and Beneficiary Costs Associated With Intensive End-of-Life Medical Services.** *JAMA Network Open.* 2019; **2**(9): e1912161.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gilissen J, Pivodic L, Wendrich-van Dael A, et al.: **Implementing advance care planning in routine nursing home care: The development of the theory-based ACP+ program.** *PLoS One.* 2019; **14**(10): e0223586.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Gomes B, Higginson JJ, Calanzani N, et al.: **Preferences for place of death if faced with advanced cancer: A population survey in England, Flanders, Germany, Italy, The Netherlands, Portugal and Spain.** *Ann Oncol.* 2012; **23**(8): 2006–2015.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Howard M, Bernard C, Klein D, et al.: **Older patient engagement in advance care planning in Canadian primary care practices: Results of a multisite survey.** *Can Fam Physician.* 2018; **64**(5): 371–377.
[PubMed Abstract](#) | [Free Full Text](#)
- Jimenez G, Tan WS, Virk AK, et al.: **Overview of Systematic Reviews of Advance Care Planning: Summary of Evidence and Global Lessons.** *J Pain Symptom Manage.* 2018; **56**(3): 436–459.e25.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Kelley AS, Wenger NS, Sarkisian CA: **Opinions: End-Of-Life Care Preferences and Planning of Older Latinos.** *J Am Geriatr Soc.* 2010; **58**(6): 1109–1116.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kelly BD: **The Assisted Decision-Making (Capacity) Act 2015: what it is and why it matters.** *Ir J Med Sci.* 2017; **186**(2): 351–356.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Khandelwal N, Kross EK, Engelberg RA, et al.: **Estimating the Effect of Palliative Care Interventions and Advance Care Planning on ICU Utilization: A Systematic Review.** *Crit Care Med.* 2015; **43**(5): 1102–1111.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lewis E, Cardona-Morrell M, Ong KY, et al.: **Evidence still insufficient that advance care documentation leads to engagement of healthcare professionals in end-of-life discussions: A systematic review.** *Palliat Med.* 2016; **30**(9): 807–824.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Lin CP, Evans CJ, Koffman J, et al.: **The conceptual models and mechanisms of action that underpin advance care planning for cancer patients: A systematic review of randomised controlled trials.** *Palliat Med.* 2019; **33**(1): 5–23.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lizarondo L, Stern C, Carrier J, et al.: **Chapter 8: Mixed methods systematic reviews.** In: E. Aromataris & Z. Munn (Eds.), *Joanna Briggs Institute Reviewer's Manual*. The Joanna Briggs Institute. 2019.
[Reference Source](#)
- Michie S, van Stralen MM, West R: **The behaviour change wheel: A new method for characterising and designing behaviour change interventions.** *Implement Sci.* 2011; **6**(1): 42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Moher D, Shamseer L, Clarke M, et al.: **Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement.** *Syst Rev.* 2015; **4**(1): 1.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Mullaney C, Gleeson M, Quinn G, et al.: **Living Well with a Chronic Condition: Framework for Self-management Support.** *National Framework and Implementation Plan for Self-management Support for Chronic Conditions: COPD, Asthma, Diabetes and Cardiovascular disease.* 2016.
[Reference Source](#)
- Nguyen KH, Sellars M, Agar M, et al.: **An economic model of advance care planning in Australia: A cost-effective way to respect patient choice.** *BMC Health Serv Res.* 2017; **17**(1): 797.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Osther K, Killoran P, Shegog R, et al.: **Death in the Digital Age: A Systematic Review of Information and Communication Technologies in End-of-Life Care.** *J Palliat Med.* 2016; **19**(4): 408–420.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Payne C, Brown MJ, Guerin S, et al.: **EMTRek: An Evidence-based Model for**

the Transfer & Exchange of Research Knowledge—Five Case Studies in Palliative Care. *SAGE Open Nurs.* 2019; 5: 1–8.
[Publisher Full Text](#)

Peck V, Valiani S, Tanuseputro P, et al.: Advance care planning after hospital discharge: qualitative analysis of facilitators and barriers from patient interviews. *BMC Palliat Care.* 2018; 17(1): 127.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Pilch M: Appendices to [Facilitators and barriers to stakeholder engagement in Advance Care Planning for older adults in community settings: A hybrid systematic review protocol]. *OSF.* 2020.
<http://www.doi.org/10.17605/OSF.IO/6TS34>

Pinquart M, Düberstein PR: Depression and cancer mortality: A meta-analysis. *Psychol Med.* 2010; 40(11): 1797–1810.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Prendergast TJ: Advance care planning: Pitfalls, progress, promise. *Crit Care Med.* 2001; 29(7): 34–39.
[PubMed Abstract](#) | [Publisher Full Text](#)

Ramsaroop S, Reid M, Adelman R: Completing an advance directive in the primary care setting: what do we need for success? *J Am Geriatr Soc.* 2007; 55(2): 277–283.
[PubMed Abstract](#) | [Publisher Full Text](#)

Rietjens JAC, Sudore RL, Connolly M, et al.: Definition and recommendations for advance care planning: an international consensus supported by the European Association for Palliative Care. *Lancet Oncol.* 2017; 18(9): e543–e551.
[PubMed Abstract](#) | [Publisher Full Text](#)

Risk J, Mohammadi L, Rhee J, et al.: Barriers, enablers and initiatives for uptake of advance care planning in general practice: a systematic review and critical interpretive synthesis. *BMJ Open.* 2019; 9(9): e030275.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Sandelowski M, Voils CI, Barroso J: Defining and Designing Mixed Research Synthesis Studies. *Res Sch.* 2006; 13(1): 29.
[PubMed Abstract](#) | [Free Full Text](#)

Sharpe T, Alsahlane A, Ward KD, et al.: Systematic Review of Clinician-Reported Barriers to Provision of Smoking Cessation Interventions in Hospital Inpatient Settings. *J Smok Cessat.* 2018; 13(4): 233–243.
[Publisher Full Text](#)

Silveira MJ, Kim SYH, Langa KM: Advance Directives and Outcomes of Surrogate Decision Making before Death. *N Engl J Med.* 2010; 362(13): 1211–1218.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Simon J, Porterfield P, Bouchal SR, et al.: “Not yet” and “Just ask”: Barriers and facilitators to advance care planning—a qualitative descriptive study of the perspectives of seriously ill, older patients and their families. *BMJ Support Palliat Care.* 2015; 5(1): 54–62.
[PubMed Abstract](#) | [Publisher Full Text](#)

Solis GR, Mancera BM, Shen MJ: Strategies used to facilitate the discussion of advance care planning with older adults in primary care settings: A literature review. *J Am Assoc Nurse Pract.* 2018; 30(5): 270–279.
[PubMed Abstract](#) | [Publisher Full Text](#)

Street M, Ottmann G, Johnstone M, et al.: Advance care planning for older people in Australia presenting to the emergency department from the community or residential aged care facilities. *Health Soc Care Community.* 2014; 23(5): 513–522.
[PubMed Abstract](#) | [Publisher Full Text](#)

Sudore RL, Boscardin J, Feuz MA, et al.: Effect of the PREPARE website vs an easy-to-read advance directive on advance care planning documentation and engagement among veterans A randomized clinical trial. *JAMA Intern Med.* 2017; 177(8): 1102–1109.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Sudore RL, Heyland DK, Lum HD, et al.: Outcomes That Define Successful Advance Care Planning: A Delphi Panel Consensus. *J Pain Symptom Manage.* 2018; 55(2): 245–255.e8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Sudore RL, Schickedanz AD, Landefeld CS, et al.: Engagement in multiple steps of the advance care planning process: A descriptive study of diverse older adults. *J Am Geriatr Soc.* 2008; 56(6): 1006–1013.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Tam-Tham H, King-Shier KM, Thomas CM, et al.: Prevalence of barriers and facilitators to enhancing conservative kidney management for older adults in the primary care setting. *Clin J Am Soc Nephrol.* 2016; 11(11): 2012–2021.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Tilburgs B, Vernooij-Dassen M, Koopmans R, et al.: Barriers and facilitators for GPs in dementia advance care planning: A systematic integrative review. *PLoS One.* 2018; 13(6): 1–21.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Van Der Steen JT, Van Soest-Poortvliet MC, Onwuteaka-Philipsen BD, et al.: Factors associated with initiation of advance care planning in dementia: A systematic review. *J Alzheimers Dis.* 2014; 40(3): 743–757.
[PubMed Abstract](#) | [Publisher Full Text](#)

Van Scoy LJ, Green MJ, Reading JM, et al.: Can Playing an End-of-Life Conversation Game Motivate People to Engage in Advance Care Planning? *Am J Hosp Palliat Care.* 2017; 34(8): 754–761.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Wright A, Zhang B, Ray A, et al.: Associations between end-of-life discussions, patient mental health, medical care near death, and caregiver bereavement adjustment. *JAMA.* 2008; 300(14): 1665–1673.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Zhang B, Wright AA, Huskamp HA, et al.: Health Care Costs in the Last Week of Life. *Arch Intern Med.* 2009; 169(5): 480.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:  

Version 2

Reviewer Report 17 June 2021

<https://doi.org/10.21956/hrbopenres.14513.r29658>

© 2021 Lee S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Susan Fiona Lee 

Nursing and Midwifery, Nursing and Health Sciences, Monash University, Frankston, Vic, Australia

I have read the comments and revisions and find them to be thoughtful and having addressed my few concerns. The review adds to our understanding of the complexity of advance care planning.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Palliative care, end of life care, decision making, advance care planning.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 01 September 2020

<https://doi.org/10.21956/hrbopenres.14182.r27813>

© 2020 Lee S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Susan Fiona Lee 

Nursing and Midwifery, Nursing and Health Sciences, Monash University, Frankston, Vic, Australia

Thank you for the opportunity to review this protocol for a novel systematic review. This mixed method review seeks to examine barriers and facilitators to stakeholder engagement in Advance Care Planning (ACP) in community settings. The context is well argued in relation to the need for

earlier ACP. The review uses a novel hybrid approach by first searching for systematic reviews and following this, the subsequently published individual articles. Risk of bias assessments are to be conducted and rigour will be maintained through the use of independent selection of studies, risk of bias assessment and data extraction by two reviewers. A further novel approach is used in the analysis involving a mixed-method research synthesis, and findings will be mapped against the existing ACP outcomes framework.

The need for this review is clearly justified and it is a reasonably well thought out process. I recommend revisiting two areas.

1. The hybrid approach does not include the common third search strategy of grey literature on the basis of a lack of peer review. I presume these studies will also not be extracted from the systematic reviews. As critique is part of the rigour of the review, studies that have not been peer reviewed, along with those that have been reviewed, undergo risk of bias assessment. I believe the exclusion of grey literature searching and studies should be reviewed.
2. I found some confusion in the treatment of the systematic reviews discovered in phase 1. It appears from the search strategy that the systematic reviews are treated as a data base of eligible studies and the studies will be extracted and added to the phase 2 study list. However, in the data management section and analysis, the results of the reviews are extracted and used in the synthesis, as in an umbrella review. This must be clarified prior to data extraction.

As a final point, I did note that the search terms did not include any synonyms for surrogate. I recommend 'substitute*' and 'alternate' to ensure the scope of this group. There also appears a typographical error in the first paragraph of the Data extraction and management section where JBM is referred to. I believe it should be JBI.

I wish you well with the review and look forward to reading the final outcomes which should have practical implications for the field.

Is the rationale for, and objectives of, the study clearly described?

Yes

Is the study design appropriate for the research question?

Partly

Are sufficient details of the methods provided to allow replication by others?

Partly

Are the datasets clearly presented in a useable and accessible format?

Not applicable

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Palliative care, end of life care, decision making, advance care planning.

I confirm that I have read this submission and believe that I have an appropriate level of

expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 04 Jun 2021

Monika Pilch, Trinity College Dublin, University of Dublin, Dublin, D2, Ireland

We are grateful for the time taken by reviewers in providing a careful analysis of our protocol. We appreciate your positive comments and constructive suggestions to further improve our manuscript. We have revised the paper and have made a point-by-point reply to your comments, as follows:

1. Thank you for the opportunity to review this protocol for a novel systematic review. This mixed method review seeks to examine barriers and facilitators to stakeholder engagement in Advance Care Planning (ACP) in community settings. The context is well argued in relation to the need for earlier ACP. The review uses a novel hybrid approach by first searching for systematic reviews and following this, the subsequently published individual articles. Risk of bias assessments are to be conducted and rigour will be maintained through the use of independent selection of studies, risk of bias assessment and data extraction by two reviewers. A further novel approach is used in the analysis involving a mixed-method research synthesis, and findings will be mapped against the existing ACP outcomes framework.

The need for this review is clearly justified and it is a reasonably well thought out process

Thank you for these positive comments.

2. The hybrid approach does not include the common third search strategy of grey literature on the basis of a lack of peer review. I presume these studies will also not be extracted from the systematic reviews. As critique is part of the rigour of the review, studies that have not been peer reviewed, along with those that have been reviewed, undergo risk of bias assessment. I believe the exclusion of grey literature searching and studies should be reviewed.

As stated in response to the first reviewer (comment 6), we decided to focus on peer-reviewed literature only. This has now been explicitly stated in the text (p. 11). As explained above, the inclusions of grey literature in this review would be beyond the scope of this PhD project. However, we will acknowledge this limitation in the discussion and will comment on the impact of this approach on findings.

3. I found some confusion in the treatment of the systematic reviews discovered in phase 1. It appears from the search strategy that the systematic reviews are treated as a data base of eligible studies and the studies will be extracted and added to the phase 2 study list. However, in the data management section and analysis, the results of the reviews are extracted and used in the synthesis, as in an umbrella review. This must be clarified prior to data extraction

Thank you for this very helpful observation. We have now revised the second paragraph in the

"Data extraction and management section" and specified that only information relevant to the processing of included primary studies will be extracted from the included systematic reviews:

"The following information will be extracted from the reviews: (a) review details (authors, year of publication); (b) search details (sources searched, range of included studies, number of studies included in the review); (c) details of quality appraisal (appraisal tool used and appraisal rating), if available." (p. 15-16)

4. As a final point, I did note that the search terms did not include any synonyms for surrogate. I recommend 'substitute*' and 'alternate' to ensure the scope of this group.

We appreciate this suggestion. As explained in response to Reviewer 1 (comment 5), our search strategy was informed by an information specialist and, at the time of receiving this feedback, searches had been completed. Due to the large workload and lack of new resources, the inclusion of new terms would not be feasible and we need to decline to conduct the search again. However, we will acknowledge the omission of the term as a study limitation in the report from this review.

5. There also appears a typographical error in the first paragraph of the Data extraction and management section where JBM is referred to. I believe it should be JBI.

Thank you, this has now been corrected. (p. 14)

6. I wish you well with the review and look forward to reading the final outcomes which should have practical implications for the field.

Thank you. We look forward to sharing the report from the study.

Competing Interests: No competing interests were disclosed.

Reviewer Report 10 August 2020

<https://doi.org/10.21956/hrbopenres.14182.r27677>

© 2020 Simon J. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Jessica E. Simon 

Department of Oncology, University of Calgary, Calgary, AB, Canada

Is the rationale for, and objectives of, the study clearly described?

This is an important systematic review addressing the issue of engagement in advance care planning (ACP) by community dwelling adults over 50 years of age. This is a key population who may benefit from participating in ACP with potential broader impacts on population health and health system resource use. The theoretical framework selected for the analysis has the advantage

of allowing the barriers and facilitators to engagement in ACP by community dwelling adults over 50 years of age to be understood in terms of behavioural change. The study should result in understanding how Capability, Motivation, and Opportunity at the individual, interpersonal, provider, and system levels of influence uptake of ACP by these adults.

The paper provides an excellent background with a current consensus definition of ACP and relevant, succinct literature reviewed.

One notable publication that was missing from the background section is Lovell and Yates (2014)¹. The authors might want to reconsider their sentence, *"..no study has synthesised available evidence to identify patterns across older adults, irrespective of their health status or condition, while taking stakeholder perspective and level of influence into account."* The review above does explore the global contextual factors for ACP uptake that this protocol will be seeking to understand, although it does not use an organizing theory such as Michie's COM-B model. Consider changing the sentence to, *"...few studies have synthesized available evidence to identify patterns across older adults, irrespective of their health status or condition and these have used an organizing behaviour change theory and taken all stakeholder perspectives and level of influence into account."*

One very minor grammatical edit could be considered. In a couple of places this phrase is used: "As readiness to ACP engagement..." . To increase ease of reading perhaps consider changing to "As readiness for ACP engagement..." or, "As readiness to engage in ACP..."

Is the study design appropriate for the research question?

The study design and search strategy is well described, using a hybrid systematic review with two phases. I am curious whether adding "public" as a setting to the search terms might be useful to identify additional references with healthy adults that the term "community" might not pick up. The authors do not comment specifically on whether "grey literature" references from the phase 1 systematic reviews will be included in phase 2 but these would seem an additional important source of information.

Given the authors stated aim that, "The findings of this review may inform the development of an innovative and evidence-based interventions for ACP" I'm inclined to suggest considering an addition to your data extraction table to include whether the study evaluated an intervention strategy and what that intervention was and which COM-B factor it addressed may enable their analysis which seeks to "differentiate individual-level, service-level, and system-level factors that facilitate and/or impede stakeholders engagement in ACP" and assist the authors in moving into describing what intervention functions and policy categories have been published as the next step in applying Michie's Behaviour Change Wheel.

Are sufficient details of the methods provided to allow replication by others?

Yes, except that the timeframe proposed for the phase 1 systematic review search and any included/excluded languages are not provided. Please add this detail.

Are the datasets clearly presented in a useable and accessible format? Yes the search terms are clearly presented and accessible.

Yes, this is well done.

References

1. Lovell A, Yates P: Advance Care Planning in palliative care: a systematic literature review of the

contextual factors influencing its uptake 2008-2012. *Palliat Med.* 2014; **28** (8): 1026-35 [PubMed Abstract](#) | [Publisher Full Text](#)

Is the rationale for, and objectives of, the study clearly described?

Yes

Is the study design appropriate for the research question?

Yes

Are sufficient details of the methods provided to allow replication by others?

Yes

Are the datasets clearly presented in a useable and accessible format?

Yes

Competing Interests: Dr Simon is a physician consultant for Advance Care Planning and Goals of Care, Alberta Health Services, Calgary Zone.

Reviewer Expertise: Advance care planning and palliative care.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 04 Jun 2021

Monika Pilch, Trinity College Dublin, University of Dublin, Dublin, D2, Ireland

We are grateful for the time taken by reviewers in providing a careful analysis of our protocol. We appreciate your positive comments and constructive suggestions to further improve our manuscript. We have revised the paper and have made a point-by-point reply to your comments, as follows:

1. This is an important systematic review addressing the issue of engagement in advance care planning (ACP) by community dwelling adults over 50 years of age. This is a key population who may benefit from participating in ACP with potential broader impacts on population health and health system resource use. The theoretical framework selected for the analysis has the advantage of allowing the barriers and facilitators to engagement in ACP by community dwelling adults over 50 years of age to be understood in terms of behavioural change. The study should result in understanding how Capability, Motivation, and Opportunity at the individual, interpersonal, provider, and system levels of influence uptake of ACP by these adults.

Thank you for these positive comments.

2. The paper provides an excellent background with a current consensus definition of ACP and relevant, succinct literature reviewed.

Thank you for this positive comment.

3. One notable publication that was missing from the background section is Lovell and Yates (2014)¹. The authors might want to reconsider their sentence, “..no study has synthesised available evidence to identify patterns across older adults, irrespective of their health status or condition, while taking stakeholder perspective and level of influence into account.” The review above does explore the global contextual factors for ACP uptake that this protocol will be seeking to understand, although it does not use an organizing theory such as Michie’s COM-B model. Consider changing the sentence to, “...few studies have synthesized available evidence to identify patterns across older adults, irrespective of their health status or condition and these have used an organizing behaviour change theory and taken all stakeholder perspectives and level of influence into account.”

*Thank you for this comment. We think the reviewer meant to state that these reviews have **not** used an organising behaviour change theory? To increase clarity, we applied the suggestion and changed the text in the relevant section (p. 6) to the following:*

*“Although several studies have been conducted to explore factors associated with the uptake of ACP among older adults in community settings; and these have been synthesised in previous systematic reviews (e.g., De Vleminck et al., 2013; Ramsaroop et al., 2007; Risk et al., 2019; Solis et al., 2018), **few studies have** synthesised available evidence to identify patterns across older adults, irrespective of their health status or condition, while taking stakeholder perspective and level of influence into account. A synthesis utilising the BCW framework is necessary to identify influences within the contextual layers of health behaviour among older adults living in a community, while highlighting individual, interpersonal, provider, and system levels of influence.” (p. 6)*

4. One very minor grammatical edit could be considered. In a couple of places this phrase is used: “As readiness to ACP engagement...” . To increase ease of reading perhaps consider changing to “As readiness for ACP engagement...” or, “As readiness to engage in ACP...”

Thank you, these changes have now been applied on pages 5 and 6.

5. The study design and search strategy is well described, using a hybrid systematic review with two phases. I am curious whether adding “public” as a setting to the search terms might be useful to identify additional references with healthy adults that the term “community” might not pick up.

We appreciate this suggestion. Our search strategy was informed by an information specialist and, at the time of receiving this feedback, searches had been completed. Due to the large workload and lack of new resources, the inclusion of new terms would not be feasible and we need to decline to conduct the search again. However, we will acknowledge the omission of the term as a study limitation in the report from this review.

6. The authors do not comment specifically on whether “grey literature” references from the phase 1 systematic reviews will be included in phase 2 but these would seem an additional important source of information.

Thank you for this comment. Prior to preparing this protocol, we completed a scoping search. The outcome of the scoping suggested a considerable amount of evidence available. Although we recognise the importance of "grey literature", we also had to consider the limited scope of research, which is a part of a PhD project, and the need to complete it within the required timeframe. Therefore, we made a decision to exclude "grey literature" in our systematic review. This has now been explicitly stated in the text, by adding the following sentence:

"Although we acknowledge the importance of grey literature, due to envisaged amount of existing evidence on the topic, grey literature will not be considered in this review." (p. 11)

7. Given the authors stated aim that, "The findings of this review may inform the development of an innovative and evidence-based interventions for ACP" I'm inclined to suggest considering an addition to your data extraction table to include whether the study evaluated an intervention strategy and what that intervention was and which COM-B factor it addressed may enable their analysis which seeks to "differentiate individual-level, service-level, and system-level factors that facilitate and/or impede stakeholders engagement in ACP" and assist the authors in moving into describing what intervention functions and policy categories have been published as the next step in applying Michie's Behaviour Change Wheel.

Thank you for this insightful comment. The aim of this review is to identify facilitators and barriers to ACP engagement, however, we appreciate potential benefits of including this additional information. We now specified that, if a study evaluated an intervention, we will include information about the name and type of a strategy used. Additionally, where relevant information is available, we will identify specific COM-B factors associated with these interventions. We included this plan in the protocol and specified (in the data extraction subsection) that "If a study evaluated an intervention, we will also extract the name of the strategy used. If relevant information is reported in a primary study, we will identify COM-B factors addressed in that intervention". (p. 15)

8. Are sufficient details of the methods provided to allow replication by others?
Yes, except that the timeframe proposed for the phase 1 systematic review search and any included/excluded languages are not provided. Please add this detail.

We have now specified the timeframe for the phase 1 systematic review. Specifically, we added this underlined part to the sentence in the "Search Strategy section.

"A comprehensive search for systematic reviews will be conducted in each database from its inception (Phase 1) and this process will be completed in June 2020". (p. 11)

With regards to the timeframe, we plan to include only studies reported in English. This information was included on page 8 of the original protocol submission. "Only reviews and primary studies reported in English and published in peer-reviewed journals will be considered for inclusion." (p. 8)

Competing Interests: No competing interests were disclosed.
