




# High Prevalence of Obesity in Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-CoV-2) Requiring Invasive Mechanical Ventilation

Arthur Simonnet<sup>1\*</sup>, Mikael Chetboun<sup>2\*</sup>, Julien Poissy<sup>1</sup>, Violeta Raverdy<sup>1</sup> <sup>2</sup>, Jerome Noulette<sup>2</sup>, Alain Duhamel<sup>3</sup>, Julien Labreuche<sup>3</sup>, Daniel Mathieu<sup>1</sup>, Francois Pattou<sup>1,2,4</sup> , and Merce Jourdain<sup>1,2</sup> ,  
on behalf of the LICORN and the Lille COVID-19 and Obesity study group

**Objective:** The COVID-19 pandemic is rapidly spreading worldwide, notably in Europe and North America where obesity is highly prevalent. The relation between obesity and severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) has not been fully documented.

**Methods:** This retrospective cohort study analyzed the relationship between clinical characteristics, including BMI, and the requirement for invasive mechanical ventilation (IMV) in 124 consecutive patients admitted in intensive care for SARS-CoV-2 in a single French center.

**Results:** Obesity (BMI >30) and severe obesity (BMI >35) were present in 47.6% and 28.2% of cases, respectively. Overall, 85 patients (68.6%) required IMV. The proportion of patients who required IMV increased with BMI categories ( $P < 0.01$ ,  $\chi^2$  test for trend), and it was greatest in patients with BMI >35 (85.7%). In multivariate logistic regression, the need for IMV was significantly associated with male sex ( $P < 0.05$ ) and BMI ( $P < 0.05$ ), independent of age, diabetes, and hypertension. The odds ratio for IMV in patients with BMI >35 versus patients with BMI <25 was 7.36 (1.63–33.14;  $P = 0.02$ ).

**Conclusions:** The present study showed a high frequency of obesity among patients admitted in intensive care for SARS-CoV-2. Disease severity increased with BMI. Obesity is a risk factor for SARS-CoV-2 severity, requiring increased attention to preventive measures in susceptible individuals.

Obesity (2020) 28, 1195–1199.

## Introduction

Patients at risk for severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) have been characterized as having preexisting diseases, such as hypertension, cardiovascular disease, diabetes, chronic respiratory disease, or cancer (1). Surprisingly, BMI has been rarely mentioned among the significant clinical risk factors for SARS-CoV-2 reported in early clinical reports from China (2), Italy (3), or the United States (4). Obesity was, however, described as an independent predisposition factor for severe H1N1 pulmonary infection (5). Moreover, abdominal obesity is associated with impaired ventilation of the base of the lungs, resulting in reduced oxygen saturation of

## Study Importance

### What is already known?

- ▶ A novel coronavirus causing severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) emerged in China and has spread globally, creating a worldwide pandemic.
- ▶ Obesity has been previously recognized as an independent predisposition factor for severe H1N1 pulmonary infection.
- ▶ Information about the clinical characteristics of infected patients who require intensive care has been limited, and the relationship between obesity and SARS-CoV-2 is unclear.

### What does this study add?

- ▶ Obesity was unexpectedly frequent in a cohort of patients admitted in intensive care for SARS-CoV-2.
- ▶ Disease severity was associated with increased BMI categories, being maximal in patients with BMI  $\geq 35$ .
- ▶ The need for invasive mechanical ventilation was associated with severe obesity and was independent of age, sex, diabetes, and hypertension.

### How might these results change the focus of clinical practice?

- ▶ Patients with obesity should avoid any COVID-19 contamination by enforcing all prevention measures during the current pandemic.
- ▶ Patients with severe obesity should be monitored more closely.
- ▶ Future studies should explore the mechanisms behind the association of SARS-CoV-2 with obesity.

<sup>1</sup> Department of Intensive Care, Centre Hospitalier Universitaire Lille, Lille, France. Correspondence: Francois Pattou (francois.pattou@univ-lille.fr) and Merce Jourdain (mercedes.jourdain@univ-lille.fr) <sup>2</sup> University of Lille, Inserm, Centre Hospitalier Universitaire Lille, Lille Pasteur Institute, U1190, European Genomic Institute for Diabetes, Lille, France <sup>3</sup> Centre Hospitalier Universitaire Lille, U2694 METRICS, University of Lille, Lille, France <sup>4</sup> Integrated Center for Obesity, Centre Hospitalier Universitaire Lille, Lille, France.

\*Arthur Simonnet and Mikael Chetboun contributed equally to this work.

© 2020 The Obesity Society. Received: 8 April 2020; Accepted: 8 April 2020; Published online 10 June 2020. doi:10.1002/oby.22831

blood (6). Furthermore, the abnormal secretion of adipokines and cytokines such as tumor necrosis factor- $\alpha$  and interferon characterize a chronic low-grade inflammation characteristic of abdominal obesity, which may impair immune response (7) and have effects on the lung parenchyma and bronchi (8). Altogether, it appears likely that obesity per se may be an independent risk factor for SARS-CoV-2 (9). COVID-19 pandemic is now rapidly spreading worldwide, notably in Europe and North America where obesity is highly prevalent (10). Exploring the relationship between obesity and the severity of the disease is therefore of major clinical importance. In this retrospective cohort, we aimed to investigate the association between BMI and clinical characteristics and the need for invasive mechanical ventilation in patients admitted to intensive care for SARS-CoV-2.

## Methods

### Study design and patients

In this single-center retrospective cohort study, we enrolled all consecutive patients admitted to intensive care for SARS-CoV-2 in Roger Salengro Hospital at "Centre Hospitalier Universitaire de Lille" (CHU Lille, France) between February 27, 2020, and April 5, 2020. CHU Lille is the tertiary referral center for 4.5 million inhabitants from two French departments (Nord and Pas de Calais). All patients were diagnosed with COVID-19 pneumonia according to World Health Organization interim guidance (11), with SARS symptoms characterized by dyspnea, increased respiratory frequency, decreased blood oxygen saturation, and need for oxygen support therapy for at least 6 L/min. Throat swab samples were obtained from all patients at admission and tested using real-time reverse transcriptase-polymerase chain reaction assays as previously described to identify SARS-CoV-2 infection (12). The study participants were also compared with a historical control group composed of 306 patients admitted to intensive care at our institution for a non-SARS-CoV-2 related severe acute respiratory disease during the year 2019.

### Data collection

A trained team of physicians reviewed and collected epidemiological data, past medical history, treatments, clinical data, and outcomes for all consecutive patients from their admission to April 6, 2020. This observational study was based on medical records, in strict compliance with the French reference methodology MR-004, established by the French National Commission on Informatics and Liberties, and approved by the institutional data protection authority of CHU Lille. Patient confidentiality was protected by assigning an anonymous identification code, and the electronic data were stored in a locked, password-protected computer.

### Study outcomes

The primary outcome of this study was the prevalence of patients receiving invasive mechanical ventilation (IMV) following admission to intensive care. The use of IMV was determined when oxygen therapy ( $\geq 10$  L/min) with target saturation pressure in O<sub>2</sub> (90-94%) was ineffective and when respiratory rate was above 25/min, with signs of acute respiratory failure despite maximal oxygen therapy. The following patient characteristics were analyzed: sex, age, weight, height, BMI, and history of diabetes, hypertension, and dyslipidemia. Patients were classified according to their BMI into the following four categories: lean (from 18.5 to  $<25$  kg/m<sup>2</sup>), overweight (from 25 to  $<30$

kg/m<sup>2</sup>), moderate obesity (from 30 to  $<35$  kg/m<sup>2</sup>), and severe obesity ( $\geq 35$  kg/m<sup>2</sup>).

### Statistical analysis

All results were expressed as median (interquartile range [IQR]) for continuous variables and as frequency (percentage) for categorical variables. Continuous variables were compared between groups with the *t* test or Mann-Whitney *U* test according to their distribution. Categorical variables were compared with the  $\chi^2$  (or Fisher exact) test. The association between BMI and the need for IMV as well as between pre-defined patient characteristics, including age, sex, history of diabetes, and hypertension, and the need for IMV was first assessed by univariate logistic regressions analysis. All variables were then included in multivariable logistic regression analysis to control the association of BMI with IMV for well-known predictors.  $P < 0.05$  was considered significant. All statistical analyses were performed with SAS Studio Statistics (version 3.71; SAS Institute Inc., Cary, North Carolina) and Prism GraphPad (version 8.1.2; GraphPad, San Diego, California).

## Results

A total of 124 patients were admitted in intensive care for SARS-CoV-2 during the study period and were all enrolled in the present study. At the time of analysis (April 6), 60 patients (48%) had been discharged alive from intensive care, 18 (15%) had died, and 46 remained hospitalized in intensive care. SARS-CoV-2 study participants were predominantly males (73%), and their median (IQR) age was 60 (51-70) years. Other clinical characteristics are detailed in Table 1. As illustrated in Figure 1, the distribution of BMI categories in SARS-CoV-2 participants differed markedly from the historical control group of patients admitted to intensive care for a non-SARS-CoV-2 severe acute respiratory disease (Figure 1A) ( $P < 0.001$ ; Fisher exact test for trend). Obesity (BMI  $\geq 30$ ) and severe obesity (BMI  $\geq 35$ ) were significantly more frequent among SARS-CoV-2 participants than in these non-SARS-CoV-2 controls (47.6% vs. 25.2% and 28.2% vs. 10.8%, respectively). The median (IQR) BMI in SARS-CoV-2 participants was higher than in non-SARS-CoV-2 controls (29.6 [26.4-36.5] vs. 24.0 [18.9-29.3], respectively;  $P < 0.0001$ , *t* test). In contrast, sex distribution and age were not significantly different from participants in non-SARS-CoV-2 controls (63.5% males,  $P = 0.08$  vs. SARS-CoV-2 patients, Fisher exact test; 63 [51-71] years,  $P = 0.97$  vs. SARS-CoV-2 patients, *t* test).


At the time of analysis, 85 out of the 124 study participants (68.6%) had required IMV, including 62 at admission, 13 at day 1, 4 at day 2, and the remaining 6 within 7 days. Their median (IQR) BMI was 31.1 (27.3-37.5) compared with 27.0 (25.3-30.8) in the 39 (31.4%) patients who did not require IMV ( $P < 0.001$ ). At the time of writing, 85 out of the 124 study participants (68.6%) had required IMV. Their BMI was higher than in the 39 (31.4%) patients who did not require IMV (31.1 [27.3-37.5] vs. 27 [25.3-30.8], respectively;  $P < 0.001$ , *t* test). As illustrated in Figure 1, the distribution of BMI categories differed markedly between the two subgroups ( $P < 0.01$ , Fisher exact test for trend); obesity (BMI  $> 30$ ) and severe obesity (BMI  $\geq 35$ ) were more frequent among patients who required IMV than among those who did not (56.4% vs. 28.2% and 35.3% vs. 12.8%, respectively) (Figure 1B).

In multivariate logistic regression analysis, the relation between male sex and BMI categories and the need for IMV remained significant after adjustment for age, diabetes, and hypertension (Table 2).

**TABLE 1** Baseline characteristics of 124 patients admitted in intensive care for SARS-CoV-2 who required invasive mechanical ventilation and those who did not

	All patients, <i>n</i> = 124	Invasive mechanical ventilation, <i>n</i> = 85	No invasive mechanical ventilation, <i>n</i> = 39
Male	90 (73)	64 (75)	26 (67)
Age (y)	60 (51-70)	60 (51-69)	60 (50-72)
Height (cm)	172 (166-178)	172 (166-178)	172 (165-180)
Weight (kg)	88 (80-108)	95 (81-112)	81 (75-94)
BMI (kg/m <sup>2</sup> )	29.6 (26.4-36.4)	31.1 (27.3-37.5)	27 (25.3-30.8)
Diabetes	28 (23)	23 (27)	5 (13)
Hypertension	60 (49)	48 (56)	12 (32)
Dyslipidemia	34 (28)	24 (28)	10 (26)

Results are expressed as median (IQR) for continuous variables and as frequency (percentage) for categorical variables.



**Figure 1** (A) Distribution of BMI categories in patients admitted during the study period in intensive care for SARS-CoV-2 (*n* = 124) and in patients admitted during the previous year in intensive care for a non-COVID-19 related severe pulmonary condition (*n* = 306) ( $\chi^2$  test for trend). (B) Distribution of BMI categories in patients who required mechanical ventilation (*n* = 85) and those who did not (*n* = 39) (*t* test). (C) Proportions of patients requiring mechanical ventilation during their stay in intensive care, according to BMI categories ( $\chi^2$  test for trend).

## Discussion

Our main finding was the unexpected high frequency of obesity among patients admitted to intensive care for SARS-CoV-2. Overall, 47.5% presented with obesity (BMI  $\geq 30$ ), including class II obesity (BMI 35-39.9) in 13.7% and class III obesity (BMI  $\geq 40$ ) in 14.5%. This distribution of BMI categories was markedly different than the distribution observed in control participants admitted during the previous years in

intensive care for severe acute pulmonary condition in the same institution. In these non-SARS-CoV-2 patients, the prevalence of obesity was only 25.8%, similar to the prevalence observed in the general population from Nord and Pas de Calais, when adjusted for age and sex (13). Importantly, we also showed that the need for IMV, a robust proxy for the severity of SARS-CoV-2, gradually increased with body mass categories, reaching nearly 90% in patients with BMI  $\geq 35$ . The main strength of the present study is its cohort design and the enrollment

**TABLE 2** Univariate and multivariate logistic regression analysis of the association between patient clinical characteristics and the need for invasive mechanical ventilation

	Univariate		Multivariate	
	Odds ratio (95% CI)	P	Odds ratio (95% CI)	P
Age (y)	1.00 (0.97-1.02)	0.73	1.00 (0.97-1.04)	0.87
Male	1.52 (0.67-3.49)	0.32	2.83 (1.02-7.85)	0.046
Diabetes vs. no	2.45 (0.85-7.03)	0.10	1.60 (0.44-5.83)	0.48
Hypertension vs. no	2.81 (1.25-6.3)	0.012	2.29 (0.89-5.84)	0.08
Dyslipidemia vs. no	1.10 (0.47-2.61)	0.83	0.68 (0.24-1.97)	0.48
BMI categories		0.023		0.049
25-30 vs. <25	1.72 (0.56-5.23)	0.22	1.69 (0.52-5.48)	0.22
30-35 vs. <25	3.38 (0.9-12.72)	0.45	3.45 (0.83-14.31)	0.48
≥35 vs. <25	6.75 (1.76-25.85)	0.015	7.36 (1.63-33.14)	0.021

Multivariate logistic regression analysis included all variables.

of all consecutive patients admitted in intensive care for SARS-CoV-2 during the study period. This single-center study also has limitations, such as its retrospective nature and the limited number of patients enrolled. The distinct role of other important predictors such as diabetes and hypertension (2,3), which did not appear to be independent from obesity in our cohort, might become significant in future studies enrolling larger numbers of patients. We had only a few younger patients (less than 30 years) and very old patients (more than 80 years) in our sample. Furthermore, we could not analyze the effect of BMI on the mortality rate because too few events were observed. Of note, many patients were still hospitalized at the time of this analysis. Taken together, our data demonstrate a distinct relation of obesity with the severity of SARS-CoV-2. Obesity is generally recognized as a risk factor for severe infection (7), as illustrated by the more severe disease of longer duration observed in patients with obesity during the influenza A H1N1 epidemic (7,14,15). Pulmonary function studies have also demonstrated a restrictive pattern and reduced lung volumes in individuals with obesity (16,17). Moreover, these pulmonary features evolved favorably in association with weight loss following bariatric surgery (18). Moreover, obesity and the metabolic syndrome are thought to increase type 2 inflammation, which may have effects on the lung parenchyma and bronchi (8). In addition, raised interleukin 6 levels are associated with obesity and/or metabolic syndrome (19). The abnormal secretion of adipokines and cytokines such as tumor necrosis factor-alpha and interferon characterize a chronic low-grade inflammation in abdominal obesity and they may induce an impaired immune response (7). Determining the cause of the relation between obesity and SRAS-CoV-2 goes beyond the scope of the present study and should now be the aim of future translational studies.

In conclusion, this cohort study showed that obesity is a factor in disease severity of SARS-CoV-2, having the greatest impact in patients with BMI ≥ 35. Patients with obesity and especially those with severe obesity should take extra measures to avoid COVID-19 contamination by enforcing prevention during the current pandemic. **O**

## Acknowledgments

The data sets generated during and/or analyzed during the current study are not publicly available, as they are subject to national data protection

laws and restrictions imposed by the institutional data protection authority of CHU Lille to ensure data privacy of the study participants.

## LICORN and Lille COVID-19 and Obesity study group

Enagnon Kazali Alidjinou (1)

Laurence Bocket (1)

Pauline Boddaert (2)

Robert Caizzo (3, 4)

Morgan Caplan (2)

Nicolas Cousin (2)

Thibault Duburcq (2)

Arthur Durand (2)

Ahmed El kalioubie (2)

Raphael Favory (2)

Bruno Garcia (2)

Patrick Girardie (2)

Julien Goutay (2)

Marion Houard (2)

Emmanuelle Jaillette (2)

Nicolas Kostuj (2)

Geoffrey Ledoux (2)

Daniel Mathieu (2)

Anne Sophie Moreau (2)

Olivier Nigeon (2)

Christopher Niles (2)

Saad Nseir (2)

Thierry Onimus (2)

Erika Parmentier (2)

Sebastien Préau (2)

Laurent Robriquet (2)

Anahita Rouze (2)

Sophie Six (2)

Claire Tinez (1)

Aurelia Toussaint (2)

Hélène Verkindt (3, 4)

1. CHU Lille, Department of Virology, Center of Biology and Pathology, 59000, Lille, France

2. CHU Lille, Department of Intensive Care, 59000, Lille, France

3. Inserm, Univ Lille, CHU Lille, Lille Pasteur Institute, European Genomic Institute for Diabetes, U1190, 59000, Lille France
4. CHU Lille, Integrated Center for Obesity, 59000, Lille, France

**Disclosure:** The authors declared no conflict of interest.

**Funding agencies:** This work was supported by grants from the European Commission (FEDER 12003944), Agence National de la Recherche (European Genomic Institute for Diabetes, E.G.I.D., ANR-10-LABX-46), and Foundation Coeur et Arteres (FCA R15112EE). The funding sources had part in patient recruitment and data collection.

## References

1. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020;395:1054-1062.
2. Li Q, Guan X, Wu P, et al. Early transmission dynamics in Wuhan, China, of novel coronavirus-infected pneumonia. *N Engl J Med* 2020;382:1199-1207.
3. Grasselli G, Zangrillo A, Zanella A, et al. Baseline characteristics and outcomes of 1591 patients infected with SARS-CoV-2 admitted to ICUs of the lombardy region, Italy. *JAMA* 2020;323:1574-1581.
4. Bhatraju PK, Ghassemieh BJ, Nichols M, et al. Covid-19 in critically ill patients in the Seattle region—case series. *N Engl J Med* 2020;382:2012-2022.
5. Kerkhove MDV, Vandemaële KAH, Shinde V, et al. Risk factors for severe outcomes following 2009 influenza A (H1N1) infection: a global pooled analysis. *PLoS Med* 2011;8:e1001053. doi:10.1371/journal.pmed.1001053
6. Peters U, Dixon AE. The effect of obesity on lung function. *Expert Rev Respir Med* 2018;12:755-767.
7. Huttunen R, Syrjänen J. Obesity and the risk and outcome of infection. *Int J Obes* 2013;37:333-340.
8. Zhang X, Zheng J, Zhang L, et al. Systemic inflammation mediates the detrimental effects of obesity on asthma control. *Allergy Asthma Proc* 2018;39:43-50.
9. Ryan DH, Ravussin E, Heymsfield S. COVID 19 and the patient with obesity – the editors speak out. *Obesity (Silver Spring)* 2020;28:847.
10. World Health Organization. *Overweight and obesity*. Geneva, Switzerland: World Health Organization; 2012.
11. World Health Organization. Clinical management of severe acute respiratory infection (SARI) when COVID-19 is suspected: interim guidance. <https://apps.who.int/iris/handle/10665/331446>. Published March 13, 2020. Accessed April 7, 2020.
12. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020;395:497-506.
13. Observatoire régional de la santé du social. Nutrition santé dans les Hauts-de-France – Surpoids, obésité et insuffisance pondérale. <http://www.or2s.fr/index.php/10-news/771-nutrition-sante-obesite>. Published June 2018. Accessed April 7, 2020.
14. Wang W, Chen H, Li Q, et al. Fasting plasma glucose is an independent predictor for severity of H1N1 pneumonia. *BMC Infect Dis* 2011;11:104. doi:10.1186/1471-2334-11-104
15. Honce R, Schultz-Cherry S. Impact of obesity on influenza a virus pathogenesis, immune response, and evolution. *Front Immunol* 2019;10:1071. doi:10.3389/fimmu.2019.01071
16. Watson RA, Pride NB, Thomas EL, et al. Reduction of total lung capacity in obese men: comparison of total intrathoracic and gas volumes. *J Appl Physiol* 2010;108:1605-1612.
17. Jones RL, Nzekwu M-MU. The effects of body mass index on lung volumes. *Chest* 2006;130:827-833.
18. Copley SJ, Jones LC, Soneji ND, et al. Lung parenchymal and tracheal CT morphology: evaluation before and after bariatric surgery. *Radiology* 2020;294:669-675.
19. Peters MC, McGrath KW, Hawkins GA, et al. Plasma interleukin-6 concentrations, metabolic dysfunction, and asthma severity: a cross-sectional analysis of two cohorts. *Lancet Respir Med* 2016;4:574-584.