

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

8-Chloro-4-oxo-4H-chromene-3-carbaldehyde

Yoshinobu Ishikawa

School of Pharmaceutical Sciences, University of Shizuoka, 52-1 Yada, Suruga-ku, Shizuoka 422-8526, Japan

Correspondence e-mail: ishi206@u-shizuoka-ken.ac.jp


Received 28 May 2014; accepted 28 May 2014

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.028; wR factor = 0.075; data-to-parameter ratio = 15.2.

In the title compound, $\text{C}_{10}\text{H}_5\text{ClO}_3$, a chlorinated 3-formylchromone derivative, all atoms are essentially coplanar (r.m.s. deviation = 0.032 Å for the non-H atoms), with the largest deviation from the least-squares plane [0.0598 (14) Å] being for a pyran-ring C atom. In the crystal, molecules are linked through stacking interactions along the b axis [shortest centroid-centroid distance between the pyran and benzene rings = 3.566 (2) Å].

Related literature

For related structures, see: Ishikawa & Motohashi (2013); Ishikawa (2014). For the synthesis of the precursor of the title compound, see: Fumagalli *et al.* (2012). For van der Waals radii; see: Bondi (1964). For halogen bonding, see: Auffinger *et al.* (2004); Metrangolo *et al.* (2005); Wilcken *et al.* (2013); Sirimulla *et al.* (2013).



Experimental

Crystal data

$\text{C}_{10}\text{H}_5\text{ClO}_3$
 $M_r = 208.60$
 Triclinic, $P\bar{1}$

$a = 6.9436$ (15) Å
 $b = 7.1539$ (17) Å
 $c = 9.165$ (2) Å

$\alpha = 102.049$ (19)°
 $\beta = 103.403$ (17)°
 $\gamma = 100.650$ (19)°
 $V = 419.89$ (18) Å³
 $Z = 2$

Mo $K\alpha$ radiation
 $\mu = 0.43$ mm⁻¹
 $T = 100$ K
 $0.38 \times 0.25 \times 0.10$ mm

Data collection

Rigaku AFC-7R diffractometer
 Absorption correction: ψ scan
 (North *et al.*, 1968)
 $T_{\min} = 0.902$, $T_{\max} = 0.958$
 2376 measured reflections
 1932 independent reflections

1750 reflections with $F^2 > 2\sigma(F^2)$
 $R_{\text{int}} = 0.011$
 3 standard reflections every 150 reflections
 intensity decay: -0.039%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.075$
 $S = 1.09$
 1932 reflections

127 parameters
 H-atom parameters constrained
 $\Delta\rho_{\max} = 0.34$ e Å⁻³
 $\Delta\rho_{\min} = -0.26$ e Å⁻³

Data collection: *WinAFC Diffractometer Control Software* (Rigaku, 1999); cell refinement: *WinAFC Diffractometer Control Software*; data reduction: *WinAFC Diffractometer Control Software*; program(s) used to solve structure: *SIR2008* (Burla *et al.*, 2007); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *CrystalStructure* (Rigaku, 2010); software used to prepare material for publication: *CrystalStructure*.

The University of Shizuoka is acknowledged for instrumental support.

Supporting information for this paper is available from the IUCr electronic archives (Reference: ZL2591).

References

- Auffinger, P., Hays, F. A., Westhof, E. & Ho, P. S. (2004). *Proc. Natl Acad. Sci. USA*, **101**, 16789–16794.
 Bondi, A. (1964). *J. Phys. Chem.* **68**, 441–451.
 Burla, M. C., Caliandro, R., Camalli, M., Carrozzini, B., Cascarano, G. L., De Caro, L., Giacovazzo, C., Polidori, G., Siliqi, D. & Spagna, R. (2007). *J. Appl. Cryst.* **40**, 609–613.
 Fumagalli, L., Pallavicini, M., Budriesi, R., Gobbi, M., Straniero, V., Zagami, M., Chiodini, G., Bolchi, C., Chiarini, A., Micucci, M. & Valoti, E. (2012). *Eur. J. Med. Chem.* **58**, 184–191.
 Ishikawa, Y. (2014). *Acta Cryst.* **E70**, o514.
 Ishikawa, Y. & Motohashi, Y. (2013). *Acta Cryst.* **E69**, o1416.
 Metrangolo, P., Neukirch, H., Pilati, T. & Resnati, G. (2005). *Acc. Chem. Res.* **38**, 386–395.
 North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
 Rigaku (1999). *WinAFC Diffractometer Control Software*. Rigaku Corporation, Tokyo, Japan.
 Rigaku (2010). *CrystalStructure*. Rigaku Corporation, Tokyo, Japan.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Sirimulla, S., Bailey, J. B., Vegesna, R. & Narayan, M. (2013). *J. Chem. Inf. Model.* **53**, 2781–2791.
 Wilcken, R., Zimmermann, M. O., Lange, A., Joerger, A. C. & Boeckler, F. M. (2013). *J. Med. Chem.* **56**, 1363–1388.

supplementary materials

Acta Cryst. (2014). E70, o743 [doi:10.1107/S1600536814012483]

8-Chloro-4-oxo-4*H*-chromene-3-carbaldehyde

Yoshinobu Ishikawa

1. Comment

Halogen bonds have been found to occur in organic, inorganic, and biological systems, and have recently attracted much attention in medicinal chemistry, chemical biology and supramolecular chemistry (Auffinger *et al.*, 2004, Metrangolo *et al.*, 2005, Wilcken *et al.*, 2013, Sirimulla *et al.*, 2013). We have recently reported the crystal structures of chlorinated 3-formylchromone derivatives 6,8-dichloro-4-oxochromene-3-carbaldehyde (Ishikawa & Motohashi, 2013, Fig.2 (top)) and 6-chloro-4-oxo-4*H*-chromene-3-carbaldehyde (Ishikawa, 2014, Fig.2 (middle)). It was found that a halogen bond is formed for 6,8-dichloro-4-oxochromene-3-carbaldehyde between the formyl oxygen atom and the chlorine atom at the 8-position, but none is formed for 6-chloro-4-oxo-4*H*-chromene-3-carbaldehyde between the formyl oxygen atom and the chlorine atom at the 6-position. As part of our interest in this type of chemical bonding, we herein report the crystal structure of a monochlorinated 3-formylchromone derivative 8-chloro-4-oxo-4*H*-chromene-3-carbaldehyde. The objective of this study is to reveal whether halogen bond(s) can be formed in the crystal of the title compound with the chlorine atom at 8-position and without a halogen atom at 6-position.

The mean deviation of the least-square planes for the non-hydrogen atoms is 0.0316 Å, and the largest deviation is 0.0598 (14) Å for C1. These mean that these atoms are essentially coplanar. In the crystal, the molecules are stacked with their inversion-symmetry equivalent along the *b*-axis direction [centroid–centroid distance between the pyran and benzene rings of the 4*H*-chromene units = 3.566 (2) Å, symmetry operator *i*: $-x + 1, -y + 1, -z + 2$], as shown in Fig.1.

The distance between the chlorine atom and the formyl oxygen atom of the translation-symmetry equivalent [C11...O3ⁱⁱ = 3.301 (2) Å, *ii*: $x, y, z + 2$] is nearly equal to the sum of their van der Waals radii [3.27 Å] (Bondi, 1964), as shown at the bottom of Fig.2. Thus, it is concluded that there is no halogen bond in the title compound. On the other hand, the angles of C–Cl...O (157.15 (6)°) and Cl...O=C (129.24 (10)°) are close to those of 6,8-dichloro-4-oxochromene-3-carbaldehyde, (C–Cl...O (160.4 (3)°) and Cl...O=C (138.7 (4)°), Fig.2(top)). Thus, the significance of the vicinal electron-withdrawing substituent in forming of a halogen bond (Wilcken *et al.*, 2013) is crystallographically validated from the fact that halogen bonding is observed in the dichlorinated 3-formylchromone, but is not observed in the monochlorinated ones. These results should be invaluable for rational drug design.

2. Experimental

2-Hydroxy-3-chloroacetophenone was prepared according to a literature method (Fumagalli *et al.*, 2012). To a solution of 2-hydroxy-3-chloroacetophenone (11.1 mmol) in *N,N*-dimethylformamide (30 ml) was added dropwise POCl₃ (27.7 mmol) for 5 min at 0 °C. After the mixture was stirred for 16 h at room temperature, water (50 ml) was added. The precipitates were collected, washed with water, and dried *in vacuo* (yield: 72%). ¹H NMR (400 MHz, DMSO-*d*₆): δ = 7.58 (t, 1H, *J* = 7.8 Hz), 8.07 (d, 1H, *J* = 7.8 Hz), 8.10 (d, 1H, *J* = 7.8 Hz), 9.03 (s, 1H), 10.12 (s, 1H). DART-MS calcd for [C₁₀H₅ClO₃ + H⁺]: 209.001, found 209.014. Single crystals suitable for X-ray diffraction were obtained by slow evaporation of a chloroform solution of the title compound at room temperature.

3. Refinement

The C(sp^2)-bound hydrogen atoms were placed in geometrical positions [C–H 0.95 Å, $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$], and refined using a riding model.





Figure 1

A packing view of the title compound, with displacement ellipsoids drawn at the 50% probability level. Hydrogen atoms are shown as small spheres of arbitrary radius.

**Figure 2**

Sphere models of the crystal structures of 6,8-dichloro-4-oxochromene-3-carbaldehyde (top), 6-chloro-4-oxo-4*H*-chromene-3-carbaldehyde (middle), and the title compound (bottom).

8-Chloro-4-oxo-4*H*-chromene-3-carbaldehyde*Crystal data* $C_{10}H_5ClO_3$ $M_r = 208.60$ Triclinic, $P\bar{1}$

Hall symbol: -P 1

 $a = 6.9436 (15) \text{ \AA}$ $b = 7.1539 (17) \text{ \AA}$ $c = 9.165 (2) \text{ \AA}$ $\alpha = 102.049 (19)^\circ$ $\beta = 103.403 (17)^\circ$ $\gamma = 100.650 (19)^\circ$ $V = 419.89 (18) \text{ \AA}^3$ $Z = 2$ $F(000) = 212.00$ $D_x = 1.650 \text{ Mg m}^{-3}$ Mo $K\alpha$ radiation, $\lambda = 0.71069 \text{ \AA}$

Cell parameters from 25 reflections

 $\theta = 15.1\text{--}17.5^\circ$ $\mu = 0.43 \text{ mm}^{-1}$

$T = 100$ K
Plate, yellow

$0.38 \times 0.25 \times 0.10$ mm

Data collection

Rigaku AFC-7R
diffractometer
 ω - 2θ scans
Absorption correction: ψ scan
(North *et al.*, 1968)
 $T_{\min} = 0.902$, $T_{\max} = 0.958$
2376 measured reflections
1932 independent reflections

1750 reflections with $F^2 > 2\sigma(F^2)$
 $R_{\text{int}} = 0.011$
 $\theta_{\text{max}} = 27.5^\circ$
 $h = -5 \rightarrow 9$
 $k = -9 \rightarrow 9$
 $l = -11 \rightarrow 11$
3 standard reflections every 150 reflections
intensity decay: -0.039%

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.075$
 $S = 1.09$
1932 reflections
127 parameters
0 restraints
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0344P)^2 + 0.1939P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 0.34 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.26 \text{ e } \text{\AA}^{-3}$

Special details

Refinement. Refinement was performed using all reflections. The weighted R -factor (wR) and goodness of fit (S) are based on F^2 . R -factor (gt) are based on F . The threshold expression of $F^2 > 2.0 \sigma(F^2)$ is used only for calculating R -factor (gt).

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C11	0.28721 (5)	0.37563 (5)	1.29528 (4)	0.02580 (11)
O1	0.21886 (13)	0.15266 (13)	0.97744 (10)	0.0187 (2)
O2	0.63950 (15)	0.10011 (16)	0.73295 (11)	0.0279 (3)
O3	0.04173 (16)	-0.20711 (16)	0.52591 (12)	0.0297 (3)
C1	0.16551 (19)	0.04262 (19)	0.82917 (14)	0.0189 (3)
C2	0.29522 (19)	0.01959 (19)	0.74169 (14)	0.0184 (3)
C3	0.51287 (19)	0.11867 (19)	0.80487 (14)	0.0184 (3)
C4	0.76947 (18)	0.35541 (19)	1.04110 (15)	0.0180 (3)
C5	0.81834 (19)	0.47543 (19)	1.19027 (15)	0.0189 (3)
C6	0.66911 (19)	0.48195 (18)	1.26931 (14)	0.0183 (3)
C7	0.47153 (19)	0.37036 (19)	1.19772 (14)	0.0176 (3)
C8	0.56861 (18)	0.24423 (18)	0.96597 (14)	0.0161 (3)
C9	0.41997 (18)	0.25375 (17)	1.04458 (14)	0.0157 (3)
C10	0.2152 (2)	-0.1091 (2)	0.58050 (15)	0.0236 (3)
H1	0.0257	-0.0238	0.7828	0.0227*
H2	0.8725	0.3481	0.9891	0.0216*
H3	0.3054	-0.1145	0.5169	0.0284*
H4	0.9536	0.5537	1.2392	0.0226*
H5	0.7035	0.5632	1.3724	0.0220*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl1	0.01652 (16)	0.0338 (2)	0.02036 (17)	-0.00018 (12)	0.00735 (12)	-0.00404 (13)
O1	0.0116 (4)	0.0215 (5)	0.0159 (5)	-0.0032 (4)	0.0015 (4)	-0.0009 (4)
O2	0.0227 (5)	0.0370 (6)	0.0190 (5)	-0.0015 (5)	0.0093 (4)	0.0010 (4)
O3	0.0252 (6)	0.0309 (6)	0.0200 (5)	-0.0059 (5)	0.0003 (4)	-0.0027 (4)
C1	0.0158 (6)	0.0182 (6)	0.0159 (6)	-0.0030 (5)	-0.0003 (5)	0.0014 (5)
C2	0.0187 (6)	0.0174 (6)	0.0138 (6)	-0.0019 (5)	0.0009 (5)	0.0026 (5)
C3	0.0182 (6)	0.0197 (6)	0.0144 (6)	-0.0004 (5)	0.0032 (5)	0.0041 (5)
C4	0.0134 (6)	0.0200 (6)	0.0183 (6)	-0.0005 (5)	0.0038 (5)	0.0047 (5)
C5	0.0132 (6)	0.0200 (6)	0.0185 (6)	-0.0012 (5)	0.0004 (5)	0.0039 (5)
C6	0.0173 (6)	0.0177 (6)	0.0147 (6)	0.0006 (5)	0.0005 (5)	0.0006 (5)
C7	0.0147 (6)	0.0196 (6)	0.0165 (6)	0.0019 (5)	0.0044 (5)	0.0026 (5)
C8	0.0147 (6)	0.0169 (6)	0.0140 (6)	-0.0002 (5)	0.0025 (5)	0.0039 (5)
C9	0.0116 (6)	0.0154 (6)	0.0157 (6)	-0.0011 (5)	0.0004 (5)	0.0026 (5)
C10	0.0248 (7)	0.0257 (7)	0.0138 (6)	-0.0017 (6)	0.0024 (5)	0.0014 (5)

Geometric parameters (\AA , $^\circ$)

Cl1—C7	1.7243 (16)	C4—C8	1.4029 (16)
O1—C1	1.3475 (15)	C5—C6	1.397 (2)
O1—C9	1.3763 (14)	C6—C7	1.3817 (17)
O2—C3	1.2250 (19)	C7—C9	1.4006 (17)
O3—C10	1.2061 (16)	C8—C9	1.393 (2)
C1—C2	1.347 (2)	C1—H1	0.950
C2—C3	1.4658 (17)	C4—H2	0.950
C2—C10	1.4836 (17)	C5—H4	0.950
C3—C8	1.4797 (17)	C6—H5	0.950
C4—C5	1.3815 (18)	C10—H3	0.950
Cl1...O1	2.8973 (12)	C3...H1	3.2929
O1...C3	2.8719 (19)	C3...H2	2.6746
O2...C1	3.574 (2)	C3...H3	2.7084
O2...C4	2.8604 (17)	C4...H5	3.2636
O2...C10	2.9089 (18)	C6...H2	3.2648
O3...C1	2.8120 (17)	C7...H4	3.2634
C1...C7	3.5981 (19)	C8...H4	3.2730
C1...C8	2.7591 (18)	C9...H1	3.1860
C2...C9	2.7695 (18)	C9...H2	3.2689
C4...C7	2.783 (2)	C9...H5	3.2672
C5...C9	2.7806 (18)	C10...H1	2.5482
C6...C8	2.7921 (18)	H1...H3	3.4825
Cl1...O2 ⁱ	3.4989 (15)	H2...H4	2.3282
Cl1...O3 ⁱⁱ	3.3012 (15)	H4...H5	2.3459
Cl1...C5 ⁱⁱⁱ	3.4247 (16)	Cl1...H1 ⁱⁱ	2.8415
O1...O1 ⁱⁱ	3.5617 (16)	Cl1...H2 ⁱⁱⁱ	3.4669
O1...O2 ⁱ	3.5683 (17)	Cl1...H4 ⁱⁱⁱ	2.8395
O1...C3 ⁱ	3.5282 (19)	Cl1...H5 ^x	2.9688
O1...C4 ^{iv}	3.5456 (19)	O1...H1 ⁱⁱ	3.2499

O1...C5 ^{iv}	3.359 (2)	O1...H2 ⁱⁱⁱ	3.0086
O1...C8 ⁱ	3.5096 (19)	O1...H4 ^{iv}	3.3704
O2...C11 ⁱ	3.4989 (15)	O2...H1 ^{viii}	2.9439
O2...O1 ⁱ	3.5683 (17)	O2...H3 ^v	2.4269
O2...C7 ⁱ	3.534 (2)	O2...H4 ^{xi}	3.3161
O2...C9 ⁱ	3.591 (2)	O3...H1 ^{vi}	3.5460
O2...C10 ^v	3.267 (2)	O3...H4 ^{vii}	2.6830
O3...C11 ⁱⁱ	3.3012 (15)	O3...H5 ^{vii}	2.5041
O3...O3 ^{vi}	3.2307 (19)	O3...H5 ⁱ	3.5184
O3...C5 ^{vii}	3.2551 (18)	C1...H2 ⁱⁱⁱ	3.5714
O3...C6 ^{vii}	3.1687 (17)	C1...H2 ⁱ	3.5400
O3...C6 ⁱ	3.560 (2)	C1...H4 ^{iv}	3.2889
O3...C10 ^{vi}	3.295 (2)	C2...H5 ^{iv}	3.3614
C1...C4 ⁱ	3.371 (3)	C3...H3 ^v	3.4629
C1...C5 ^{iv}	3.472 (3)	C3...H5 ^{iv}	3.4282
C1...C8 ⁱ	3.542 (3)	C4...H1 ⁱ	3.4694
C2...C6 ^{iv}	3.553 (3)	C4...H2 ^{xi}	3.0614
C2...C7 ⁱ	3.552 (3)	C5...H2 ^{xi}	3.2259
C2...C9 ⁱ	3.578 (2)	C6...H3 ⁱ	3.5976
C3...O1 ⁱ	3.5282 (19)	C10...H4 ^{vii}	3.3912
C3...C6 ^{iv}	3.460 (3)	C10...H5 ⁱ	3.4945
C3...C7 ⁱ	3.515 (3)	H1...C11 ⁱⁱ	2.8415
C3...C9 ⁱ	3.304 (3)	H1...O1 ⁱⁱ	3.2499
C4...O1 ^{iv}	3.5456 (19)	H1...O2 ⁱⁱⁱ	2.9439
C4...C1 ⁱ	3.371 (3)	H1...O3 ^{vi}	3.5460
C4...C7 ^{iv}	3.570 (3)	H1...C4 ⁱ	3.4694
C4...C9 ^{iv}	3.459 (3)	H1...H2 ⁱⁱⁱ	3.4159
C5...C11 ^{viii}	3.4247 (16)	H1...H2 ⁱ	3.4928
C5...O1 ^{iv}	3.359 (2)	H1...H3 ^{vi}	3.5853
C5...O3 ^{ix}	3.2551 (18)	H1...H4 ^{iv}	3.3899
C5...C1 ^{iv}	3.472 (3)	H2...C11 ^{viii}	3.4669
C5...C9 ^{iv}	3.521 (2)	H2...O1 ^{viii}	3.0086
C6...O3 ^{ix}	3.1687 (17)	H2...C1 ^{viii}	3.5714
C6...O3 ⁱ	3.560 (2)	H2...C1 ⁱ	3.5400
C6...C2 ^{iv}	3.553 (3)	H2...C4 ^{xi}	3.0614
C6...C3 ^{iv}	3.460 (3)	H2...C5 ^{xi}	3.2259
C6...C8 ^{iv}	3.538 (2)	H2...H1 ^{viii}	3.4159
C6...C10 ⁱ	3.387 (3)	H2...H1 ⁱ	3.4928
C7...O2 ⁱ	3.534 (2)	H2...H2 ^{xi}	2.4762
C7...C2 ⁱ	3.552 (3)	H2...H4 ^{xi}	2.7931
C7...C3 ⁱ	3.515 (3)	H3...O2 ^v	2.4269
C7...C4 ^{iv}	3.570 (3)	H3...C3 ^v	3.4629
C7...C8 ^{iv}	3.427 (3)	H3...C6 ⁱ	3.5976
C8...O1 ⁱ	3.5096 (19)	H3...H1 ^{vi}	3.5853
C8...C1 ⁱ	3.542 (3)	H3...H3 ^v	3.0081
C8...C6 ^{iv}	3.538 (2)	H3...H4 ^{vii}	3.2450
C8...C7 ^{iv}	3.427 (3)	H3...H5 ⁱ	3.5572
C8...C9 ⁱ	3.560 (2)	H4...C11 ^{viii}	2.8395
C8...C9 ^{iv}	3.600 (2)	H4...O1 ^{iv}	3.3704

C9...O2 ⁱ	3.591 (2)	H4...O2 ^{xi}	3.3161
C9...C2 ⁱ	3.578 (2)	H4...O3 ^{ix}	2.6830
C9...C3 ⁱ	3.304 (3)	H4...C1 ^{iv}	3.2889
C9...C4 ^{iv}	3.459 (3)	H4...C10 ^{ix}	3.3912
C9...C5 ^{iv}	3.521 (2)	H4...H1 ^{iv}	3.3899
C9...C8 ⁱ	3.560 (2)	H4...H2 ^{xi}	2.7931
C9...C8 ^{iv}	3.600 (2)	H4...H3 ^{ix}	3.2450
C10...O2 ^v	3.267 (2)	H5...C11 ^x	2.9688
C10...O3 ^{vi}	3.295 (2)	H5...O3 ^{ix}	2.5041
C10...C6 ⁱ	3.387 (3)	H5...O3 ⁱ	3.5184
C11...H5	2.8072	H5...C2 ^{iv}	3.3614
O2...H2	2.5915	H5...C3 ^{iv}	3.4282
O2...H3	2.6355	H5...C10 ⁱ	3.4945
O3...H1	2.4818	H5...H3 ⁱ	3.5572
C1...H3	3.2782		
C1—O1—C9	118.02 (11)	C4—C8—C9	119.19 (11)
O1—C1—C2	125.01 (11)	O1—C9—C7	117.14 (12)
C1—C2—C3	120.64 (11)	O1—C9—C8	122.57 (10)
C1—C2—C10	119.09 (11)	C7—C9—C8	120.29 (11)
C3—C2—C10	120.26 (13)	O3—C10—C2	123.51 (15)
O2—C3—C2	123.90 (11)	O1—C1—H1	117.494
O2—C3—C8	122.20 (11)	C2—C1—H1	117.497
C2—C3—C8	113.90 (12)	C5—C4—H2	119.817
C5—C4—C8	120.37 (13)	C8—C4—H2	119.815
C4—C5—C6	120.11 (11)	C4—C5—H4	119.941
C5—C6—C7	120.09 (11)	C6—C5—H4	119.944
C11—C7—C6	120.35 (10)	C5—C6—H5	119.956
C11—C7—C9	119.75 (10)	C7—C6—H5	119.956
C6—C7—C9	119.90 (13)	O3—C10—H3	118.246
C3—C8—C4	121.00 (13)	C2—C10—H3	118.242
C3—C8—C9	119.81 (10)		
C1—O1—C9—C7	178.77 (11)	C8—C4—C5—C6	2.1 (3)
C1—O1—C9—C8	-0.80 (18)	C8—C4—C5—H4	-177.9
C9—O1—C1—C2	1.7 (2)	H2—C4—C5—C6	-177.9
C9—O1—C1—H1	-178.3	H2—C4—C5—H4	2.1
O1—C1—C2—C3	-0.5 (3)	H2—C4—C8—C3	-1.5
O1—C1—C2—C10	179.02 (12)	H2—C4—C8—C9	178.9
H1—C1—C2—C3	179.5	C4—C5—C6—C7	-0.9 (2)
H1—C1—C2—C10	-1.0	C4—C5—C6—H5	179.1
C1—C2—C3—O2	178.13 (14)	H4—C5—C6—C7	179.1
C1—C2—C3—C8	-1.5 (2)	H4—C5—C6—H5	-0.9
C1—C2—C10—O3	-5.7 (3)	C5—C6—C7—C11	179.30 (12)
C1—C2—C10—H3	174.3	C5—C6—C7—C9	-1.3 (2)
C3—C2—C10—O3	173.77 (13)	H5—C6—C7—C11	-0.7
C3—C2—C10—H3	-6.2	H5—C6—C7—C9	178.7
C10—C2—C3—O2	-1.3 (3)	C11—C7—C9—O1	2.11 (18)
C10—C2—C3—C8	179.06 (12)	C11—C7—C9—C8	-178.31 (9)

O2—C3—C8—C4	3.0 (3)	C6—C7—C9—O1	-177.27 (12)
O2—C3—C8—C9	-177.39 (13)	C6—C7—C9—C8	2.3 (2)
C2—C3—C8—C4	-177.42 (12)	C3—C8—C9—O1	-1.2 (2)
C2—C3—C8—C9	2.21 (19)	C3—C8—C9—C7	179.27 (11)
C5—C4—C8—C3	178.54 (12)	C4—C8—C9—O1	178.46 (12)
C5—C4—C8—C9	-1.1 (2)	C4—C8—C9—C7	-1.1 (2)

Symmetry codes: (i) $-x+1, -y, -z+2$; (ii) $-x, -y, -z+2$; (iii) $x-1, y, z$; (iv) $-x+1, -y+1, -z+2$; (v) $-x+1, -y, -z+1$; (vi) $-x, -y, -z+1$; (vii) $x-1, y-1, z-1$; (viii) $x+1, y, z$; (ix) $x+1, y+1, z+1$; (x) $-x+1, -y+1, -z+3$; (xi) $-x+2, -y+1, -z+2$.