

Review

Open Access

Expected values for pedometer-determined physical activity in older populations

Catrine Tudor-Locke*¹, Teresa L Hart² and Tracy L Washington²

Address: ¹Walking Behavior Laboratory, Pennington Biomedical Research Center, Baton Rouge, LA 70808, USA and ²Department of Exercise and Wellness, Arizona State University, Mesa, AZ 85212, USA

Email: Catrine Tudor-Locke* - Tudor-Locke@pbr.edu; Teresa L Hart - harttl@uwm.edu; Tracy L Washington - washington.tracy@gmail.com

* Corresponding author

Published: 25 August 2009

Received: 5 February 2009

International Journal of Behavioral Nutrition and Physical Activity 2009, **6**:59 doi:10.1186/1479-5868-6-59

Accepted: 25 August 2009

This article is available from: <http://www.ijbnpa.org/content/6/1/59>

© 2009 Tudor-Locke et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

The purpose of this review is to update expected values for pedometer-determined physical activity in free-living healthy older populations. A search of the literature published since 2001 began with a keyword (pedometer, "step counter," "step activity monitor" or "accelerometer AND steps/day") search of PubMed, Cumulative Index to Nursing & Allied Health Literature (CINAHL), SportDiscus, and PsychInfo. An iterative process was then undertaken to abstract and verify studies of pedometer-determined physical activity (captured in terms of steps taken; distance only was not accepted) in free-living adult populations described as ≥ 50 years of age (studies that included samples which spanned this threshold were not included unless they provided at least some appropriately age-stratified data) and not specifically recruited based on any chronic disease or disability. We identified 28 studies representing at least 1,343 males and 3,098 females ranging in age from 50–94 years. Eighteen (or 64%) of the studies clearly identified using a Yamax pedometer model. Monitoring frames ranged from 3 days to 1 year; the modal length of time was 7 days (17 studies, or 61%). Mean pedometer-determined physical activity ranged from 2,015 steps/day to 8,938 steps/day. In those studies reporting such data, consistent patterns emerged: males generally took more steps/day than similarly aged females, steps/day decreased across study-specific age groupings, and BMI-defined normal weight individuals took more steps/day than overweight/obese older adults. The range of 2,000–9,000 steps/day likely reflects the true variability of physical activity behaviors in older populations. More explicit patterns, for example sex- and age-specific relationships, remain to be informed by future research endeavors.

Introduction

The recently released U.S. Physical Activity Guidelines <http://www.health.gov/paguidelines> were informed by an advisory committee report that concluded that, for older adults, in addition to the well-know health benefits of a physically active lifestyle, "strong evidence indicates that being physically active is associated with higher levels of functional health, a lower risk of falling, and better cognitive functioning" [1]. Although we have a collective under-

standing that physical activity declines with age [2], we know little about the actual physical activity patterns of older adults. Accurate and precise quantification of physical activity behaviors is elemental to epidemiologists, physiologists, and behavioral scientists, as well as clinicians engaged in efforts to extend quality years of life and ultimately compress morbidity. In addition, researchers and practitioners must be prepared with practicable and appropriate assessment tools, including becoming com-

petent in the interpretation of their data, if they are to successfully address measurement challenges related to surveillance, screening, intervention, and program evaluation.

Motion sensor technology has evolved quickly in recent years, producing a wide variety of commercially available body worn instrumentation, including accelerometers and pedometers, capable of objectively detecting physical activity behaviors. Although accelerometers can offer minute-by-minute estimates of physical activity volume and intensity, their relatively high costs, data management demands, and tandem requisites for personnel time and technical expertise stand as a barrier to widespread implementation outside formal research studies. In contrast, simple and affordable pedometers are generally considered more practical for individual, clinical, and population level applications. In addition, pedometer outputs correlate highly with that of different accelerometers [3]. Although pedometers are not designed to directly detect physical activity intensity, they do provide a feasible and practical means of capturing daily physical activity volume (especially walking) expressed as a summary output of steps/day. Walking for exercise increases with age and it is also a fundamental behavior in daily life activities [4]. An indicator of volume is likely sufficient when studying older populations, who do not typically engage in high levels of physical activity intensity. Based on accelerometer data, Troiano et al. [5] reported that 50–59 year olds average 1.1 minutes daily in vigorous intensity activities, and 60–69 years olds and those over 70 years of age, respectively average 0.4 and 0.1 minutes in vigorous activity daily.

Pedometers are considered analogous to computer hardware (e.g., keyboards, monitors, disks, etc.); without the matching software (e.g., expected values, etc.) they are of limited utility [6]. Expected values are normative or benchmark values necessary for interpreting change and comparison purposes [7]. In 2001, we compiled expected values of free-living pedometer-determined physical activity in adults [8]. At that time (and based on only 4 studies [9-12]) we concluded that healthy older adults (≥ 50 years without chronic disease) accumulated 6,000–8,500 steps/day. Since then, pedometry as a scholarly and applied practice has burgeoned and the related scientific literature is again ripe for a comparative examination. Therefore, the purpose of this review is to present expected values for pedometer-determined physical activity published since the original review paper, pertinent to free-living healthy older populations.

Methods for Literature Review

A search of the literature (current and verified as of January 05, 2009) began with a keyword (pedometer, "step counter," "step activity monitor" or "accelerometer AND

steps/day") search of PubMed, Cumulative Index to Nursing & Allied Health Literature (CINAHL), SportDiscus, and PsychInfo. A publication date limit was set from 2001, coinciding with the year of publication of the original review [8]. All authors then participated in an iterative process of identifying and verifying studies of free-living pedometer-determined physical activity (captured in terms of steps taken; distance only was not accepted) in older adult populations described as ≥ 50 years of age (studies that included samples which spanned this threshold were not included unless they provided at least some appropriately age-stratified data) and not specifically recruited based on any chronic disease or disability. We excluded one study that presented pedometer data only in a figure form that required interpretation to extract data [13]; however, we decided to include a study by Moreau et al. [14] that appeared to report rounded values for steps/day rather than exact numbers. We included the single report with the largest sample size [15] reporting steps/day from the Nakanojo Study papers that used pedometers to assess physical activity in older adults. We also excluded two other papers [16,17] after confirming with authors (Strath and Swartz, personal communication) their overlap with other papers from the same group. Finally, since the explicit focus of this review was on healthy older populations we also excluded two free-living studies [18,19] and a case study [20] of older adults living in assisted living facilities. The majority of these residents walk at speeds less than 0.6 meters/second [21], a speed which produces lower gravitational forces than are typically detected by research-quality pedometers [19,22,23].

Once the studies were assembled, we collated them according to the original reference, country where study was conducted, sample descriptions (including sex, age descriptors as available, and verbatim additional descriptions of the sample, if provided), verbatim descriptions of pedometer brands used, monitoring frame (i.e., number of days worn), whether the instrument was sealed or unsealed (explicitly stated or implicit from requiring participants to self-record data) to the participant, and mean (or median if mean not reported) and standard deviation or standard error of steps/day for the total sample and/or stratified groups, as available. We also looked for any data treatment descriptions including identifying extreme values or addressing missing data. In the case of intervention or other longitudinal-type studies, only the baseline data were considered. Although in previous reviews [24,25], an overall median steps/day was imputed considering those studies originally reporting mean values, we determined that it was overly reductionistic to attempt to represent all older adults by a single value for pedometer-determined physical activity. Instead we offer the studies' data as published and only capture their range as an indicator of distribution.

Findings

We identified 28 unique studies of pedometer-determined free-living physical activity in healthy older adult populations representing at least 1,343 males and 3,098 females (another 568 older adults were not identified by sex in the original studies [26-29]) ranging in age from 50–94 years. All are assembled by year of publication [see Additional file 1]. Inconsistencies in presentation across table columns reflect underlying inconsistencies in reporting across studies. Eighteen studies (64%) were conducted in the United States, 5 in Japan (18%), 2 in Australia (7%), and 1 each in Thailand and Belgium. A final study was a secondary analysis of data assembled from a number of different countries. Eighteen (or 64%) of the studies clearly identified using a Yamax pedometer model (i.e., Yamasa, Yamax, Digi-walker). Monitoring frames (where reported) ranged from 3 days to 1 year; the modal length of time was 7 days (17 studies, or 61%). Only 5 studies (15%) indicated that they used a sealed pedometer (1 study unsealed the pedometer one week, and sealed it the next). Mean pedometer-determined physical activity ranged from a low of 2,015 steps/day (in one sample of 85+ year olds) [30] to a high of 8,938 steps/day in a sample described as normal weight males, aged 51–88 years [31].

Four studies offered pedometer data by sex [31-34]. In all but one case (males vs. females 50–59.9 years of age [34]) males took more steps/day than females, ranging from a low difference of 497 steps/day [33] to a high of 1,450 steps/day [32] for similarly aged groups. A clear decline in steps/day within study-specific age groupings was consistently apparent in the four studies that provided age-stratified pedometer data [27,30,31,34]. Two studies examined pedometer data by BMI-defined weight status categories [31,35]; individuals classified as normal weight ($BMI \leq 25 \text{ kg/m}^2$) took between 1,659 [31] and 4,060 steps/day [35] more than individuals classified as overweight/obese. Three studies [28,36,37] also provided at least some pedometer data stratified by race/ethnicity. In each study, samples identified as White or Caucasian took more steps/day than other race/ethnic groupings considered.

Conclusion

This review represents an extension of our earlier efforts to inform expected values for free-living pedometer-determined physical activity, previously focused on general adult populations [8] and youth (including children and adolescents) [25]. Older adults embody a special population that can derive additional benefits from a physically active lifestyle [1]. The data assembled here, and the amalgamation of expected values ultimately distilled, are useful to clinicians and researchers alike. As previously stated, accurate quantification of physical activity behaviors is an

important requisite for surveillance, screening, intervention and program evaluation, as well as more basic research questions.

In our first attempt to assemble expected values for different populations [8], we concluded that healthy older adults take between 6,000–8,500 steps/day. This earlier conclusion was based on only four studies with limited sample sizes. We acknowledged at that time that expected values should be refined as more pedometer studies emerged. Although we did impute overall median values for other similar reviews (considering underlying studies reporting original mean values) [25], we believed that, in the case of older adults as a broadly defined population, a single value for steps/day fails to capture the fact that older adults range in ability and habit with age as well as other factors. In recognition of this phenomenon, we also conclude that the range of 2,000–9,000 steps/day more likely reflects the true variability of physical activity behaviors in healthy older populations. We believe that it is prudent to designate rounded anchors for the ranges of such expected values (to not overstate precision), and the use of 1,000 steps/day increment is rationale [31]. One thousand steps are approximately equivalent to 10 minutes of brisk walking in healthy adults [38].

Pedometer-determined physical activity has been shown to be inversely related to age [39]. In our previous review of pedometer-determined physical activity for youth [25], we were able to construct a graph of sex- and age-specific expected values from age 6 to 18 years. Herein we observed a clear decline in steps/day in those studies reporting age-stratified data in older adults [27,30,31,34], however, the study-specific age groupings were broad and ultimately were not comparable between studies precluding an opportunity to draw more explicit conclusions about expected values for different ages of older adults. In order to extend and complete the step/day curve that has been initiated in young populations [25], researchers should continue to provide both sex- and age-specific data to the extent that it is possible in their original publications.

Most studies of free-living pedometer-determined older adult physical activity have been conducted in the United States, but none of the assembled studies can be considered nationally representative. The most frequently used pedometer was different models of the Yamax brand, manufactured in Japan and available through a variety of distributors around the world. In a review of literature focused on youth pedometer studies [40], we found that 30 of 34 (88%) studies identified used the Yamax pedometer. The Yamax pedometer is considered to be a criterion pedometer, against which others may be compared [41]. However, its reduced sensitivity to low force ambulation

is well known [42,43], and there are concerns that these pedometers miss steps in gait-impaired elderly [23]. Other pedometers and accelerometers offer reduced sensitivity thresholds, capable of detecting such low force movements [19,44,45]. However, the inevitable sensitivity-specificity trade-off can result in the over-detection of erroneous steps [44,45] and complicates comparison and amalgamation of data between studies, the central focus of this review. Further, instrumentation that is greatly discrepant from established research quality pedometers would require its own normative data for interpretation purposes. Even if low force ambulation is missed, however, the steps that are detected are considered to be above 0.35 G (i.e., the manufacturer-reported force threshold for the Yamax pedometer) [46], offering an unyielding standard that ultimately simplifies comparisons of the same measurement across populations and studies. Although it remains plausible that at least some age-related decline in steps/day is complicated by slower gaits with advanced age and disability, the pedometer-determined pattern displayed herein is congruent with what we know from other assessments of physical activity in these populations, including doubly labeled water [47,48]. Ultimately, the choice of the instrument used is a function of the research question, participant burden, and resources available to the researcher or practitioner.

As noted above, reported monitoring frames ranged from 3 days to 1 year although the most frequently reported length of time was 7 days. Three days is considered sufficient to capture habitual pedometer-determined physical activity (defined as the mean of 7 days) in healthy adult populations [49,50]. Sedentary populations with chronic illnesses appear to require even fewer days to obtain a stable estimate of habitual physical activity [51,52]. Likewise, Rowe et al. [53] studied older adults (over 60 years of age) and reported that between-day reliability was high (ICC = .90 for Yamax steps and .87 for ActiGraph steps) and that likely only 2 days are sufficient to obtain a stable estimate. We conservatively refrained from conducting any analyses to examine the relationship between monitoring frame and step count since it would likely be confounded by study differences in sample characteristics, sample sizes, instrumentation, assessment protocols, etc.

Relatively few studies used sealed pedometers to blind participants to their step counts. Since research participants can typically view unsealed pedometer data displays and observe how many steps they are accumulating, there is a concern that certain individuals may set out to accrue more steps, either because they are trying to please the researcher or because they are naturally motivated to achieve. Marshall et al. [54] examined this question in older adults by comparing data collected under unsealed and subsequently sealed conditions. Participants accumulated 400 more steps/day ($p = 0.02$) in the unsealed con-

dition. The authors concluded that although the difference in steps/day was statistically significant, it might not be considered clinically significant. Put in context, this magnitude of difference attributed to reactivity is substantially less than what has come to be expected of successful pedometer-based interventions [55,56].

There was largely no reporting whatsoever of data treatment regarding missing data or extreme values. King et al. [57] indicated that "when a participant returned her activity diary with missing data, she was asked to start a new diary and wear the pedometer for another week." Fukukawa et al. [58] indicated that "to estimate the participants' usual walking activity, we discarded the maximum and minimum daily records from the entire data." Tudor-Locke et al. [34] wrote that "Data for any single day indicating < 1,000 steps were removed and values > 30,000 steps on any single day were truncated (i.e., replaced with 30,000 steps)." No other study described any other treatment for missing data or extreme values.

We located only two longitudinal studies of older adult pedometer-determined physical activity at this time [34,58]. Fukukawa et al. [58] collected two waves of data from 314 older adults (65–79 years of age) separated by two years. Values increased 1,483 steps/day over time. The participants were not part of an intervention and the authors did not offer any explanation for the observed increase. Another sample that included older adults (e.g., 50–71 years of age) was followed before and one year after they moved residences within Perth, Australia [34]. An overall small decrease in steps/day was evident with mean values ranging from -100 steps in 50–59.9 year old males and -776 steps in 60+ year old males to -407 in 50–59.9 year old females and -408 steps in 60+ year old females. Spearman rank order correlations were computed to assess relative stability of behavior over time. Males and females ranging in age from 50–59.9 years were moderately stable over one year ($r = 0.533$ and 0.645 , respectively). Males aged 60+ years were also moderately stable ($r = 0.542$), however, the oldest females were less stable over time ($r = 0.304$).

It is important to stress that the expected values assembled herein do not envisage what older adults "should" be taking (i.e., an "indicator," "cut point," or "threshold" value). A recent analysis identified preliminary steps/day cut points for adults that best discriminated between BMI-defined normal weight and overweight/obesity [31]. Simultaneous consideration of a number of different indices produced estimates of 11,000 steps/day for males age 51–88 years (based on $n = 220$), 10,000 steps/day for females aged 50–59.9 years ($n = 366$), and 8,000 for females aged 60–94 years ($n = 214$). These values are obviously much higher than most of the expected values collected herein [see Additional file 1], and reflect the fact

that expected values are frequently lower than criterion-based values [31]. Rowe et al. [53] conducted Receiver Operator Curve analysis to identify steps/day associated with achieving 30 minutes of continuous moderate-to-vigorous physical activity (typically associated with public health recommendations for physical activity) by older adults (mean age 74 years). An estimate of 7,000 to 8,000 steps/day was determined. It is important to emphasize that the first study concerned itself with steps associated with a healthy BMI and the second one focused on time in moderate-to-vigorous physical activity; these are distinct measurement criteria used to evaluate "how many steps are enough?" There are no other studies at this time that have attempted to address this question for older adults in particular.

In summary, expected values for pedometer-determined physical activity are necessary to enable research and program planning and aid in interpretation of similar data. The literature focused on older adults has expanded since an initial review published in 2001 [8], presenting a timely opportunity to amass comparable parameters and refine expected values. We located 28 studies of pedometer-determined free-living physical activity in healthy older adults ranging in age from 50–94 years. Steps/day ranged from 2,000–9,000 steps/day, effectively capturing the diversity of age. There is limited data portraying sex- and age-specific values for steps/day, however. These are necessary if we are to complete a fully evolved "life step curve," expanding on earlier versions of a youth step curve [25]. This shall hopefully be addressed as more pedometer-based research is accumulated and expected values continue to be cultivated.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

CTL conceived the review and lead its implementation and writing; TLW led the electronic search and TLA administered the database. All authors took part in identifying and abstracting articles and editing the final version. All authors read and approved the final manuscript.

Additional material

Additional file 1

Table 1. Expected values for pedometer-determined physical activity in healthy older adults. Compiled pedometer studies abstracted from the scientific literature.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1479-5868-6-59-S1.doc>]

References

- Physical Activity Guidelines Advisory Committee: *Physical Activity Guidelines Advisory Committee Report, 2008* Washington, D.C.: U.S. Department of Health and Human Services; 2008.
- Taylor AH, Cable NT, Faulkner G, Hillsdon M, Narici M, Bij AK Van Der: **Physical activity and older adults: a review of health benefits and the effectiveness of interventions.** *J Sports Sci* 2004, **22(8)**:703-725.
- Tudor-Locke C, Williams JE, Reis JP, Pluto D: **Utility of pedometers for assessing physical activity: convergent validity.** *Sports Med* 2002, **32(12)**:795-808.
- Tudor-Locke C, Ham SA: **Walking behaviors reported in the American Time Use Survey 2003–2005.** *J Phys Act Health* 2008, **5(5)**:633-647.
- Troiano RP, Berrigan D, Dodd KW, Masse LC, Tilert T, McDowell M: **Physical activity in the United States measured by accelerometer.** *Med Sci Sports Exerc* 2008, **40(1)**:181-188.
- Tudor-Locke C: **Taking steps toward increased physical activity: using pedometers to measure and motivate.** *Research Digest* 2002, **3(17)**:1-8.
- Myers AM: *Program Evaluation for Exercise Leaders* Champaign, IL: Human Kinetics; 1999.
- Tudor-Locke C, Myers AM: **Methodological considerations for researchers and practitioners using pedometers to measure physical (ambulatory) activity.** *Res Q Exerc Sport* 2001, **72(1)**:1-12.
- Bassey EJ, Dallosso HM, Fentem PH, Irving JM, Patrick JM: **Validation of a simple mechanical accelerometer (pedometer) for the estimation of walking activity.** *Eur J Appl Physiol Occup Physiol* 1987, **56(3)**:323-330.
- Bassey EJ, Bendall MJ, Pearson M: **Muscle strength in the triceps surae and objectively measured customary walking activity in men and women over 65 years of age.** *Clin Sci* 1988, **74(1)**:85-89.
- Voorrips LE, Ravelli AC, Dongelmans PC, Deurenberg P, Van Staveren WA: **A physical activity questionnaire for the elderly.** *Med Sci Sports Exerc* 1991, **23(8)**:974-979.
- Visser M, De Groot LC, Deurenberg P, Van Staveren WA: **Validation of dietary history method in a group of elderly women using measurements of total energy expenditure.** *Br J Nutr* 1995, **74(6)**:775-785.
- Kitagawa J, Omasu F, Nakahara Y: **Effect of daily walking steps on ultrasound parameters of the calcaneus in elderly Japanese women.** *Osteoporos Int* 2003, **14(3)**:219-224.
- Moreau KL, Degarmo R, Langley J, McMahon C, Howley ET, Bassett DR Jr, Thompson DL: **Increasing daily walking lowers blood pressure in postmenopausal women.** *Med Sci Sports Exerc* 2001, **33(11)**:1825-1831.
- Yoshiuchi K, Nakahara R, Kumano H, Kuboki T, Togo F, Watanabe E, Yasunaga A, Park H, Shephard RJ, Aoyagi Y: **Yearlong physical activity and depressive symptoms in older Japanese adults: cross-sectional data from the Nakanajo study.** *Am J Geriatr Psychiatry* 2006, **14(7)**:621-624.
- Swartz AM, Strath SJ, Miller NE, Cashin SE, Cieslik LJ: **Glucose control and walking in a multiethnic sample of older adults.** *Gerontology* 2007, **53(6)**:454-461.
- Swartz AM, Strath SJ, Parker SJ, Miller NE: **The impact of body-mass index and steps per day on blood pressure and fasting glucose in older adults.** *J Aging Phys Act* 2008, **16**:188-200.
- Croteau KA, Richeson NA, Vines SW, Jones DB: **Effects of a pedometer-based physical activity program on older adults' motility-related self-efficacy and physical performance.** *Activ Adapt Aging* 2004, **28(2)**:19-33.
- Bergman RJ, Bassett DR Jr, Muthukrishnan S, Klein DA: **Validity of 2 devices for measuring steps taken by older adults in assisted-living facilities.** *J Phys Act Health* 2008, **5(Suppl 1)**:S166-175.
- Richeson NE, Vines SW, Jones DB, Croteau KA: **Pedometer as a minimal intervention to improve physical performance indicators for an older adult: An interdisciplinary health team approach and case study.** *Am J Rec Ther* 2003, **2(2)**:21-26.
- Giuliani CA, Gruber-Baldini AL, Park NS, Schrodt LA, Rokoske F, Sloane PD, Zimmerman S: **Physical performance characteristics of assisted living residents and risk for adverse health outcomes.** *Gerontologist* 2008, **48(2)**:203-212.

22. Bassett DR Jr, Ainsworth BE, Leggett SR, Mathien CA, Main JA, Hunter DC, Duncan GE: **Accuracy of five electronic pedometers for measuring distance walked.** *Med Sci Sports Exerc* 1996, **28(8)**:1071-1077.
23. Cyarto EV, Myers AM, Tudor-Locke C: **Pedometer accuracy in nursing home and community-dwelling older adults.** *Med Sci Sports Exerc* 2004, **36(2)**:205-209.
24. Tudor-Locke C, Washington TL, Hart TL: **Expected values for steps/day in special populations.** *Prev Med* 2009, **49(1)**:3-11.
25. Tudor-Locke C, McClain JJ, Hart TL, Sisson SB, Washington TL: **Expected values for pedometer-determined physical activity in youth.** *Res Q Exerc Sport* 2009, **80(2)**:164-174.
26. Tudor-Locke C, Ham SA, Macera CA, Ainsworth BE, Kirtland KA, Reis JP, Kimsey CD Jr: **Descriptive epidemiology of pedometer-determined physical activity.** *Med Sci Sports Exerc* 2004, **36(9)**:1567-1573.
27. Wyatt HR, Peters JC, Reed GW, Barry M, Hill JO: **A Colorado statewide survey of walking and its relation to excessive weight.** *Med Sci Sports Exerc* 2005, **37(5)**:724-730.
28. Strath S, Swartz A, Parker S, Miller N, Cieslik L: **Walking and metabolic syndrome in older adults.** *J Phys Act Health* 2007, **4(4)**:397-410.
29. Payn T, Pfeiffer KA, Hutto B, Vena JE, LaMonte MJ, Blair SN, Hooker SP: **Daily steps in midlife and older adults: relationship with demographic, self-rated health, and self-reported physical activity.** *Res Q Exerc Sport* 2008, **79(2)**:128-132.
30. Croteau KA, Richeson NA: **A matter of health: Using pedometers to increase the physical activity of older adults.** *Activ Adapt Aging* 2005, **30(2)**:37-47.
31. Tudor-Locke C, Bassett DR Jr, Rutherford WJ, Ainsworth BE, Chan CB, Croteau K, Giles-Corti B, Le Masurier G, Moreau K, Mrozek J, Oppert JM, Raustorp A, Strath SJ, Thompson D, Whitt-Glover MC, Wilde B, Wojcik JR: **BMI-referenced cut points for pedometer-determined steps per day in adults.** *J Phys Act Health* 2008, **5(Suppl 1)**:S126-139.
32. Zhang JG, Ohta T, Ishikawa-Takata K, Tabata I, Miyashita M: **Effects of daily activity recorded by pedometer on peak oxygen consumption (VO₂peak), ventilatory threshold and leg extension power in 30- to 69-year-old Japanese without exercise habit.** *Eur J Appl Physiol* 2003, **90(1-2)**:109-113.
33. Shimizu K, Kimura F, Akimoto T, Akama T, Kuno I: **Effect of free-living daily physical activity on salivary secretory IgA in elderly.** *Med Sci Sports Exerc* 2007, **39(4)**:593-598.
34. Tudor-Locke C, Giles-Corti B, Knuiaman M, McCormack G: **Tracking of pedometer-determined physical activity in adults who relocate: results from RESIDE.** *Int J Behav Nutr Phys Act* 2008, **5**:39.
35. Yamakawa K, Tsai CK, Haig AJ, Miner JA, Harris MJ: **Relationship between ambulation and obesity in older persons with and without low back pain.** *Int J Obes Relat Metab Disord* 2004, **28(1)**:137-143.
36. King WC, Belle SH, Brach JS, Simkin-Silverman LR, Soska T, Kriska AM: **Objective measures of neighborhood environment and physical activity in older women.** *Am J Prev Med* 2005, **28(5)**:461-469.
37. Swartz A, Strath S, Parker S, Miller N, Cieslik L: **Ambulatory activity and body mass index in white and non-white older adults.** *J Phys Act Health* 2007, **4(3)**:294-304.
38. Tudor-Locke C, Sisson SB, Collova T, Lee SM, Swan PD: **Pedometer-determined step count guidelines for classifying walking intensity in a young ostensibly healthy population.** *Can J Appl Physiol* 2005, **30(6)**:666-676.
39. Tudor-Locke C, Williams JE, Reis JP, Pluto D: **Utility of pedometers for assessing physical activity: construct validity.** *Sports Med* 2004, **34(5)**:281-291.
40. Tudor-Locke C, McClain JJ, Hart TL, Sisson SB, Washington TL: **Pedometer methods for assessing free-living youth.** *Res Q Exerc Sport* 2009, **80(2)**:175-184.
41. Schneider PL, Crouter SE, Bassett DR: **Pedometer measures of free-living physical activity: comparison of 13 models.** *Med Sci Sports Exerc* 2004, **36(2)**:331-335.
42. Le Masurier GC, Lee SM, Tudor-Locke C: **Motion sensor accuracy under controlled and free-living conditions.** *Med Sci Sports Exerc* 2004, **36(5)**:905-910.
43. Crouter SE, Schneider PL, Karabulut M, Bassett DR Jr: **Validity of 10 electronic pedometers for measuring steps, distance, and energy cost.** *Med Sci Sports Exerc* 2003, **35(8)**:1455-1460.
44. Le Masurier GC, Tudor-Locke C: **Comparison of pedometer and accelerometer accuracy under controlled conditions.** *Med Sci Sports Exerc* 2003, **35(5)**:867-871.
45. Karabulut M, Crouter SE, Bassett DR Jr: **Comparison of two waist-mounted and two ankle-mounted electronic pedometers.** *Eur J Appl Physiol* 2005, **95(4)**:335-343.
46. Tudor-Locke C, Ainsworth BE, Thompson RW, Matthews CE: **Comparison of pedometer and accelerometer measures of free-living physical activity.** *Med Sci Sports Exerc* 2002, **34(12)**:2045-2051.
47. Frisard MI, Fabre JM, Russell RD, King CM, DeLany JP, Wood RH, Ravussin E: **Physical activity level and physical functionality in nonagenarians compared to individuals aged 60-74 years.** *J Gerontol A Biol Sci Med Sci* 2007, **62(7)**:783-788.
48. Westterterp KR, Meijer EP: **Physical activity and parameters of aging: a physiological perspective.** *J Gerontol A Biol Sci Med Sci* 2001, **56(Spec No 2)**:7-12.
49. Tudor-Locke C, Burkett L, Reis JP, Ainsworth BE, Macera CA, Wilson DK: **How many days of pedometer monitoring predict weekly physical activity in adults?** *Prev Med* 2005, **40(3)**:293-298.
50. Felton GM, Tudor-Locke C, Burkett L: **Reliability of pedometer-determined free-living physical activity data in college women.** *Res Q Exerc Sport* 2006, **77(3)**:304-308.
51. Schonhofer B, Ardes P, Geibel M, Kohler D, Jones PW: **Evaluation of a movement detector to measure daily activity in patients with chronic lung disease.** *Eur Respir J* 1997, **10(12)**:2814-2819.
52. Sieminski DJ, Cowell LL, Montgomery PS, Pillai SB, Gardner AVV: **Physical activity monitoring in patients with peripheral arterial occlusive disease.** *J Cardiopulm Rehabil* 1997, **17(1)**:43-47.
53. Rowe DA, Kemble CD, Robinson TS, Mahar MT: **Daily walking in older adults: day-to-day variability and criterion-referenced validity of total daily step counts.** *J Phys Act Health* 2007, **4(4)**:434-446.
54. Marshall AL: **Should all steps count when using a pedometer as a measure of physical activity in older adults?** *J Phys Act Health* 2007, **4**:305-314.
55. Richardson CR, Newton TL, Abraham JJ, Sen A, Jimbo M, Swartz AM: **A meta-analysis of pedometer-based walking interventions and weight loss.** *Ann Fam Med* 2008, **6(1)**:69-77.
56. Bravata DM, Smith-Spangler C, Sundaram V, Gienger AL, Lin N, Lewis R, Stave CD, Olkin I, Sirard JR: **Using pedometers to increase physical activity and improve health: a systematic review.** *JAMA* 2007, **298(19)**:2296-2304.
57. King WC, Brach JS, Belle S, Killingsworth R, Fenton M, Kriska AM: **The relationship between convenience of destinations and walking levels in older women.** *Am J Health Promot* 2003, **18(1)**:74-82.
58. Fukukawa Y, Nakashima C, Tsuboi S, Kozakai R, Doyo W, Niino N, Ando F, Shimokata H: **Age differences in the effect of physical activity on depressive symptoms.** *Psychol Aging* 2004, **19(2)**:346-351.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

