



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



## Letter to the Editor

### The potential significance of vitamin D binding protein polymorphism in COVID-19



With interest, we read the paper by Nasiri et al. (Nasiri et al., 2021), which investigated the relationship between vitamin D and the prognosis in Coronavirus Disease 2019 (COVID-19) patients. A significant association between the hospital stay and lower serum vitamin D levels was demonstrated. Although several confounders were taken into account, we would like to focus on the potential important influence of vitamin D binding protein (DBP) polymorphisms on the reported results.

DBP is a polymorphic protein with three major phenotypes, DBP1F [rs7041-T (ASP), rs4588-C (Thr)], DBP1S [rs7041-G (ASP), rs4588-C (Thr)], and DBP2 [rs7041-T (ASP), rs4588-A (Lys)], and more than 120 variants. The majority (85-90%) of 25-hydroxyvitamin D in the circulation is bound to DBP, whereas 10-15% is loosely bound to albumin, and < 1% circulates in its free form (Speeckaert et al., 2014). As demonstrated in a cross-sectional study, plasma concentrations of DBP, 25-hydroxyvitamin D and 1,25-dihydroxyvitamin D are highest in DBP1-1, intermediate in DBP2-1, and lowest in DBP2-2. Multiple regression analysis showed that the DBP concentration was an independent predictor of 1,25-dihydroxyvitamin D, whereas the DBP phenotype was a significant predictor of the 25-hydroxyvitamin D concentration, even after adjustment for confounders (Lauridsen et al., 2005). Besides rs7041 and rs4588, a genome-wide meta-analysis identified additional SNPs that affect the 25-hydroxyvitamin D concentration: e.g. rs2282679 of the *DBP* gene, which is a near-perfect proxy for rs4588. rs2282679-A is typically co-inherited with rs4588-C, whereas rs2282679-C is co-inherited with rs4588-A. rs2282679-C/C allele carriers have lower vitamin D concentrations than carriers of one rs2282679-C-allele, who in turn on average have lower vitamin D levels than rs2282679-A/A individuals (Wang et al., 2010).

Investigating the influence of the DBP phenotypes in patients with a Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-CoV-2) infection, we demonstrated that DBP1 carriers might be less susceptible to infection and mortality due to COVID-19 (Speeckaert et al., 2021). Besides, in a recent paper, a positive correlation was observed between the Metabolism score (= DBP rs2282679 + CYP24A1 rs17216707) and COVID-19 disease severity, in which a deeper analysis showed that the rs2282679 polymorphism could explain most of this interesting correlation (Freitas et al., 2021). Besides the transport of vitamin D and its metabolites, DBP could exert several other key roles in COVID-19, being involved in the extracellular actin scavenger system or acting as a neutrophil chemotactic factor and a macrophage activator (Delanghe et al., 2015).

### Conflicts of interest

None disclosed.

### Ethical Approval

N/A.

### Funding sources

None disclosed.

### Acknowledgments

None.

### References

- Delanghe JR, Speeckaert R, Speeckaert MM. Behind the scenes of vitamin D binding protein: more than vitamin D binding. *Best Pract Res Clin Endocrinol Metab* 2015;29:773–86.
- Freitas AT, Calhau C, Antunes G, Araújo B, Bandeira M, Barreira S, et al. Vitamin D-related polymorphisms and vitamin D levels as risk biomarkers of COVID-19 infection severity; 2021 medRxiv preprint doi: <https://doi.org/10.1101/2021.05.14.21251111>.
- Lauridsen AL, Vestergaard P, Hermann AP, Brot C, Heickendorff L, Mosekilde L, et al. Plasma concentrations of 25-hydroxy-vitamin D and 1,25-dihydroxy-vitamin D are related to the phenotype of Gc (vitamin D-binding protein): a cross-sectional study on 595 early postmenopausal women. *Calcif Tissue Int* 2005;77:15–22.
- Nasiri M, Khodadadi J, Molaei S. Does vitamin D serum level affect prognosis of COVID-19 patients? *Int J Infect Dis* 2021 S1201971221003957.
- Speeckaert MM, De Buyzere ML, Delanghe JR. Vitamin D binding protein polymorphism and COVID-19. *J Med Virol* 2021;93:705–7.
- Speeckaert MM, Speeckaert R, van Geel N, Delanghe JR. Vitamin D binding protein. *Adv Clin Chem* 2014;63:1–57.
- Wang TJ, Zhang F, Richards JB, Kestenbaum B, van Meurs JB, Berry D, et al. Common genetic determinants of vitamin D insufficiency: a genome-wide association study. *Lancet* 2010;376:180–8.

Marijn M Speeckaert\*

Department of Nephrology, Ghent University Hospital, Ghent, Belgium  
Research Foundation-Flanders (FWO), Brussels, Belgium

Joris R. Delanghe

Department of Diagnostic Sciences, Ghent University, Ghent, Belgium

\*Corresponding author: Marijn M. Speeckaert, Ghent University Hospital, Department of Nephrology, Corneel Heymanslaan 10 9000 Ghent, Phone: ++ 32 9 332 4509; Fax: ++ 32 9 332 3847.  
E-mail address: [Marijn.Speeckaert@ugent.be](mailto:Marijn.Speeckaert@ugent.be) (M.M. Speeckaert)