



Understanding fish waste management using bibliometric analysis: A supply chain perspective

Farook Abdullah Sultan¹, Srikanta Routroy² and Maitri Thakur³

Abstract

Food loss and waste have become an issue of global significance, considering their concurrent effects on the socioeconomic and environmental facet of society. Despite this domain gaining prolific attention recently, issues hampering the effective utilization of residues from fish processing usually go unidentified in developing economies such as India. This occurs mainly owing to fragmented supply chains, inappropriate handling, discontinuous cold chains, inadequate temperature monitoring and so on, affecting quality and causing underuse. Any researcher trying to understand the prospects of utilizing these fish processing co-streams in a developing economy with the vision of improving consumption, economic sustainability, reducing discards and promoting circularity faces a lacuna. The authors address this demand in research by identifying the validity of this domain both in the global and native research community by conducting a detailed review using bibliometric analysis and content analysis. Data from Scopus with 717 documents, comprising 612 research articles from 78 countries, 1597 organizations and 2587 authors, are analysed. Results signify (i) developing a focus on hydroxyapatite production, bio-methane generation, transesterification processes, biomass and the rest raw material generated from fish processing, and (ii) reduced research on supply chain-related aspects despite their considerable importance. To comprehend this deficiency, especially in the Indian stance, barriers hindering the utilization of generated by-products are identified, and recommendations for improvements are proposed. The results will provide the struts for a circular and sustainable supply chain for processed seafood in developing economies.

Keywords

Food waste, processing co-streams, waste management, bibliometric analysis, supply chain

Received 11th November 2021, accepted 29th July 2022 by Associate Editor Rodrigo Navia.

Introduction

Food wastage (FW) has become an issue of global enormity, with 931 million tonnes of edible food being lost in 2019, accounting for a 17 % colossal volume of the global food production. Stresses such as the advent of Covid-19 have risen food insecurity counts to 108 million people in 48 countries (FSIN, 2020), increasing global pressures to regulate food waste/losses occurring in supply chains (Magalhães et al., 2020). The complications of FW gain further attention in a nation like India, which obliges a two-fold role of serving the rising global market and its self-growing population. Posing second in occupancy figures after China with a population of 1.38 billion, India incurs an annual loss of INR 440 billion from food waste/loss, sharing 40% in its autogenous food produced (NAAS, 2019). Comprising 14% of the world's undernourished people (India Food Banking Network, 2021), the problem of FW in India is undoubtedly a topic of concern in recent times. Despite the varied types of FW produced, this article only focuses on understanding the various utilizations possible from product co-streams generated from fish processing. The generated co-streams involve the fish head, viscera, trimmings,

bone and cartilage hide tails and wash water. The co-streams are categorized into rest raw materials (RRMs) and by-products based on their usage. Both being process outputs, the former is food grade, whereas the latter is used for producing feed/fertilizers (Mozuraityte et al., 2020; Penven et al., 2013). Literature also indicates an absence of proper terminologies considering generated co-streams. Furthermore, many articles focus on these co-streams as fish wastes in general (Arvanitoyannis and Kassaveti, 2008; Caruso et al., 2020). This can be due to the subjective use of this terminology for being utilizable for some processors and

¹School of Business Management, Narsee Monjee Institute of Management Studies, Hyderabad, Telangana, India

²Department of Mechanical Engineering, Birla Institute of Technology and Science, Pilani, Pilani, India

³SINTEF Ocean, Trondheim, Norway

Corresponding author:

Farook Abdullah Sultan, School of Business Management, Narsee Monjee Institute of Management Studies, Hyderabad, Telangana 509301, India.

Email: abdullahsultan1991@gmail.com

its consideration as waste by others (Adler et al., 2014). Therefore, to account for the utilization of all the co-streams generated, this article considers the term 'fish wastes' in the entire manuscript.

Indian fisheries fetch USD 7.08 billion by exporting 11% of their catch (NFDB, 2019). These exports comprise shrimp, frozen fish, squid, dried fish, live fish, chilled fish and other forms of seafood (i.e. processed seafood, surimi etc.). As of 2018, India has an installed processing capacity of 27300 MT seafood/day with 547 registered processing plants (CATR, 2018). Consumption and seafood processing generate several underutilized co-streams that generally get discarded as wastes. India constitutes around 2 MMT of fish waste, with the global fish waste summing to 130 MMT (Binsi, 2018). Seafood wastes produced in India share >4% (Binsi, 2018) amid the worldwide share of 50 million tonnes (McCarthy, 2018). Future estimations citing increment in India's seafood share from 4.1 to 6.7% by 2030 (ET Bureau, 2020) subsequently indicate an added rise in seafood waste generation. All these lay added stresses to understand the various causes deterring the utilization and commercialization of high-value seafood wastes in India, despite its dominance in research.

In the global context, seafood production has recorded a sharp rise of 20% compared to 148.1 MMT produced in 2010 (Statista, 2020). This has eventually accounted for an analogous increase in seafood processing to counter consumers' growing demand. Seafood processing generates a large amount of liquid and solid waste (Arvanitoyannis and Kassaveti, 2008) that needs to be managed due to its undesirable ecological effects caused by improper disposal (Arvanitoyannis and Tserkezou, 2014). Though valorization has proved effective in handling fish wastes through the production of fish meal, fish oil, the output of silage (Islam and Peñarubia, 2021) and organic fertilizers (Ahuja et al., 2020; Arruda et al., 2007), the products produced, however, are of low value. The advent of recent technologies nevertheless enables efficient valorizing of high-value products such as collagen, gelatine and enzymes (Kumar et al., 2018).

Fish waste management

Efficient management of fish wastes necessitates the firm espousal of The United Nations Responsible Consumption and Production Goals (i.e. Sustainable Development Goal 12) by encouraging efficient resource management. Supply chain inefficiencies are a plausible reason for the ineffective utilization of fish wastes in high valued products (Jouvenot, 2015). However, significant references cite their utilization in agriculture (Islam, 2006), pet food (De Silva and Turchini, 2008), nutraceutical and pharmaceutical industries (Hayes, 2012) and animal feed (Afreem and Ucak, 2020) with varying margin levels. Arvanitoyannis and Kassaveti (2008) reviewed the treatments, environmental impacts and potential uses of fish wastes. A thorough description of the usage of fish waste was provided by Ghaly et al. (2013). They highlighted that processing fish wastes provided a rich yield of proteins, amino acids and oils. Caruso (2015) stressed the need to impart sustainability by utilizing fish wastes by valorizing them

for making fish feed, thereby solving the ecological impacts caused due to the improper disposal of the generated by-products. Nevertheless, a large volume of literature was found on the utilization of fish wastes to produce bioactive compounds (Karkal and Kudre, 2020; Mohapatra et al., 2017; Ramanujam et al., 2016). This has been followed by a rise in review papers on managing fish wastes (Coppola et al., 2021; Hjellnes et al., 2020; Omar et al., 2019), aiding in clarifying the level of research existing in the last decade.

Research influences from India reveal noteworthy contributions from Mathew (2010), who focussed on the utilization of fish wastes and discussed its importance. Sasidharan and Mathew (2011) deliberated on the status of fish waste management in Kerala (India) by explaining the handling capacities of seafood processing industries and stressed a better management practices. Jayathilakan et al. (2012) reviewed the valorizing of animal by-products, including fish and provided an inclusive view of various waste management techniques. Sasidharan et al. (2013) magnified the role played by private players in the Indian waste disposal scenario and underscored the absence of authority control. Rejula and Mohanty (2018) further emphasized developing a structured fisheries waste management technique in India by initiating technification via public-private partnerships and technology commercialization via policy-level initiatives. A similar indication was delivered by Binsi (2018), who weighed the importance of the upgradation of existing valorizing technologies and suggested changes to enhance the valorization of secondary products produced from fish waste.

Among the various studies on fish waste valorization, many studies address the various valorized fish by-products produced. Despite such proficient research, it is witnessed that the existence of high demand products generated is generally meagre and localized to Scandinavian regions (Olsen et al., 2014). It is concluded that a significant gap exists between seafood waste management and other waste management techniques, with the latter having a substantial share of research done, especially in the Indian stance (Sharma et al., 2020; Vanapalli et al., 2021). Considering all the above-discussed factors, a comprehensive review is essential, further succumbing to the nonexistence of studies focusing on sustainable seafood supply chains, especially in India. All of these indicate a discontinuity in the literature on seafood waste management. Hence, the authors try to bridge this research gap using bibliometric analysis by undergoing a detailed understanding of managing fish waste and comprehending research progress in this domain.

Bibliometric analysis

Bibliometric analysis proves to be an effective review system to demonstrate the general state and trajectory of research by providing a concise perspective compared to a conventional reviewing technique (Costa et al., 2017; Qaiser et al., 2017). Introduced in 1969 by Pritchard (Pritchard, 1969), this technique statistically evaluates published literature to measure researchers' degree of inspiration/adoption of a research topic. The added

uniqueness of this technique also lies in its ability to capture the temporal evolution of multiple parameters (Zhu et al., 2019). The validity of adopting bibliometric studies as literature analysing tool is evident from recent research on food security (Xie et al., 2021), blockchain technology (Ante et al., 2021), big data analytics (Inamdar et al., 2020), manufacturing (De Oliveira et al., 2019), internet of things (Bouzembrak et al., 2019), food waste research (Zhang et al., 2018) and so on.

Research novelty

The novelty of this research lies in its twofold approach of understanding the various uses of fish by-products (globally) and highlighting factors inhibiting the effective utilization of fish wastes generated from processing (Indian perspective). To the author's best knowledge, no study to date has focussed both quantitatively and visually on the domain of fish waste. This research explores the inherent information generally undermined in a conventional review, such as bibliographic coupling and co-occurrence network analysis, and reveals critical research hotspots. This study is additionally made comprehensive by carrying out a content analysis. The research aims to provide a geocentric outcome in the Indian seafood supply chain context by carrying out a barrier assessment.

The research findings will benefit the research fraternity by providing a coherent picture for framing a foundation to conduct valuable research on managing fish waste. The suggested implications being of its kind owing will also positively pave the way for bureaucrats, decision-makers and industry-based practitioners to implement measures for new product development, plan operations, engage in effective policymaking and develop effective marketing strategies; deliberating towards sustainable processing in the Indian context by enhancing circularity. Our research follows the stages proficiently analysed by Li et al. (2021) and aims to address the following research questions (RQs) on fish waste management:

RQ1: What is the present status of research in the global context?

RQ2: What research contexts/hotspots are being explored in literature?

RQ3: Is research in this domain domineered, or is it collaborative (w.r.t authors, countries and organizations)?

RQ4: Identify issues hampering the valorization of fish wastes in the Indian perspective?

Methods

Search strategy

This article adopts a mixed review method assuming both quantitative bibliometric analysis and qualitative content analysis (Du et al., 2021). Primarily espousing a seven-step approach, the initial three steps of the methodology deal with retrieving, sorting and screening relevant information from the selected database followed by analysing, explaining and proposing improvements

in the next three steps prior to concluding the derived outcomes. The methodological procedure adopted in this research (refer Figure 1) coincides with that followed by Jia and Jiang (2018), confirming its validity.

Data curation

Data curation is initiated using the Scopus database. The Scopus database was chosen due to the high scientific quality and coverage compared to other available databases (Mongeon et al., 2016; Pham-Duc et al., 2020). A keyword-oriented search is initially adopted, targeting literature on fish waste and associated management techniques using the keywords 'fish waste utilization' OR 'fish waste management' AND 'supply chain management'. The top 70 articles obtained are initially reviewed to expand the keyword search string used. This further expanded the keyword search to words such as '*fish waste(s)*' OR '*fish waste utilization*' OR '*rest raw material*' OR '*fish by-product*' OR '*fish waste management*'. When searching relevant literature, the keywords mentioned above provided additional keywords such as '*bio-economy in fisheries*' OR '*circular economy in fisheries*'. The search string revolves around '*fish waste(s)*' OR '*fish waste utilization*', OR '*rest raw material*' OR '*fish by-product*' OR '*fish waste management*' OR '*bio-economy in fisheries*' OR '*circular economy in fisheries*' AND '*supply chain management*'. This derived a total of 975 documents till December 2020, including contributions as early as 1917. Selection criteria involve the inclusion criteria of published literature only in English ($n=934$) and neglects books, book series and trade journals, reducing the total count to 888. An added initiative adopted in this work is the manual screening (filtering) of literature to identify and eliminate listed grey literature. This further reduced the total article count to 717 documents comprising 612 research articles, 82 conference papers and 23 review papers authored by 2587 authors from 1597 organizations and 87 countries subjected to further analysis.

Analysis

Bibliometric analysis principally comprises the application of quantitative techniques (such as citation analysis) on bibliometric data (i.e. publication and citation data). The resultant data in *CSV format* from the Scopus website is analysed using performance analysis and science mapping techniques. Performance analysis is a preliminary analysis used to analyse contributions by source, author, affiliation, country, document type, subject area and funding. Being qualitative, this analysis forms a base for any review undertaken (Donthu et al., 2021). Detailed outcomes are revealed using science mapping techniques showing relations between scientific literature and contributing authors w.r.t existing collaborations, analysing publication patterns (at institutional and geographical level) and aid in proposing new research directions (Li et al., 2021). This analysis capacitates understanding the exploratory perception w.r.t co-authorship, co-occurrence, citation, bibliographic coupling and co-citation links (Van Eck and




Figure 1. Research flow.

Waltman, 2013). To evaluate this analysis, vital information such as author details, citation information, title, journal, DOI, references, keywords and research text is collected. All results related to the various science mapping analysis used in this research is

presented by applying network analysis using VOS viewer V1.6.16 software (refer to the section ‘Network analysis’). VOS viewer is a tool used for creating and exploring maps depicting network data and functions. This is done based on a normalized

term co-occurrence matrix and a similarity measure used to calculate the association strength between terms by employing an integrated approach of mutual mapping and clustering (Van Eck and Waltman, 2019). Table 1 details the various metrics/techniques used in this research for conducting the bibliometric analysis.

Network analysis. Though a general bibliometric study provides adequate information regarding document/source/author/organization/country, the relation between the mentioned factors goes un-noted. Network analysis presents this relation by circular nodes and connecting lines. Visualizations in the VOS viewer software are represented using the distance-based approach where the distance between the visualized nodes (document/source/author, etc.) indicates the relatedness (i.e. shorter the distance, more vital the relatedness) and the size of the nodes denotes article importance (i.e. citation, number of documents, etc.).

The analysis is initiated by applying the default association strength normalization (Eck and Waltman, 2009). Once normalized, the network is constructed. This is followed by the positioning of nodes using the default ‘visualization of similarities mapping technique’ (Van Eck et al., 2010). Evaluation is carried out using total link strength (TLS) and links. ‘TLS’ measures the extant links between the compared research w.r.t researcher/country/organization (Lancho-Barrantes and Cantú-Ortiz, 2019), whereas links denote the extent of the relation existing between two items. Upon completion, the assignment of nodes to clusters is carried out and represented by unique colours. The clustering operation purposes of generating thematic or social clusters that aid in understanding the development of a research field. A cluster, in general, refers to a set of items included in developed network analysis and is identified using cluster numbers. Clustering in VOS viewer is performed by default using its inherent smart local moving algorithm (Van Eck and Waltman, 2014). The final set of clusters obtained, dependent on resolution, calls for optimal resolutions during network analysis according to the clarity of the derived clusters. The various Acronyms used in this research are tabulated in the appendix section.

Content analysis. Content analysis provides information regarding the existing intellectual structure in the literature analysed (Song et al., 2021) and indicates the correlation between articles by clustering (Costa et al., 2017). This analysis focuses on understanding valuable information available in textual data in a structured and systematic manner. Content analysis is carried out in this research by selecting research articles based on the top cited articles, done by the procedure adopted by Du et al. (2021).

Impediment assessment of sustainable fish waste management in India

Based on the results obtained, impediments are suggested with a geo-localized perspective. This section deals with the ambiguities understood from the bibliometric and content analysis and proceeds by posing recommendations that need to be

incorporated. The research team surveyed academic literature focusing on perishable foods to understand the antecedents that influence stakeholders in adopting sustainable improvements in the Indian stance. This derived 75 barriers identified from an extensive survey of the literature. A screening questionnaire is developed to filter the identified barriers to avoid discrepancies in the barriers identified. The screening questionnaire filtered the identified barriers to a total of 52 barriers. The identified barriers were further validated and screened through interactions conducted with prime stakeholders of the supply chain (fishers, aggregators, preprocessing centres and processing centres). This finally concluded to 48 finalized and categorized barriers (refer to the section ‘Impediment assessment of sustainable fish waste management in India’ and Supplementary Material for more details). The identified barriers are validated via interactions conducted with 74 stakeholders along the supply chain, with the mode of interaction primarily being offline (face to face) and through telephonic conversation with the top management of processing centres. The developed screening questionnaire is as mentioned below.

1. Are the barriers related to the objective discussed in this research?
2. Do the barriers denote sufficient information?
3. Are the barriers coherent in the perishable food supply chain?
4. Are the barriers linked with facets of sustainability?

Interviews were conducted with various stakeholders such as fishermen, aggregators, preprocessing centres and processing centres along Sassoon dock in Mumbai (India). Responses were collected between September 2019 and April 2021. Respondent selection threshold was kept to a minimum experience criterion of 1 year in their respective fields throughout stakeholder validation. Table 2 summarizes the details of stakeholders who participated in the validation process. This broad perceptive study conducted throughout the supply chain helped the research team gain an adequate idea regarding the coherency of the identified barriers in the Indian stance of seafood supply chains and also aided in identifying the echelon (member) influenced by these impediments in the supply chain. This is finally concluded using the improvements derived from stakeholder interactions required for the sustainable operation of the seafood supply chain in India.

Findings

Statistical overview

An evolution centric study of published literature reveals an exponential rise in article distribution and their corresponding citations from 1917 to December 2020 (Figure 2). A scrutiny of the assessed research articles reveals distribution across 147 sources. Further, break up discloses a sharp rise of 87.58% in publications contributions from 2002 among the entire range of documents assessed. Research contributions also denote a drop/decrease at irregular intervals. This possibly occurs owing

Table 1. Metric/technique adopted for bibliometric analysis.

Analysis type	Metric(s)/technique	Explanation	Purpose	Data source/particulars	
Performance analysis	<i>Metric(s) adopted</i>				
	Total publication	Total number of research documents published	Insight regarding research activity in a domain	Scopus Database	
	Year-wise categorization	Year-wise categorization of published literature	Insight regarding year-wise developments in research activity in a research domain		
	Author contributions	Total number of research document(s) by the author(s)	Insight regarding author contributions in a research domain		
	Co-author contributions	Total number of research documents by two or more authors	Insight regarding contributions from two or more authors author in a research domain		
Citation data	Total number of citations of a research document	Provides insight regarding the number of times an article is followed			
Science mapping	<i>H</i> index	The measure of the influence of an article. Denotes the number of publications cited at least 'H' times	The comparative ratio of publications to citations showcases the cumulative impact of an author's contribution	Document sourced analysis that requires the use of following critical information	
	<i>Technique(s) adopted</i>				
	Citation analysis	Analyse relatedness between research documents based on the number of times these are cited	Identifying the most influential publication in a domain		<ul style="list-style-type: none"> • Author • Citation • Title • Journal • DOI • References
	Co-citation analysis	Analyse relatedness between research items based on the number of times these are cited together	Understand the development of foundational research themes		Document sourced analysis that requires the use of references
	Co-authorship analysis	Analyse relatedness of items based on the number of co-authored documents	Reveal the level of inherent relation existing in research between authors and their affiliations, countries		This analysis requires authors their affiliations and their respective domicile (countries)
Bibliographic coupling	Bibliographic coupling	Analyse the relatedness of items based on the number of references they share, assuming similarity in research content	Forms as an indicator of a robust research foundation by forming an insight into prospective research in a domain	Document sourced analysis that requires the following critical information	
				<ul style="list-style-type: none"> • Author • Citation • Title • Journal • DOI • References 	
				Keyword sourced analysis that requires the following critical information	
				<ul style="list-style-type: none"> • Title • Abstract • Research Text • Author/index keywords 	
				Visualise research hotspots and emerging research trends	
Co-occurrence analysis	Analyse the relatedness of items in which the terms are found to occur together				

Table 2. Survey response summary.

Stakeholder	Role	Members	Response (n)	Response share (%)
Fishermen	The starting point of the supply chain and purpose by providing raw material (whole fish) for the functioning of the supply chain. Highly skilled fishers are the skipper/captain of the fishing vessel and are the prime decision-makers during fishing.	Fishers	18	24.32
Aggregators	Collect fish from numerous fishers and carry bulk auctions to disperse the possessed fish.	Aggregator	15	20.27
Preprocessing centre	Engage in a manual de-heading operation involving the separation of de-headed fish and RRM. Once de-headed, whole fish is sent for additional processing, whereas the generated RRM is either sent for valorization to fish waste processing plants or is discarded.	In charge Top management	16 8	21.62 10.81
Processing centres	Whole fish processing plants engage in added processing of received raw material	Manager	8	10.81
		Top management	4	5.41
	Fish waste processing plants engage in the valorization of received RRM into fish meal/fish oil or any other value derived product	Manager	5	6.76
		Top management	4	5.41
		Total responses	74	100%

to the time in publication and adoption of new concepts/ideas/ techniques in the academic fraternity, hence suitably correlating to the fact stated by Van Meeteren et al. (2016).

Source wise contribution. Assessment reveals significant contributions in areas of agricultural and biological sciences (27%) environmental science (15%), with minor contributions in biochemistry, genetics, microbiology and energy and so on (Figure 3). A similarity in the diversification of publications is also visible in publication sources, irrespective of the existing diversification in scopes (Refer Table 3) Scrutiny indicates an increased fondness towards journals such as *Bioresource Technology* (34%), *Journal of Cleaner Production* (18%), *Aquaculture* (15%), *Renewable Energy* (12%), *Journal of The Science of Food and Agriculture* (10%), *Waste Management* (7%) and other publishing sources (<1%), with the former three constituting around 40% among the top 10 sources. A year-wise citation assessment reveals maximum likeliness towards *Bioresource Technology* and *Journal of Cleaner Production*, with average sources of these journals exceeding their counterparts, especially in the last decade. However, in terms of total citations, *Bioresource Technology*, *Food Chemistry* and *Aquacultural Engineering* lead the total citation frontier together, having the lion’s share of 51.28% among the top 10 publishing sources. This makes these sources a preferable basis for modern research in fish waste.

Author and country-wise contribution. Maximum contributions were from Peter J. Bechtel and Narayan Bhaskar (eight documents each) (Table 4). Analysing the geo-local contributions reveal that India leads the list in global contributions (99 documents), followed by the United States (87 documents) and Brazil (51 documents) (Figure 4). It is worth noting that the United States spots the largest share of citations despite holding the second position in the percentage of overall publications. The adaptability of published literature (in terms of citation) seems higher for developed countries (80%), whereas the volume of publications supersedes for developing nations (53%).

Affiliation share. One hundred and sixty organizations were found to have contributed to the domain of fish wastes. The maximum contribution was cited from SINTEF Ocean, Norway, solely contributing 12 publications (1.96%). Universidad Federal do Rio Grande (Brazil), University of Alaska Fairbanks (United States) and Norges teknisk-naturvitenskapelige Universitet (Norway) each contribute 11 publications (1.79%) in the domain of fish waste management. Organizations from developing nations contribute 40% of the leading contributors, evidently showcasing maximum contributions from organizations located in developed countries. Brazil (Universidad Federal do Rio Grande – 1.79% and UNESP Universidade Estadual Paulista – 1.31%) tops the list in terms of a developing nation contributing to the domain of fish waste, followed by India (Central Food Technological Research Institute – 1.47%) and Malaysia (Universiti Sains Malaysia – 1.31%). A citation-based ranking on the




Figure 2. Publication citations and articles (1917-2020).





Figure 3. Subject focussed assessment.

Table 3. Top 10 sources for publications (till December 2020).

Source title	Publisher	Publications	IF	H index	Quartile of JCR
<i>Bioresource Technology</i>	Elsevier	20	7.539	273	Q 1
<i>Aquaculture</i>	Elsevier	16	3.224	164	Q 1
<i>Journal of Cleaner Production</i>	Elsevier	15	7.246	173	Q 1
<i>Waste and Biomass Valorization</i>	Springer	12	2.851	35	Q 2
<i>IOP Conference Series</i>	IOP Science	10	-	18	-
<i>Waste Management</i>	Elsevier	9	5.448	145	Q 1
<i>Renewable Energy</i>	Elsevier	9	6.274	174	Q 1
<i>Journal of Food Science and Technology</i>	Springer	9	2.705	55	Q 2
<i>Aquacultural Engineering</i>	Elsevier	9	2.638	67	Q 1
<i>Journal of The Science of Food and Agriculture</i>	Wiley	8	2.614	131	Q 1

Table 4. Top 10 contributing authors (till December 2020).

Name of the author	Organization	H index	No. of publications
Peter J. Bechtel	USDA ARS Southern Regional Research Center, United States	39	8
Narayan Bhaskar	Central Food Technological Research Institute, India	25	8
José A. Siles	Department of Inorganic Chemistry and Chemical Engineering, Spain	22	7
M. A. Martín Santos	Universidad de Córdoba, Spain	30	6
Rasa Šližyt	SINTEF Ocean, Norway	15	6
Frank George Guimarães Cruz	Universidade Federal do Amazonas, Brazil	5	5
Masayuki Furuichi	Kyushu University, Japan	17	5
Wilson A. Lennard	University of South Australia	4	5
María Elvira López-Mosquera	Universidad de Santiago de Compostela, Spain	11	5
Vilásia Guimarães Martins	Universidade Federal do Rio Grande, Brazil	13	5

**Figure 4.** Global distribution of publication density (numbers represent publications).

organizational deliverable, however, deliberates that the University of Alaska Fairbanks and USDA REE Agricultural Research Service (both belonging to the United States) ranks as the most cited organization, followed by Central Food Technological Research Institute (India) and SINTEF Ocean (Norway). Hence, the outcomes mentioned above satisfy RQ1 by presenting the status of research on fish waste management in the global context.


Co-occurrence analysis

Co-occurrence/keyword analysis primarily reveals the core topics addressed by the research community in recent times. Figure 5(a) and (b) signifies the keyword network aiding in understanding the interlinks between research using keyword specifics and representing a keyword cloud. These interlinks/lines represent the strengths and relevance of the analysed keywords (Mishra et al., 2021). In Figure 5(c), letter sizes indicate keyword occurrence frequency among the assessed literature, representing the most popular terms.


Keyword assessments reveal the importance of deriving biodiesel (Yahyaee et al., 2013) and biogas (Salam et al., 2009) from fish waste as an ecologically sustainable fuel alternative. Nevertheless, the generation of these value-added products using anaerobic digestion makes this digestion technique also a renowned term used by researchers (Bücker et al., 2020). Researchers have shown the ability to efficiently generate biodiesels using the transesterification process (Jung et al., 2019), improving the recovery of value-added products. Production of animal feed from fish by-products has also seen rising adoptions recently (Afreeen and Ucak, 2020). This can be cited in aquaculture's rising adoption in populous nations like India and China (Mo et al., 2018). Fish waste (such as skin and bone) generated from fish processing, being rich sources of collagen, has found increased usage via extractions and characterization, thereby expanding the scope of collagen extraction from fish waste (Mahboob, 2015). Fish wastes have also found profound usage in fish oil extraction for leather tanning and the production of hydrolysed oils (Saranya et al., 2020; Nascimento et al., 2015). Another well-renowned method is the process of

fermentation, majorly used for silage production (Lindgren and Pleje, 1983) and animal feed production (Faid et al., 1997). Recent times have seen this process as a potential technique for handling environmental issues caused by fish wastes/fish by-products (Marti-Quijal et al., 2020). Other products produced

involve generating high-value compounds such as bioactive peptides and antifungal compounds. Scrutiny also reveals recent trends towards recovering oils and fats using enzymatic hydrolysis (Fadhil et al., 2017). Author keyword analysis primarily reveals significant contributions and combinations



(a)



(b)

Figure 5. (Continued)




Figure 5. Keyword network with (a) clustered index keywords (note: threshold criteria of minimum four keywords, resolution=0.8), (b) trends in keyword usage (note: threshold criteria of minimum four keywords, resolution=0.8), and (c) cloud of frequency of keywords.

Table 5. Top keyword occurrence.

Keyword	Occurrences	TLS	Average year
Fish waste(s)	134	154	2014
Biodiesel	29	43	2016
Aquaculture	27	52	2015
Anaerobic digestion	17	24	2016
Collagen	16	25	2014
Biogas	15	18	2016
Fish oil	15	33	2016
Fermentation	14	19	2009
Fish by-products	12	16	2008
Transesterification	12	34	2018
Enzymatic hydrolysis	11	33	2017

TLS: total link strength.

existing between fish waste(s)–biodiesel, fish waste(s)–aquaculture and fish waste(s)–anaerobic digestion.

The top 10 identified keywords based on the frequency of occurrence and TLS is reported in Table 5, signifying the average year of occurrence. Keyword valuation carried out considering the last four years’ research (i.e. 2017–2020) indicates the rising emphasis on hydroxyapatite, bio-methane, waste management, biofuel, transesterification, biomass and RRM in the domain of fish waste(s). These indicate interest among researchers, encouraging the addition of prodigious work and satisfying RQ2.

Hydroxyapatite has recently found increasing interest among research groups. This is evident from the synthesizing studies by Yamamura et al. (2018) from fish wastes. This extraction modus has even reached prodigious heights of having its own review work done by Omar et al. (2019), clearly citing its rich influence on the scientific community.

In general, both author and index keywords indicate the prima foci of research, with index keywords highlighting an increasing focus on concepts that author keywords that fail to detail. To analyse and identify the research hotspots evident via analysing index keywords, we categorize the various keywords into clusters using the innate software functionality. Cluster analysis reveals the categorization of keywords into six clusters. Despite variations among keywords between clusters, there happen to be visible coherences existing between analysed clusters. Outputs from the cluster analysis have been detailed in Table 6.

Network analysis

This analysis is initiated by conducting bibliographic coupling, citation, co-citation and co-authorship analysis to understand individual interrelations between authors, countries and organizations. This will enable in understanding the undertaken research collaborations and citation dependencies across various strata

Table 6. Cluster wise categorization.

Identity	Cluster name	Discussion
Cluster 1	Aquaculture	The most discussed topic under this cluster were solid wastes (Bureau and Hua, 2010), waste disposal (Wu and Song, 2021b), nitrogen generation (von Ahnen et al., 2020) and the effect of bacteria and microorganisms (Schneider et al., 2006). Aquaculture, in general, has become research within itself, with prominent research being done on the usage of fish wastes as a feed product (Aifreen and Ucak, 2020).
Cluster 2	Non-human consumption (in direct forms)	This cluster comprises the extraction and valorization of fish by-products. Fish waste possesses antioxidant nature (Elvevoll, 2007), proving consumable in a valorized form. Pivotal keywords identified deals with the derivation of proteins and amino acids via hydrolysis (Lü et al., 2007) and enzymatic hydrolysis (Hathwar et al., 2011). However, increased focus is found on bioactive gene compounds (Hosseini and Zandi, 2017) and bioactive peptides (Coppola et al., 2021). Extensive research is evident on marine collagen and chitosan for biomedical applications (Irasorza et al., 2021) and food packaging film (Bhuiabar et al., 2019).
Cluster 3	Fish wastes characterization	The essential keywords identified involve bacteria (microorganisms), food processing, fermentation, pH, microbial activity, enzyme activity, biotechnology and food processing. Lactic acid bacteria (microorganisms) in fermentation during fish waste processing (Martí-Quijal et al., 2020) play a critical role in enzymatic activity. These procedures have become feasible considering the recent advances in biotechnology (Ramkumar et al., 2016).
Cluster 4	Food aspect of fish wastes	This cluster covers animal food/feed, dietary supplements, fish proteins and protein hydrolysate. The conversion of fish wastes into a dietary product for human and animal consumption has increased the scope of nutrition capture. Fish waste has been used as animal feed (Faid et al., 1997) due to its inherent protein content and ability to get converted into high value-added products by added processing (Araujo et al., 2021). Fish protein hydrolysate is one such product that has gained importance due to its growth and health properties (Ildowu et al., 2020). The rising use of fish waste-derived food supplements owing to its good Omega 3 constituent (Alfio et al., 2021) has amplified the consumption of fish wastes (though indirectly) among humans.
Cluster 5	Fish wastes and associated extraction techniques	Anaerobic digestion is the prime extraction method resulting from the extraction of methane, biogas and biofuels (Bücker et al., 2020; Wu and Song, 2021a, 2021b). Containing high amounts of biodegradable organics, fish wastes also prove attractive for producing activated sludge (Coppola et al., 2021).
Cluster 6	Biodiesel	The critical keywords identified are fatty acids, fish oil, transesterification and catalysis. Characterized by the fatty acids present in fish wastes, biodiesel produced from fish wastes by enzymatic catalysis (Ching-Velasquez et al., 2020) allows for reduced environmental pollution. Notable works are also found on transesterification for biodiesel synthesis (Jung et al., 2019). A similar research trend is also witnessed in bioreactors which are used for the enzymatic conversion of fish oils to biodiesel fuels (Bücker et al., 2020)

Table 7. Bibliographic coupling of top authors, countries and organizations.

Top authors	Top countries	Top organizations
Deepika Dave, Memorial University of Newfoundland, Canada (4406)	India (5610)	SINTEF Ocean, Norway (887)
Rasa Šližytė, SINTEF Ocean, Norway (3261)	Italy (4114)	Department of Integrative Marine Ecology, Stazione Zoologica Anton Dohrn, Italy (765)
Suzanne M. Budge, Dalhousie University, Canada (3146)	Spain (4061)	Department of Marine Biotechnology, Stazione Zoologica Anton Dohrn, Italy (765)
Abdel E. Ghaly, Dalhousie University, Canada (3146)	Unites States (3298)	Institute of Biochemistry and Cell Biology (Ibbc), National Research Council, Italy (765)
Su Ling Brooks, Dalhousie University, Canada (2396)	China (3197)	Institute of Biosciences and Bioresources (Ibbr), National Research Council, Italy (765)
Keiko Shirai, Universidad Autónoma Metropolitana – Unidad Iztapalapa, Mexico (2395)	Brazil (2990)	Institute of Polymers, Composites and Biomaterials, National Research Council, Italy (765)
Ioannis Sotirios Arvanitoyannis, Panepistimio Thesalias, Greece (2334)	Australia (2731)	Bapalal Vaidya Botanical Research Centre, Department of Biosciences, Veer Narmad South Gujarat University, India (752)
Narayan Bhaskar, Central Food Technological Research Institute India (2303)	Malaysia (2329)	Department of Biology, College of Science, University of Hail, Saudi Arabia (752)
Carlos Prentice, Universidade Federal do Rio Grande, Brazil (2158)	Canada (2170)	Department of Clinical Nutrition, College of Applied Medical Sciences, University of Hail, Saudi Arabia (752)
J. A. Siles, Department of Inorganic Chemistry and Chemical Engineering, Spain (2000)	Mexico (1818)	Department of Oral Radiology, College of Dentistry, University of Hail, Saudi Arabia (752)

Threshold of one document per author and five citations per document. () – value within parenthesis represents the TLS value.

Table 8. Citation analysis by documents.

Cited as	Citations	Links
Kristinsson and Rasco (2000)	702	12
Muyonga et al. (2004)	402	3
Wu (1995)	400	2
Van Rijn et al. (2006)	394	1
Nagai and Suzuki (2000)	381	2
Jayathilakan et al. (2012)	342	10
Arvanitoyannis and Kassaveti (2008)	242	6
Mshandete et al. (2004)	211	5
Álvarez et al. (2010)	207	1
Sathivel et al. (2003)	202	1

hence satisfying RQ3. Representations are primarily interpreted using circular nodes and connecting lines, revealing the number of citations and co-citations achieved.

Bibliographic coupling. This analysis achieved by coupling bibliographic data aids in identifying adopted collaborations to solve a particular research problem (i.e. fish waste management in our case). Ranked using the TLS values (mentioned in parenthesis), this analysis denotes the link between two different documents that cite the same document (i.e. they share the exact references). Documents have a strong bibliographic coupling depending on the significant coexistence of references existing between publications. Bibliographic coupling is performed within top authors, top countries and top organizations. Table 7 reveals the outcomes from the bibliographic coupling. Valuable contributions are evident from developed countries despite a significant share from developing nations such as India, China, Brazil and Malaysia

with India topping the list of developing nations offering substantial contributions to the domain.

Citation analysis. Majorly used as an indicator for measuring research influence (Bonilla et al., 2015), a citation analysis measures the scientific quality of a research/researcher/organization/country. Despite older papers gaining large citation counts compared to recent contributions (Bornmann and Williams, 2013), the likelihood of being discovered decreases due to outdated research outputs. This gets further evident from Table 8, where research by Kristinsson and Rasco (2000) has garnered higher citations than Wu (1995). To further explore the influence of research, citation analysis is adopted across documents, countries and organizations (Figure 6).

A document intended citation analysis (Table 8) reveals the contributions of Kristinsson and Rasco (2000), Muyonga et al. (2004) and Wu (1995) to be the most cited documents




Figure 6. Citation analysis: (a) citation by authors, (b) citation by organization, and (c) citation by country.

(citation > 400) among the top 10 research articles addressed, with only one research document in 2010–2020 by Jayathilakan et al. (2012). An author wise citation analysis reveals derivatives similar to the document wise citation analysis w.r.t the contributions from Kristinsson and Rasco (2000). Themed on Fish protein hydrolysates, Kristinsson’s research to date still is of significant influence to the academic world.

A similar analysis focusing on countries reveals the United States to be the most cited country with 2928 citations, followed by India and Spain. Organization wise citation assessment shows that the University of Massachusetts, Washington State University and the University of Washington (all from the United States) are the most cited organization, constituting 14.34% of the mentioned documents among the top 10 highly cited organizations worldwide (Figure 6). As a reiterative technique, a bibliometric analysis of the organization’s contribution revealed similar outcomes validating the contributions achieved.

Co-citation analysis. The analysis is an indicator of the relationship existing between two documents. This analysis forms ideal for establishing thematic relevance between research fields if two recent research articles quote two other older articles. Therefore, it creates a co-citation value of two, thereby establishing scientific contributions based on demonstrated scientific knowledge (Small, 1973).

An insight into the top 10 authors by co-citation reveals that around 6 authors have a citation above 100, with 1 author exceeding 200 sources (Table 9). Network analysis further reveals a working prominence in works by Soottawat Benjakul with Fereidoon Shahidi, Asbjörn Gildberg, Hördur G. Kristinsson and Moncef Nasri. Combined works are mainly evident with Fereidoon Shahidi, who have often worked together (Benjakul et al., 2012). Soottawat Benjakul is also found to be contributing with Chantachum et al. (2000), Klomklao et al. (2009), Thiansilakul et al. (2007), Phanturat et al. (2010) and Nalinanon et al. (2011). A similar collaboration is also witnessed among Asbjörn Gildberg, Rasa Šližytė and Hördur G. Kristinsson (Thorkelsson et al., 2009). To gain a more profound directive, a reference-based co-citation is also carried out (Table 10). A large amount of co-citation was also found to exist between three prominent publishing sources, that is, *Food Chemistry* (867 citations), *Aquaculture* (792 citations) and *Bioresource Technology* (709 citations).

However, an analysis is carried out to understand the correlation between the top authors and top references (by co-citation). Results reveal augmented contributions from Benjakul, S., Shahidi, F., Gildberg, A., Kristinsson, H.G. and Visessanguan, W., who have had noted impacts in terms of contribution and referencing. Among the top 10 references (by co-citation), 4 were published by the top authors (six authors) enlisted by co-citation.

Table 9. Top 10 authors by co-citation.

Author	Organization	Region	Citations	Link	TLS	Cluster
Benjakul, S.	Prince of Songkla University	Thailand	285	878	30574	7
Kim, S. K.	Korea Maritime and Ocean University	South Korea	170	786	24092	4
Fereidoon Shahidi	Memorial University of Newfoundland	Canada	168	888	20670	4
Asbjörn Gildberg	NOFIMA	Norway	127	753	11982	2
Hördur G. Kristinsson	Haskoli Islands	Iceland	119	752	12787	4
Moncef Nasri	Ecole Nationale d'Ingenieurs de Sfax	Tunisia	115	800	12419	3
Turid rustad	Norges teknisk-naturvitenskapelige universitet	Norway	96	868	9719	2
Montero, Pilar G.	Universidad de Granada	Spain	93	726	11098	7
Wonnop Visessanguan	Thailand National Center for Genetic Engineering and Biotechnology	Thailand	92	729	10498	7
Narayan Bhaskar	Central Food Technological Research Institute	India	91	728	5531	2

TLS: total link strength.

Table 10. Top 10 references by co-citation.

Cited reference	Citations	TLS	Cluster	Link
Mshandete et al. (2004)	10	28	3	26
Liaset et al. (2000)	5	19	3	18
Chalamaiah et al. (2012)	5	13	3	13
Kristinsson and Rasco (2000)	5	13	3	13
Ghaly et al. (2013)	5	8	3	7
Crab et al. (2007)	5	5	4	5
Nges et al. (2012)	5	2	3	1
Sarmadi et al. (2010)	4	38	3	36
Villamil et al. (2017)	4	33	3	31
Arvanitoyannis and Kassaveti (2008)	4	32	3	32

TLS: total link strength.

Mshandete (Anthony Manoni Mshandete) is a Professor of Biotechnology at the Nelson Mandela African Institute of Science and Technology. The second most co-cited reference belongs to Bjoern Liaset, who works at the Directorate of fisheries at the Institute of Nutrition, Norway. Furthermore, the network analysis reveals that 9 references (all belonging to cluster 3) were co-cited among the 10 references found via co-citation, except Crab et al. (2007). This represents an unbroken link being progressed by the research loop that requires insight.

Co-authorship analysis. The analysis assists in revealing the scholarly collaborations given the contributing authors, countries and organizations (Uddin et al., 2012). Setting an initial threshold of one document per author with a minimum number of five citations, the contributions of 1364 authors out of the 2587 authors assessed are presented. The primary analysis of the authors shows the formation of seven clusters (Figure 7(a)), with each cluster posing its contribution. Strong collaboration is visible in the cluster among all authors in cluster 1 (seven authors) from China. Cluster 2 (six authors) emanated dominance from Liu Y. (Kochi University, Japan) and five other authors. Cluster 3 (seven authors) had maximum networking impacts from Luo G. and Tan H. from Shanghai Ocean University, China. Cluster 4 (seven authors) is one of a kind cluster focussing only on

Indian contributors. Maximum productivity of Madhu D. and Sharma Y.C. belonging to the IIT, Banaras Hindu University, India, dominate their author cluster. Relatively, cluster 5 (five authors) is dominated by Liu Z., with all author collaborations from China. Comprising five authors, cluster 6 involves scholarly impacts from Fang J. (Yellow Sea Fisheries Research Institute Chinese Academy of Fishery Science, China) and Jansen H.M. (Wageningen University & Research, Wageningen, the Netherlands). Finally, cluster 7 (four authors) is led by the works of Jiang Z. University of Agder, Norway, and Li I. Universitetet i Stavanger, Norway. However, it is also witnessed that most of the authors had a geographical influence while networking making them collaborate among authors available in proximate localities.

To further explore the level of co-authoring existing, an analysis based on countries and organizations is carried out (Figure 7(b)). Cluster wise categorization reveals classification into 11 clusters with South Korea (cluster 1 has ten countries), Indonesia (cluster 2 has seven countries), Brazil (cluster 3 has seven countries), Germany (cluster 4 has six countries), Norway (cluster 5 has five countries), Spain (cluster 6 has five countries), India (cluster 7 has five countries), France (cluster 8 has five countries), United States (cluster 9 has four countries), China (cluster 10 has three countries) and Italy (cluster 11 has three




Figure 7. Network analysis: (a) author collaboration network (note: threshold criteria of 1, resolution=0.8), (b) collaboration network between countries (note: threshold criteria of 1, resolution=0.8), and (c) collaboration network between organizations (note: threshold criteria of 1, resolution=0.8).

countries) leading each cluster. Analysis of node size further reveal magnified influence from the United States, India and Spain. Posing a robust well-connected network, the representations indicate more citations when the publications originate from these nations. An organization-oriented assessment ($n=1597$) reveals unique collaborations between organizations in the Scandinavian region (Figure 7(c)). The outcomes derived in two clusters showcase a total of 8 organizations. The derivatives attained direct organizational research impacts, making the Nordic regions a prolific domain for fish waste-related research.

Qualitative content analysis

Results denote the clustering of research articles based on citation by references. Categorizing the articles into 4 clusters, as evident in Section 3.3.3, an exploratory review is proceeded to understand and uncover the research directions. Clustering reveals that 15 articles are categorized in four clusters after satisfying a threshold criterion of 4 citations (local citations). Table 11 provides a holistic view of the various clusters identified and provides a concise description of each cluster.

Table 11. Results from clustering with theoretical direction.

Cluster	Articles	Research direction	Description	References
Cluster 1	5	Extraction technique	Co-products utilization techniques	Chantachum et al. (2000); Sarmadi and Ismail (2010); He et al. (2013); Ghaly et al. (2013)
Cluster 2	4	Protein generation	Fish protein hydrolysates	Liaset et al. (2000); Karim and Bhat (2009); Chalamaiah et al. (2012); Villamil et al. (2017)
Cluster 3	4	Management of Fish wastes	Anaerobic digestion	Mshandete et al. (2004); Chen et al. (2008); Arvanitoyannis and Kassaveti (2008); Nges et al. (2012); Kristinsson and Rasco (2000); Ghaly et al. (2013)
Cluster 4	2	Nutrient sources	Nutrient source	

Cluster 1: Extraction technique. The direction adopted in this cluster denotes a developing pattern initiating from extraction procedures adopted throughout the analysed articles. Chantachum et al. (2000) explain the procedural extraction of fish oil from fish discards. Though this article stands alone from the other themes discussed, the reference to using the wet rendering method for extraction makes this article related to the other research outcomes discussed under this cluster. Different significances include discussing enzymatic extraction procedures correlating with the other texts discussed further. This paper's importance is supported by the latest works on fish oil extraction using enzymatic hydrolysis (Araujo et al., 2021; Zhang et al., 2021). He et al. (2013) focused on protein extraction using protein hydrolysates. The authors also suggest the hydrolysis technique using enzymes to be a better adoption owing to the milder reactions witnessed with higher quality and functionality of the derived product. However, the authors cite high expenses in technology, confirming its effectiveness in improving RRM use in processing industries. Stressing on the characteristic of fish protein extracted, Sarmadi and Ismail (2010) highlighted the presence of anti-oxidative properties in the food proteins. Ghaly et al. (2013) elaborated analogous information and the various value-added products derivable using enzymatic hydrolysis.

Cluster 2: Protein generation. Cluster 2 relates to the generation of proteins from fish discards and encompasses review papers, indicating the positive growth of this cluster focusing on protein hydrolysates (Idowu et al., 2020). Liaset et al. (2000) focused on producing and characterizing proteins from salmon and codfish frames. Keeping the derivatives obtained as struts, Karim and Bhat (2009) understood the dire need for an alternative source for gelatin (i.e. protein form) which traditionally had a mammalian origin. The author's prime focus was to assess the market acceptability for the new product with competitive and similar properties to existing alternatives. Chalamaiah et al. (2012) published a review highlighting the generation of fish protein hydrolysates. Apart from focusing on hydrolysates and compositions of amino acids, the authors also emphasized the dual benefit of being both anti-oxidative and a nutritious alternative for humans and aquaculture feeds. A more recent addition to this cluster is the contributions of Villamil et al. (2017) which stressed the adverse challenges such as preserving product quality and developing cheap operative procedures for recovering proteins in agricultural, cosmetic, pharma and food industries.

Cluster 3: Management of fish wastes. Cluster 3 deals with the anaerobic digestion of fish wastes and highlights the acceptance of this technique even in recent times. Recent works involve the sustainable digestion of fish wastes using anaerobic digestion by Nazurally (2018), reducing sludge production while producing biogas using anaerobic digestion (Choi, 2020). However, analysis reveals trailing literature confined to four research articles discussed in this cluster. Mshandete et al. (2004) explored the possibility of producing biogas using an anaerobic digestion process with fish waste and sisal pulp. Despite this article being a profound analysis, the intent to use an anaerobic digestion process for biogas generation makes this research article imperative to other research articles discussed under this cluster. Considering the laid focus on anaerobic digestion, Chen et al. (2008) examined the various underperforming factors that reduce the efficacy of the anaerobic digestion process. Arvanitoyannis and Kassaveti (2008) published a reputed article on treatments, impacts and issues of using fish industry wastes. Though this directly does not address the digestion process, the article touches on the concept of biogas production and a list of other renowned management methods that make this article coherent. Nges et al. (2012) anaerobically digested fish waste and fish sludge to study the properties and characteristics of the methane produced.

Cluster 4. Nutrient sources. Nutrient derivative, a commonly discussed topic under this section, has been extensively discussed, focusing on the derivation of proteins from fish wastes. Articles such as Alfio et al. (2021) emphasize the sustainable recovery of Omega 3 fatty acids from fish wastes; Coppola et al. (2021) concentrated on the various value-added products derivable from fish wastes. Radziemska et al. (2019) applied the use of fish waste for compost. Arvanitoyannis and Tserkezou (2014) focused on the generalized fish waste management techniques, found to be extensively based on articles from this cluster. Kristinsson and Rasco (2000) discuss the various features of fish protein hydrolysates, including the production, application and comparison with other food derived protein hydrolysates. The other article discussed under this cluster that extracts and expands the outcomes derived includes the article by Ghaly et al. (2013). Besides focusing on fish protein hydrolysates, the authors try to cover various value-added products. The article also deeply discusses the different methods of extractions available and the cons of each technique focusing on the biological aspect.




Figure 8. Barriers to fish waste valorization (Indian context).

Impediment assessment of sustainable fish waste management in India

Outcomes from bibliometric analysis conclude the undermined focus among researchers on supply chain-related aspects in fish waste management and meagre commercialization of fish waste valorized products despite noticeable research contributions. The limited focus on supply chain aspects of fish waste management envisages gaining concentrated focus. Furthermore, the capitalization of valorized co-streams generated from fish processing has only been witnessed in published literature from nations such as Iceland and Norway (Olsen et al., 2014). This could be possible due to the integrated nature of supply chains in these countries that permit the rapid utilization of fish wastes that is prone to high-quality degradation. Literature additionally reveals dynamically reducing quality (Amos et al., 2007), high valorizing costs (Olsen et al., 2014), availability of preferable alternatives (Karim and Bhat, 2009) and reduced RRM & by-product yield (Gildberg and Stenberg, 2001) to hamper the effective valorization of fish wastes generated. This holds essential considering the dependent nature of perishable food on handling and

storage conditions (Jensen et al., 2010), thereby necessitating the need to undertake supply chain-related research, particularly in this domain, especially in nations that do not have vertically integrated supply chain structure in fisheries.

To accentuate the stresses hampering the adoption of sustainable practices in the Indian context, a barrier assessment validated by stakeholder perception is carried out. The existing fish waste management scenario in India is detailed in the Supplementary Material of the manuscript. Forty-eight barriers identified from literature after appropriate stakeholder validations are classified into eight categories: government issues, local participation, public participation, training/skill development, quality, infrastructure, processing and cost (Refer Figure 8). The various barrier categories have been elucidated in the Supplementary Material due to space constraints and summarized in Table 12, hence satisfying RQ4 of this research.

The way forward?

Assessments reveal the need for regulatory bodies to play dominant roles involving the immediate need to frame seafood waste

Table 12. Barrier wise identification of issues and proposed remedies.

Barrier category	Issues	Proposed remedial measure
Local participation	Investment risks Lack of technical assistance Product marketability Fear of workforce redundancy Reduced eco-literacy Reduced top management willingness	Stakeholder education and training Government should act as a linking block between research organizations and industries Cost-benefit analysis should be done before any investments
Public participation	Achieve strategic competitiveness Complex decision-making processes Demand planning	Understand consumer acceptability Involve public opinion during decision making Consumer awareness programs using community drives and advertisements to design appropriate strategies Economic stimulus Involvement of local municipal authorities can induce public involvement
Quality	Maintain controlled temperatures Delayed processing Broken cold chain Initial investments	Refine internal organizational infrastructure Make definition of quality uniform across the supply chain Administrative callouts regarding planned expenditure Use of government benefits such as tax exemptions (80IB,35AD) and excise duty exemptions Government supported financial assistance @ zero-interest loans
Processing	Business sustainability Dynamic seafood availability Inadequate infrastructure	Create dedicated RRM pools Adopt state-level procurement and processing strategies Control quality Regulate logistics and control operational costs State-owned enterprises should pave structured operating roadmaps
Infrastructure	Infrastructure restructuring Managing operation capacities Demand planning in unpredictable market scenarios Cold chain Localization Logistics network Fragmented supply chain Co-ordination	Supply chain networked via a common platform. Possible technological instigation such as blockchain technologies. Agile logistics should be introduced to make raw material utilization swift Introduce supply chain localization Localizing processors within a particular zone will influence transportation costs forming a hub and spoke network Aggregated RRM can be processed in localized processing units saving global emissions Reduce procurement costs, fuel costs and raw material inventory costs
Costs	Cost crunches Financial assistance	Interest-free credits, financial assistance (such as tax exemptions) Government financial aids
Training/skill development	Gain technical skills post improvement adoptions	—
Government initiatives	—	Frame strict legislatures Account wastes generated from seafood processors Levy waste taxes for by-product discards Penalize non-practicing industries Setup a centralized authority body for managing fish waste Promote industry-research collaborations Introduce incentives comprising tax evasions Financial aid for purchasing required infrastructure Pave structured guidelines/framework for managing RRM

RRM: rest raw material.

management plans requiring extensive contributions from local municipal authorities and NGOs. A significant share of the concerned bodies should focus on educating investors regarding sustainable business opportunities in the domain of by-product valorization in India. Apart from the mere framing of regulations, an active region-level monitoring committee need to be appointed that is centrally controlled by a national-level monitoring committee. Knowledge sharing sessions need to be arranged between NGOs, government institutions, and central and internationally acclaimed seafood research laboratories with a scope of venturing into collaborations. A long-term government-run initiative might not be an alternative to this issue, hence requiring the informal sector to step into the domain to gain substantial market benefits by producing and exporting valorized products to international markets. Focus also need to be laid on training and capacity building on every stakeholder in the seafood supply chain, considering the conglomerated effect in moving towards a sustainable Indian seafood supply chain.

The implications of research derived from this study imply applying sustainable practices in the Indian seafood sector, as mentioned by Sultan et al. (2021). The outcomes derived mainly hold high importance due to the extensive possibilities of exploring viable markets to understand and foresee issues before facilitating breakthrough innovations in India's valorized seafood by-products.

Conclusion and recommendation

This research aims to record the various developments in managing fish waste. In general, processes include the conversion of fish wastes into low-value products such as fish meals, fish oils, and other high-value products such as fish protein hydrolysate, bio-gas/bio-methane production, collagen extraction, and various other by-products. This research first uses bibliometric analysis to address the current study status, research contexts explored, and research hotspots in the global scenario. Secondly, the explorative research further finds possible reasons and proposes developments for improving fish waste utilization in India by providing localized implication strategy(s). The limited availability of high-value valorized co-stream products as a dominant market substitute among the conventional products in the Indian market instigates the authors to explore the variable causes for the underutilization of fish wastes. The generalized conclusion derived from this research are as follows:

- Research in managing fish wastes has improved in the last decade, with a significant share of work carried out on biological, chemical and characteristic aspects of various laboratory-derived by-products.
- Geo-local contributions reveal twofold outcomes. Despite dominance in research contributions to be evident from India, favoured research adaptability was found to be among developed countries (80%) denoting preference to research from developed nations (RQ1).
- Keyword evaluations considering the last quadrennial (i.e. from 2017 to 2020) reveal prolific studies on hydroxyapatite, bio-methane, waste management, biofuel, transesterification, biomass and RRM in the domain of fish waste(s) (RQ2).
- Strong collaboration is visible among clusters with geographical influences playing significant roles in collaborative networks formed between authors, with most authors willing to collaborate within proximate localities (RQ3).
- Scandinavian countries significantly contribute to sustainable utilization and consumption of seafood by-products with an excessive amount of research collaborations and a wider acceptance (in citations) in the international academic community.
- Market availability of valorized fish by-products is difficult due to expensive commercialization processes, fewer interested investors and better properties shown by existing products.
- The supply chain development with a high level of vertical integration and modern technology can be a potential solution for the enhanced utilization of processing derived fish co-streams. With modern technology, countries like Iceland and Norway have a vertically integrated inbound supply chain. However, it is observed that though the supply chain in India is fragmented with comparatively nascent technologies in place, not much research has been reported in the literature for this understudied area.
- Government role, local and public participation, training/skill development, quality, costing, processing and infrastructure (cold chain in specific) along the supply chain were the identified barriers hindering India's effective utilization of fish by-products (RQ4).
- The outcomes obtained can bridge the large gap in seafood quality between nations (both product and utilization wise), hence laying the struts for a circular and sustainable supply chain for processed seafood.

Future research can be proceeded by documenting upstream and downstream activities in the supply chain, cost analysis for localization and by-product valorizations to develop a vertically integrated supply chain. Evident work may be carried out on strategy development for RRM consolidation, wash water utilization, productivity improvement and marketability of value-added products.

Declaration of conflicting interests


The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The authors disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work is supported by the ReValue project funded through the INNO-Indigo programme under a DBT India grant (BT/IN/INNO-Indigo/12/NK/2017-18 dt. 21/02/18) and the Research Council of Norway project number 281262.

ORCID iDs

Farook Abdullah Sultan  <https://orcid.org/0000-0002-8757-8723>

Srikanta Routroy  <https://orcid.org/0000-0002-4592-0009>

Supplemental material

Supplemental material for this article is available online.

References

- Adler SA, Honkapää K, Saarela M, et al. (2014) Utilisation of co-streams in the Norwegian food processing industry. A multiple case study *Bioforsk Rapp. Bioforsk Rep.*, 9 Nr. 82.
- Afreen M and Ucak I (2020) Fish processing wastes used as feed ingredient for animal feed and aquaculture feed. *Survey in Fisheries Sciences* 6: 55–64.
- Ahuja I, Dauksas E, Remme JF, et al. (2020) Fish and fish waste-based fertilizers in organic farming – with status in Norway: A review. *Waste Management* 115: 95–112.
- Alfio VG, Manzo C and Micillo R (2021) From fish waste to value: An overview of the sustainable recovery of omega-3 for food supplements. *Molecules* 26: 1002.
- Álvarez JA, Otero L and Lema JM (2010) A methodology for optimising feed composition for anaerobic co-digestion of agro-industrial wastes. *Bioresource Technology* 101: 1153–1158.
- Amos B, Sector F, Einarsson H, et al. (2007) *Analysis of Quality Deterioration at Critical Steps/Points in Fish Handling in Uganda and Iceland and Suggestions for Improvement*. Uganda: United Nations University.
- Ante L, Steinmetz F and Fiedler I (2021) Blockchain and energy: A bibliometric analysis and review. *Renewable and Sustainable Energy Reviews* 137: 110597.
- Aranganathan L and SR RR (2016) Bioconversion of marine trash fish (MTF) to organic liquid fertilizer for effective solid waste management and its efficacy on Tomato growth. *Management of Environmental Quality: An International Journal* 27:93–103
- Araujo J, Sica P, Costa C, et al. (2021) Enzymatic hydrolysis of fish waste as an alternative to produce high value-added products. *Waste and Biomass Valorization* 12: 847–855.
- Arruda LFD, Borghesi R and Oetterer M (2007) Use of fish waste as silage: A review. *Brazilian Archives of Biology and Technology* 50: 879–886.
- Arvanitoyannis IS and Kassaveti A (2008) Fish industry waste: treatments, environmental impacts, current and potential uses. *International Journal of Food Science and Technology* 43: 726–745.
- Arvanitoyannis IS and Tserkezou P (2014) Fish waste management. In: Bozaris IS (ed.) *Seafood Processing: Technology, Quality and Safety*. John Wiley & Sons, Ltd. pp. 263–309.
- Benjakul S, Nalinanon S and Shahidi F (2012) Fish collagen. In: Simpson BK (ed.) *Food Biochemistry and Food Processing* (2nd ed.). Wiley-Blackwell. pp. 365–387.
- Bhuimbar MV, Bhagwat PK and Dandge PB (2019) Extraction and characterization of acid soluble collagen from fish waste: Development of collagen-chitosan blend as food packaging film. *Journal of Environmental Chemical Engineering* 7: 102983.
- Binsi PK (2018) *Overview of waste generation in fish and shellfish processing industry*. Cochin: ICAR-Central Institute of Fisheries Technology.
- Bonilla CA, Merigó JM and Torres-Abad C (2015) Economics in Latin America: A bibliometric analysis. *Scientometrics* 105: 1239–1252.
- Bornmann L and Williams R (2013) How to calculate the practical significance of citation impact differences? An empirical example from evaluative institutional bibliometrics using adjusted predictions and marginal effects. *Journal of Informetrics* 7: 562–574.
- Bouzembrak Y, Klüche M, Gavai A, et al. (2019) Internet of things in food safety: Literature review and a bibliometric analysis. *Trends in Food Science and Technology* 94: 54–64.
- Bücker F, Marder M, Peiter MR, et al. (2020) Fish waste: an efficient alternative to biogas and methane production in an anaerobic mono-digestion system. *Renewable Energy* 147: 798–805.
- Bureau DP and Hua K (2010) Towards effective nutritional management of waste outputs in aquaculture, with particular reference to salmonid aquaculture operations. *Aquaculture Research* 41: 777–792.
- Caruso G (2015) Fishery wastes and by-products: A resource to be valorised. *Journal of Fisheries Science* 9: 80–83.
- Caruso G, Floris R, Serangeli C, et al. (2020) Fishery wastes as a yet undiscovered treasure from the sea: Biomolecules sources, extraction methods and valorization. *Marine Drugs* 18: 622.
- Centre for Advance Trade Research (2018) Frozen fish export from India continue to see astounding growth. [Online]. Available at: <https://www.tpci.in/blogs/frozen-fish-export-from-india-continue-to-see-astounding-growth/> (accessed 18 May 2021).
- Chalamaiah M, Hemalatha R and Jyothirmayi T (2012) Fish protein hydrolysates: Proximate composition, amino acid composition, antioxidant activities and applications: A review. *Food Chemistry* 135: 3020–3038.
- Chantachum S, Benjakul S and Sriwirat N (2000) Separation and quality of fish oil from precooked and non-precooked tuna heads. *Food Chemistry* 69: 289–294.
- Chen Y, Cheng JJ and Creamer KS (2008) Inhibition of anaerobic digestion process: A review. *Bioresource Technology* 99: 4044–4064.
- Ching-Velasquez J, Fernández-Lafuente R, Rodrigues RC, et al. (2020) Production and characterization of biodiesel from oil of fish waste by enzymatic catalysis. *Renewable Energy* 153: 1346–1354.
- Choi HJ (2020) Acid-fermented fish by-products broth: an influence to sludge reduction and biogas production in an anaerobic co-digestion. *Journal of Environmental Management* 262: 110305.
- Coppola D, Lauritano C, Palma Esposito F, et al. (2021) Fish waste: From problem to valuable resource. *Marine Drugs* 19: 116.
- Costa DF, de Melo Carvalho F, de Melo Moreira BC, et al. (2017) Bibliometric analysis on the association between behavioral finance and decision making with cognitive biases such as overconfidence, anchoring effect and confirmation bias. *Scientometrics* 111: 1775–1799.
- Crab R, Avnimelech Y, Defoirdt T, et al. (2007) Nitrogen removal techniques in aquaculture for a sustainable production. *Aquaculture* 270: 1–14.
- De Oliveira RI, Sousa SO and De Campos FC (2019) Lean manufacturing implementation: Bibliometric analysis 2007–2018. *The International Journal of Advanced Manufacturing Technology* 101: 979–988.
- De Silva SS and Turchini GM (2008) Towards understanding the impacts of the pet food industry on world fish and seafood supplies. *Journal of Agricultural and Environmental Ethics* 21: 459–467.
- Donthu N, Kumar S, Mukherjee D, et al. (2021) How to conduct a bibliometric analysis: An overview and guidelines. *Journal of Business Research* 133: 285–296.
- Du HS, Xu J, Li Z, et al. (2021) Bibliometric mapping on sustainable development at the base-of-the-pyramid. *Journal of Cleaner Production* 281: 125290.
- Eck NJV and Waltman L (2009) How to normalize cooccurrence data? An analysis of some well-known similarity measures. *Journal of the American Society for Information Science and Technology* 60: 1635–1651.
- Eck NJV and Waltman L (2014). Visualizing bibliometric networks. In *Measuring scholarly impact*. Cham: Springer, pp. 285–320
- Elvevoll EO (2007) *Fish waste and functional foods*. Norway: Norwegian College of Fishery Science, Department of Marine Biotechnology, University of Tromsø.
- ET Bureau (2020) India's share in global seafood trade set to rise from 4.1% to 6.7% by 2030. [Online]. Available at: <https://m.economicstimes.com/news/economy/foreign-trade/indias-share-in-global-seafood-trade-set-to-rise-from-4-1-to-6-7-by-2030/articleshow/74010401.cms> (accessed 19 April 2021).
- Fadhil AB, Ahmed AI and Salih HA (2017) Production of liquid fuels and activated carbons from fish waste. *Fuel* 187: 435–445.
- Faid M, Zouiten A, Elmarrakchi A, et al. (1997) Biotransformation of fish waste into a stable feed ingredient. *Food Chemistry* 60: 13–18.
- Food Security Information Network (FSIN) (2020) Global report on food crisis. [Online]. Available at: <https://www.fsplatform.org/report/global-report-food-crises-2020/> (accessed 5 May 2021).
- Ghaly AE, Ramakrishnan VV, Brooks MS, et al. (2013) Fish processing wastes as a potential source of proteins, amino acids and oils: A critical review. *Journal of Microbial and Biochemical Technology* 5: 107–129.
- Gildberg A and Stenberg E (2001) A new process for advanced utilisation of shrimp waste. *Process Biochemistry* 36: 809–812.
- Hathwar SC, Bijinu B, Rai AK, et al. (2011) Simultaneous recovery of lipids and proteins by enzymatic hydrolysis of fish industry waste

- using different commercial proteases. *Applied Biochemistry and Biotechnology* 164: 115–124.
- Hayes M (2012) Chitin, chitosan and their derivatives from marine rest raw materials: Potential food and pharmaceutical applications. In: *Marine Bioactive Compounds*. Boston, MA: Springer, pp. 115–128.
- He S, Franco C and Zhang W (2013) Functions, applications and production of protein hydrolysates from fish processing co-products (FPCP). *Food Research International* 50: 289–297.
- Hjellnes V, Rustad T and Falch E (2020) The value chain of the white fish industry in Norway: History, current status and possibilities for improvement – a review. *Regional Studies in Marine Science* 36: 101293.
- Hosseini SF and Zandi M (2017) Derivation bioactive compounds from seafood processing by-products. *Utilization and Cultivation of Aquatics* 6: 61–73.
- Idowu AT, Igichon OO, Idowu S, et al. (2021) Bioactivity potentials and general applications of fish protein hydrolysates. *International Journal of Peptide Research and Therapeutics* 27(1): 109–118.
- Idowu AT, Igichon OO, Idowu S, et al. (2020) Bioactivity potentials and general applications of fish protein hydrolysates. *International Journal of Peptide Research and Therapeutics*: 1–10.
- Inamdar Z, Raut R, Narwane VS, et al. (2020) A systematic literature review with bibliometric analysis of big data analytics adoption from period 2014 to 2018. *Journal of Enterprise Information Management*. DOI: 10.1108/JEIM-09-2019-0267
- Irastorza A, Zarandona I, Andonegi M, et al. (2021) The versatility of collagen and chitosan: From food to biomedical applications. *Food Hydrocolloids* 116: 106633.
- Islam MS (2006) Use of bioslurry as organic fertilizer in Bangladesh agriculture. In: *Prepared for the presentation at the international workshop on the use of bioslurry domestic biogas programme*, Bangkok, Thailand, 2006, pp. 3–16.
- Islam MJ and Peñarubia OR (2021) Seafood waste management status in Bangladesh and potential for silage production. *Sustainability* 13: 2372.
- Jayathilakan K, Sultana K, Radhakrishna K, et al. (2012) Utilization of byproducts and waste materials from meat, poultry and fish processing industries: a review. *Journal of Food Science and Technology* 49: 278–293.
- Jensen TK, Nielsen J, Larsen EP, et al. (2010) The fish industry—toward supply chain modeling. *Journal of Aquatic Food Product Technology* 19: 214–226.
- Jia F and Jiang Y (2018) Sustainable global sourcing: A systematic literature review and bibliometric analysis. *Sustainability* 10: 595.
- Jouvenot L (2015) *Utilisation of rest raw materials from the fish industry: Business opportunities and logistics requirements*. Master's Thesis, NTNU, Norway.
- Jung JM, Oh JI, Park YK, et al. (2019) Biodiesel synthesis from fish waste via thermally-induced transesterification using clay as porous material. *Journal of Hazardous Materials* 371: 27–32.
- Karim AA and Bhat R (2009) Fish gelatin: Properties, challenges, and prospects as an alternative to mammalian gelatins. *Food Hydrocolloids* 23: 563–576.
- Karkal SS and Kudre TG (2020) Valorization of fish discards for the sustainable production of renewable fuels. *Journal of Cleaner Production* 275: 122985.
- Klomkloao S, Kishimura H, Nonami Y, et al. (2009) Biochemical properties of two isoforms of trypsin purified from the intestine of skipjack tuna (*Katsuwonus pelamis*). *Food Chemistry* 115: 155–162.
- Kristinsson HG and Rasco BA (2000) Fish protein hydrolysates: production, biochemical, and functional properties. *Critical Reviews in Food Science and Nutrition* 40: 43–81.
- Kumar V, Muzaddadi AU, Mann S, et al. (2018) Utilization of fish processing waste: A waste to wealth approach. In: *Emerging Post-Harvest Engineering and Technological Interventions for Enhancing Farmer's Income*, vol 04. Compend ICAR Summer School, pp. 127–131.
- Lancho-Barrantes BS and Cantú-Ortiz FJ (2019) Science in Mexico: A bibliometric analysis. *Scientometrics* 118: 499–517.
- Li Z, Chen Z, Yang N, et al. (2021) Trends in research on the carbon footprint of higher education: A bibliometric analysis (2010–2019). *Journal of Cleaner Production* 289: 125642.
- Liaset B, Lied E and Espe M (2000) Enzymatic hydrolysis of by-products from the fish-filleting industry; chemical characterisation and nutritional evaluation. *Journal of the Science of Food and Agriculture* 80: 581–589.
- Lindgren S and Pleje M (1983) Silage fermentation of fish or fish waste products with lactic acid bacteria. *Journal of the Science of Food and Agriculture* 34: 1057–1067.
- Lü F, He PJ, Shao LM, et al. (2007) Effects of ammonia on hydrolysis of proteins and lipids from fish residues. *Applied Microbiology and Biotechnology* 75: 1201–1208.
- Magalhães VS, Ferreira LMD, da Silva César A, et al. (2020) Food loss and waste in the Brazilian beef supply chain: an empirical analysis. *The International Journal of Logistics Management* 32: 214–236.
- Mahboob S (2015) Isolation and characterization of collagen from fish waste material-skin, scales and fins of *Catla catla* and *Cirrhinus mrigala*. *Journal of Food Science and Technology* 52: 4296–4305.
- Marti-Quijal FJ, Remize F, Meca G, et al. (2020) Fermentation in fish and by-products processing: an overview of current research and future prospects. *Current Opinion in Food Science* 31: 9–16.
- Mathew PT (2010) Fish waste utilisation in India, Proceedings of the National Seminar on Conservation and Sustainability of Coastal Living Resources of India, 2009.
- McCarthy JJ (2018) Nearly 50 million tons of fish are wasted every year: UN. [Online]. Available at: <https://www.globalcitizen.org/en/content/world-wastes-50-million-tons-of-fish-each-year/> (accessed 11 April 2021).
- Mishra M, Sudarsan D, Santos CAG, et al. (2021) An overview of research on natural resources and indigenous communities: A bibliometric analysis based on Scopus database (1979–2020). *Environmental Monitoring and Assessment* 193: 1–17.
- Mo WY, Man YB and Wong MH (2018) Use of food waste, fish waste and food processing waste for China's aquaculture industry: needs and challenge. *Science of the Total Environment* 613: 635–643.
- Mohapatra S, Sarkar B, Samantaray DP, et al. (2017) Bioconversion of fish solid waste into PHB using *Bacillus subtilis* based submerged fermentation process. *Environmental Technology* 38: 3201–3208.
- Mongeon P and Paul-Hus A (2016) The journal coverage of Web of Science and Scopus: A comparative analysis. *Scientometrics* 106: 213–228.
- Mozuraityte R, Perote A, Kumari A, et al. (2020) Regulations on the use of rest raw materials from seafood processing in EU and India – ReValue Project Deliverable 4.2 – 2020: 00471 A. Publikasjoner fra CRISTin - SINTEF Ocean.
- Mshandete A, Kivaisi A, Rubindamayugi M, et al. (2004) Anaerobic batch co-digestion of sisal pulp and fish wastes. *Bioresource Technology* 95: 19–24.
- Muyonga JH, Cole CGB and Duodu KG (2004) Characterisation of acid soluble collagen from skins of young and adult Nile perch (*Lates niloticus*). *Food Chemistry* 85: 81–89.
- NAAS (2019) Saving the harvest: Reducing the food loss and waste. Policy Brief No. 5. [Online]. Available at: <http://naas.org.in/documents/Saving%20the%20Harvest.pdf> (accessed 8 May 2021).
- Nagai T and Suzuki N (2000) Isolation of collagen from fish waste material—skin, bone and fins. *Food Chemistry* 68: 277–281.
- Nalinanon S, Benjakul S, Kishimura H, et al. (2011) Functionalities and antioxidant properties of protein hydrolysates from the muscle of ornate threadfin bream treated with pepsin from skipjack tuna. *Food Chemistry* 124: 1354–1362.
- Nascimento VLVD, Bermúdez VMS, Oliveira ALLD, et al. (2015) Characterization of a hydrolyzed oil obtained from fish waste for nutritional application. *Food Science and Technology* 35: 321–325.
- National Fisheries Development Board (NFDB)(2019) Fisheries in India. [Online]. Available at: <http://nfdb.gov.in/> (accessed 18 May 2021).
- Nazurully N (2018) Anaerobic digestion of fish waste and seagrass/macroalgae: Potential sustainable waste management for tropical Small Island Developing States. *Journal of Material Cycles and Waste Management* 20: 1724–1735.
- Nges IA, Mbatia B and Björnsson L (2012) Improved utilization of fish waste by anaerobic digestion following omega-3 fatty acids extraction. *Journal of Environmental Management* 110: 159–165.
- Olsen RL, Toppe J and Karunasagar I (2014) Challenges and realistic opportunities in the use of by-products from processing of fish and shellfish. *Trends in Food Science and Technology* 36: 144–151.
- Omar S, Muhamad MS, Te Chuan L, et al. (2019) A review on lead sources, occurrences, health effects, and treatment using hydroxyapatite (HAP) adsorbent made from fish waste. *Water, Air, and Soil Pollution* 230: 1–21.
- Penven A, Pérez-Gálvez R and Bergé JP (2013) By-products from fish processing: Focus on French industry. *Utilization of Fish Waste*: 232.

- Pham-Duc B, Tran T, Trinh TPT, et al. (2020) A spike in the scientific output on social sciences in Vietnam for recent three years: Evidence from bibliometric analysis in Scopus database (2000–2019). *Journal of Information Science*. Epub ahead of print 30 November 2020. DOI: 10.1177/0165551520977447.
- Phanturat P, Benjakul S, Visessanguan W, et al. (2010) Use of pyloric caeca extract from bigeye snapper (*Priacanthus macracanthus*) for the production of gelatin hydrolysate with antioxidative activity. *LWT - Food Science and Technology* 43: 86–97.
- Pritchard A (1969). Statistical bibliography or bibliometrics. *Journal of Documentation* 25: 348–349.
- Qaiser FH, Ahmed K, Sykora M, et al. (2017) Decision support systems for sustainable logistics: A review and bibliometric analysis. *Industrial Management and Data Systems* 117: 1376–1388.
- Radziemska M, Vaverková MD, Adamcová D, et al. (2019) Valorization of fish waste compost as a fertilizer for agricultural use. *Waste and Biomass Valorization* 10: 2537–2545.
- Ramanujam PK, Aranganathan L and Rajasree SRR (2016) Bioconversion of marine trash fish (MTF) to organic liquid fertilizer for effective solid waste management and its efficacy on tomato growth. *Management of Environmental Quality: An International Journal* 27: 93–103.
- Ramkumar A, Sivakumar N and Victor R (2016) Fish waste-potential low cost substrate for bacterial protease production: A brief review. *The Open Biotechnology Journal* 10: 335–341.
- Rejula K and Mohanty AK (2018) *Innovative extension approaches for fish waste management*. Cochin: ICAR-Central Institute of Fisheries Technology.
- Salam B, Islam M and Rahman MT (2009) Biogas from anaerobic digestion of fish waste. In: *Proceedings of the international conference on mechanical engineering*, Dhaka, Bangladesh, 2009, vol. 26, pp. 1–3.
- Saranya R, Selvi AT, Jayapriya J, et al. (2020) Synthesis of fat liquor through fish waste valorization, characterization and applications in tannery industry. *Waste and Biomass Valorization* 11: 6637–6647.
- Sarmadi BH and Ismail A (2010) Antioxidative peptides from food proteins: A review. *Peptides* 31: 1949–1956.
- Sasidharan A and Mathew S (2011). Current status of fish waste management in Kerala, India. *Waste Management* 31(8): 1898–1900.
- Sasidharan A, Baiju KK and Mathew S (2013) Seafood processing waste management and its impact on local community in Cochin Corporation, India. *International Journal of Environment and Waste Management* 12: 422–441.
- Schneider O, Sereti V, Machiels MA, et al. (2006) The potential of producing heterotrophic bacteria biomass on aquaculture waste. *Water Research* 40: 2684–2694.
- Sharma HB, Vanapalli KR, Cheela VS, et al. (2020) Challenges, opportunities, and innovations for effective solid waste management during and post COVID-19 pandemic. *Resources, Conservation and Recycling* 162: 105052.
- Small H (1973) Co-citation in the scientific literature: A new measure of the relationship between two documents. *Journal of the American Society for Information Science* 24: 265–269.
- Song Y, Wu L and Ma F (2021) A study of differences between all-author bibliographic coupling analysis and all-author co-citation analysis in detecting the intellectual structure of a discipline. *The Journal of Academic Librarianship* 47: 102351.
- Statista (2020) Value of fish and fishery products exported from India from financial year 2011 to 2019, with an estimate for 2020. [Online]. Available at: <https://www.statista.com/statistics/623927/export-value-of-fish-and-fishery-products-india/> (accessed 31 May 2021).
- Sultan FA, Routroy S and Thakur M (2021) A simulation-based performance investigation of downstream operations in the Indian Surimi Supply Chain using environmental value stream mapping. *Journal of Cleaner Production* 286: 125389.
- Thiansilakul Y, Benjakul S and Shahidi F (2007) Compositions, functional properties and antioxidative activity of protein hydrolysates prepared from round scad (*Decapterus maruadsi*). *Food Chemistry* 103: 1385–1394.
- Thorkelsson G, Slizyte R, Gildberg A, et al. (2009) Fish proteins and peptide products: processing methods, quality and functional properties. *Marine Functional food*: 115–133.
- Uddin S, Hossain L, Abbasi A, et al. (2012) Trend and efficiency analysis of co-authorship network. *Scientometrics* 90: 687–699.
- Vanapalli KR, Sharma HB, Ranjan VP, et al. (2021) Challenges and strategies for effective plastic waste management during and post COVID-19 pandemic. *Science of the Total Environment* 750: 141514.
- Van Eck NJ and Waltman L (2013) *VOS viewer manual*. Leiden: Universteit Leiden.
- Van Eck NJ and Waltman L (2014). Visualizing bibliometric networks. In: *Measuring Scholarly Impact*. Cham: Springer, pp. 285–320.
- Van Eck NJ and Waltman L (2019) VOSviewer manual. Available at: https://www.vosviewer.com/documentation/Manual_VOSviewer_1.6.13.pdf (accessed 25 January 2022).
- Van Eck NJ, Waltman L, Dekker R, et al. (2010). A comparison of two techniques for bibliometric mapping: Multidimensional scaling and VOS. *Journal of the American Society for Information Science and Technology* 61: 2405–2416.
- Van Meeteren M, Poorthuis A, Derudder B, et al. (2016) Pacifying Babel's tower: A scientometric analysis of polycentricity in urban research. *Urban Studies* 53: 1278–1298.
- Van Rijn J, Tal Y and Schreier HJ (2006) Denitrification in recirculating systems: Theory and applications. *Aquacultural Engineering* 34: 364–376.
- Villamil O, Váquiro H and Solanilla JF (2017) Fish viscera protein hydrolysates: Production, potential applications and functional and bioactive properties. *Food Chemistry* 224: 160–171.
- von Ahnen M, Pedersen PB and Dalsgaard J (2020) Improving denitrification in an aquaculture wetland using fish waste: A case study. *Ecological Engineering* 143: 105686.
- Wu RSS (1995) The environmental impact of marine fish culture: Towards a sustainable future. *Marine Pollution Bulletin* 31: 159–166.
- Wu Y and Song K (2021a) Anaerobic co-digestion of waste activated sludge and fish waste: Methane production performance and mechanism analysis. *Journal of Cleaner Production* 279: 123678.
- Wu Y and Song K (2021b) Source, treatment, and disposal of aquaculture solid waste: A review. *Journal of Environmental Engineering* 147: 03120012.
- Xie H, Wen Y, Choi Y, et al. Global trends on food security research: A bibliometric analysis. *Land* 10: 119.
- Yahyaee R, Ghobadian B and Najafi G (2013) Waste fish oil biodiesel as a source of renewable fuel in Iran. *Renewable and Sustainable Energy Reviews* 17: 312–319.
- Yamamura H, da Silva VHP, Ruiz PLM, et al. (2018) Physico-chemical characterization and biocompatibility of hydroxyapatite derived from fish waste. *Journal of the Mechanical Behavior of Biomedical Materials* 80: 137–142.
- Zhang M, Gao M, Yue S, et al. (2018) Global trends and future prospects of food waste research: A bibliometric analysis. *Environmental Science and Pollution Research* 25: 24600–24610.
- Zhang Y, Sun Q, Liu S, et al. (2021) Extraction of fish oil from fish heads using ultra-high pressure pre-treatment prior to enzymatic hydrolysis. *Innovative Food Science and Emerging Technologies* 70: 102670.
- Zhongming Z, Linong L, Xiaona Y, et al. (2021) UNEP Food Waste Index Report 2021.
- Zhu S, Jin W and He C (2019) On evolutionary economic geography: A literature review using bibliometric analysis. *European Planning Studies* 27: 639–660.

Appendix. Acronyms table.

Acronym	Full form
DOI	Digital object identifier
FW	Food wastage
H index	Hirsch index
JCR	Journal citation reports
RQs	Research questions
RRM	Rest raw materials
SDG	Sustainable development goal
TLS	Total link strength
UN	United nations
IF	Impact Factor