




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Editorial

Thromboprophylaxis in Patients With Cancer and COVID-19

Interest in thromboprophylaxis has increased since the onset of the SARS-CoV-2 pandemic. It quickly became clear that patients hospitalized for COVID-19 should receive pharmacological thromboprophylaxis with low-molecular-weight heparin (LMWH), unless contraindicated (in line with long-established clinical practice guidelines on antithrombotic prophylaxis in patients hospitalized for an acute medical process). However, early reports of a high rate of thrombotic events in some cohorts of patients hospitalized for COVID-19 despite the use of standard prophylactic doses of LMWH prompted various scientific societies and expert groups to propose the use of higher doses of LMWH in some patient subgroups.^{1,2} At the time, the evidence supporting these recommendations was scant, and, as might be expected, more intense thromboprophylaxis was associated with an increase in hemorrhagic complications.³

Results from randomized clinical trials comparing various intensities of antithrombotic prophylaxis (standard, intermediate or therapeutic doses of LMWH) in hospitalized patients with COVID-19 have helped answer some, but not all, questions. In COVID-19 patients in intensive care units, the use of higher than standard LMWH prophylactic doses does not appear to provide clinical benefit,⁴ but in patients admitted to conventional hospital wards the results are more heterogeneous. In a meta-analysis of aggregated data, therapeutic doses of LMWH compared with standard prophylactic doses were associated with a reduction in thromboembolic events, but also with an increase in major bleeding (although lower in absolute terms than the reduction in thrombosis) and no significant differences in mortality.⁵ On the basis of these data, several guidelines were updated to propose the use of therapeutic doses of LMWH versus standard prophylaxis in selected COVID-19 patients in conventional hospital wards with no additional hemorrhagic risk factors.^{6–8} Meta-analyses of individual patient data will soon be available, and these reports will probably help identify patient subgroups that benefit most from this strategy. Other pertinent topics addressed in the guidelines are the prolongation of prophylaxis after discharge (not generally recommended) and prophylaxis in non-hospitalized patients (not consistently recommended).

The latest version of the international clinical practice guidelines for the treatment and prevention of venous thromboembolism (VTE) in cancer patients, sponsored by the International Thrombosis and Cancer Initiative (ITAC),⁹ includes for the first time a section on the treatment and prevention of VTE in patients with cancer and COVID-19. In these cases, the panelists recommend the same approach used in cancer patients without COVID-19. The

argument for this recommendation is two-fold. Firstly, there is no evidence that the incidence of VTE in hospitalized patients with cancer and COVID-19 is higher than in non-cancer patients with COVID-19 (although it may have been more appropriate to compare the incidence of VTE in hospitalized cancer patients with and without COVID-19). Secondly, few specific data are available in the oncological population on the risk and benefit of different prophylactic strategies in patients with COVID-19. In fact, several of the clinical trials mentioned above excluded patients with active cancer and all of them excluded patients with primary or secondary central nervous system involvement.

Cancer patients are a special population with a high thrombotic and hemorrhagic risk due to multiple factors.¹⁰ Hospitalization is a recognized risk factor for VTE in patients with active cancer.¹¹ For this reason, routine prophylaxis with LMWH is recommended in all hospitalized patients with cancer in the absence of any contraindication, except for patients admitted solely for the administration of cancer treatment and who are not bedridden. Nevertheless, 3–8% of patients develop an episode of VTE during hospitalization or within days after discharge, despite thromboprophylaxis.¹²

Whence the dilemma: should standard prophylactic doses be administered as suggested by the ITAC guidelines for cancer patients or therapeutic doses as suggested by the more general guidelines? (Table 1). The simple but less useful answer is that patients should be assessed on a case-by-case basis, taking into account the thrombotic/hemorrhagic risk balance in each case. However, although there are no published data, the changing characteristics of prevalent strains of the virus, widespread vaccination, and advances in disease management may mean that the thrombotic risk associated with COVID-19 is not now as crucial as it was in the initial waves of the pandemic. If this is indeed the case, the validity of the results of clinical trials that were mostly conducted at the beginning of the pandemic could be questioned, since in absolute terms, the use of therapeutic doses of LMWH would have a lower impact on the prevention of thrombotic events. Given this uncertainty and the increased hemorrhagic risk of cancer patients, it does not seem unreasonable to be prudent and suggest that, in general, patients with cancer and COVID-19 admitted to a conventional hospital ward should receive standard prophylaxis with LMWH.

Even less evidence is available for specific recommendations on thromboprophylaxis in cancer and COVID-19 in patients who do not require hospitalization. Many patients starting cancer treatment in routine practice do not undergo a thrombotic risk assessment, a situation that may lead to the underuse of outpatient

<https://doi.org/10.1016/j.arbres.2022.08.006>

0300-2896/© 2022 SEPAR. Published by Elsevier España, S.L.U. All rights reserved.

Table 1

Summary of current recommendations/suggestions in different clinical practice guidelines on thromboprophylaxis in patients with COVID-19 (July 2022).

	CHEST guidelines	ASH guidelines	ISTH guidelines	ITAC (cancer) guidelines
Critical patients	Standard prophylaxis	Standard prophylaxis	Standard prophylaxis	na
Acute patients (conventional hospital ward)	LMWH therapeutic doses ^a	LMWH therapeutic doses ^a	LMWH therapeutic doses ^a	Standard prophylaxis
Prolongation of prophylaxis after discharge	Not recommended	Not recommended	Consider in selected patients	Not recommended
Non-hospitalized patients	na	na	Consider sulodexide ^b	According to individual risk ^c

^a Selected patients with low hemorrhagic risk.^b Patients at high risk of disease progression, to reduce risk of hospitalization.^c Follow the same recommendations as in cancer patients without COVID-19, na, not applicable.

thromboprophylaxis. In cancer patients with COVID-19, thromboembolic risk should be assessed, if it had not been done previously (there are several validated scales for this purpose).¹³ Pharmacological thromboprophylaxis is probably indicated in a significant proportion of patients, irrespective of a COVID-19 diagnosis.

Emerging concepts such as personalized medicine and precision medicine are also applicable in the field of thromboembolic disease. Evidence-based guidelines are very useful, but they inevitably fail to address several circumstances commonly encountered in daily practice. Although these knowledge gaps are gradually closing thanks to scientific advances, there remains ample scope for practicing the art of medicine.

Conflicts of interest

The authors declare that they have no conflict of interests related with the contents of this manuscript.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi: 10.1016/j.arbres.2022.08.006](https://doi.org/10.1016/j.arbres.2022.08.006).

References

- Mackman N, Antoniak S, Wolberg AS, Kasthuri R, Key NS. Coagulation abnormalities and thrombosis in patients infected with SARS-CoV-2 and other pandemic viruses. *Arterioscler Thromb Vasc Biol.* 2020;40:2033–44. [http://dx.doi.org/10.1161/ATVBAHA.120.314514](https://doi.org/10.1161/ATVBAHA.120.314514).
- Sociedad Española de Trombosis y Hemostasia (SETH). Recomendaciones de tromboprolifaxis y tratamiento antitrombótico en pacientes con COVID-19. Fecha de publicación: 21 de abril de 2022. Available from: <https://www.covid-19.seth.es/recomendaciones-de-tromboprolifaxis-y-tratamiento-antitrombotico-en-pacientes-con-covid-19/>.
- Demelo-Rodríguez P, Farfán-Sedano AI, Pedrajas JM, Llamas P, Sigüenza P, Jaras MJ, et al. Bleeding risk in hospitalized patients with COVID-19 receiving intermediate- or therapeutic doses of thromboprophylaxis. *J Thromb Haemost.* 2021;19:1981–9. [http://dx.doi.org/10.1111/jth.15400](https://doi.org/10.1111/jth.15400).
- REMAP-CAP Investigators; ACTIV-4a Investigators; ATTACC Investigators. Therapeutic anticoagulation with heparin in critically ill patients with Covid-19. *N Engl J Med.* 2021;385:777–89. [doi:10.1056/NEJMoa2103417](https://doi.org/10.1056/NEJMoa2103417).
- Valeriani E, Porfidia A, Ageno W, Spoto S, Pola R, Di Nisio M. High-dose versus low-dose venous thromboprophylaxis in hospitalized patients with COVID-19: a systematic review and meta-analysis. *Intern Emerg Med.* 2022;27:1–9. [http://dx.doi.org/10.1007/s11739-022-03004-x](https://doi.org/10.1007/s11739-022-03004-x).

- Moores LK, Tritschler T, Brosnahan S, Carrier M, Collen JF, Doerschug K, et al. Thromboprophylaxis in patients with COVID-19: a brief update to the CHEST guideline and expert panel report. *Chest.* 2022;162:213–25. [http://dx.doi.org/10.1016/j.chest.2022.02.006](https://doi.org/10.1016/j.chest.2022.02.006).
- Schulman S, Sholzberg M, Spyropoulos AC, Zarychanski R, Resnick HE, Bradbury C, et al. ISTH guidelines for antithrombotic treatment in COVID-19. *J Thromb Haemost.* 2022. [http://dx.doi.org/10.1111/jth.15808](https://doi.org/10.1111/jth.15808).
- Cuker A, Tseng EK, Nieuwlaat R, Angchaisuksiri P, Blair C, Dane K, et al. American Society of Hematology living guidelines on the use of anticoagulation for thromboprophylaxis in patients with COVID-19: January 2022 update on the use of therapeutic-intensity anticoagulation in acutely ill patients. *Blood Adv.* 2022. [http://dx.doi.org/10.1182/bloodadvances.2022007561](https://doi.org/10.1182/bloodadvances.2022007561), blood-advances.2022007561.
- Farge D, Frere C, Connors JM, Khorana AA, Kakkar A, Ay C, et al. 2022 international clinical practice guidelines for the treatment and prophylaxis of venous thromboembolism in patients with cancer, including patients with COVID-19. *Lancet Oncol.* 2022;23:e334–47. [http://dx.doi.org/10.1016/S1470-2045\(22\)00160-7](https://doi.org/10.1016/S1470-2045(22)00160-7).
- Gervaso L, Dave H, Khorana AA. Venous and arterial thromboembolism in patients with cancer: *JACC: CardioOncology* state-of-the-art review. *JACC CardioOncol.* 2021;3:173–90. [http://dx.doi.org/10.1016/j.jacc.2021.03.001](https://doi.org/10.1016/j.jacc.2021.03.001).
- Figueroa R, Alfonso A, López-Picazo J, Gil-Bazo I, García-Mouriz A, Hermida J, et al. Insights into venous thromboembolism prevention in hospitalized cancer patients: lessons from a prospective study. *PLOS ONE.* 2018;13. [http://dx.doi.org/10.1371/journal.pone.0200220](https://doi.org/10.1371/journal.pone.0200220), e0200220.
- Figueroa R, Alfonso A, López-Picazo J, Gil-Bazo I, García-Mouriz A, Hermida J, et al. Improvement of appropriate pharmacological prophylaxis in hospitalised cancer patients with a multiscreen e-alert system: a single-centre experience. *Clin Transl Oncol.* 2019;21:805–9. [http://dx.doi.org/10.1007/s12094-018-1986-1](https://doi.org/10.1007/s12094-018-1986-1).
- Moik F, Ay C, Pabinger I. Risk prediction for cancer-associated thrombosis in ambulatory patients with cancer: past, present and future. *Thromb Res.* 2020;191 Suppl. 1:S3–11. [http://dx.doi.org/10.1016/S0049-3848\(20\)30389-3](https://doi.org/10.1016/S0049-3848(20)30389-3).

Ramón Lecumberri^{a,b,*}, María Marcos-Jubilar^a, Carolina Guillén^c
^a Servicio de Hematología, Clínica Universidad de Navarra, Pamplona, Spain

^b CIBER-CV, Instituto de Salud Carlos III, Madrid, Spain

^c Servicio de Hematología, Clínica Universidad de Navarra, Madrid, Spain

Corresponding author.

E-mail address: rlcumber@unav.es (R. Lecumberri).